

(This is a sample cover image for this issue. The actual cover is not yet available at this time.)

This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.


In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/copyright>



Contents lists available at SciVerse ScienceDirect

NeuroImage

journal homepage: www.elsevier.com/locate/ynimg

Single pulse TMS induced disruption to right and left parietal cortex on addition and multiplication[☆]

Elena Salillas^{a,*}, Carlo Semenza^{b,c}, Demis Basso^{d,e}, Tomaso Vecchi^f, Michael Siegal^{g,h,**}

^a Basque Center on Cognition Brain and Language, Donostia, Spain

^b Department of Neuroscience, University of Padova, Italy

^c IRCCS Ospedale S Camillo, Lido di Venezia, Italy

^d Faculty of Education, Free University of Bozen, Italy

^e CeNCA, Centro di Neuroscienze Cognitive Applicate, Rome, Italy

^f Department of Psychology, University of Pavia, Italy

^g Department of Psychology, University of Trieste, Italy

^h Department of Psychology, University of Sheffield, UK

ARTICLE INFO

Article history:

Received 23 June 2011

Revised 15 September 2011

Accepted 31 October 2011

Available online 4 November 2011

Keywords:

Mathematics

Intraparietal sulcus

Verbal codes

ABSTRACT

Whether or not mathematical operations are dependent on verbal codes in left hemisphere areas – particularly the left intraparietal sulcus – remains an issue of intense debate. Using single pulse transcranial magnetic stimulation directed at horizontal and ventral regions of the left and right intraparietal sulcus, we examined disruption to reaction times in simple addition and multiplication. Results indicate that these two operations differ in the pattern of lateralization across time for the two areas studied. These show that computational efficiency is not specifically dependent on left hemisphere regions and, in particular, that efficiency in multiplication is dependent on the ventral region of the intraparietal sulcus in the right hemisphere considered to be critical for motion representation and automatization.

© 2011 Elsevier Inc. All rights reserved.

Introduction

In mapping mathematical abilities in the brain, a still very coarse distinction is made, despite recent progress, between the respective functions of the left and of the right hemisphere. Traditionally, since acalculia mostly derives, in several varieties, from left hemisphere lesions, mathematical cognition has been linked in large measure to the left hemisphere and to linguistic functions (Jackson and Warrington, 1986). Number words have been considered to provide a verbal code that facilitates exact calculation (Dehaene et al., 1999; Spelke and Tsivkin, 2001). In this respect, activation in left hemisphere (LH) brain regions has been found in fMRI studies in operations relying

on memory retrieval (Dehaene et al., 1999, 2003; Pinel and Dehaene, 2010; Stanescu-Cosson et al., 2000; Zhou et al., 2007). In particular, operations such as addition and multiplication involve activation of the left horizontal portion of the intraparietal sulcus (IHIPS) – an area seen to be critical in the representation of quantities. Although the left angular gyrus (IANG) can also be activated in operations as addition or multiplication, it has been seen to concern linguistic processing more than the specific processing of quantity, and the role of the IANG in calculation appears dependent on the IHIPS (Dehaene et al., 2003, 2004, p. 219). By this account, emphasis has been placed on the predominance of a left hemisphere network for calculation. In sum, although the processing of number magnitude has been proposed to reside bilaterally in HIPS, it has been proposed that the IANG jointly with parietal areas is of prime importance for retrieving results in arithmetic problems. (Dehaene et al., 2003, 2004).

Within this context, our first research question concerns whether *both right and left parietal regions are critical for exact numerical processing*. There are reasons to further explore the contribution of the right hemisphere and non-language-related processing during these tasks. First, not all neuroimaging studies have found more activation in left hemisphere language regions on calculation tasks (Pesenti et al., 2000; Zago et al., 2001). There are investigations that have shown right posterior parietal activations (Dehaene, 2009; Hubbard et al., 2005; Knops et al., 2009a; McCrink et al., 2007) and these activations have been attributed to the need of a visuospatial medium for arithmetic. Second, there is evidence from neuropsychological data

[☆] This research was supported by European Union Marie Curie Action Contract “NUMBRA” (504927) and by a MIUR grant 2005 and Italian PRIN 2007 (2007XFM93B_005) to C.S. M.S. is supported by a European Union 6th Framework Marie Curie Chair and a grant from the Fondazione Benefica Kathleen Foreman-Casali. T.V. is supported by the Italian PRIN2006 and in part by Fondazione CARIPLO. D.B. is supported by the fund FIRB RBNE018ET9_003. E.S. is supported by the Marie Curie FP7-PEOPLE-2010-IEF programme.

* Correspondence to: E. Salillas, Basque Center on Cognition, Brain and Language (BCBL), Paseo Mikeletegi 69 2°, 20009 Donostia, Spain.

** Correspondence to: M. Siegal, Department of Psychology, University of Sheffield, Western Bank, Sheffield S10 2TP, UK.

E-mail addresses: e.salillas@bcbl.eu (E. Salillas), m.siegal@sheffield.ac.uk (M. Siegal).

showing dissociations between linguistic and mathematical abilities in developmental and acquired language disorders (Landerl et al., 2004; Varley et al., 2005), and patient studies have documented that acalculia can occur without damage to LH language areas (Granà et al., 2006; Hartje, 1987) or that simple calculation can be preserved in persons with aphasia (Rossor et al., 1995; Warrington, 1982; Whalen et al., 2002).

Lesion studies are indeed important for discovering whether a given anatomical structure is necessary to carry over a given task. However, a structure may contribute to performing a task and yet be not strictly necessary. Results from the use of TMS techniques imply causal relationships, overcoming the limitations of correlation approaches in neuroimaging and the lack of clearly identifiable neural loci in the study of patients. The virtual lesion approach provided by TMS can therefore help to establish the importance of left and right parietal areas in calculation. Use of these techniques should facilitate investigation of the extent to which both left and right IPS are fundamental for addition and/or multiplication.

Therefore the present study used single-pulse TMS to explore the role of bilateral IPS in calculation. Our focus was on two areas: *HIPS* and *VIPS*. Stimulation was directed at the *right and left HIPS* in order to compare whether interference to operations of addition and multiplication would be confined to the *LHIPS* in a LH network dedicated to calculation, as consistent with a verbally-mediated account (Dehaene et al., 2003, 2004), or to the *HIPS* bilaterally as would be consistent with an account in which verbal mediation is not specifically necessary and sufficient to calculation (Bloom and Keil, 2001; Campbell and Epp, 2004; Gelman and Butterworth, 2005).

We also examined the effects of TMS on the *ventral portion of the intraparietal sulcus* bilaterally (*rVIPS* and *IVIPS*). Recently, fMRI activation accompanying numerical processing in adults in the *HIPS* has been reported to extend adjacently to the *rVIPS* region (Cantlon et al., 2006). Moreover, applying TMS to the ventral portion of the *rIPS* (mean Talairach coordinates, $x = 22.0$, $y = 68.6$, $z = 39.8$) impairs processing of number magnitude (Cohen Kadosh et al., 2007). Although this location differs from the posterior superior parietal lobe studied by Andres et al. (2011) in a TMS study of calculation, it is in the region studied by Salillas et al. (2009) who showed that TMS inhibition of *VIPS* results in impaired efficiency in both motion perception and number comparison. As Salillas et al. (2009) have suggested, *VIPS* may be recruited within a network sustaining number comparison to complement *HIPS* by facilitating use of a mental number line. In order for a person to decide whether a number is higher or lower than a reference number, the focus of attention has to “move” along the mental number line in a way similar to that in which sensory motion is computed. This process may be carried by adjacent motion sensitive areas such as *VIPS* forming the basis of visuospatial operations implied in arithmetic (Knops et al., 2009b; Salillas et al., 2009). Therefore, concerning our first research question, if right *IPS* is essential for efficiency in exact calculation, then no differences across hemispheres should be obtained. Only a main effect of site of stimulation or visual field should be found. In this regard, we predicted a joint involvement of ventral and horizontal *IPS* in calculation as shown through TMS induced disruption to these regions.

In contrasting addition and multiplication, our study was intended to address a second research question concerning *whether multiplication is more verbally-mediated than addition*. We examined the neural basis of computations involving simple addition and multiplication — operations that have both been considered to be especially dependent on verbally-coded facts (Lemer et al., 2003). Nevertheless, simple addition is usually done with the use of procedural strategies whereas multiplication is thought to rely more in memorization (e.g., Dehaene and Cohen, 1997; Roussel et al., 2002). These strategies may lead to differences in the form of final representations (e.g., Siegler and Shipley, 1995; Siegler and Shrager, 1984). Thus as has often been reported (e.g. Cohen et al., 2000; Dehaene and Cohen, 1997; Delazer

and Benke, 1997; Lemer et al., 2003; Pesenti et al., 1994; van Harskamp et al., 2002), mental representations of multiplication facts may have greater reliance on verbal memory and hence greater left hemisphere involvement than those for addition facts.

A third research question involved examination of *the temporal pattern of left and right parietal contributions to calculation*. The study of the time course of *IPS* involvement in calculation can be achieved using single pulse TMS while targeting left or right parietal areas. However, to date, research has been limited to studies of repetitive TMS effects on number processing that have mainly concerned magnitude comparisons (Andres et al., 2005; Cohen Kadosh et al., 2007; Göbel et al., 2001; Knops et al., 2006; Sandrini et al., 2004). Recently, Andres et al. (2011) have used *rTMS* to address lateralization effects in the *HIPS* during simple arithmetic. They have reported that *rTMS* directed at the bilateral *HIPS* during simple subtraction and multiplication influences calculation, thus questioning the proposal of a left lateralized network for exact arithmetic.

The use of single pulse TMS at different stimulus onset asynchronies (SOAs) in our investigation was aimed to determine whether lateralization is constant across time or whether left and right *IPS* areas are necessary at different moments (i.e. visuospatial processes are used before a left lateralized verbal recovery of the answer occurs). Stimulation was delivered at one of four SOA intervals (150, 200, 250 and 300 ms). These SOAs were chosen because they have been shown to be critical time points where arithmetic processes are detected in ERP studies. Specifically, similar arithmetic problems elicit ERP components starting as soon as 250 ms after problem presentation, and this effect lasts until 280, 350 or 400 ms (e.g., Galfano et al., 2009; Stanesco-Cosson et al., 2000). Since no study has addressed the time course in lateralization due to restrictions in the imaging techniques used, our investigation of SOA effects was exploratory.

In the two experiments reported here, lateralization was examined in two ways: through the presentation of lateralized stimuli and through the selective stimulation of a hemisphere at a time. In **Experiment 1**, we compared stimulation in the contralateral visual field to TMS stimulation with the interhemispheric sulcus as the control site. In **Experiment 2**, stimulation in the contralateral field was compared to ipsilateral stimulation as the control site. These two approaches allowed us to give consistency to the results and a more fine grained timing of processes could be observed in **Experiment 2**. Lateralized presentation was intended to maximize the projections of the stimuli contralaterally and the involvement of the TMS stimulation to each hemisphere. In **Experiment 1**, the critical comparison concerned RTs after contralateral stimulation vs. central stimulation. Therefore, maximal effects were expected since disruption was compared against a neutral condition. As in **Experiment 2**, the critical comparison concerned the visual field that was ipsilateral vs. contralateral to the stimulated hemisphere in tracking how and when ipsilateral stimuli presentation differs with the maximum expected disruption after contralateral presentation. Thus this experiment was expected to yield weaker effects due to inter-hemispheric transfer although the comparison between visual fields entailed a finer investigation of interhemispheric differences.

Experiment 1

Method

The procedure was approved by the Ethics Committee of the Department of Psychology of the University of Pavia.

Participants

Twelve right-handed undergraduates participated in the study (mean age: 23 yrs. range 21 to 26; 3 males and 9 females; average + 91% laterality index scores as indicated from the Oldfield (1971)). All of them provided informed consent before the experiment.

Design and stimuli

The participants were given two types of tasks: addition and multiplication consisting of one-digit operands and operators from 3 to 9. The items were classified as easy and difficult after observing differences in reaction times in a preliminary study involving a different group of four undergraduates who were given the calculation items using the same paradigm as in the main TMS experiments, except that the operations appeared centrally on the screen. During this item classification phase, each of the 28 addition or multiplication items was presented seven times. Each item was then classified as easy or hard on the basis of mean RTs to a button press after arriving to a solution. Easy tasks were defined using the median of RTs as those with RTs of 1042 ms or less for addition and with RTs of 1290 ms or less for multiplication employing the same experimental paradigm as that used during TMS. These criteria served to divide the items in two halves, with 50% above and below those cut-offs. As the multiplication item classification closely replicated that of Campbell and Graham (1985; see Table 1), RTs proved to be good indices of item difficulty. The use of a varied pool of pairs was intended to maximize the novelty of each pair of operands and solutions to examine the association of the studied areas with problem size variation. Items were divided into easy and hard in order to maximize effects for hard items.

Procedure

The task consisted of the presentation of two lateralized digits to the right visual field or the left visual field in different blocks. Participants were asked to look at a central fixation point while attending parafoveally to the problems. After mentally arriving at a solution for each addition or multiplication task, participants pressed a button with their right hand. They were told to be careful and not to press the button until they really knew the solution. Afterwards they were asked to verbalize the solution; this was a control task in order to assure participants were really calculating. The position of the larger digit in each equation was randomized across trials with the restrictions that the larger digit did not appear first more than twice and the digits could not appear in

consecutive equations as commuted pairs (e.g., $3 + 6$, $6 + 3$). In order to project the digits to the hemisphere to which TMS was applied, the stimuli were presented lateralized with respect to a central fixation asterisk (the 4.6° visual angle of the mathematical sign at the center of the equation). The addition and multiplication equations were presented in three blocks each for each visual field in a total of 12 blocks. The order of the blocks was fully counterbalanced. Within each block, participants were given 112 trials. For each trial, RTs between the interval from the initial display of the equation and button pressing were recorded. Participants were tested in two sessions on different days. In the first session, they received three blocks for addition and three blocks for multiplication. One session included TMS stimulation of the rHIPS, rVIPS, and the control area while, in the other session, TMS was delivered to lHIPS, lVIPS, and the control area. Half the participants began with the first session, while the other half began with the second session. The order of stimulation sites within each session was randomized.

Following the procedure used by Salillas et al. (2009), single pulse TMS was applied at 110% of the phosphene threshold (mean $MT = 37.4 \pm 4.3$), with a figure-of-eight coil and a Magstim 200 stimulator. Stimulation was delivered (see Fig. 1) either to Talairach coordinates in (a) the rHIPS or lHIPS (± 43 , -48 , 47), to (b) the right or left VIPS (± 24 , -76 , 30) or to (c) the interhemispheric sulcus (central/control site: 0 , -76 , 30). These points were identified using the SoftTactic Evolution Navigator system. This system works in absence of radiological images on the basis of digitized skull landmarks (nasion, inion and two preauricular points) from which it maps 40 uniformly-distributed points in the scalp. Talairach coordinates of cortical sites underlying coil locations are then estimated for each subject MRI constructed stereotaxic template. The coil, perpendicular to the scalp surface, was positioned with the handle pointing up either with the wings parallel to the coronal plane when stimulating VIPS, or with the wings 30° from the sagittal plane when stimulating HIPS. The coil position on the stimulation sites was continuously monitored using the navigation software coupled with the Fastrak Polhemus system

Table 1

Items used in the experiment for addition and multiplication (D = Difficulty; H = Hard; E = Easy). For multiplication, the disruption index average is provided based on the RT difference when rVIPS, lVIPS, rHIPS or lHIPS is disrupted by TMS minus the RT when the control-central site is disrupted. Item = number of the item. Num1 and num2 = numbers presented for addition and multiplication items. Res = correct calculation result.

ADDITION					MULTIPLICATION								
Item	Num1	Num2	Res	D	Item	Num1	Num2	Res	D	rVIPS	rHIPS	lVIPS	lHIPS
1	3	3	6	E	1	3	3	9	E	64,14	2,01	288,80	185,08
2	3	4	7	E	2	3	4	12	E	132,20	140,34	105,05	84,82
3	3	5	8	E	3	3	5	15	E	41,46	-94,28	233,54	108,47
4	3	6	9	E	4	3	6	18	E	289,62	182,39	116,75	219,97
5	3	7	10	E	5	3	7	21	E	-88,85	-125,91	199,99	195,52
6	3	8	11	H	6	3	8	24	E	68,91	66,69	218,00	166,60
7	3	9	12	H	7	3	9	27	E	200,13	54,25	192,85	131,93
8	4	4	8	E	8	4	4	16	E	188,31	32,24	341,42	123,11
9	4	5	9	E	9	4	5	20	E	159,09	-91,20	246,22	110,84
10	4	6	10	E	10	4	6	24	H	191,06	122,28	63,30	565,45
11	4	7	11	H	11	4	7	28	H	268,89	-47,76	262,14	108,69
12	4	8	12	H	12	4	8	32	H	130,38	169,79	230,73	193,96
13	4	9	13	H	13	4	9	36	H	261,60	-19,50	297,67	261,42
14	5	5	10	E	14	5	5	25	E	-30,42	-68,02	349,88	-9,55
15	5	6	11	E	15	5	6	30	H	103,16	11,23	174,35	59,20
16	5	7	12	H	16	5	7	35	H	58,89	1,89	250,23	18,91
17	5	8	13	H	17	5	8	40	H	-2,33	-19,33	204,79	183,72
18	5	9	14	H	18	5	9	45	H	224,09	-52,84	203,78	196,95
19	6	6	12	E	19	6	6	36	E	48,69	-1,49	194,35	-16,06
20	6	7	13	H	20	6	7	42	H	215,66	-63,23	202,73	354,30
21	6	8	14	H	21	6	8	48	H	197,57	97,55	18,21	158,58
22	6	9	15	H	22	6	9	54	H	467,25	27,57	603,31	308,74
23	7	7	14	E	23	7	7	49	E	220,79	204,89	278,82	436,92
24	7	8	15	H	24	7	8	56	H	492,32	-110,98	453,80	668,78
25	7	9	16	H	25	7	9	63	H	992,76	338,77	629,66	519,31
26	8	8	16	E	26	8	8	64	H	222,57	47,93	-114,84	74,84
27	8	9	17	H	27	8	9	72	H	628,49	256,24	303,63	529,02
28	9	9	18	E	28	9	9	81	E	265,23	185,35	152,73	135,75




Fig. 1. Sites of TMS-induced interference. a) Right and left VIPS b) right and left HIPS c) central control regions d) bidimensional representation of the TMS stimulation sites obtained through the FasTrak Polhemus neuronavigation system.

(1 d). Stimulation was delivered at one of four stimulus onset asynchrony (SOA) intervals. These were 150, 200, 250 or 300 ms after the operation appeared. Single pulse stimulation was delivered after the operation was presented at one of these SOAs randomly.

Results

One participant was excluded from the analysis on the basis on an abnormally high error rate (9 to 16%). The verbal response errors of the other 12 participants were rare (1.8% of answers in multiplication; 1.1% in addition). Two separate 2 (difficulty: easy vs. difficult) \times 2 (hemisphere: RH vs LH) \times 3 (site: HIPS, VIPS, central) ANOVAs on the proportion of errors in addition and multiplication respectively indicated that errors were associated with item difficulty. More errors were produced in difficult items both for multiplication (3% vs. 0.7%: $F(1,10) = 24.5$; $p = 0.001$; $\eta^2 = 0.71$) and for addition (1.5% vs. 0.6%: $F(1,10) = 13.4$; $p = 0.004$; $\eta^2 = 0.57$). Errors were not associated with TMS location. Reaction times (RTs) for each operation were examined in two separate 2 (difficulty: easy vs. hard) \times 2 (hemisphere: RH vs. LH) \times 3 (site: HIPS, VIPS, central) \times 4 SOA (150/200/250/300) ANOVAs. Subsequent RT analyses were carried out using correct responses.

For addition, there were no significant main or interaction effects involving hemisphere or SOA (all $F_s < 1$). However, there was a significant difficulty \times site interaction, $F(2,20) = 4.12$; $p = 0.03$; $\eta^2 = 0.29$ (Fig. 2a). The effects of site were confined to difficult items: $F(2,20) = 3.65$; $p = 0.04$ $\eta^2 = 0.27$. TMS induced disruption in VIPS did not affect performance compared to the control, $F(1,10) = 1.92$; $p = 0.19$; $\eta^2 = 0.16$. By contrast, disruption directed at the HIPS bilaterally increased reaction times: $F(1,10) = 6.891$; $p = 0.02$; $\eta^2 = 0.41$.

For multiplication, there was a significant main effect for site: $F(2,20) = 3.40$; $p = 0.05$; $\eta^2 = 0.25$, as well as a significant difficulty \times site interaction effect: $F(2,20) = 5.35$; $p = 0.018$; $\eta^2 = 0.35$, and a difficulty \times hemisphere \times site interaction effect, $F(2,20) = 5.48$; $p = 0.02$; $\eta^2 = 0.35$ (Fig. 2b). Again significant effects were confined to the difficult items, $F(2,20) = 4.23$; $p = 0.03$; $\eta^2 = 0.30$. Interference to VIPS bilaterally compared to the relevant central control sites significantly increased RTs, for rVIPS, $F(1,10) = 6.62$; $p = 0.03$; $\eta^2 = 0.40$; for lVIPS, $F(1,10) = 6.75$; $p = 0.03$; $\eta^2 = 0.40$. Although disruption directed at lHIPS compared to the relevant control site also resulted in significantly increased RTs, $F(1,10) = 6.14$; $p = 0.03$; $\eta^2 = 0.38$, disruption of the rHIPS did not, $F(1,10) < 1$, $\eta^2 < 0.01$. In both analyses, the SOA main effect was significant, addition: $F(3,30) = 4.31$; $p = 0.012$; $\eta^2 = 0.30$; multiplication: $F(3,30) = 6.42$; $p = 0.006$; $\eta^2 = 0.39$. Addition was significantly more impaired with SOAs of 250 ms ($M = 1161.3$, $SD = 152.9$) and 300 ms ($M = 1114.63$, $SD = 137.4$) compared to an SOA of 150 ms ($M = 1155.6$, $SD = 145.8$), $F(1,10) > 5.10$; $p > 0.88$; $\eta^2 > 0.33$. Multiplication was significantly more impaired with SOAs of

250 ms ($M = 1419.1$, $SD = 176.4$) and 300 ms ($M = 1421.1$, $SD = 172.1$) compared to SOAs of 150 ms ($M = 1369.2$, $SD = 174.4$) and 200 ms ($M = 1381.8$, $SD = 169.8$), $F_s(1,10) > 6.41$; $p > 0.002$; $\eta^2 > 0.35$. However, there were no significant interaction effects involving SOA.

Size effects in rVIPS. The effects of TMS-induced disruption on the rVIPS for multiplication were related to solution size. Regression analyses of solution size using the difference between RTs for each site stimulated and control sites as the dependent measure revealed that solution size significantly decreased calculation efficiency only in the case of TMS induced disruption to the rVIPS ($t = 3.39$; $\beta = 0.7$; $p < 0.005$) for the difficult multiplication items. There were no significant relationships for other sites of disruption or for either the easy or difficult addition items or for the easy multiplication items. For either addition or multiplication, the order of the larger and smaller numbers in the operation (e.g., $3 + 6$ vs. $6 + 3$; 3×6 vs. 6×3) did not significantly influence RTs.




Fig. 2. RTs as a function of site of stimulation for Experiment 1. RTs (ms) for addition (a) and multiplication (b) under TMS directed at the VIPS, HIPS, and central control regions in the RH and LH for the easy and hard items with 95% confidence intervals.

The goal of [Experiment 2](#) was to compare the effects of ipsilateral vs. contralateral stimulation for both VIPS and HIPS sites — a contrast aimed to provide fine grained comparisons between hemispheres. In addition, structural MRIs were used to localize the sites in order to facilitate precise localization of the sites with respect to that in [Experiment 1](#). As the contralateral hemisphere serves to process preferentially lateralized stimuli, our hypothesis was that contralateral TMS stimulation would impair calculation efficiency compared to ipsilateral stimulation. We expected a similar pattern of results as in [Experiment 1](#) such that (1) stimulation to the HIPS bilaterally either to the RH or LH would impair efficiency in addition and that (2) stimulation to the VIPS bilaterally either to the RH or LH would impair efficiency in multiplication. Further, we hypothesized that multiplication would be impaired by stimulation directed at the IHIPS compared to the rHIPS. Thus although [Experiment 2](#) aimed again to characterize the lateralization pattern for exact calculation, our comparison of ipsilateral versus contralateral stimulation allowed for a more precise look at the interplay of the two hemispheres along time.

Experiment 2

Method

The procedure was approved by the Ethics Committee of the Department of General Psychology of the University of Padua.

Participants

These were 10 right-handed undergraduates (mean age: 24 yrs, range 23 to 27; 3 males and 7 females; average +83.3% laterality index as indicated on the [Oldfield \(1971\)](#)). All provided informed consent before the experiment. None had been participants in [Experiment 1](#).

Stimuli

The material was identical to [Experiment 1](#), with the exception that only difficult items for both operations were used since these produced the strongest effects in [Experiment 1](#).

Procedure

The session consisted of 16 blocks as a combination of site of stimulation (HIPS/VIPS), hemisphere (left/right), visual field for the stimulus (ipsilateral/contralateral) and operation. The order of blocks was randomized between participants and none of them was given the same order. The procedure was otherwise identical to that of [Experiment 1](#).

As in [Experiment 1](#), single pulse TMS was applied at 110% of the phosphene threshold (mean MT = 55.3 ± 3.6), with a figure-of-eight coil and a Magstim Rapid² stimulator. Stimulation was delivered either to VIPS and HIPS, either to left and right hemispheres following the same coil positions used in the previous experiment. The TMS coil was placed on the skull of each subject and continuously monitored during the whole experiment with theBrainsight stereotaxic neuronavigator (Rogue Research Inc, Montreal, Canada) coupled with a Polaris Vicra infrared camera system (NDI, Waterloo, Canada) using individual MRI images of the participants. T1-weighted MR scans were obtained from each participant using a Signa 3 T system GE Healthcare, Milwaukee, WI, USA ($1.3 \times 1.3 \times 1.3$ mm, sagittal acquisition). The four sites were localized through optical inspection of anatomical MRI images considering for VIPS the ventral part of the caudal IPS and for HIPS the site between the two gyri separated by the anterior part of the IPS, 1 cm posterior to the postcentral gyrus (see resulting average coordinates in [Table 2](#)). Stimulation was again delivered at one of four SOAs intervals: 150, 200, 250 and 300 ms after each item had been presented.

Results

Participants' verbal response errors were infrequent (2% for addition and 3% for multiplication) and did not follow any systematic pattern.

Table 2

Mean Talairach coordinates and standard deviations for TMS stimulation in Experiment 2.

	X	Y	Z	sdX	sdY	sdZ
IVIPS	−26.6	−84.3	33.3	2.6	3.8	4.3
rVIPS	26.5	−83.8	33.1	3.1	2.7	4.8
IHIPS	−44.1	−49.6	46.2	2.9	4.2	3.2
rHIPS	44.9	−50.0	48.4	2.0	2.5	2.6

RTs are shown in [Table 2](#). Two separate 2 (site: VIPS/HIPS) \times 2 (hemisphere: left/right) \times 2 (visual field: ipsilateral/contralateral) \times 4 (SOA: 150/200/250/300) ANOVAs were performed for each operation.

For addition, only the site \times hemisphere \times visual field \times SOA interaction was significant ($F(3, 27) = 6.62$; $p = 0.002$; $\eta^2 = 0.42$). There were no other significant main or interaction effects. We then proceeded to analyze each stimulation site separately to determine the extent of lateralization effects.

For HIPS, the hemisphere \times visual field \times SOA interaction was not significant, ($F(3,27) = 1.93$; $p = 0.14$; $\eta^2 = 0.18$) indicating, as in [Experiment 1](#), the absence of significant effects for hemisphere. For VIPS, the hemisphere \times visual field \times SOA interaction effect was significant ($F(3,27) = 6.38$; $p = 0.005$; $\eta^2 = 0.41$). However, there were no simple main or interaction effects for either hemisphere or for any of the SOAs (see [Fig. 3](#) and [Table 2](#)).




Fig. 3. RTs as a function of site of stimulation for [Experiment 2](#). RTs (ms) for addition (a) and multiplication (b) under TMS directed to HIPS and VIPS in the RH and LH for ipsilateral or contralateral stimuli presentation. RTs are plotted with 95% confidence interval.

For *multiplication*, no main or interaction effects were significant, although the site \times hemisphere \times visual field marginal interaction approached significance ($F(1,9) = 3.54$; $p = 0.09$; $\eta^2 = 0.28$). A separate analysis for each site revealed a significant hemisphere \times visual field \times SOA interaction ($F(3,27) = 3.48$; $p = 0.03$; $\eta^2 = 0.28$) for IHIPS. Similar to [Experiment 1](#), simple effects tests showed a visual field effect restricted to IHIPS. However, this effect was only significant at the 300 ms SOA ($F(1,9) = 5.27$; $p = 0.047$; $\eta^2 = 0.37$). Again as in [Experiment 1](#), the analysis for VIPS showed a significant visual field main effect ($F(1,9) = 6.35$; $p = 0.03$; $\eta^2 = 0.41$) but also there was a significant visual field \times hemisphere interaction ($F(1,9) = 6.35$; $p = 0.03$; $\eta^2 = 0.41$), with the effect of visual field restricted to the rVIPs ($F(1,9) = 7.92$; $p = 0.02$; $\eta^2 = 0.47$). This significant effect extended throughout 150 ms to 250 ms SOAs (150 ms: $t = 2.51$, $p = 0.03$; false discovery rate (FDR) correction $p = 0.049$; 200 ms: $t = 2.76$, $p = 0.02$; FDR $p = 0.049$; 250 ms: $t = 2.44$, $p = 0.037$; FDR $p = 0.049$), becoming marginal at 300 ms ($t = 1.99$, $p = 0.07$; FDR $p = 0.07$) ([Table 3](#)).

Size effects in rVIPs. A regression analysis between the different sites/visual field/SOAs and the size of the result in an analysis by items showed a significant relationship between rVIPs disruption and the size of the result. The difference of RTs after contralateral minus ipsilateral stimulation was predicted by the problem size ($t = 2.28$; $p = 0.04$; $\beta = 0.5$). This relationship became larger when only those SOAs with significant effects were considered (150 ms to 250 ms) ($t = 2.56$; $p = 0.02$; $\beta = 0.6$).

Discussion

Both left and right IPS were implicated in efficiency of exact calculation

Overall, results from [Experiments 1 and 2](#) converge in showing that disruption of IPS in either hemisphere compared to control sites or contrasting ipsi vs. contralateral visual presentation results in loss of efficiency in the form of increased RTs. This result occurs both in addition and multiplication.

Addition differs from multiplication

For *addition*, disruption of the HIPS in either hemisphere compared to control sites results in loss of efficiency in the form of increased RTs. rHIPS disruption is consistent with the use of the analog magnitude representation jointly with verbal coding strategies ([Göbel et al.](#),

[2001](#)). By contrast, for *multiplication*, significantly increased RTs resulted only from disruption of the IHIPS – and not the rHIPS. However, disruption to the rVIPs was as effective as IHIPS disruption in increasing RTs on multiplication items. The absence of a widespread increase across SOAs in RTs following disruption to the IVIPS in [Experiment 2](#) points to a weaker involvement of this site compared to the rVIPs. Solution size predicts the extent of TMS disruption to the rVIPs. This result indicates an overlap between number processing and the motion representation and automatization of cognitive function that has been shown to occur in this region ([Cohen Kadosh et al., 2007](#); [Orban et al., 2003](#)). As we have noted, it has been shown in a related study that TMS induced interference in VIPs affects both motion perception and numerical comparison ([Salillas et al., 2009](#)). Efficient calculation is based on the automatic retrieval of multiplication facts. This process involves scanning correct products as well as cohort candidate solutions ([Campbell, 1994](#); [Galfano et al., 2003](#)). For example, the selection of 56 as the correct product to 7×8 involves scanning cohort solutions such as the table errors 49 and 54. Impairment to this automatized scanning process through TMS-induced disruption to the rVIPs points to the presence of a RH network that supports multiplication – one that is retained by persons with LH lesions ([Varley et al., 2005](#)) and that may be impaired by RH damage ([Granà et al., 2006](#)). Thus, although both addition and multiplication imply bilateral IPS, specific recruited areas for each operation differ, and rHIPS is only important for addition as proposals based on learning processes suggest ([Siegler and Shipley, 1995](#); [Siegler and Shrager, 1984](#)).

Patterns of lateralization along time

Involvement of both hemispheres appears in fact to be sustained across time. Lateralization tends to appear when the process of result retrieval is disrupted through stimulation on IHIPS at a late onset time (300 ms), however. This lateralization in IHIPS is found for multiplication at this SOA but again, a sustained disruption of the retrieval process is found while disrupting rVIPs from 150 ms SOAs until a later lateralization to IHIPS. The tendency for a late lateralization to the IHIPS suggests involvement of visuospatial processes early during calculation. Other studies have focused on lateralization during fact retrieval using repetitive TMS. Left lateralization has been found using this technique and focusing on left angular gyrus (IANG: [Göbel et al., 2001](#)) with a very different experimental paradigm. While a lateralization was found on IANG, stimulating IPS has led to a bilateral pattern in the recent report of [Andres et al. \(2011\)](#). Our investigation is the first to

Table 3
Effects of SOA in [Experiment 2](#). RTs and (SD) depending on site of stimulation, hemisphere, visual field for each SOA.

	Addition	Contralateral		Ipsilateral	
	SOA	LH	RH	LH	RH
HIPS	150	1220.9 (584.3)	1257.5 (588.5)	1180.8 (576.9)	1191.1 (503.3)
	200	1329.5 (616.2)	1295.4 (565.3)	1093.8 (410.5)	1209.6 (558.7)
	250	1233.6 (565.9)	1255.7 (536.6)	1147.2 (474.1)	1284.7 (609.1)
	300	1304.7 (593.5)	1380.6 (576.4)	1143.2 (464.1)	1215.8 (518.3)
VIPs	150	1312.9 (531.6)	1136.1 (486.8)	1144.4 (366.6)	1205.7 (512.3)
	200	1217.7 (546.3)	1260.8 (533.5)	1256.7 (446.8)	1228.5 (508.1)
	250	1219.8 (509.1)	1141.7 (473.3)	1212.3 (437.5)	1269.5 (513.5)
	300	1207.4 (44.404)	1170.1 (512.4)	1237.2 (387.3)	1217.9 (471.4)
	Multiplication	Contralateral		Ipsilateral	
	SOA	LH	RH	LH	RH
HIPS	150	1603.9 (382.8)	1462.9 (470.8)	1396.8 (453.3)	1423.7 (499.6)
	200	1514.4 (502.9)	1422.8 (586.7)	1422.2 (445.3)	1437.1 (649.5)
	250	1527.9 (349.1)	1485.1 (586.9)	1411.8 (527.9)	1389.7 (675.1)
	300	1586.8 (508.7)	1404.8 (472.2)	1324.3 (476.1)	1590.3 (639.5)
VIPs	150	1327.5 (488.6)	1455.7 (509.4)	1356.5 (278.6)	1176.6 (424.4)
	200	1391.5 (609.3)	1506.7 (445.3)	1368.9 (521.8)	1176.3 (422.7)
	250	1441.4 (547.4)	1575.7 (663.9)	1451.1 (479.2)	1185.3 (424.9)
	300	1371.1 (513.9)	1445.2 (530.9)	1396.8 (447.1)	1250.5 (397.6)

address patterns of lateralization in the IPS using single pulse TMS with different SOAs and provides a description of the temporal course of the bilateral involvement. In our study, we included an observation of the role of VIPS in the automatization of solution search, indicating that use of repetitive stimulation in periods up to 300 ms after stimulus may not reveal possible rapid changes in the use of right and left areas of IPS or even LANG. This temporal sensitivity is lacking both in neuroimaging studies using fMRI and of course in patient data.

A methodological limitation for the present study comes from the anatomical localization used to identify the target regions that may have added a source of variability, as suggested by Rosenberg-Lee et al. (2011). However, we used a single pulse paradigm that, with respect to rTMS studies, decreased the probability that the stimulation would spread to other regions, thus limiting variability in the TMS methodology.

In sum, our investigation demonstrates for the first time using evidence from TMS that simple addition and multiplication calculations involve different RH and LH substrates. For addition, disruption to both rHIPS and IHIPS resulted in impaired efficiency. For multiplication, efficiency was unaffected by disruption to the rHIPS. However, disruption of the rVIPS resulted in a loss of efficiency and increased RTs and these effects were significantly related to problem size. Although adults – at least in Western cultures – may have achieved efficient, automatized calculation through a verbally-mediated route during childhood (Rivera et al., 2005; Tang et al., 2006), verbal mediation may no longer be sufficient for calculation in the mature cognitive system and numerical processing may also rely on visual representation of numerals and arithmetic facts.

References

- Andres, M., Seron, X., Olivier, E., 2005. Hemispheric lateralization of number comparison. *Brain Res.* 25, 283–290.
- Andres, M., Pelgrims, B., Michaux, N., Olivier, E., Pesenti, M., 2011. Role of distinct parietal areas in arithmetic: an fMRI-guided TMS study. *NeuroImage* 54, 3048–3056.
- Bloom, P., Keil, F.C., 2001. Thinking through language. *Mind Lang.* 16, 351–367.
- Campbell, J.I.D., 1994. Architectures for numerical cognition. *Cognition* 53, 1–44.
- Campbell, J.I.D., Epp, L.J., 2004. An encoding-complex approach to numerical cognition in Chinese-English bilinguals. *Can. J. Exp. Psychol.* 58, 229–244.
- Campbell, J.I.D., Graham, D.J., 1985. Mental multiplication skill: structure, process and acquisition. *Can. J. Psychol.* 39, 338–366.
- Cantlon, J.F., Brannon, E.M., Carter, E.J., Pelphrey, K.A., 2006. Functional imaging of numerical processing in adults and 4-y-old children. *PLoS Biology* 4, e125.
- Cohen Kadosh, R., Cohen Kadosh, K., Schuhmann, T., Kaas, A., Goebel, R., Henik, A., Sack, A., 2007. Virtual dyscalculia induced by parietal-lobe TMS impairs automatic magnitude processing. *Curr. Biol.* 17, 1–5.
- Cohen, L., Dehaene, S., Chochon, F., Leh'eric, S., Naccache, L., 2000. Language and calculation within the parietal lobe: a combined cognitive, anatomical and fMRI study. *Neuropsychologia* 38, 1426–1440.
- Dehaene, S., 2009. Origins of mathematical intuitions. *Ann. NY Acad. Sci.* 1156, 232–259.
- Dehaene, S., Cohen, L., 1997. Cerebral pathways for calculation: double dissociation between rote verbal and quantitative knowledge of arithmetic. *Cortex* 33, 219–250.
- Dehaene, S., Spelke, E., Pinel, P., Stanescu, R., Tsivkin, S., 1999. Sources of mathematical thinking: behavioral and brain-imaging evidence. *Science* 284, 970–974.
- Dehaene, S., Piazza, M., Pinel, P., Cohen, L., 2003. Three parietal circuits for number processing. *Cognitive Neuropsych.* 20, 487–506.
- Dehaene, S., Molko, N., Cohen, L., Wilson, A.J., 2004. Arithmetic and the brain. *Curr. Opin. Neurobiol.* 14, 218–224.
- Delazer, M., Benke, T., 1997. Arithmetic facts without meaning. *Cortex* 33, 697–710.
- Galfano, G., Rusconi, E., Umiltà, C., 2003. Automatic activation of multiplication facts: evidence from the nodes adjacent to the product. *Q. J. Exp. Psychol.* 56A, 31–61.
- Galfano, G., Penolazzi, B., Vervaeck, I., Angrilli, A., Umiltà, C., 2009. Event-related brain potentials uncover activation dynamics in the lexicon of multiplication facts. *Cortex* 45 (10), 1167–1177.
- Gelman, R., Butterworth, B., 2005. Numbers and language: how are they related? *Trends Cogn. Sci.* 9, 6–10.
- Göbel, S., Walsh, V., Rushworth, F.S., 2001. The mental number line and the human angular gyrus. *NeuroImage* 14, 1278–1289.
- Granà, A., Hofer, R., Semenza, C., 2006. Acalculia from a right hemisphere lesion: dealing with “where” in multiplication procedures. *Neuropsychologia* 44, 2972–2986.
- Hartje, W., 1987. The effect of spatial disorders on arithmetical skills. In: Deloche, G., Seron, X. (Eds.), *Mathematical Disabilities*. Erlbaum, Hillsdale NJ, pp. 121–125.
- Hubbard, E.M., Piazza, M., Pinel, P., Dehaene, S., 2005. Interactions between number and space in parietal cortex. *Nat. Rev. Neurosci.* 6, 435–448.
- Jackson, M., Warrington, E.K., 1986. Arithmetic skills in patients with unilateral cerebral lesions. *Cortex* 22, 611–620.
- Knops, A., Nuerk, H., Sparing, R., Foltys, H., Willmes, K., 2006. On the functional role of human parietal cortex in number processing: how gender mediates the impact of a “virtual lesion” induced by rTMS. *Neuropsychologia* 44, 2270–2283.
- Knops, A., Thirion, B., Hubbard, E.M., Michel, V., Dehaene, S., 2009a. Recruitment of an area involved in eye movements during mental arithmetic. *Science* 324, 1583–1585.
- Knops, A., Viarouge, A., Dehaene, S., 2009b. Dynamic representations underlying symbolic and nonsymbolic calculation: evidence from the operational momentum effect. *Atten. Percept. Psychophys.* 71, 803–821.
- Landerl, K., Bevan, A., Butterworth, B., 2004. Developmental dyscalculia and basic numerical capacities: a study of 8–9-year-old students. *Cognition* 93, 99–125.
- Lemer, C., Dehaene, S., Spelke, E., Cohen, L., 2003. Approximate quantities and exact number words: dissociable systems. *Neuropsychologia* 41, 1942–1958.
- McCrink, K., Dehaene, S., Dehaene-Lambertz, G., 2007. Moving along the number line: operational momentum in nonsymbolic arithmetic. *Percept. Psychophys.* 69, 1324–1333.
- Oldfield, R.C., 1971. The assessment and analysis of handedness: the Edinburgh inventory. *Neuropsychologia* 9, 97–113.
- Orban, G.A., Fize, D., Peuskens, H., Denys, K., Nelissen, K., Snaert, S., Todd, J.T., Vanduffel, W., 2003. Similarities and differences in motion processing between the human and macaque brain: evidence from fMRI. *Neuropsychologia* 41, 1757–1768.
- Pesenti, M., Seron, X., van der Linden, M., 1994. Selective impairment as evidence for mental organisation of arithmetical facts: BB, a case of preserved subtraction? *Cortex* 30, 661–671.
- Pesenti, M., Thioux, M., Seron, X., De Volder, A., 2000. Neuroanatomical substrate of Arabic number processing, numerical comparison and simple addition: a PET study. *J. Cogn. Neurosci.* 12, 461–479.
- Pinel, P., Dehaene, S., 2010. Beyond hemispheric dominance: brain regions underlying the joint lateralization of language and arithmetic to the left hemisphere. *J. Cogn. Neurosci.* 22, 48–66.
- Rivera, S.M., Reiss, A.L., Eckert, M.A., Menon, V., 2005. Developmental changes in mental arithmetic: evidence for increased functional specialization in the left inferior parietal cortex. *Cereb. Cortex* 15, 1779–1790.
- Rosenberg-Lee, M., Chang, T.T., Young, C.B., Wu, S., Menon, V., 2011. Functional dissociations between four basic arithmetic operations in the human posterior parietal cortex. A cytoarchitectonic mapping study. *Neuropsychologia* 49, 2592–2608.
- Rossor, M.N., Warrington, E.K., Cipolotti, L., 1995. The isolation of calculation skills. *J. Neurol.* 242, 78–81.
- Roussel, J., Fayol, M., Barrouillet, P., 2002. Procedural vs. direct retrieval strategies in arithmetic: a comparison between additive and multiplicative problem solving. *Eur. J. Cogn. Psychol.* 14, 61–104.
- Salillas, E., Basso, D., Baldi, M., Semenza, C., Vecchi, T., 2009. Motion on numbers: Transcranial Magnetic Stimulation on VIPS alters both numerical and motion processes. *J. Cogn. Neurosci.* 21, 2129–2138.
- Sandrini, M., Rossini, P.M., Miniussi, C., 2004. The differential involvement of inferior parietal lobule in number comparison: an rTMS study. *Neuropsychologia* 42, 1902–1909.
- Siegler, R.S., Shipley, C., 1995. Variation, selection, and cognitive change. In: Simon, T., Halford, G. (Eds.), *Developing Cognitive Competence: New Approaches to Process Modeling*. Erlbaum, Hillsdale, NJ, pp. 31–76.
- Siegler, R.S., Shrager, J., 1984. Strategy choices in addition and subtraction: how do children know what to do? In: Sophian, C. (Ed.), *The Origins of Cognitive Skills*. Erlbaum, Hillsdale, NJ, pp. 229–293.
- Spelke, E.S., Tsivkin, S., 2001. Language and number: a bilingual training study. *Cognition* 78, 45–88.
- Stanescu-Cosson, R., Pinel, P., van de Moortele, P.F., Le Bihan, D., Cohen, L., Dehaene, S., 2000. Understanding dissociations in dyscalculia: a brain imaging study of the impact of number size on the cerebral networks for exact and approximate calculation. *Brain* 123, 2240–2255.
- Tang, Y., Zhang, W., Chen, K., Feng, S., Ji, Y., Shen, J., Reiman, E.M., Liu, Y., 2006. Arithmetic processing in the brain shaped by cultures. *Proc. Natl. Acad. Sci. U. S. A.* 103, 10775–10780.
- van Harskamp, N.J., Rudge, P., Cipolotti, L., 2002. Are multiplication facts implemented by the left supramarginal and angular gyri? *Neuropsychologia* 40, 1786–1793.
- Varley, R.A., Klessinger, N.J.C., Romanowski, C.A.J., Siegal, M., 2005. Agrammatic but numerate. *Proc. Natl. Acad. Sci. U. S. A.* 102, 3519–3524.
- Warrington, E.K., 1982. The fractionation of arithmetical skills: a single case study. *Q. J. Exp. Psychol. Proc. Natl. Acad. Sci. U. S. A.* 34, 31.
- Whalen, J., McCloskey, M., Lindemann, M., Bouton, G., 2002. Representing arithmetic table facts in memory: evidence from acquired impairments. *Cogn. Neuropsychol.* 19, 505.
- Zago, L., Pesenti, M., Mellet, E., Crivello, F., Mazoyer, B., Tzourio-Mazoyer, N., 2001. Neural correlates of simple and complex mental calculation. *NeuroImage* 13, 314–327.
- Zhou, X., Chen, C., Zang, Y., Dong, Q., Chen, C., Qiao, S., Gong, Q., 2007. Dissociated brain organization for single-digit addition and multiplication. *NeuroImage* 35, 871–880.