Università degli Studi di Padova

Padua Research Archive - Institutional Repository

Conformity and controversies in the diagnosis, staging and follow-up evaluation of canine nodal lymphoma: a systematic review of the last 15 years of published literature

Original Citation:

Availability:

This version is available at: 11577/3194308 since: 2017-11-06T08:50:04Z

Publisher:

Published version:

DOI: 10.1111/vco.12244

Terms of use:

Open Access

This article is made available under terms and conditions applicable to Open Access Guidelines, as described at <http://www.unipd.it/download/file/fid/55401> (Italian only)

(Article begins on next page)

Conformity and consensus in the diagnosis, staging and follow-up evaluation of canine nodal lymphoma: a systematic review of the last 15 years of published literature

Journal:	<i>Veterinary and Comparative Oncology</i>
Manuscript ID	Draft
Manuscript Type:	Original Article
Keywords:	dog, multicentric lymphoma, staging, efficacy assessment, systemic review

SCHOLARONE™
Manuscripts

View Only

**Conformity and controversies in the diagnosis, staging and follow-up
evaluation of canine nodal lymphoma: a systematic review of the last 15
years of published literature**

Marconato L,^{1*} Polton GA,² Sabattini S,³ Dacasto M,⁴ Garden OA,⁵ Grant I,⁶
Hendrickx T,⁷ Henriques J,⁸ Lubas G,⁹ Morello E,¹⁰ Stefanello D,¹¹ Comazzi S,¹¹
on behalf of the European Canine Lymphoma Network

- 1 Centro Oncologico Veterinario, Sasso Marconi (Bologna), Italy
- 2 North Downs Specialist Referrals, Bletchingley, UK
- 3 Department of Veterinary Medical Sciences, University of Bologna, Italy
- 4 University of Padua, Department of Comparative Biomedicine and Food
Science, Legnaro (Padua), Italy
- 5 Immune Regulation Laboratory, Department of Clinical Science and Services,
Royal Veterinary College, London, UK and Queen Mother Hospital for Animals,
Royal Veterinary College, Hatfield, UK
- 6 Small Animal Clinical Sciences, School of Veterinary Medicine, University of
Glasgow, Glasgow, UK
- 7 Dierenkliniek Sanimalia, Diepenbeek, Belgium
- 8 Centro Veterinário Berna, OnevetGroup, Lisboa, Portugal
- 9 Department of Veterinary Sciences, University of Pisa, Pisa, Italy
- 10 Department of Veterinary Sciences, University of Torino, Grugliasco (Turin), Italy
- 11 Department of Veterinary Sciences and Public Health, University of Milan, Italy

*** Corresponding author:**

Laura Marconato, DVM, DECVIM-CA (Oncology)

Centro Oncologico Veterinario

Via San Lorenzo 1/4

40037 Sasso Marconi, Italy

marconato@centroncologicovet.it

Abbreviated title

Canine lymphoma staging and efficacy assessment

Keywords

Dog, multicentric lymphoma, staging, efficacy assessment, systematic review

Presented in part at the 13-ICML, Lugano, Switzerland, June 20th, 2015.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

Abstract

Diagnostic methods used in the initial and post-treatment evaluation of canine lymphoma are heterogeneous and can vary within countries and institutions. Accurate reporting of clinical stage and response assessment is crucial in determining the treatment efficacy and predicting prognosis. This study comprises a systematic review of all available canine multicentric lymphoma studies published over a period of 15 years. Data concerning clinical stage evaluation and response assessment procedures were extracted and compared. Sixty-five studies met the eligibility criteria. The survey results expose variations in diagnostic criteria and treatment response assessment in canine multicentric lymphoma. Variations in staging procedures performed and recorded led to an unquantifiable heterogeneity among patients in and between studies, making it difficult to compare treatment efficacies. Awareness of this inconsistency of procedure and reporting may help in the design of future clinical trials.

Introduction

Variation in diagnostic criteria and inconsistencies in staging procedures in veterinary cancer patients have important consequences for patient selection in clinical studies and will often preclude meaningful comparison of published data between studies. Standardizing staging and treatment response assessment criteria are therefore critical to the successful performance of clinical trials and to subsequent evaluations and comparisons of study outcomes.

Canine lymphoma is a heterogeneous group of diseases that exhibit distinct biological behaviors according to histological subtype and extent of systemic distribution.^{1,2} In addition to histopathological classification, clinical stage is one of the most important prognostic factors and may therefore represent a key variable in dictating treatment with respect to drug choice. Indeed, modern methods for diagnosis and staging of human lymphoma have improved in parallel with the spectrum of therapeutic options in recent years.³

Modern lymphoma classifications are based on the rationale of defining clinico-pathological disease entities, enabling greater insight into the biological mechanisms that underlie specific diseases and the clinical consequences in terms of progression patterns and responses to different treatments.^{4,5} The ultimate goal is to develop treatment protocols that are specifically tailored to the characteristics of the individual disease entity.⁶

Much emphasis has lately been placed on the morphological subtype of disease.^{1,4,7} Whereas morphological subtype is not expected to change in response to therapy, for accurate evaluation of treatment response, a complete knowledge of lymphoma extension prior to therapy makes it possible to

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

87 accurately re-stage dogs at the end of therapy and thus to define the quality of
88 response. Standardized methods for staging are essential to make critical
89 assessments and comparisons between different therapeutic strategies;
90 incomplete or inconsistent staging work-up impedes comparison of study
91 results.

92 Currently, controversies exist regarding the extent of staging work-up that
93 needs to be carried out at initial presentation and after completion of
94 chemotherapy to assess treatment response. Over the years, much of this
95 controversy arose from the assumption that an extensive staging work-up, while
96 it might result in stage migration, did not influence prognosis or therapy.⁸

97 Recent progress in the field of canine lymphoma is not limited to improvements
98 in determining morphological subtype. Refinements have also been made in
99 molecular diagnosis and detection of minimal residual disease (MRD). A
100 prognostic impact of the presence of MRD as detected by thymidine kinase
101 assay⁹ or PARR (PCR for Antigen Receptor Rearrangement) testing¹⁰ has been
102 demonstrated. While progress has been made in the publication of consensus
103 guidelines concerning the standardization of lymph node assessment by
104 physical examination (VCOG, Veterinary Cooperative Oncology Group), in the
105 light of such recent progress, it can now be considered very likely that these
106 guidelines would tend to overstate complete remission rates and understate
107 progression rates.¹¹

108
109 In order to continue the current trajectory of progress in our understanding and
110 management of canine lymphoma, and to be able to retrospectively evaluate
111 and compare between clinical studies, it is clear that there is a need for greater

accuracy in the staging of lymphoma at first presentation and the assessment of treatment response.

In this systematic review, data that report various staging methods in canine lymphoma are summarized. The main aim was to determine to what extent different approaches to evaluate treatment efficacy were comparable. In conclusion, we will make some recommendations for further studies that may help address significant unresolved clinical issues surrounding the disease.

Methods

Literature search and study selection processes

A literature search limited to manuscripts published from January 1999 to December 2014 was performed. The search was limited to a 15 year period to ensure the studies represented contemporary diagnostic procedures and management options.

A systematic MEDLINE search of articles was conducted by using the following search terms: “lymphoma” AND “dog” OR “canine” AND “treatment” OR “therapy” OR “chemotherapy” OR “immunotherapy” OR “adoptive therapy” AND “prognosis” OR “outcome” OR “assessment” OR “survival” OR “progression” OR “remission” OR “relapse” OR “disease-free”. The following were inclusion criteria for the studies to be selected: the article was published in English; the full text was available for review; the number of cases was more than 5; and finally the study was published in a peer-reviewed journal. Eligible studies for inclusion in the final data analysis were those evaluating the efficacy of first-line protocols for canine multicentric lymphoma. Exclusion criteria were studies

describing dogs with extranodal lymphoma, dogs undergoing rescue treatment, or dogs for which treatment efficacy was not recorded.

After the initial search, article titles and abstracts were first evaluated for relevance and potential exclusion, then the studies included for manuscript review were subjected to full article review. The resulting list was therefore screened for non-research articles, duplicates, case reports and irrelevant references.

Two authors were assigned to a time-period as follows: 1999-2001 MD and OAG, 2002-2004 IG and JH, 2005-2007 TH and GL, 2008-2011 LM and EM, 2012-2014 DS and GAP. Selected papers were independently reviewed on the basis of the selected criteria by the two authors for each assigned time period and a consensus on the requested information was reached.

Data extraction

Studies were selected based on completeness of data and inclusion criteria only. From eligible articles, the following data were extracted: study characteristics (authors, nationality, publishing year, journal), study design (prospective versus retrospective, randomized versus non-randomized, controlled versus non-controlled), recruitment period, recruiting practices/institutions, disease (all histotypes versus B-cell lymphoma versus T-cell lymphoma versus specific histotype), number of enrolled dogs, staging work-up (including complete blood count and serum biochemical profile, urinalysis, thoracic radiographs, abdominal radiographs, abdominal ultrasound, fine-needle aspirate of liver and spleen, bone marrow aspirate, flow cytometry to quantify peripheral blood and bone marrow infiltration, others), diagnosis (histological review with or without immunohistochemistry, cytological review, flow

cytometry), type of chemotherapeutic protocol (drugs used, duration), type of remission assessment (physical examination and subjective assessment of lymph node size reduction/ enlargement, with or without confirmative cytology, flow cytometry, PARR), duration of first remission, and survival time.

Any uncertainty about the inclusion of data from any article was resolved with a consensus meeting. No attempt was made to contact authors for additional information.

Descriptive analysis

Descriptive analysis was performed to present the proportion of studies with each characteristic. Given the small sample size and heterogeneous study methodologies, no statistical comparisons were performed.

Agreement by the Editors and Participants of the European Canine Lymphoma Network

The European Canine Lymphoma Network (ECLN) is a network created in 2009 with the aim of establishing cooperation among different institutions working on canine lymphoma across the fields of diagnosis and therapy.¹² The definition of common guidelines and approaches is one of the main goals of ECLN. This review was submitted to the 25 Editors and Participants of Workgroup 2. The review was planned to be submitted to a peer-reviewed journal only if at least 75% of the participants agreed on its content.

Results

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The initial search yielded over 508 references, many of which were not specifically relevant to our topic. After the exclusion of irrelevant studies, 63 articles that appeared relevant to our aim and that met all study criteria were identified and fully reviewed.^{4,10,13-72} The main characteristics of the included studies are summarized in Table 1.

Among these studies, 40 (63.5%) were from the USA, 9 (14.3%) were from Italy, 3 (4.8%) were from Germany, 3 (4.8%) were from Brazil, 2 (3.2%) were from Japan, 2 (3.2%) were from UK, 2 (3.2%) were from The Netherlands, 1 (1.6%) was from France, and 1 (1.6%) was from Poland.

Forty-five (71.4%) studies were conducted in single centres, 7 (11.1%) were multicentre studies and 6 (9.5%) were undertaken by two centres. The number of recruiting practices was not stated in 5 (7.9%) studies.

Thirty-seven (58.7%) studies were conducted prospectively, 7 of which were randomized controlled trials comparing chemotherapy alone with chemotherapy and steroids, chemotherapy alone and chemo-immunotherapy, chemotherapy alone and chemotherapy plus total body hyperthermia, chemotherapy plus control diet and chemotherapy plus experimental diet, or two different chemotherapy protocols. Three studies were Phase 1 clinical trials. Twenty-five (39.7%) studies were retrospective and the design of 1 (1.6%) study was unclear.

The median number of dogs per study was 46 (mean, 58; range, 7-456; IQR, interquantile range, 63).

Forty-nine (77.8%) studies included all lymphoma histotypes; 10 (15.9%) studies focused on B-cell lymphomas (3 specifically on high-grade B-cell

lymphoma and 2 on diffuse large B-cell lymphoma, DLBCL); 4 (63.5%) focused on T-cell lymphoma (1 specifically on high-grade T-cell lymphoma) (Figure 1). Depending on the study, dogs had diagnostic assessment of disease by cytological review only (n=12; 19%), histological review only (n=17; 27%), cytology or histology (n=20; 31.7%), cytology and histology (n= 13; 20.6%), whereas the diagnostic method was not stated in 1 (1.6%) study. Immunophenotype of disease was determined either by flow cytometry or by immunohistochemistry or by PARR in 47 (74.6%) of the 63 studies examined. Immunophenotype of disease was determined in all dogs in 29 (46%) studies, in the majority to half of the cases in 5 (7.9%) studies, and only occasionally (less than 50% of cases) in 13 (20.6%) studies.

The evaluation of disease extent differed significantly among studies. In 52 (82.5%) studies, staging work-up was described, while in 11 (17.5%) studies the tests performed to assess disease extent were not mentioned. Considering the 52 studies in which staging was described, the most commonly suggested tests included a complete blood cell count (CBC), serum biochemical profile (83.1%), and/or urinalysis (100%), thoracic radiographs (82.7%), abdominal radiographs (28.8%), abdominal ultrasound examination with or without fine-needle aspiration of liver and spleen regardless of their sonographic appearance (59.6%), and bone marrow evaluation (73.1%). In some studies, the following tests were also performed: serology for infectious diseases (3.8%), and echocardiography and/or electrocardiography (9.6%).

For the purpose of this analysis, staging procedures were grouped in the following categories: minimum work-up (including a CBC and serum biochemical profile and/or radiography or ultrasound, and/or bone marrow

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

evaluation; 25 [39.7%] studies) or full staging (including a CBC and serum biochemical profile, thoracic radiography, abdominal ultrasound, and bone marrow evaluation; 27 [42.8%] studies) (Figure 2). When specifically focusing on studies in which a full-staging was suggested, tests were not always performed on all dogs.

The most commonly used first-line treatment protocols included vincristine, cyclophosphamide, doxorubicin, and prednisone, with or without other drugs, radiation therapy or immunotherapy (CHOP-based protocols; 45 studies, 71.4%). Fourteen (22.2%) papers evaluated the efficacy of other drugs or combinations of drugs. The adopted protocol was not described in 4 (6.3%) studies.

The duration of the chemotherapeutic protocols was described in 49 (77.8%) studies, and not reported in 8 (12.7%) studies. In 6 (9.5%) studies, the duration of the protocol depended on treatment response and was therefore variable. When described, the median duration of the chemotherapeutic protocol was 19 weeks (range, 4 to 130 weeks; IQR, 12).

Regarding treatment efficacy, if response to treatment was generically described as “regression of measurable tumours”, it was assumed that peripheral lymph nodes were at least measured. Thus, for the purpose of this review, this type of remission assessment was grouped into the category “subjective or radiological/sonographic measurement of peripheral lymph nodes”. The methods for assessing treatment response varied greatly among studies. In 41 (65.1%) studies, treatment response was based on subjective or radiological/sonographic measurement of peripheral lymph nodes; in none of

1
2
3 268 them, confirmative nodal cytology was described as mandatory. In 2 (3.2%)
4
5 269 studies, a complete end-staging was carried out, including bloodwork,
6
7 270 urinalysis, imaging and confirmative cytology. In 9 (14.3%) studies, minimal
8
9 271 residual disease analysis was carried out, including flow cytometry and/or
10
11 272 PARR. Finally, the methods were not described in 11 (17.5%) studies (Figure
12
13 273 3).
14
15
16 274

17
18 275 The endpoint remission duration was described in 57 (90.5%) studies; the
19
20 276 endpoint survival was reported in slightly fewer studies (n=52; 82.5%).
21
22 277
23
24 278

27 279 **Discussion**

28
29 280 For each dog with suspected multicentric lymphoma, the overall goal is a timely
30
31 281 diagnosis and administration of appropriate therapy. Needless to say, accurate
32
33 282 staging influences management decisions and predicts prognosis for cancer
34
35 283 patients in general. Also, clinical staging procedures allow determination of a
36
37 284 patient's response to therapy. Finally, clinical stage evaluations serve an
38
39 285 important role in allowing the comparison of treatments between studies.
40
41 286

42
43 287 The purpose of this work was to review the last 15 years of published literature
44
45 288 to determine to what extent different approaches to evaluate treatment efficacy
46
47 289 in the first-line setting were comparable. To the authors' knowledge, there are
48
49 290 no other published systematic reviews assessing the methods used for staging
50
51 291 canine lymphoma at diagnosis and post treatment.
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

292 This systematic review identified a total of 63 articles that satisfied the search
293 criteria. The total number of dogs in the current systematic review is relatively
294 large, with a median 46 dogs per study.

295 Based on the results of the current review, certain points of controversy were
296 found.

297

298 First, there was significant variability across studies concerning histotypes. The
299 greatest majority of studies have been severely hampered by the admixture of a
300 variety of lymphoma subtypes in the analysis of outcome, making it difficult to
301 assess the clinical efficacy of any given treatment. Indeed, it has been well
302 documented that canine lymphomas encompass a group of types of tumors,
303 with different biologic behaviors, patterns of chemosensitivity and treatment
304 responses.^{1,2} Thus, clinical trial results need to be interpreted in the context of
305 the distribution of histologic subtypes treated. This, in turn, complicates the
306 assessment of chemotherapy efficacy, making it impossible, in studies
307 describing mixed lymphoma subtypes, to determine whether high or low
308 response rates are due to the specific treatment or to the specific population
309 under study. Only 2 out of the 63 studies evaluated a single lymphoma subtype.

310

311 Second, there were striking differences in the criteria for the diagnosis and the
312 extent of the staging procedures. These differences inevitably have an
313 unquantifiable influence on the patients' final outcome, and preclude meaningful
314 comparisons between studies. Briefly, in 17.5% of the studies the staging work-
315 up was not described. Furthermore, almost half of the studies relied on a
316 minimum work-up. Unfortunately, to date no single diagnostic algorithm
317 sufficiently addresses the complexity and variation in disease patterns of canine

lymphoma. Furthermore, local expertise and financial resources can also influence the approach taken. Doubtless, the different opinions concerning the minimum criteria for the diagnosis of canine lymphoma do result in differences in patient selection for different chemotherapeutic protocols and therefore do bias treatment outcome.

Third, the comparability of efficacy between studies was also hampered by differences in response assessment criteria employed.

The importance of response assessment criteria is well described in the literature: recently, the VCOG developed a consensus document, dictating guidelines to standardize definition of normal lymph node size, when and how responses should be assessed, and definitions for response categories and endpoints.⁸ However, cytological and molecular diagnostic techniques allow one to state that the VCOG guidelines would tend to overstate complete remission rates and understate progression rates.⁸ Indeed, most of the limitations of this document reside in the inter- and intra-observer variability of physical examination, rendering the guidelines not suited for end-staging; furthermore, they do not allow assessment of MRD. A recent study has indeed shown presence of MRD by PARR despite clinical remission in 9 of 12 (75%) dogs with diffuse large B-cell lymphoma.⁹ As a matter of fact, despite the ease and practicality of lymph node measurement, the VCOG guidelines have not been validated in clinical and therapeutic studies.

According to the results obtained here, a good proportion of studies (17.5%) did not describe the methods used for evaluating treatment response at all. The majority of studies relied on subjective or radiological measurement of peripheral lymph nodes, whereas few studies defined treatment response

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

344 based on MRD evaluation. Although the induction of clinical remission is
345 associated with clinical benefit, RECIST criteria are restricted to measuring
346 tumor size, being insensitive to changes in tumor load in other matrices (such
347 as peripheral blood, bone marrow and abdominal organs), and may therefore
348 overestimate the anti-tumor treatment effect.

349
350 In this study it was shown that the absence of accurate diagnostic work-up
351 during the initial and the end-staging may be one of the confounding factors
352 leading to controversial results and different rates of success of antitumoral
353 treatment in the different studies. Clearly, standardization of staging techniques,
354 both initially and after treatment, is needed to decrease, if not eliminate,
355 variability due to selection bias. Until the validity and reliability of measurement
356 tools are ensured, it cannot be accurately determined which of the published
357 treatment protocols will benefit lymphoma dogs. Awareness of these effects for
358 patient selection and for treatment outcome may help in the design of future
359 clinical trials. These trials will require international collaboration and should
360 ideally be designed following multidisciplinary clinical input and include dogs
361 classified according to histological guidelines to ensure homogeneous
362 enrolment.

363
364 The participants in the Clinical Working Group of ECLN make the following
365 concluding observations and recommendations.

366 While the shortcomings of retrospective studies are familiar to all, such clinical
367 studies describing historical actions to real patients will always be of value to
368 our understanding of treatment and disease.

When clinical information concerning canine nodal lymphoma is gained prospectively, thought must be given to the utility of that information for the scientific community at large.

For all dogs enrolled in prospective studies, optimal diagnosis, clinical stage evaluation and response evaluation criteria should comprise as a minimum:

Diagnosis: WHO classification of lymphoma type and/or flow cytometry and cytomorphological analysis to define B/T immunophenotype and morphological subtype within the limits of what is possible using those diagnostic modalities. For the histopathological diagnosis of lymphoma, lymph node excision biopsies (lymphadenectomy) rather than core biopsies are regarded as standard of care.

Clinical Stage: Complete blood count and smear evaluation; thoracic and abdominal imaging (x-ray, ultrasound, CT/MRI as appropriate); cytology of splenic and hepatic aspirates, and bone marrow evaluation prior to initiation of therapy.

Response Evaluation: Two to four weeks following administration of final chemotherapy treatment for discontinuous protocols or four to six months after initiation of therapy for continuous protocols: complete blood count and smear evaluation; thoracic and abdominal imaging (x-ray, ultrasound, CT/MRI as appropriate); cytology of splenic and hepatic aspirates, bone marrow evaluation, and MRD monitoring.

It is recognized that these observations and recommendations are pertinent in the present; future discoveries and trends should lead to their modification. By achieving conformity as suggested, such progress, it is hoped, will be made faster.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

395

For Peer Review Only

References

1. Valli VE, San Myint M, Barthel A, Bienzle D, Caswell J, Colbatzky F, *et al.* Classification of canine malignant lymphomas according to the World Health Organization criteria. *Veterinary Pathology* 2011; **48**: 198-211.
2. Aresu L, Martini V, Rossi F, Vignoli M, Sampaolo M, Aricò A, *et al.* Canine indolent and aggressive lymphoma: clinical spectrum with histologic correlation. *Veterinary and Comparative Oncology* 2013. doi: 10.1111/vco.12048.
3. Marconato L. The staging and treatment of multicentric high-grade lymphoma in dogs: a review of recent developments and future prospects. *The Veterinary Journal* 2011; **188**: 34-38.
4. Valli VE, Kass PH, San Myint M and Scott F. Canine lymphomas: association of classification type, disease stage, tumor subtype, mitotic rate, and treatment with survival. *Veterinary Pathology* 2013; **50**: 738-748.
5. Frantz AM, Sarver AL, Ito D, Phang TL, Karimpour-Fard A, Scott MC, *et al.* Molecular profiling reveals prognostically significant subtypes of canine lymphoma. *Veterinary Pathology* 2013; **50**: 693-703.
6. Klopfleisch R. Personalised medicine in veterinary oncology: One to cure just one. *The Veterinary Journal* doi: 10.1016/j.tvjl.2015.01.004.
7. Ponce F, Marchal T, Magnol JP, Turinelli V, Ledieu D, Bonnefont C, *et al.* A morphological study of 608 cases of canine malignant lymphoma in France with a focus on comparative similarities between canine and human lymphoma morphology. *Veterinary Pathology* 2010; **47**: 414-433.

8. Flory AB, Rassnick KM, Stokol T, Scrivani PV and Erb HN. Stage migration in dogs with lymphoma. *Journal of Veterinary Internal Medicine* 2007; **21**: 1041-1047.
9. von Euler H, Einarsson R, Olsson U, Lagerstedt AS and Eriksson S. Serum thymidine kinase activity in dogs with malignant lymphoma: a potent marker for prognosis and monitoring the disease. *Journal of Veterinary Internal Medicine* 2004; **18**: 696-702.
10. Aresu L, Aricò A, Ferraresso S, Martini V, Comazzi S, Riondato F, *et al.* Minimal residual disease detection by flow cytometry and PARR in lymph node, peripheral blood and bone marrow, following treatment of dogs with diffuse large B-cell lymphoma. *The Veterinary Journal* 2014; **200**: 318-324.
11. Piek CJ, Rutteman GR and Teske E. Evaluation of the results of a L-asparaginase-based continuous chemotherapy protocol versus a short doxorubicin-based induction chemotherapy protocol in dogs with malignant lymphoma *Veterinary Quarterly* 1999; **21**:44-49.
12. Comazzi S, Marconato L, Argyle DJ, Aresu L, Stirn M, Grant IA, *et al.* The European canine lymphoma network: a joining initiative to generate consensus guidelines for the diagnosis and therapy in canine lymphoma and research partnership. *Veterinary and Comparative Oncology* 2015; **13**:494-497.
13. Larue SM, Fox MH, Ogilvie GK, Page RL, Getzy DM, Thrall DE, *et al.* Tumour cell kinetics as predictors of response in canine lymphoma treated with chemotherapy alone or combined with whole body hyperthermia. *International Journal of Hyperthermia* 1999; **15**:475-486.

14. Phillips BS, Kass PH, Naydan DK, Winthrop MD, Griffey SM and Madewell BR. Apoptotic and proliferation indexes in canine lymphoma. *Journal of Veterinary Diagnostic Investigation* 2000; **12**:111-117.
15. Ogilvie GK, Fettman MJ, Mallinckrodt CH, Walton JA, Hansen RA, Davenport DJ, et al. Effect of fish oil, arginine, and doxorubicin chemotherapy on remission and survival time for dogs with lymphoma: a double-blind, randomized placebo-controlled study. *Cancer* 2000; **88**:1916-1928.
16. Chun R, Garrett LD and Vail DM. Evaluation of a high-dose chemotherapy protocol with no maintenance therapy for dogs with lymphoma. *Journal of Veterinary Internal Medicine* 2000; **14**:120-124.
17. Boyce KL and Kitchell BE. Treatment of canine lymphoma with COPLA/LVP. *Journal of the American Animal Hospital Association* 2000; **36**:395-403.
18. Moore AS, Cotter SM, Rand WM, Wood CA, Williams LE, London CA, et al. Evaluation of a discontinuous treatment protocol (VELCAP-S) for canine lymphoma. *Journal of Veterinary Internal Medicine* 2001; **15**:348-354.
19. Dobson JM, Blackwood LB, McInnes EF, Bostock DE, Nicholls P, Hoather TM, et al. Prognostic variables in canine multicentric lymphosarcoma. *Journal of Small Animal Practice* 2001; **42**:377-384.
20. Garrett LD, Thamm DH, Chun R, Dudley R and Vail DM. Evaluation of a 6-month chemotherapy protocol with no maintenance therapy for dogs with lymphoma. *Journal of Veterinary Internal Medicine* 2002; **16**:704-709.

21. Jagielski D, Lechowski R, Hoffmann-Jagielska M and Winiarczyk S. A retrospective study of the incidence and prognostic factors of multicentric lymphoma in dogs (1998-2000). *Journal of Veterinary Medicine. A, Physiology, Pathology and Clinical Medicine* 2002; **49**:419-424.
22. Mutsaers AJ, Glickman NW, DeNicola DB, Widmer WR, Bonney PL, Hahn KA, et al. Evaluation of treatment with doxorubicin and piroxicam or doxorubicin alone for multicentric lymphoma in dogs. *Journal of the American Veterinary Medical Association* 2002; **220**:1813-1817.
23. Morrison-Collister KE, Rassnick KM, Northrup NC, Kristal O, Chretien JD, Williams LE, et al. A combination chemotherapy protocol with MOPP and CCNU consolidation (Tufts VELCAP-SC) for the treatment of canine lymphoma. *Veterinary and Comparative Oncology* 2003; **1**:180-90.
24. Moore AS, Imondi AR, de Souza PL and Wood CA. Intravenous administration of 9-aminocamptothecin to dogs with lymphoma. *Veterinary and Comparative Oncology* 2003; **1**:86-93.
25. Ponce F, Magnol JP, Ledieu D, Marchal T, Turinelli V, Chalvet-Monfray K, et al. Prognostic significance of morphological subtypes in canine malignant lymphomas during chemotherapy. *The Veterinary Journal* 2004; **167**:158-166.
26. Ricci Lucas SR, Pereira Coelho BM, Marquezi ML, Franchini ML, Miyashiro SI, et al. Carmustine, vincristine, and prednisone in the treatment of canine lymphosarcoma. *Journal of the American Animal Hospital Association* 2004; **40**:292-299.
27. Williams LE, Johnson JL, Hauck ML, Ruslander DM, Price GS and Thrall DE. Chemotherapy followed by half-body radiation therapy for canine lymphoma. *Journal of Veterinary Internal Medicine* 2004; **18**:703-709.

28. Gustafson NR, Lana SE, Mayer MN and LaRue SM. A preliminary assessment of whole-body radiotherapy interposed within a chemotherapy protocol for canine lymphoma. *Veterinary and Comparative Oncology* 2004; **2**:125-131.
29. MacDonald VS, Thamm DH, Kurzman ID, Turek MM and Vail DM. Does L-asparaginase influence efficacy or toxicity when added to a standard CHOP protocol for dogs with lymphoma? *Journal of Veterinary Internal Medicine* 2005; **19**:732-736.
30. Simon D, Nolte I, Eberle N, Abbrederis N, Killich M and Hirschberger J. Treatment of dogs with lymphoma using a 12-week, maintenance-free combination chemotherapy protocol. *Journal of Veterinary Internal Medicine* 2006; **20**:948-954.
31. Turner AI, Hahn KA, Rusk A, Gamblin RM, Cosgrove SB, Griffice K, et al. Single agent gemcitabine chemotherapy in dogs with spontaneously occurring lymphoma. *Journal of Veterinary Internal Medicine* 2006; **20**:1384-1388.
32. Siedlecki CT, Kass PH, Jakubiak MJ, Dank G, Lyons J and Kent MS. Evaluation of an actinomycin-D-containing combination chemotherapy protocol with extended maintenance therapy for canine lymphoma. *Canadian Veterinary Journal* 2006; **47**:52-59.
33. Turek MM, Thamm DH, Mitzey A, Kurzman ID, Huelsmeyer MK, Dubielzig RR, et al. Human granulocyte-macrophage colony-stimulating factor DNA cationic-lipid complexed autologous tumour cell vaccination in the treatment of canine B-cell multicentric lymphoma. *Veterinary and Comparative Oncology* 2007; **5**:219-231.

34. Hosoya K, Kisseberth WC, Lord LK, Alvarez FJ, Lara-Garcia A, Kosarek CE, et al. Comparison of COAP and UW-19 protocols for dogs with multicentric lymphoma. *Journal of Veterinary Internal Medicine* 2007; **21**:1355-1363.
35. Kaiser CI, Fidel JL, Roos M and Kaser-Hotz B. Reevaluation of the University of Wisconsin 2-year protocol for treating canine lymphosarcoma. *Journal of the American Animal Hospital Association* 2007; **43**:85-92.
36. Gavazza A, Lubas G, Valori E and Gugliucci B. Retrospective survey of malignant lymphoma cases in the dog: clinical, therapeutical and prognostic features. *Veterinary Research Communications* 2008; **32**:S291-S293.
37. Marconato L, Bonfanti U, Stefanello D, Lorenzo MR, Romanelli G, Comazzi S, et al. Cytosine arabinoside in addition to VCAA-based protocols for the treatment of canine lymphoma with bone marrow involvement: does it make the difference? *Veterinary and Comparative Oncology* 2008; **6**:80-89.
38. Merlo A, Rezende BC, Franchini ML, Monteiro PR and Lucas SR. Serum amyloid A is not a marker for relapse of multicentric lymphoma in dogs. *Veterinary Clinical Pathology* 2008; **37**:79-85.
39. Rebhun RB, Lana SE, Ehrhart EJ, Charles JB and Thamm DH. Comparative analysis of survivin expression in untreated and relapsed canine lymphoma. *Journal of Veterinary Internal Medicine* 2008; **22**:989-95.
40. Simon D, Moreno SN, Hirschberger J, Moritz A, Kohn B, Neumann S, et al. Efficacy of a continuous, multiagent chemotherapeutic protocol versus

- a short-term single-agent protocol in dogs with lymphoma. *Journal of the American Veterinary Medical Association* 2008; **232**:879-885.
41. Gavazza A, Sacchini F, Lubas G, Gugliucci B and Valori E. Clinical, laboratory, diagnostic and prognostic aspects of canine lymphoma: A retrospective study. *Comparative Clinical Pathology* 2009; **18**:291-299.
42. Miller AG, Morley PS, Rao S, Avery AC, Lana SE and Olver CS. Anemia is associated with decreased survival time in dogs with lymphoma. *Journal of Veterinary Internal Medicine* 2009; **23**:116-122.
43. Brodsky EM, Maudlin GN, Lachowicz JL and Post GS. Asparaginase and MOPP treatment of dogs with lymphoma. *Journal of Veterinary Internal Medicine* 2009; **23**:578-584.
44. Daters AT, Mauldin GE, Mauldin GN, Brodsky EM and Post GS. Evaluation of a multidrug chemotherapy protocol with mitoxantrone based maintenance (CHOP-MA) for the treatment of canine lymphoma. *Veterinary and Comparative Oncology* 2010; **8**:11-22.
45. Lori JC, Stein TJ and Thamm DH. Doxorubicin and cyclophosphamide for the treatment of canine lymphoma: a randomized, placebo-controlled study. *Veterinary and Comparative Oncology* 2010; **8**:188-195.
46. Marconato L, Crispino G, Finotello R, Mazzotti S and Zini E. Clinical relevance of serial determinations of lactate dehydrogenase activity used to predict recurrence in dogs with lymphoma. *Journal of the American Veterinary Medical Association* 2010; **236**:969-974.
47. Rassnick KM, Bailey DB, Malone EK, Intile JL, Kiselow MA, Flory AB, et al. Comparison between L-CHOP and an L-CHOP protocol with interposed treatments of CCNU and MOPP (L-CHOP-CCNU-MOPP) for

- lymphoma in dogs. *Veterinary and Comparative Oncology* 2010; **8**:243-53.
48. Sorenmo K, Overley B, Krick E, Ferrara T, LaBlanc A and Shofer F. Outcome and toxicity associated with a dose-intensified, maintenance-free CHOP-based chemotherapy protocol in canine lymphoma: 130 cases. *Veterinary and Comparative Oncology* 2010; **8**:196-208.
49. Yamazaki J, Takahashi M, Setoguchi A, Fujino Y, Ohno K and Tsujimoto H. Monitoring of minimal residual disease (MRD) after multidrug chemotherapy and its correlation to outcome in dogs with lymphoma: a proof-of-concept pilot study. *Journal of Veterinary Internal Medicine* 2010; **24**:897-903.
50. Zenker I, Meichner K, Steinle K, Kessler M and Hirschberger J. Thirteen-week dose-intensifying simultaneous combination chemotherapy protocol for malignant lymphoma in dogs. *The Veterinary Record* 2010; **167**:744-748.
51. Sato M, Yamazaki J, Goto-Koshino Y, Takahashi M, Fujino Y, Ohno K, et al. Evaluation of cytoreductive efficacy of vincristine, cyclophosphamide, and Doxorubicin in dogs with lymphoma by measuring the number of neoplastic lymphoid cells with real-time polymerase chain reaction. *Journal of Veterinary Internal Medicine* 2011; **25**:285-291.
52. Marconato L, Stefanello D, Valenti P, Bonfanti U, Comazzi S, Roccabianca P, et al. Predictors of long-term survival in dogs with high-grade multicentric lymphoma. *Journal of the American Veterinary Medical Association* 2011; **238**:480-485.
53. Perry JA, Thamm DH, Eickhoff J, Avery AC and Dow SW. Increased monocyte chemotactic protein-1 concentration and monocyte count

- independently associate with a poor prognosis in dogs with lymphoma. *Veterinary and Comparative Oncology* 2011; **9**:55-64.
54. Flory AB, Rassnick KM, Erb HN, Garrett LD, Northrup NC, Selting KA, et al. Evaluation of factors associated with second remission in dogs with lymphoma undergoing retreatment with a cyclophosphamide, doxorubicin, vincristine, and prednisone chemotherapy protocol: 95 cases (2000-2007). *Journal of the American Veterinary Medical Association* 2011; **238**:501-506.
55. Rebhun RB, Kent MS, Borroffka SA, Frazier S, Skorupski K and Rodriguez CO. CHOP chemotherapy for the treatment of canine multicentric T-cell lymphoma. *Veterinary and Comparative Oncology* 2011; **9**:38-44.
56. Sorenmo KU, Krick E, Coughlin CM, Overley B, Gregor TP, Vonderheide RH, et al. CD40-activated B cell cancer vaccine improves second clinical remission and survival in privately owned dogs with non-Hodgkin's lymphoma. *PLoS One* 2011; **6**:e24167.
57. O'Connor CM, Sheppard S, Hartline CA, Huls H, Johnson M, Palla SL, et al. Adoptive T-cell therapy improves treatment of canine non-Hodgkin lymphoma post chemotherapy. *Scientific Reports* 2012; **2**:249.
58. Silver M, Rusk A, Phillips B, Beck E, Jankowski M, Philibert J, et al. Evaluation of the oral antimitotic agent (ABT-751) in dogs with lymphoma. *Journal of Veterinary Internal Medicine* 2012; **26**:349-354.
59. Willcox JL, Pruitt A and Suter SE. Autologous peripheral blood hematopoietic cell transplantation in dogs with B-cell lymphoma. *Journal of Veterinary Internal Medicine* 2012; **26**:1155-1163.

60. Vail DM, Husbands BD, Kamerling SG, Simpson H, Kurzman ID and McDonnell A. Phase I study to determine the maximal tolerated dose and dose-limiting toxicities of orally administered idarubicin in dogs with lymphoma. *Journal of Veterinary Internal Medicine* 2012; **26**:608-613.
61. Gentilini F, Turba ME and Forni M. Retrospective monitoring of minimal residual disease using hairpin-shaped clone specific primers in B-cell lymphoma affected dogs. *Veterinary Immunology and Immunopathology* 2013; **153**:279-288.
62. Sato M, Yamzaki J, Goto-Koshino Y, Takahashi M, Fujino Y, Ohno K, et al. The prognostic significance of minimal residual disease in the early phases of chemotherapy in dogs with high-grade B-cell lymphoma. *The Veterinary Journal* 2013; **195**:319-324.
63. Marconato L, Martini V, Aresu L, Sampaolo M, Valentini F, Rinaldi V, et al. Assessment of bone marrow infiltration diagnosed by flow cytometry in canine large B cell lymphoma: prognostic significance and proposal of a cut-off value. *The Veterinary Journal* 2013; **197**:776-781.
64. Elliott JW, Cripps P, Marrington AM, Grant IA and Blackwood L. Epirubicin as part of a multi-agent chemotherapy protocol for canine lymphoma. *Veterinary and Comparative Oncology* 2013; **11**:185-198.
65. Burton JH, Garrett-Mayer E and Thamm DH. Evaluation of a 15-week CHOP protocol for the treatment of canine multicentric lymphoma. *Veterinary and Comparative Oncology* 2013; **11**:306-315.
66. Zandvliet M, Rutteman GR and Teske E. Prednisolone inclusion in a first-line multidrug cytostatic protocol for the treatment of canine lymphoma does not affect therapy results. *The Veterinary Journal* 2013; **197**:656-661.

- 649 67. Warry EE, Willcox JL and Suter SE. Autologous peripheral blood
650 hematopoietic cell transplantation in dogs with T-cell lymphoma. *Journal*
651 *of Veterinary Internal Medicine* 2014; **28**:529-537.
- 652 68. Avery PR, Burton J, Bromberek JL, Seelig DM, Elmslie R, Correa S, et
653 al. Flow cytometric characterization and clinical outcome of CD4+ T-cell
654 lymphoma in dogs: 67 cases. *Journal of Veterinary Internal Medicine*
655 2014; **28**:538-546.
- 656 69. Marconato L, Frayssinet P, Rouquet N, Comazzi S, Leone VF, Laganga
657 P, et al. Randomized, placebo-controlled, double-blinded
658 chemoimmunotherapy clinical trial in a pet dog model of diffuse large B-
659 cell lymphoma. *Clinical Cancer Research* 2014; **20**:668-677.
- 660 70. Mutz M, Boudreaux B, Kearney M, Stroda K, Gaunt S and Shiomitsu K.
661 Prognostic value of baseline absolute lymphocyte concentration and
662 neutrophil/lymphocyte ratio in dogs with newly diagnosed multi-centric
663 lymphoma. *Veterinary and Comparative Oncology* 2015; **13**:337-347.
- 664 71. Lucas SR, Maranhão RC, Guerra JL, Coelho BM, Barboza R and Pozzi
665 DH. Pilot clinical study of carmustine associated with a lipid
666 nanoemulsion in combination with vincristine and prednisone for the
667 treatment of canine lymphoma. *Veterinary and Comparative Oncology*
668 2015; **13**:184-193.
- 669 72. Childress MO, Fulkerson CM, Lahrman SA, Weng HY. Inter- and intra-
670 rater reliability of calliper-based lymph node measurement in dogs with
671 peripheral nodal lymphomas. *Veterinary and Comparative Oncology*
672 2014 Nov 16. doi: 10.1111/vco.12125.
- 673

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Captions to figures

Figure 1. Pie chart showing the distribution of disease types in dogs enrolled in the 63 studies. (DLBCL: diffuse large B-cell lymphoma).

Figure 2. Pie chart showing the staging methods according to the 63 studies.

Figure 3. Pie chart showing the methods used to assess treatment response according to the 63 studies. (LN: lymph node).

Table 1. Characteristics of included studies (chronologic order).

Author (n)	Study type	Disease	Staging methods				Diagnosis methods			Remission assessment			
			B+U	IM	BM	Other*	C	H	PH	LN meas	C	B+IM	MRD
Piek et al 1999 (117)	R	All histotypes	NR				x	(or) x	no	x	no	no	no
Larue et al 1999 (42)	P	All histotypes	NR				no	x	no	x	no	no	no
Phillips et al 2000 (41)	P	All histotypes	x	x	x	no	no	x	x	x	no	no	no
Ogilvie et al 2000 (32)	P	All histotypes	x	x	x	no	no	x	no	x	no	no	no
Chun et al 2000 (49)	P	All histotypes	x	x	x	no	no	x	OCC	NR			
Boyce et al 2000 (75)	NR	All histotypes	x	x	OCC	no	x	(or) x	no	x	no	no	no
Moore et al 2001 (82)	R	All histotypes	x	OCC	OCC	no	x	(or) x	no	x	no	no	no

Dobson et al 2001 (49)	P	All histotypes	x	x	OCC	no	no	x	x	x	no	no	no
Garrett et al 2002 (53)	P	All histotypes	x	x	x	no	x	OCC	no	NR			
Jagielski et al 2002 (43)	R	All histotypes	x	x	x	no	no	x	no	x	no	no	no
Mutsaers et al 2002 (33)	P	All histotypes	x	x	x	no	no	x	no	x	no	x (IM only)	no
Morrison- Collister et al 2003 (94)	R	All histotypes	X	x	x	no	no	x	MOST	x	no	x	no
Moore et al 2003 (10)	P	All histotypes	X	x	x	no	no	x	MOST	x	no	no	no
Ponce et al 2004 (57)	R	All histotypes	x	x	x	no	x	x	x	x	no	no	no
Ricci Lucas et al 2004 (7)	P	All histotypes	x	x	x	no	x	x	no	x	no	no	no
Williams et al	P	All	x	x	OCC	x	x	(or)	x	x	no	x	no

2004 (52)		histotypes						x					
Gustafson et al 2004 (8)	P	All histotypes	x	x	x	no	x	x	x	x	no	no	no
MacDonald et al 2005 (115)	R	All histotypes	x	x	x	no	no	x	x	NR			
Simon et al 2006 (77)	P	All histotypes	x	x	OCC	x	x	(or) x	MOST	x	no	no	no
Turner et al 2006 (21)	P	All histotypes	x	x	x	no	no	x	no	x	no	no	no
Siedlecki et al 2006 (39)	R	All histotypes	x	x	OCC	no	x	(or) x	OCC	x	no	no	no
Turek et al 2007 (52)	P	B-cell	x	x	x	no	no	x	x	x	no	no	no
Hosoya et al 2007 (101)	R	All histotypes	x	x	OCC	no	x	(or) x	OCC	x	no	no	no
Kaiser et al 2007 (96)	R	All histotypes	x	MOST	OCC	no	x	(or) x	no	x	no	no	no
Gavazza et al	R	All	x	no	x	x	x	no	OCC	NR			

2008 (114)		histotypes											
Marconato et al 2008 (17)	P	All histotypes	x	x	x	x	x	no	x	x	x	x	x
Merlo et al 2008 (20)	P	All histotypes	x	x	no	no	x	no	no	x	no	no	no
Rebhun et al 2008 (31)	R	All histotypes	x	x	x	no	no	x	x	x	no	x	no
Simon et al 2008 (106)	P	All histotypes	x	x	OCC	x	x	x	OCC	NR			
Gavazza et al 2009 (114)	R	All histotypes	x	no	x	x	x	no	OCC	x	no	no	no
Miller et al 2009 (84)	R	All histotypes	NR				x	(or) x	OCC	NR			
Brodsky et al 2009 (50)	R	T-cell	x	x	OCC	no	x	(or) x	x	x	no	no	no
Daters et al 2010 (65)	P	All histotypes	x	x	x	no	x	x	no	x	no	no	no
Lori et al 2010	P	All	x	OCC	OCC	no	x	(or)	OCC	x	no	no	no

(32)		histotypes						x					
Marconato et al 2010 (50)	P	All histotypes	x	x	x	x	x	no	x	x	x	x	no
Rassnick et al 2010 (66)	P	All histotypes	x	x	x	no	no	x	x	x	no	x	no
Sorenmo et al 2010 (119)	R	All histotypes	x	x	x	x	x	no	OCC	x	OC C	NR	no
Yamazaki et al 2010 (17)	P	All histotypes	NR				x	no	x	x	no	no	x
Zenker et al 2010 (17)	P	All histotypes	x	x	OCC	no	x	OCC	no	x	no	no	no
Sato et al 2011 (29)	P	B-cell high grade	x	x	no	no	x	no	x	x	no	no	no
Marconato et al 2011 (127)	R	All histotypes	x	x	x	x	x	no	x	x	no	no	no
Perry et al 2011 (26)	R	All histotypes	x	no	OCC	x	no	x	MOST	NR			
Flory et al 2011	R	All	x	OCC	OCC	no	x	(or)	OCC	NR			

(95)		histotypes						x					
Rebhun et al 2011 (24)	R	T-cell (intermediate or high grade)	x	x	x	no	x	(or) x	x	x	no	no	no
Sorenmo et al 2011 (83)	P	B-cell	x	x	x	no	x	x	x	x	x	x	no
O'Connor et al 2012 (8)	P	B-cell	NR				x	x	x	NR			
Silver et al 2012 (19)	P	All histotypes	x	x	no	no	NR			x	no	no	no
Willcox et al 2012 (19)	P	B-cell	NR				x	(or) x	x	no	no	no	x
Vail et al 2012 (19)	P	All histotypes	x	OCC	OCC	no	no	x	MOST	x	no	no	no
Gentilini et al 2013 (8)	R	B-cell	NR				x	no	x	x	no	no	x
Sato et al 2013	P	B-cell high	x	x	no	no	x	no	x	x	no	no	x

(36)		grade											
Valli et al 2013 (456)	R	All histotypes	NR				no	x	x	NR			
Marconato et al 2013 (46)	P	B-cell high grade	x	x	x	x	x	OCC	x	x	x	x	x
Elliott et al 2013 (97)	R	All histotypes	x	MOST	OCC	OCC	x	(or) x	OCC	x	no	no	no
Burton et al 2013 (31)	R	All histotypes	NR				x	(or) x	OCC	x	no	no	no
Zandvliet et al 2013 (81)	P	All histotypes	x	OCC	x	OCC	x	no	x	x	no	no	no
Warry et al 2014 (14)	P	T-cell high grade	NR				x	(or) x	x	no	no	no	x
Avery et al 2014 (67)	R	T-cell	OCC	OCC	OCC	no	x	(or) x	x	NR			
Marconato et al 2014 (19)	P	DLBCL	x	x	x	no	x	x	x	x	x	no	x
Aresu et al 2014	P	DLBCL	x	x	x	no	x	x	x	x	x	x	x

(14)													
Mutz et al 2015** (77)	R	All histotypes	x	x	OCC	no	x	(or) x	OCC	x	no	x	no
Lucas et al 2015** (15)	P	All histotypes	x	x	x	no	x	x	x	x	no	no	no
Childress et al 2015** (15)	P	All histotypes	NR				x	(or) x	no	x	no	no	no

n: number of dogs, P: prospective, R: retrospective, B+U: blood and urinalysis, IM: imaging (thoracic radiography and/or abdominal radiography and/or abdominal ultrasound), BM: bone marrow evaluation, C: cytology, H: histology, PH: phenotype assessment, LN meas: subjective or radiological measurement of peripheral lymph nodes, MRD: minimal residual disease

NR: not reported, OCC: occasionally (<50% of cases), MOST: most cases (>50%)

* Other: infectious disease serology or cardiac evaluation or fine-needle aspiration of liver and spleen regardless of their sonographic appearance

** The papers published in 2015 were available for early view already when this review was started and were therefore included in the analysis.

Figure 1. Pie chart showing the distribution of disease types in dogs enrolled in the 63 studies. (DLBCL: diffuse large B-cell lymphoma).
167x149mm (300 x 300 DPI)

Figure 2. Pie chart showing the staging methods according to the 63 studies.
167x149mm (300 x 300 DPI)

Figure 3. Pie chart showing the methods used to assess treatment response according to the 63 studies. (LN: lymph node). 167x149mm (300 x 300 DPI)