Università degli Studi di Padova

Padua Research Archive - Institutional Repository

Colonic metaplasia of the neo-terminal ileum in Crohn's Disease

Original Citation:

Availability:

This version is available at: 11577/3223650 since: 2017-03-21T12:36:40Z

Publisher:

Published version:

DOI: 10.1016/j.dld.2016.12.032

Terms of use:

Open Access

This article is made available under terms and conditions applicable to Open Access Guidelines, as described at <http://www.unipd.it/download/file/fid/55401> (Italian only)

(Article begins on next page)

Image of the Month

Colonic metaplasia of the neo-terminal ileum in Crohn's Disease

Livia Biancone^{a,*}, Laura Albertoni^b, Marta Ascolani^a, Claudia Mescoli^b, Massimo Rugge^b^a Department of Systems Medicine, Unit of Gastroenterology, University of Rome "Tor Vergata", Rome, Italy^b Department of Medicine, DIMED, Pathology Unit, University of Padova, Padova, Italy

Mucosa of the neo-terminal ileum after ileo-colonic resection for Crohn's Disease (CD), showing sulfomucin-type metaplasia. The High-Iron-Diamine (HID) histochemical reaction documents the neo-expression of sulfomucins in ileal columnar/goblet enterocytes (Fig. 1).

In a previous study [1] we already documented in CD ileum the metaplastic transformation of the native ileal mucosa into colonic type sulfomucin-rich mucosa (i.e. colonic metaplasia).

Submucosal and myenteric plexitis in proximal margins of ileocolonic resection specimens have been associated with CD recurrence [2,3]. Differently, the clinico-pathological usefulness of colonic metaplasia as early predictor of clinical and endoscopic CD recurrence is undefined.

Panels A–C

Ileal mucosa (biopsy sample obtained 6 months after ileo-colonic resection): *Panel A*. In the routine hematoxylin-eosin stain, the original structure of the ileal villi is well recognizable and only mild distortion of the crypts and lymphocytic flogosis is observable. Hematoxylin-eosin, original magnification 10×. *Panel B*. The original structure of the ileal villi is well recognizable (sialomucin secreting goblet cells bordering the villous structure [light blue]). Some of the crypts are bordered by intense-brown stained epithelia (containing sulfomucins: i.e. sulfomucin-secreting metaplasia). High Iron Diamine; original magnification 10×. *Panel C*. At a higher magnification, the sulfomucin secreting goblet cells (intense-brown) bordering the crypts structure (light blue) are more assessable. High Iron Diamine; original magnification 25×.

Panels D–F

Ileal mucosa (biopsy sample obtained 12 months after ileo-colonic resection): *Panel D*. In the routine hematoxylin-eosin stain, the original structure of the ileal villi is almost lost and a more intense flogosis (but no granulocytes) and crypt distortion are observed. Hematoxylin-eosin, original magnification 10×. *Panel E*. The sulfomucin-secreting cells (intense brown) appear along the remaining villous structure, beyond crypts (sulfomucin-secreting metaplasia involving the whole villous structure). High Iron Diamine; original magnification 25×. *Panel F*. At a higher magnification, the sulfomucin secreting crypts (intense-brown) predominate on sialomucin-secreting (light blue) goblet cells (sulfomucin-secreting metaplasia involving the whole epithelium). High Iron Diamine; original magnification 40×.

Fig. 1.

tion are observed. Hematoxylin-eosin, original magnification 10×. *Panel E*. The sulfomucin-secreting cells (intense brown) appear along the remaining villous structure, beyond crypts (sulfomucin-secreting metaplasia involving the whole villous structure). High Iron Diamine; original magnification 25×. *Panel F*. At a higher magnification, the sulfomucin secreting crypts (intense-brown) predominate on sialomucin-secreting (light blue) goblet cells (sulfomucin-secreting metaplasia involving the whole epithelium). High Iron Diamine; original magnification 40×.

References

- [1] Ascolani M, Mescoli C, Palmieri G, et al. Colonic phenotype of the ileum in Crohn's Disease: a prospective study before and after ileo-colonic resection. *Inflammatory Bowel Disease* 2014;20:1555–61.
- [2] Sokol H, Polin V, Lavegne-Slove A, et al. Plexitis as a predictive factor of early postoperative clinical recurrence in Crohn's Disease. *Gut* 2009;58:1218–25.
- [3] Ferrante M, De Hertogh G, Hlavaty T, et al. The value of myenteric plexitis to predict early postoperative Crohn's Disease recurrence. *Gastroenterology* 2006;130:1595–606.

* Corresponding author at: Department of Systems Medicine, Unit of Gastroenterology, University of Rome "Tor Vergata", Via Montpellier 1, 00133 Rome, Italy. Fax: +39 06 20903738.

E-mail address: biancone@med.uniroma2.it (L. Biancone).

<http://dx.doi.org/10.1016/j.dld.2016.12.032>

1590-8658/© 2017 Editrice Gastroenterologica Italiana S.r.l. Published by Elsevier Ltd. All rights reserved.