

# Marine Biotechnology

## The Voltage Dependent Anion Channel (VDAC) of Pacific oysters *Crassostrea gigas* is upaccumulated during infection by the ostreid herpesvirus-1 (OsHV-1): an indicator of the Warburg effect. --Manuscript Draft--

<b>Manuscript Number:</b>	MBTE-D-17-00092R1
<b>Full Title:</b>	The Voltage Dependent Anion Channel (VDAC) of Pacific oysters <i>Crassostrea gigas</i> is upaccumulated during infection by the ostreid herpesvirus-1 (OsHV-1): an indicator of the Warburg effect.
<b>Article Type:</b>	Original Paper
<b>Funding Information:</b>	
<b>Abstract:</b>	Voltage-Dependent Anion Channel (VDAC) is a key mitochondrial protein. VDAC drives cellular energy metabolism by controlling the influx and efflux of metabolites and ions through the mitochondrial membrane, playing a role in its permeabilization. This protein exerts a pivotal role during the White Spot Syndrome Virus (WSSV) infection in shrimp, through its involvement in a particular metabolism that plays in favor of the virus, the Warburg effect. The Warburg effect corresponds to an atypical metabolic shift toward an aerobic glycolysis that provides energy for rapid cell division and resistance to apoptosis. In the Pacific oyster <i>Crassostrea gigas</i> , the Warburg effect occurs during infection by Ostreid herpesvirus (OsHV-1). At present, the role of VDAC in the Warburg effect, OsHV-1 infection and apoptosis is unknown. Here we developed a specific antibody directed against <i>C. gigas</i> VDAC. This tool allowed us to quantify the tissue-specific expression of VDAC, to detect VDAC oligomers, and to follow the amount of VDAC in oysters deployed in the field. We showed that oysters sensitive to a mortality event in the field presented an accumulation of VDAC. Finally, we propose to use VDAC quantification as a tool to measure the oyster susceptibility to OsHV-1 depending on its environment.
<b>Corresponding Author:</b>	Lizenn DELISLE Ifremer Centre de Bretagne FRANCE
<b>Corresponding Author Secondary Information:</b>	
<b>Corresponding Author's Institution:</b>	Ifremer Centre de Bretagne
<b>Corresponding Author's Secondary Institution:</b>	
<b>First Author:</b>	Lizenn Delisle
<b>First Author Secondary Information:</b>	
<b>Order of Authors:</b>	Lizenn Delisle
	Marine Fuhrmann
	Claudie Quéré
	Marianna Pauletto
	Vianney Pichereau
	Fabrice Pernet
	Charlotte Corporeau
<b>Order of Authors Secondary Information:</b>	
<b>Author Comments:</b>	You will find in the summary PDF file, downloaded with the figures of the article, a document for the reviewer 2 (titled: document for reviewer 2), it is not to add in the body of the article but it is provided in addition to the answer to the reviewer 2.

	<p>On December 4th, I modified the table 1 in the manuscript and in the attached files. I reversed the rows and columns to improve the readability of the figure whose quality was poor.</p>
<b>Response to Reviewers:</b>	<p>Brest, December 1, 2017 Dear Editor,</p> <p>Please find enclosed our revised manuscript entitled “The Voltage-Dependent Anion Channel (VDAC) of Pacific oysters <i>Crassostrea gigas</i> is up-accumulated during infection by the ostreid herpes virus-1 (OsHV-1): an indicator of the Warburg effect” submitted for publication in Marine Biotechnology.</p> <p>In this paper, we validate a specific antibody to quantify VDAC in oyster whole body or tissues as an indicator of susceptibility to infection. This study represents a first step toward the analysis of VDAC functioning and mechanisms during an invertebrate Warburg effect, that will be addressed in further fundamental research projects. We carefully addressed all comments made by the two reviewers. In particular, in our revised manuscript, we add new information concerning the occurrence of the Warburg effect in the oyster <i>C. gigas</i>. In particular, a highly relevant reference published in 2017 is now provided, which confirms by metabolomics that <i>C. gigas</i> metabolism shifts toward the Warburg effect when it is infected by OsHV-1 (Young et al., 2017). We hope these results will help to convince the reviewer of the occurrence of the Warburg effect in infected oysters as published in Corporeau et al. (2014) and already demonstrated in shrimp. We hope the manuscript now presents a more convincing case for why VDAC up-accumulation is an indicator of the Warburg effect in infected oyster. As recommended, we also improved technical findings, figure legends, and copy editing. The answers to each point brought up by reviewers are developed below.</p> <p>Thank you in advance for considering this revised manuscript.</p> <p>Sincerely,</p> <p>Lizenn Delisle.</p> <p>The Voltage Dependent Anion Channel (VDAC) of Pacific oysters <i>Crassostrea gigas</i> is up-accumulated during infection by the ostreid herpesvirus-1 (OsHV-1): an indicator of the Warburg effect.</p> <p>Response to the editor Based on the advice received, I have decided that your manuscript could be reconsidered for publication should you be prepared to incorporate major revisions. When preparing your revised manuscript, you are asked to carefully consider the reviewer comments which can be found below, and submit a list of responses to the comments.</p> <p>In this paper we aim to develop a tool to quantify the VDAC protein in oyster, in whole body animals or tissues. This development is a first step toward the study of VDAC functioning and precise mechanisms of invertebrate Warburg effect during oyster infection and mortalities. As recommended, we improved interpretation and discussion of the data to respond to major comments, and we submit a list of responses to each reviewer’s comment (see below).</p> <p>You are kindly requested to also check the website for possible reviewer attachment(s). While submitting, please check the filled in author data carefully and update them if applicable - they need to be complete and correct in order for the revision to be processed further. You will need to log in to the journal and check the submission details as described below.</p> <p>Done.</p> <p>Response to reviewers</p> <p>Reviewer #1: This article developed a specific antibody and quantified the amount of oyster VDAC.</p>

The authors showed a correlation between oyster susceptibility to OsHV-1 and VDAC amount, which indicated the feasibility of using VDAC quantification as a tool to measure the oyster susceptibility to OsHV-1. This is an interesting work, but the following recommendations need to be considered before the manuscript can be accepted for publication.

1. The authors found VDAC is up-accumulated in oysters exposed to OsHV-1 in the field. They considered the up-accumulation as an indicator of the Warburg effect. However, VDAC is not only involved in Warburg effect, but also a key protein regulating mitochondrial membrane permeabilization and apoptosis.

Thanks to the reviewer comment, we better describe the generic role of VDAC in normal physiology as well as in the Warburg effect in cancer cells (see Introduction, lines 73-74 and 82-85).

2. Therefore, the authors should at least measure the level of glycolysis by measuring activities of glycolytic enzymes, such as G6PDH, Triose phosphate isomerase and fructose 1,6-biphosphatase. As little has been done to study the role of oyster VDAC in Warburg effect, it would be great if the authors could test whether there exists an interaction between VDAC and glycolytic enzyme hexokinase, an important component of Warburg effect in cancer cells. Only when oyster VDAC has been demonstrated to be involved in the Warburg effect could the up-accumulation of it be an indicator of Warburg effect

To address the comments in our revised manuscript, we have added some information concerning the occurrence of the Warburg effect in the oyster *C. gigas*. In particular, a highly pertinent reference from 2017 is now provided, which confirms by metabolomics that *C. gigas* metabolism shifts toward the Warburg effect when oysters are infected by OsHV-1 (Young et al., 2017). Taken with the observation of this effect in infected oysters (Corporeau et al., 2014) and shrimp (Chen et al., 2011; Su et al., 2014), we think these new metabolomics results provide convincing evidence of the occurrence of the Warburg effect in infected oysters. Below we provide details about improvements to our manuscript which we hope make a more convincing case that VDAC up-accumulation is an indicator of the Warburg effect in infected oysters:

i) Given the reviewer major concerns, we have changed slightly the focus of the manuscript. Instead of trying to demonstrate the Warburg effect in oyster, we rather rely on the fact that the Warburg effect is a known required mechanism for infection and mortalities in *C. gigas*, as ascertained by the recent publication of Young et al. (2017). We try to be clearer in the revised manuscript that we would like to provide a biochemical tool to quantify VDAC, rather to describe the precise mechanisms of action of VDAC and the Warburg effect in oyster (see Introduction, lines 91-94 and 101). This is the reason we didn't perform enzymatic assays, biochemical studies, or immuno-precipitation.

ii) In fact, the study of VDAC functioning and the precise mechanisms of the Warburg effect in *C. gigas* need further investigation, in samples that will be designed for this purpose. We are developing fundamental research projects aiming to compare the precise mechanisms of VDAC action between oyster and other model organisms (mouse/yeast). These projects will use metabolic measurements and enzymatic assays to describe the dynamic of the metabolic shift toward the Warburg effect. The Cg-VDAC antibody developed here will be used to identify VDAC partners by immunoprecipitation studies. In the work presented here, we wanted to validate the specific anti-VDAC antibody in oyster whole body animals or tissues, as a first step toward the study of VDAC functioning and mechanisms of the invertebrate Warburg effect.

#### c. Minor comments

Line 63: a comma should be added between "2006)" and "Kaposi's Done.

Line 69-71: Results in these two references did not link the accumulation of the protein VDAC with metabolic shift toward the Warburg effect in shrimp. Correct the expression or add some new references.

As recommended, we corrected the expression (see lines 69-71).

Line 70 & 84: "Han-Ching Wang" should be "Wang", please also modify it at the Reference section

Done.

Line 245: "Any" should be "No"

Done

Line 265: "Fig 2C" should be "Fig 2c"

Done

Line 367-368: Results in this reference did not show the role of VDAC in infection stages. Correct the expression or change the references. Line 369-370: Results in this reference did not demonstrate the interaction between VDAC and HK during WSSV infection in shrimp. Correct the expression.

As recommended, we tried to clarify the text (see lines 386-393).

Reviewer #2:

The Voltage-Dependent Anion Channel (VDAC) is a key mitochondrial protein related to cellular energy metabolism. The manuscript MBTE-D-17-00092 entitled "The Voltage Dependent Anion Channel (VDAC) of Pacific oysters *Crassostrea gigas* is up-accumulated during infection by the ostreid herpesvirus-1 (OsHV-1): an indicator of the Warburg effect" by Lizenn Delisle and colleagues developed a specific antibody directed against *Crassostrea gigas* VDAC. Then they used this tool to quantify the tissue-specific expression of VDAC, to detect VDAC oligomers, and to follow the amount of VDAC in oysters deployed in the field. The results suggested that oysters sensitive to a mortality event in the field presented an accumulation of VDAC. The topic is interest and the paper is technically sound, but it requires a few controls with different technical approaches to fully validate the data. Although the manuscript is well written and the figures are clear. Nevertheless, the figure legends need to be improved. The description of the figures in the text are incomplete.

1. In Figure 1, Immuno-detection on western-blot with 10 to 30 µg of protein lysates and using anti-Cg-VDAC as primary antibody. VDAC was detected at 30 kDa (arrow), 45 kDa (line), 60 kDa and 90 kDa (asterisks). But in Figure 2, VDAC was detected at 30 kDa (arrows), 45 kDa (line), 60 kDa, 90 kDa or 120 kDa (asterisks) depending on the tissue. In Figure 3, VDAC was detected in spots at 30 kDa (arrows), 45 kDa (line) or 60 kDa (asterisks) depending on the tissue. The author needs to provide a reasonable explanation.

As recommended, we tried to clarify these points (see lines 254-256; 261-263, 292-296, 355-371). The bands corresponding to VDAC oligomers detected in gels can vary between immunoblots for some technical reasons: i) signals can be masked due to high background level of chemiluminescence, or ii) oligomers can be disrupted due to the conditions of protein extract denaturation before immunoblots:

i) Figures 1-3 presented in the manuscript show the Image analysis of immunoblots obtained through G:Box software with the background signal removed (see lines 191-193). The level of the background signal was higher with whole body samples (Fig.1) than tissue samples (Fig.2). This could explain why we cannot detect the size of oligomers at 120 kDa in whole body samples (Fig. 1), which does not necessarily mean that the oligomer at 120 kDa does not exist. In fact, SDS PAGE immunoblots were more resolute in tissue samples (enrichment in specific tissue proteins) than whole body samples (mix of all proteins from the animal, less specific signal detected).

ii) In Figure 3, two-dimensional electrophoresis was performed on tissue samples. The 2D immunoblots are much more thorough in terms of how proteins are treated than SDS PAGE immunoblots. The sample denaturation using 2D (urea, thiourea, CHAPS, DTT, iodoacetamide, mercaptoethanol, heat, SDS) is more complete and efficient than in monodimensional electrophoresis (mercaptoethanol, heat, SDS). Thus, we can consider that oligomers at size 90 and 120 kDa detected in SDS PAGE using tissue protein extracts (Fig. 2) were undetectable in 2D (Fig. 3) because of destructure, and that might be due to strong denaturation of tissue protein extracts in 2D.

2. Page 13th and line 281-283. Textual material 'VDAC was detected in a spot at 30 kDa (pI= 8 and 8.3) with a shift at around 32 kDa (pI=7.6) that might be due to heart-specific post-translational modifications such as phosphorylation.' I suggest the authors to provide some experimental results confirm the speculative conclusions.

Thanks to the reviewer, we clarified this point and tried to be less conclusive. We decided to rewrite this paragraph to be less speculative in its conclusions. As published in Martel et al. 2014, oligomerization and a lot of post-translational modifications are now described for VDAC in a lot of species and control VDAC functioning. Martel et al. (2014) describe the many VDAC post-translational modifications that can occur, such as phosphorylation, acetylation, S-nitrosylation, all of which may influence the interactome and the activity of VDAC (Martel et al., 2014). In our case, further analysis should be carried out to verify the identities of the putative post-translational modifications or oligomerization found at 45, 60, 90 and 120 kDa in western blotting. Further studies must be designed to investigate the post-translational modifications and oligomerization of Cg VDAC, and we cannot address this question in our samples.

3. The Figure 4 only represented the relative level of VDAC (30 kDa) in oysters. I think needs to provide all VDAC, such as 45 kDa, 60 kDa and 90 kDa, even 120 kDa.

In Figure 4, the electrophoretic profile of VDAC in SDS-PAGE using whole body protein extracts from animals coming from the field presented a major band at 30 kDa, and other sizes (muscle 45 kDa or oligomers) remained less detectable. We present to the reviewer one image of blot obtained from whole body proteins of animals from the field. We can see that VDAC is mainly detected at 30 kDa.

We chose to analyze the link between mortality and relative level of VDAC at 30 kDa for statistical analysis. Indeed, as shown in Figure 2, the statistical analysis between relative level of VDAC at 30 kDa and citrate synthase activity was conclusive, meaning that the analysis of VDAC at 30 kDa can be informative, even if it doesn't represent all the VDAC oligomers. We clarified this point in the discussion (see lines 395-399) and in figure legends (see lines 259; 262-263; 304-305).

3 Other more, the author needs to provide the reference protein electrophoretogram in Figure 4d.

We present to the reviewer in attached file, one blot and its corresponding Ponceau staining, which confirms equal amount of total protein loaded onto gels. As recommended, we improved figure legends (see lines 279-280; 322-324). We hope our response can answer the reviewer's concerns.

[Click here to view linked References](#)

1  
2  
3  
4  
5  
6 1 The Voltage Dependent Anion Channel (VDAC) of Pacific  
7  
8 2 oysters *Crassostrea gigas* is upaccumulated during infection by  
9  
10 3 the ostreid herpesvirus-1 (OsHV-1): an indicator of the Warburg  
11  
12 4 effect.  
13  
14 5

15 6 Lizenn Delisle<sup>1\*</sup>, Marine Fuhrmann<sup>1</sup>, Claudie Quéré<sup>1</sup>, Marianna Pauletto<sup>2</sup>, Vianney Pichereau<sup>3</sup>,  
16  
17 7 Fabrice Pernet<sup>1</sup> and Charlotte Corporeau<sup>1</sup>.  
18

19  
20 8 <sup>1</sup> Ifremer, UMR 6539 CNRS/UBO/IRD/Ifremer, Laboratoire des sciences de l'Environnement  
21  
22 9 Marin (LEMAR), 29280 Plouzané, France.  
23

24  
25 10 <sup>2</sup> Department of Comparative Biomedicine and Food Science. University of Padova, Viale  
26  
27 11 dell'Università 16, 35020 Legnaro, Padova, Italy.  
28

29  
30 12 <sup>3</sup> Université de Bretagne Occidentale, UMR 6539 CNRS/UBO/IRD/Ifremer, Laboratoire des  
31  
32 13 sciences de l'Environnement Marin (LEMAR), 29280 Plouzané, France  
33

34  
35 14 \* **Corresponding author:** Lizenn Delisle, Centre Ifremer de Bretagne, CS 10070, 29280 Plouzané,  
36  
37 15 France. Tél: +33 2 98 22 43 86. Fax: + 33 2 98 22 46 53. E-mail: Lizenn.Delisle@ifremer.fr  
38  
39

40 16 **Keywords:** Voltage dependent anion channel, Warburg effect, *Crassostrea gigas*, Ostreid herpes  
41  
42 17 virus.  
43  
44

## 45 46 47 18 **ABSTRACT**

48  
49 19 Voltage-Dependent Anion Channel (VDAC) is a key mitochondrial protein. VDAC drives cellular  
50  
51 20 energy metabolism by controlling the influx and efflux of metabolites and ions through the  
52  
53 21 mitochondrial membrane, playing a role in its permeabilization. This protein exerts a pivotal role  
54  
55 22 during the White Spot Syndrome Virus (WSSV) infection in shrimp, through its involvement in a  
56  
57 23 particular metabolism that plays in favor of the virus, the Warburg effect. The Warburg effect  
58  
59  
60  
61  
62  
63  
64  
65

corresponds to an atypical metabolic shift toward an aerobic glycolysis that provides energy for rapid cell division and resistance to apoptosis. In the Pacific oyster *Crassostrea gigas*, the Warburg effect occurs during infection by Ostreid herpesvirus (OsHV-1). At present, the role of VDAC in the Warburg effect, OsHV-1 infection and apoptosis is unknown. Here we developed a specific antibody directed against *C. gigas* VDAC. This tool allowed us to quantify the tissue-specific expression of VDAC, to detect VDAC oligomers, and to follow the amount of VDAC in oysters deployed in the field. We showed that oysters sensitive to a mortality event in the field presented an accumulation of VDAC. Finally, we propose to use VDAC quantification as a tool to measure the oyster susceptibility to OsHV-1 depending on its environment.

## INTRODUCTION

Since 2008, massive mortality events of young oysters *C. gigas* have been reported in France (Miossec et al. 2009; EFSA 2010; Barbosa Solomieu et al. 2015; Pernet et al. 2016). These mortality events are associated with the infection of oysters with a newly described genotype ( $\mu$ Var) of Ostreid herpesvirus 1 (OsHV-1) (Segarra et al. 2010). A causal relationship between OsHV-1 and oyster mortality has been established (Schikorski et al. 2011; EFSA 2015). OsHV-1 is a double strand DNA virus which belongs to Malacoherpesviridae family (Davison 2005; Segarra et al. 2010) and is now distributed along the European coastline from Portugal to Scandinavia, and closely related variants have been detected in Australia, New Zealand and Asia (EFSA 2015; Barbosa Solomieu et al. 2015; Pernet et al. 2016). Mortalities of juveniles range from 40 to 100%, resulting in huge losses in Pacific oyster production (Pernet et al. 2012; Dégremont 2013).

To date, there is only limited data on how OsHV-1 interacts with its host, and the pathogenesis of the disease is not completely understood. The first study of this interaction, used global protein

expression profiling to investigate the oyster cellular response to OsHV-1 infection, was carried out in 2014 (Corporeau et al. 2014). This study demonstrated that during early stages of infection (2 days after injection), several biological pathways were modulated and that infected oysters exhibited an increased glycolysis and accumulation of the protein Voltage-Dependent Anion Channel (Cg-VDAC) that reflected a “Warburg effect” (Corporeau et al. 2014). Recently, Li et al. further illustrated the up-regulation of the Cg-VDAC transcript in hemolymph of OsHV-1 infected oysters (Li et al. 2016).

The Warburg effect was first described by Otto Warburg in 1930s in cancer cells (Warburg 1956), and partly results from deregulation of cellular energy pathways (Poliseno 2012). Cells experiencing the Warburg effect show a metabolic shift toward an “aerobic glycolysis”, which presents several benefits to support the high energy and macromolecular synthesis requirement in rapidly dividing cells (Pedersen 2007; Vander Heiden et al. 2009; Puyraimond-Zemmour and Vignot 2013). Several human viruses reprogram the host energy metabolism toward the Warburg effect to support the biosynthesis of viral building blocks (Mesri et al. 2014): Human Papillomavirus (HPV) (Guo et al. 2014), Human Cytomegalovirus (HCMV;  $\beta$ -Herpesvirus) (Munger et al. 2006), Kaposi’s Sarcoma Herpesvirus (KSHV) (Delgado et al. 2010) or Hepatitis C virus (Diamond et al. 2010).

To date, the Warburg effect has been shown to occur in the shrimp *Litopenaeus vannamei* as a metabolic shift that provides cellular energy and building blocks during the replication phase of the White Spot Syndrome Virus (WSSV) (Chen et al. 2011; Su et al. 2014). The WSSV infection is a lethal disease that can cause up to 100% mortality in 10 days (Chen et al. 2011; Su et al. 2014). In *Litopenaeus vannamei*, the Warburg effect is an essential component of the host-viral interaction, providing essential energy for successful WSSV viral replication (Wang et al. 2010; Su et al. 2014).



The Voltage-Dependent Anion Channel (VDAC) is a mitochondrial protein that plays a pivotal role in normal cells, but is also involved in the Warburg effect, occurring with cancer and viral infection in mammals. In normal cells, VDAC is a major membrane protein located in the mitochondrial outer membrane that controls metabolism and apoptosis, and it is considered as a multiple stress sensor, being an apoptotic checkpoint during stress and pathological conditions (Lemasters and Holmuhamedov 2006; Martel et al. 2014; Brahimi-Horn et al. 2015). The VDAC pore mediates the transport of metabolites such as ADP, ATP and NADH, ions and even larger molecules up to 4-6 kDa (Rostovtseva et al. 2002; Naghdi and Hajnóczy 2016). This channel is a key protein that drives cellular energy metabolism by controlling the influx and efflux of metabolites and ions, and participates in mitochondrial membrane permeabilization (Martel et al. 2014). VDAC acts as a platform for many proteins supporting glycolysis and prevents apoptosis by interacting with hexokinase, or members of the Bcl-2 family, respectively. VDAC is thus involved in the metabolic reprogramming of cancer cells toward the Warburg effect (Mazure 2017). In shrimp, when the expression of VDAC is silenced by RNA interference (RNAi) before WSSV infection, the mortality decreases by 50 %, the detection of WSSV DNA drop markedly, and the Warburg effect does not occur (Wang et al. 2010; Chen et al. 2011). The accumulation of VDAC during infection has also been reported in some other species including the flounder *Paralichthys olivaceus* infected with the *Scophthalmus maximus* rhabdovirus (Lü et al. 2007), and during Grass carp hemorrhagic disease in the grass carp *Ctenopharyngodon idella* (Shen et al. 2014). In *Crassostrea gigas* infected by OsHV-1, both the mRNA (Renault et al. 2011; Li et al. 2016) and the protein VDAC (Corporeau et al. 2014; Young et al. 2017) were up-accumulated during the infection processes linked with the metabolic shift toward the Warburg effect (Young et al. 2017). In this context, characterizing the Cg-VDAC protein and quantifying its accumulation becomes a key component for following OsHV-1 infection processes in oyster. VDAC was recently described

in oyster (Li et al. 2016). Like other invertebrates, oysters have only one type of VDAC. Cg-VDAC clustered into the group of VDAC 2, strongly conserved gene from cnidarians to mammals. This study showed that VDAC transcripts were expressed during all oyster developmental stages and in all tissues at adult stage.

As a first step toward the study of VDAC functioning in infection processes of *C. gigas*, the objectives of the present study are: (1) producing and validating a specific antibody directed against *C. gigas* VDAC, (2) analyzing VDAC tissue-specific expression and electrophoretic profiles, and finally (3) assessing the VDAC amount in oysters exposed to OsHV-1 in the field.

## **MATERIALS AND METHODS**

### **Experimental design**

#### **Ethics Statement**

The Pacific oyster, *C. gigas*, used in this study is a marine-cultured animal and cultured in the Ifremer facilities in Argenton (Brittany, France; 48°34'30''N, 4°36'18'' W). All of the experiments were conducted according to local and national regulations. Permission for deploying oysters outside of farming areas was issued by the French Ministry of Ecology and Sustainable Development, dept. of maritime affairs, in February 2013. For locations within farming areas, the owner of the farm gave permission to conduct the study on this site. The present field studies did not involve endangered or protected species.

#### **Expt. 1. Validation of a specific antibody directed against *C. gigas* VDAC and tissue-specific analysis of VDAC**

Specific-pathogen-free (SPF) oysters were produced according (Petton et al. 2015). Spawning occurred on 18 August 2014 (cohort NSI 01/15) in Ifremer facilities in Argenton (Brittany, France; 48°34'30''N, 4°36'18'' W). The fecundation rates were up to 90%. The embryos developed in 150L tanks at 21°C for 48h, and D-larvae were transferred to flow-through rearing systems at 25°C. After 15 days, competent larvae were collected and allowed to settle in downwellers. On 6 October 2014, oysters were transferred to Ifremer facilities in Bouin (46°57' N - 2°02' O). Before being transferred to Ifremer facilities in Argenton, oysters were deployed in farming area located in the Bay of Brest at Pointe du Chateau (48° 20' 06.19'' N, 4° 19' 06.37'' W) for 11 months since 5 Mars 2015 (Petton et al. 2013). Oysters were sampled on 15 February 2016. The fleshs of 16 oysters were pooled, flash-frozen, crushed and stored in liquid nitrogen for further validation of the antibody. The mantle, gills, digestive gland, labial palp, striated and smooth adductor muscle, heart and visceral ganglia were carefully dissected from 50 oysters on ice, immediately flash-frozen in liquid nitrogen, pooled together by tissues, crushed and stored in liquid nitrogen.

## **Expt 2. Quantification of the protein VDAC in oysters exposed to OsHV-1 in the field.**

Specific-pathogen-free (SPF) oysters were produced according to Petton *et al* (Petton et al. 2015). Animals were reared under controlled conditions until the age of 8 months (mean individual wet mass = 0.51 g). The oysters were screened for the herpesvirus by qPCR at the different stages of production, and it was undetected in all cases. These SPF oysters (also called “sentinel oysters”) were deployed at 46 sites located along an inshore-offshore gradient in the Mor-Braz area, South Brittany (France). This deployment took place before the start of a disease induced mortality event on 8 April 2013 that lasted for 171 days until 26 September 2013 (Pernet et al. in prep). At each site, 16 small mesh bags containing 85 individual oysters were grouped in one big mesh bag. These bags were attached to iron tables for the sites situated in the intertidal farming area or immersed

vertically at 2 meters depth and attached to a mooring point for the sites in the offshore area. The oysters were sampled 15 times at each site at low tide slack water  $\pm$  2 hours. Upon arrival in the laboratory, live and dead oysters were counted to evaluate survival. Individual shell length and wet mass were measured on a sub-sample of 25 live oysters per bag. The soft tissues of these oysters were removed from the shells, pooled together, dipped into liquid nitrogen and stored at  $-80^{\circ}\text{C}$  until laboratory analyses. Western blot analyses were conducted on samples collected at two inshore sites located within the oyster farming area where mortality occurred the earliest between 7 and 14 June (sites 37 and 39, **Fig 1**), at two offshore sites where mortality occurred later, from 6 to 14 July (sites 33 and 38) and at two sites where no mortality occurred (sites 32 and 36). Analyses were conducted on samples collected before (30 April and 27 May), at the onset (7 and 14 June) and during the earliest mortality event (20 June).

## **Total protein extraction**

Total protein extraction was performed using 1 g of oyster powder (flesh or tissues) that was homogenized with a Polytron® PT 2500 E (Kinematica). Proteins were solubilized during 40 min at  $4^{\circ}\text{C}$  by adding 5 mL of lysis buffer (Guévelou et al. 2013b) (150 mM NaCl, 10 mM Tris pH 7.4, 1 mM EDTA, 1 mM EGTA, 1% Triton X-100, and 0.5% Igepal; pH 7.4 at  $4^{\circ}\text{C}$ ) containing phosphatase and protease inhibitors (1% of Phosphatase inhibitor cocktail II [Sigma-Aldrich], 2% of NaPPi 250 mM, and 1 tablet of complete EDTA free protease inhibitor cocktail [Roche] in 25 mL of lysis buffer). Solubilized proteins were extracted by centrifugation at 4,000 g for 1 h at  $4^{\circ}\text{C}$  to eliminate lipids and cellular debris. The phase containing proteins was then collected and centrifuged at 10,000 g for 45 min at  $4^{\circ}\text{C}$ . Total protein content in each lysate was analyzed using the DC protein assay (Bio-Rad), in 96-well microplates (Nunc™) using a microplate reader (Bio-

Tek®Synergy™ HT). Concentration was obtained using Gen5 version 2.03 software (Bio-Tek).

The resulting lysates were divided in aliquots and stored at -80°C for further analysis.

## Antibody

The *Crassostrea gigas* VDAC protein BAF63641.1 is a 280 amino acids protein with a predicted size of 30.33 kDa and a pI at 8.25 ([http://web.expasy.org/compute\\_pi](http://web.expasy.org/compute_pi)). A polyclonal anti-Cg - VDAC antibody was produced in rabbit by Eurogentec (France) using the protocol describe in (Fabioux et al. 2009), against one peptide (104-QTGTKSGKIKTSYKM-118) located in the middle of the *C. gigas* VDAC protein sequence, in a turn and  $\beta$  structure.

The antibody was purified and analyzed by indirect Elisa against the purified peptide to compare with pre-immune serum, large bleed and final bleed (Eurogentec, France). Purified anti-CgVDAC antibody was provided in PBS-BSA 0.1% with thimerosal (0.01 %) as preservative and was diluted vol/vol in glycerol for preservation in aliquots at -20 °C.

## Immunodetection on Western-blot

Immunodetection on western-blot was done after mono-dimensional electrophoresis of total protein lysates: 10  $\mu$ g, 20  $\mu$ g or 30  $\mu$ g. Protein lysates were heated in Laemmli buffer for 10 minutes at 100°C and loaded onto 4-15% SDS-Page (Criterion® TGX™ Precast Gels Bio-Rad, Hercules, CA, USA) in parallel with broad range SDS-Polyacrylamide gel electrophoresis (PAGE) molecular weight markers (Precision Plus Protein™Dual Color Standard, Bio-Rad, Hercules, CA, USA). SDS-PAGE run at 200 V constant voltage, 40 mA, for 10 minutes then 200 V constant voltage, 80 mA, for 40 minutes. Proteins were then transferred onto a PVDF membrane (Trans-Blot®Turbo™Midi PVDF Transfer Packs, Bio-Rad, Hercules, CA, USA) using the Trans-Blot®Turbo™Transfer System (Bio-Rad, Hercules, CA, USA). For the immunodetection, we used

the anti Cg-VDAC (Eurogentec; dilution 1:5,000 in PBS-BSA 3% - Tween 1%), or the preimmune serum as the primary antibody, overnight at 4°C. Blots were revealed using a secondary horseradish peroxidase-linked goat anti-rabbit antibody (dilution 1:2500 in PBS-BSA 3% - Tween 1%) and a horseradish peroxidase detection kit (GE-Healthcare). The relative amount of protein detected was quantified using Gel Imaging for fluorescence and chemiluminescence G:Box Chemi XX6 (Syngene, Gene tools software- Syngene) with the background signal removed. The value obtained was expressed in OD/mm<sup>2</sup> and represents the band intensity. To ensure that identical amounts of total protein samples were loaded into gels, membranes were stained 5 minutes with Ponceau S (0.2% with TCA 0.3%, 5-sulfosalicylic acid 3%), then rinsed in distilled water under gentle shaking until the background signal had been removed.

## **Two-dimensional electrophoresis (2-DE) and immunodetection**

Anti-Cg-VDAC was used for immunodetection on two-dimensional electrophoresis (2-DE) followed by western-blot using mantle, gills, smooth adductor muscle or heart protein lysates. For 2-DE, 500 µg of tissue protein lysates were precipitated and desalted by adding 4 volumes of TCA 20% during 2 h at 4°C, followed by centrifugation at 12.000 g for 15 min at 4°C. Pellets were washed 20 times with 80% acetone in 0.05M Tris-HCl, pH 8. Proteins were resuspended in DeStreak rehydration solution (GE Healthcare). Protein concentrations were determined using a modified Bradford assay (Ramagli 1999) and all samples were adjusted to 200 µg in 125 µl of DeStreak rehydration solution (GE Healthcare) containing 1% IPG buffer, then samples were placed at room temperature for 1 hour before isoelectric focusing (IEF). The first dimension was conducted on Bio-Rad protean IEF Cell System <sup>TM</sup>, (Bio-Rad, Hercules, CA, USA). Samples were loaded onto each strip (Immobiline DryStrip pH 4-7, 7 cm, GE Healthcare) and passive rehydration was allowed at room temperature overnight using mineral oil to prevent sample evaporation. The

IEF was carried out at 20°C in four steps: a calibration step at 250 V constant voltage for 10 min, an active hydration step at 250V for 30 min, a continuous increase in voltage up to 3500 V over 2 h, then kept at 3500 V for 2 h. Finally, 50 V were maintained for 1 h. Before the second dimension, IPG strips were equilibrated for 15 min in a solution containing 6 M urea, 2% SDS, 30% glycerol, and 1% DTT in 0.05 M Tris-HCL, pH 8.8. The strips were then further equilibrated for 15 min in a similar buffer in which DTT was replaced with 2.5% iodoacetamide to alkylate the proteins. For the second dimension, 10% acrylamide gels were used and run on Mini protean tetra cell system (Bio-Rad, Hercules, CA, USA). The gels were loaded with broad range SDS-PAGE molecular weight markers (Precision Plus Protein™Dual Color Standard, Bio-Rad, Hercules, CA, USA). Migration was carried out at 200 V constant voltage, 40 mA, 10 min, and 200 V constant voltage, 80 mA, 1 h. After 2-DE, immunodetection on western-blot was performed using anti-Cg-VDAC as described above.

### **Citrate synthase activity**

Citrate synthase (CS; EC 2.3.3.1) activity was measured in oyster tissues using 20 µl of total protein lysate obtained as describe before. CS assay buffer contains 100 mM Tris/HCl at pH 8, 0.1 mM 5,5'-dithio-bis-[2-nitrobenzoic] acid (DNTB), 0.2 mM acetyl-coenzyme A, and 0.5 mM oxaloacetate. CS activity is measured by following the increase in TNB absorbance for 10 min at 412 nm using a Synergy HT microplate reader (BioTek). Enzyme activity was related to the total protein concentration of each sample.

### **Statistical analysis**

Statistical analyses were performed using R software (<http://www.R-project.org>). For all tests, the differences were accepted as statistically significant at the 95% of confidence level ( $p < 0.05$ ).

Linear regressions were used to investigate the relation between VDAC relative protein level (30 kDa) and the activity of citrate synthase among tissues (Expt 1). Analyses of Variance (ANOVA) followed by Tukey's post hoc test were conducted to investigate the effect of tissues (Expt 1) and time and sites (Expt 2) on VDAC relative protein levels.

## VDAC amino acid sequence comparison

In order to evaluate the amino acid sequence conservation of the VDAC epitope across bivalves, we searched the VDAC transcript through several transcriptome assemblies. Contigs putatively encoding VDAC were identified through BlastX similarity searches against the non-redundant protein sequences database (nr). Coding sequences were predicted by similarity to *C. gigas* VDAC transcript (AB262088.1) and amino acid sequences were obtained by means of ExPASy translate tool (<http://web.expasy.org/tools/translate/>). The predicted amino acid sequences are aligned to *C. gigas* VDAC protein (BAF63641.1) by using BlastP.

## RESULTS

### Validation of anti-Cg-VDAC

As demonstrated by immuno-detection on western-blot, the purified synthetic polyclonal anti-Cg-VDAC (Eurogentec) strongly and rapidly recognized a band with an apparent molecular weight of 30 kDa in oyster flesh (**Fig 1a**), which corresponds to the predicted size for VDAC protein in *C. gigas* (GI:148717311). No signal was revealed when primary antibody was replaced by pre-immune serum (**Fig 1b**). As a result, we validated anti-Cg-VDAC as a specific tool to quantify the amount of VDAC in *C. gigas*. Anti-Cg-VDAC also detected a signal at around 45, 60 and 90 kDa (**Fig 1a**). These signal likely corresponded to the size of multimeric forms of VDAC, i.e. dimers



(60 kDa) and trimers (90 kDa). The existence of monomers to tetramers and higher oligomers of VDAC has already been characterized in many other species (Hoogenboom et al. 2007).

### **Fig 1: Validation of anti-Cg-VDAC antibody.**

Immuno-detection on western-blot with 10 to 30 µg of protein lysates from whole-body protein extracts using (a) anti-Cg-VDAC as primary antibody (dilution 1:5000) or (b) by pre-immune serum. VDAC was detected at 30 kDa (arrow), 45 kDa (line), 60 kDa and 90 kDa (asterisks). The bands detected at 60 and 90 kDa in SDS PAGE immunoblots correspond to the size of VDAC oligomers.

### **Tissue-specific relative abundance of VDAC**

The 30 kDa VDAC was constitutively expressed in the eight tissues tested (**Fig 2**) in accordance with the Cg VDAC mRNA expression pattern (Li et al. 2016). Oligomers at 60 kDa, 90 and 120 kDa were also detected at low levels, depending on the tissue (**Fig 2a**). The 30 kDa VDAC was less abundant in striated and smooth adductor muscle than in other tissues (**Fig 2b**). Also the electrophoretic profile of VDAC was specific to muscle since VDAC was mainly detected at 45 kDa rather than at 30 kDa in other tissues. This could be due to muscle-specific post-translational modifications of VDAC such as glycosylation. The relative levels of VDAC among tissues were correlated with their citrate synthase activities ( $p < 0.001$ , Fig 2c).

### **Fig 2: Tissue-specific amount of VDAC and citrate synthase activity.**

(a) Immuno-detection on western-blot using anti-Cg VDAC with 10 µg of tissues protein lysates and (b) relative quantification of VDAC at 30 kDa. Tissues analyzed were: mantle, gills, digestive

gland, labial palp, striated adductor muscle, smooth adductor muscle, heart and visceral ganglia. VDAC was detected at 30 kDa (arrows), 45 kDa (line), 60 kDa, 90 kDa or 120 kDa (asterisks) depending on the tissue. The bands detected at 60, 90 and 120 kDa in SDS PAGE immunoblots corresponds to the size of VDAC oligomers. (c) Linear regression model using citrate synthase activity (mU/mg protein) as the explanatory variable and VDAC (30 kDa) relative protein level as the response variable in eight tissues ( $y = 0.086x + 1.178$ ,  $r^2 = 0.8789$ ,  $p < 0.005$ ).

## 2-DE profile of VDAC

The immunodetection on western-blot after 2-DE was performed in order to compare the 2-DE electrophoretic profiles of VDAC in four tissues: mantle, gills, smooth adductor muscle and heart (**Fig 3**). VDAC was detected in one specific train of spots at 30 kDa in the mantle (pI 6.2 to 9.6) and in the gills (pI 7.5 to pI 9.2). In the smooth muscle, the 2-DE profile revealed that VDAC was mainly detected at 45 kDa in a train of spot from pI 6.8 to 7.5 as observed in the monodimensional electrophoresis. In the heart, VDAC was detected in a spot at 30 kDa (pI= 8 and 8.3) with a shift at around 32 kDa (pI=7.6) that might be due to heart-specific post-translational modifications. In vertebrates, VDAC modifications refer mainly to acetylation, phosphorylation and S-nitrosylation. (Martel et al., 2014). In mammals and plants, VDAC has been found to be phosphorylated (Martel et al., 2014). According to what is known about the structure of VDAC, the horizontal trains on gels could correspond to different phosphorylation states of the protein, and the vertical shifts could be due to glycosylation. In the four tested tissues, the spots detected at 60 kDa or 90 kDa might indicate that some multimerization of VDAC remains visible even after protein denaturation followed by IEF.

### **Fig 3: Tissue-specific 2-DE profile of VDAC.**

Immuno-detection on western-blot using anti-Cg VDAC after 2-DE with 500 µg of (a) mantle, (b) gills, (c) smooth adductor muscle and (d) heart protein lysates. VDAC was detected in spots at 30 kDa (arrows), 45 kDa (line) and 60 kDa (asterisks) depending on the tissue. The spots detected at 60 kDa in 2-DE immunoblots correspond to the size of VDAC dimers.

### **VDAC amount in oysters during a field experiment**

The relative amounts of VDAC were followed in oysters deployed at 6 locations in the field for two months when OsHV-1 outbreaks (**Fig 4**). The sites 37 and 39 correspond to inshore sites located within the oyster farming area where OsHV-1 induced mortality occurred the earliest between 7 and 14 June (**Fig 4a**). The sites 33 and 38 were located offshore and OsHV-1 induced mortality occurred later, between 6 to 14 July (**Fig 4a**). Finally, the sites 32 and 36 were also located offshore, and there was no abnormal mortality (**Fig 4a**). The levels of VDAC at 30 kDa in oysters varied as a function of sites and time, the interaction of site and time being not significant ( $p=0.081$ ). Overall, the level of VDAC at 30 kDa increased with precocity of the mortality event (**Fig 4b, 4c, 4d**). On 30 April, the levels of VDAC at 30 kDa were similar irrespective of locations (**Fig 4d**). Then, the level of VDAC at 30 kDa increased between 30 April and 27 May and reached a plateau until 14 June, at the onset of the mortality period (**Fig 4c**).

### **Fig 4: VDAC amount in oysters in the field.**

(a) Map of 6 sampling sites, located along an inshore-offshore gradient in the Mor-Braz area, South Brittany (France). Sites 37 and 39 correspond to sites where mortalities occurred “early” (in red), sites 38 and 33 correspond to sites where mortalities occurred “late” (in yellow), and in sites 36

and 32 “no mortality” (blue) were recorded. (b) Effect of mortality level on the relative level of VDAC (30 kDa) in oysters in early mortality site (red), in late mortality site (yellow), and in site without mortality (blue). (c) Effect of time on the relative level of VDAC (30 kDa) in oysters from 30th April 2013 to 20th June 2013. (d) Representative blocs of western blot obtained the 30th April 2013 and 20th June 2013 in each site using anti-Cg VDAC antibody.

## VDAC in other marine species

We identified a contig encoding VDAC in the transcriptomes of six bivalve species: *Crassostrea rhizophorae*, *Pecten maximus*, *Ruditapes philippinarum*, *Ruditapes decussatus*, *Mytilus galloprovincialis* and *Mytilus edulis*. As compared to *C. gigas*, the entire VDAC amino-acid sequence of the six marine species listed above showed a percentage of identity ranging from 64 to 97 % (Table 1). In these species, we investigated the conservation of the epitope employed for the anti-Cg VDAC antibody synthesis, which consisted of 15 amino acids. With 10 to 14 conserved amino acids in the epitope, we could assume that anti-Cg VDAC antibody might be able to detect VDAC in these bivalve species (**Table 1**). The best record of epitope conservation was found for *C. rhizophorae* with 14 conserved amino acids.

**Table 1: VDAC in *C. gigas* and other marine species.**

	% identity Cg VDAC	Epitope	Epitope identity (number aa)
Cg VDAC	-	<u><b>QTGTKSGKIKTSYKM</b></u>	-
<i>Crassostrea gigas</i>			
Cr VDAC	97 %	<u><b>QTGTKSGKIKSSYKM</b></u>	14/15
<i>Crassostrea rhizophorae</i>			

<i>Mg</i> VDAC	67 %	<u>QTGKKQGT</u> <u>IKTGYKQ</u>	10/15
<i>Mytilus galloprovincialis</i>			
<i>Me</i> VDAC	67 %	<u>QTGKKQGT</u> <u>IKTGYKQ</u>	10/15
<i>Mytilus edulis</i>			
<i>Pm</i> VDAC	65 %	<u>QTGKKNG</u> <u>QIKTAYKM</u>	11/15
<i>Pecten maximus</i>			
<i>Rp</i> VDAC	64 %	<u>QTGKKSGK</u> <u>VKTGYKQ</u>	11/15
<i>Ruditapes philippinarum</i>			
<i>Rd</i> VDAC	64 %	<u>QTGKKSGK</u> <u>VKTGFKQ</u>	10/15
<i>Ruditapes decussatus</i>			

Comparison of VDAC amino acid sequence between *C. gigas* and others marine mollusks (in percentage of identity). The conserved amino acids from the epitope of anti-Cg VDAC are bold and underlined. Epitope identity is expressed as the number of amino acids conserved with *C. gigas* epitope.

## DISCUSSION

Owing to the purified anti-Cg VDAC antibody developed in our study, we showed that VDAC was constitutively expressed in all the tissues analyzed. This result agrees with the repartition of *Cg* VDAC transcripts in oysters (Li et al, 2016). The protein VDAC (30 kDa) was particularly accumulated in the heart, the labial palp, the ganglia, and the gills. The amount of VDAC was correlated with a high activity of citrate synthase in these tissues, a proxy of mitochondria number in tissues (Moran and Manahan 2004; Holmborn et al. 2009).

Analyses of *Cg* VDAC electrophoretic profile using both the western blot and the 2-DE western-blot might reveal the existence of dimers and trimers and their tissue-specific regulations. The detection of bands corresponding to VDAC oligomers can vary between immunoblots, due to either

a low level of band detection in a complex protein sample, or to the protocol of denaturation of protein extracts before analysis. VDAC oligomerization is well documented and VDAC is present as a dimer in rat liver (Lindén and Gellerfors 1983), as both dimers and trimers in yeast (Krause et al. 1986), and higher oligomers in plants (Hoogenboom et al., 2007). In striated and smooth muscles, VDAC was mainly detected at 45 kDa rather than at 30 kDa. In fact, specific post-translational modifications of VDAC might occur depending on the tissue, and this was confirmed by 2D immunoblots. Tissue specific regulation of VDAC likely reflects post-translational modifications, as previously reported in rats (Martel et al. 2014). These authors show that various VDAC post-translational modifications can occur, such as phosphorylation, acetylation, S-nitrosylation, and they influence the interactome and the activity of VDAC (Martel et al. 2014). In our case, further analysis should be carried out to verify the identities of the putative post-translational modifications and oligomerization found at 32, 45, 60, 90 and 120 kDa in western blotting. The anti-Cg VDAC antibody developed here might help to study the functioning of VDAC in oyster in further studies designed to investigate the post-translational modifications and oligomerization of Cg VDAC.

In *C. gigas*, the role of VDAC remains unclear. A recent study strongly supported the well conserved role of VDAC in the control of apoptosis during UV exposure and the direct interaction between VDAC and the pro-apoptotic protein Bak (Li et al. 2016). However, VDAC could be involved in the early stage of viral infection in invertebrates. This was already proposed in *C. gigas*/OsHV-1 (Corporeau et al. 2014; Young et al. 2017) and demonstrated in shrimps/WSSV. For instance, VDAC is accumulated during WSSV infection in *L. vannamei*, and it facilitates the infection process (Wang et al. 2007, 2010; Leu et al. 2013). Indeed, when the expression of VDAC is silenced, the infection process is delayed (Chen et al. 2011).

For the first time, our study shows that VDAC is up-accumulated in oysters exposed to OsHV-1 in the field. This result agrees with the accumulation of transcripts of *Cg* VDAC in the hemolymph of oysters infected with OsHV-1 (Renault et al. 2011) six hours after the viral injection (Li et al. 2016).

In marine invertebrates, knowledge on proteins playing a role in viral infection is still limited (Li et al. 2016) and studies have mainly explored the host or viral transcriptomes (Jouaux et al. 2013; Segarra et al. 2014). In shrimp, the Warburg effect is induced by viral mechanisms that alter the host metabolome through the PI3K-Akt-mTOR signaling pathways for production of energy and metabolic precursors for viral biogenesis (Su et al. 2014). In shrimp infected by WSSV, VDAC is up-regulated, and silencing of VDAC reduces WSSV-induced mortalities and virion copy number (Chen et al., 2011). As a key component of the Warburg effect, it now seems important to develop tools to study VDAC in *C. gigas* at the proteomic level, and to further evaluate its ability to bind partners, like hexokinase, depending on the infection status and the oyster environment.

For the first time, we proved the validity of the anti-*Cg* VDAC antibody as a tool to follow the amount of VDAC in oyster deployed in the field. We showed that the relative amount of VDAC at 30 kDa, as a monomer, can be informative even if it does not represent all the VDAC oligomers.

We demonstrated a higher accumulation of VDAC at 30 kDa in tissues exhibiting a higher citrate synthase activity, and in oysters from sites suffering from higher levels of mortality compared to control animals. These results likely indicate that (1) increasing amount of VDAC is related to the susceptibility of oysters to OsHV-1, and (2) disease-susceptibility of oyster and the amount of VDAC in oyster tissues are influenced by the local environment. The high amount of VDAC might reflect the ability of the oyster in the field to shift toward the Warburg effect and to replicate the virus, thus leading to death. Ongoing studies are investigating the role of environmental factors on

the interaction between VDAC, *C. gigas* metabolism and OsHV-1 replication. The antibody anti-Cg VDAC is a new tool to measure the impact of environmental factors on oyster metabolism.

## ACKNOWLEDGMENTS

We are grateful to Ifremer and the French ministry of agriculture for partly supporting this study. We acknowledge E. Harney for his help in editing English. The authors are grateful to Bruno Petton and the ifremer staff involved in oyster and algae production Argenton for their help and delivery of animals used in the study. We thank the shellfish network Resco II ([http://wwz.ifremer.fr/observatoire\\_conchylicole](http://wwz.ifremer.fr/observatoire_conchylicole)).

## CONFLICT OF INTEREST

The author declares no conflict of interest.

## REFERENCES

- Barbosa Solomieu V, Renault T, Travers M-A (2015) Mass mortality in bivalves and the intricate case of the Pacific oyster, *Crassostrea gigas*. *J Invertebr Pathol* 131:2–10 . doi: 10.1016/j.jip.2015.07.011
- Brahimi-Horn MC, Lacas-Gervais S, Adaixo R, et al (2015) Local Mitochondrial-Endolysosomal Microfusion Cleaves Voltage-Dependent Anion Channel 1 To Promote Survival in Hypoxia. *Mol Cell Biol* 35:1491–1505 . doi: 10.1128/MCB.01402-14
- Chen I-T, Aoki T, Huang Y-T, et al (2011) White Spot Syndrome Virus Induces Metabolic Changes Resembling the Warburg Effect in Shrimp Hemocytes in the Early Stage of Infection. *J Virol* 85:12919–12928 . doi: 10.1128/JVI.05385-11
- Corporeau C, Tamayo D, Pernet F, et al (2014) Proteomic signatures of the oyster metabolic response to herpesvirus OsHV-1  $\mu$ Var infection. *J Proteomics* 109:176–187 . doi: 10.1016/j.jprot.2014.06.030
- Davison AJ (2005) A novel class of herpesvirus with bivalve hosts. *J Gen Virol* 86:41–53 . doi: 10.1099/vir.0.80382-0




- 1
- 2
- 3
- 4 423 Dégremont L (2013) Size and genotype affect resistance to mortality caused by OsHV-1 in *Crassostrea*
- 5 424 *gigas*. *Aquaculture* 416–417:129–134 . doi: 10.1016/j.aquaculture.2013.09.011
- 6
- 7 425 Delgado T, Carroll PA, Punjabi AS, et al (2010) Induction of the Warburg effect by Kaposi's sarcoma
- 8 426 herpesvirus is required for the maintenance of latently infected endothelial cells. *Proc Natl Acad*
- 9 427 *Sci* 107:10696–10701 . doi: 10.1073/pnas.1004882107
- 10
- 11
- 12 428 Diamond DL, Syder AJ, Jacobs JM, et al (2010) Temporal Proteome and Lipidome Profiles Reveal Hepatitis
- 13 429 C Virus-Associated Reprogramming of Hepatocellular Metabolism and Bioenergetics. *PLoS*
- 14 430 *Pathog* 6:e1000719 . doi: 10.1371/journal.ppat.1000719
- 15
- 16
- 17 431 EFSA (2010) Scientific Opinion on the increased mortality events in Pacific oysters, *Crassostrea gigas*:
- 18 432 Oyster mortality
- 19
- 20 433 EFSA (2015) Oyster mortality: Oyster mortality. *EFSA J* 13:4122 . doi: 10.2903/j.efsa.2015.4122
- 21
- 22 434 Fabioux C, Corporeau C, Quillien V, et al (2009) In vivo RNA interference in oyster -*vasa* silencing inhibits
- 23 435 germ cell development. *FEBS J* 276:2566–2573 . doi: 10.1111/j.1742-4658.2009.06982.x
- 24
- 25
- 26 436 Guévelou E, Huvet A, Sussarellu R, et al (2013a) Regulation of a truncated isoform of AMP-activated
- 27 437 protein kinase  $\alpha$  (AMPK $\alpha$ ) in response to hypoxia in the muscle of Pacific oyster *Crassostrea*
- 28 438 *gigas*. *J Comp Physiol [B]* 183:597–611 . doi: 10.1007/s00360-013-0743-6
- 29
- 30 439 Guévelou E, Huvet A, Sussarellu R, et al (2013b) Regulation of a truncated isoform of AMP-activated
- 31 440 protein kinase  $\alpha$  (AMPK $\alpha$ ) in response to hypoxia in the muscle of Pacific oyster *Crassostrea*
- 32 441 *gigas*. *J Comp Physiol B* 183:597–611 . doi: 10.1007/s00360-013-0743-6
- 33
- 34
- 35 442 Guo Y, Meng X, Ma J, et al (2014) Human Papillomavirus 16 E6 Contributes HIF-1 $\alpha$  Induced Warburg
- 36 443 Effect by Attenuating the VHL-HIF-1 $\alpha$  Interaction. *Int J Mol Sci* 15:7974–7986 . doi:
- 37 444 10.3390/ijms15057974
- 38
- 39
- 40 445 Holmborn T, Dahlgren K, Høleton C, et al (2009) Biochemical proxies for growth and metabolism in
- 41 446 *Acartia bifilosa* (Copepoda, Calanoida). *Limnol Oceanogr Methods* 7:785–794
- 42
- 43 447 Hoogenboom BW, Suda K, Engel A, Fotiadis D (2007) The Supramolecular Assemblies of Voltage-
- 44 448 dependent Anion Channels in the Native Membrane. *J Mol Biol* 370:246–255 . doi:
- 45 449 10.1016/j.jmb.2007.04.073
- 46
- 47
- 48 450 Jouaux A, Lafont M, Blin J-L, et al (2013) Physiological change under OsHV-1 contamination in Pacific
- 49 451 oyster *Crassostrea gigas* through massive mortality events on fields. *BMC Genomics* 14:590
- 50
- 51 452 Krause J, Hay R, Kowollik CH, Brdiczka D (1986) Cross-linking analysis of yeast mitochondrial outer
- 52 453 membrane. *Biochim Biophys Acta BBA-Biomembr* 860:690–698
- 53
- 54
- 55 454 Lemasters JJ, Holmuhamedov E (2006) Voltage-dependent anion channel (VDAC) as mitochondrial
- 56 455 governor—Thinking outside the box. *Biochim Biophys Acta BBA - Mol Basis Dis* 1762:181–190 .
- 57 456 doi: 10.1016/j.bbadis.2005.10.006
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

- 1
- 2
- 3
- 4 457 Leu J-H, Lin S-J, Huang J-Y, et al (2013) A model for apoptotic interaction between white spot syndrome
- 5 458 virus and shrimp. *Fish Shellfish Immunol* 34:1011–1017 . doi: 10.1016/j.fsi.2012.05.030
- 6
- 7 459 Li Y, Zhang L, Qu T, et al (2016) Characterization of Oyster Voltage-Dependent Anion Channel 2 (VDAC2)
- 8 460 Suggests Its Involvement in Apoptosis and Host Defense. *PLOS ONE* 11:e0146049 . doi:
- 9 461 10.1371/journal.pone.0146049
- 10
- 11
- 12 462 Lindén M, Gellerfors P (1983) Hydrodynamic properties of porin isolated from outer membranes of rat
- 13 463 liver mitochondria. *Biochim Biophys Acta BBA-Biomembr* 736:125–129
- 14
- 15 464 Lü A-J, Dong C-W, Du C-S, Zhang Q-Y (2007) Characterization and expression analysis of *Paralichthys*
- 16 465 *olivaceus* voltage-dependent anion channel (VDAC) gene in response to virus infection. *Fish*
- 17 466 *Shellfish Immunol* 23:601–613 . doi: 10.1016/j.fsi.2007.01.007
- 18
- 19
- 20 467 Martel C, Wang Z, Brenner C (2014) VDAC phosphorylation, a lipid sensor influencing the cell fate.
- 21 468 *Mitochondrion* 19:69–77 . doi: 10.1016/j.mito.2014.07.009
- 22
- 23
- 24 469 Mazure NM (2017) VDAC in cancer. *Biochim Biophys Acta BBA - Bioenerg* 1858:665–673 . doi:
- 25 470 10.1016/j.bbabo.2017.03.002
- 26
- 27 471 Mesri EA, Feitelson MA, Munger K (2014) Human Viral Oncogenesis: A Cancer Hallmarks Analysis. *Cell*
- 28 472 *Host Microbe* 15:266–282 . doi: 10.1016/j.chom.2014.02.011
- 29
- 30 473 Miossec L, Le Deuff R-M, Goulletquer P (2009) Alien species alert: *Crassostrea gigas* (Pacific oyster). *ICES*
- 31 474 *Coop Res Rep* 299:
- 32
- 33
- 34 475 Moran A, Manahan D. (2004) Physiological recovery from prolonged “starvation” in larvae of the Pacific
- 35 476 oyster *Crassostrea gigas*. *J Exp Mar Biol Ecol* 306:17–36 . doi: 10.1016/j.jembe.2003.12.021
- 36
- 37 477 Munger J, Bajad SU, Collier HA, et al (2006) Dynamics of the Cellular Metabolome during Human
- 38 478 Cytomegalovirus Infection. *PLoS Pathog* 2:e132 . doi: 10.1371/journal.ppat.0020132
- 39
- 40
- 41 479 Naghdi S, Hajnóczky G (2016) VDAC2-specific cellular functions and the underlying structure. *Biochim*
- 42 480 *Biophys Acta BBA - Mol Cell Res.* doi: 10.1016/j.bbamcr.2016.04.020
- 43
- 44 481 Pedersen PL (2007) Warburg, me and Hexokinase 2: Multiple discoveries of key molecular events
- 45 482 underlying one of cancers’ most common phenotypes, the “Warburg Effect”, i.e., elevated
- 46 483 glycolysis in the presence of oxygen. *J Bioenerg Biomembr* 39:211–222 . doi: 10.1007/s10863-
- 47 484 007-9094-x
- 48
- 49
- 50 485 Pernet F, Barret J, Le Gall P, et al (2012) Mass mortalities of Pacific oysters *Crassostrea gigas* reflect
- 51 486 infectious diseases and vary with farming practices in the Mediterranean Thau lagoon, France.
- 52 487 *Aquac Environ Interact* 2:215–237 . doi: 10.3354/aei00041
- 53
- 54
- 55 488 Pernet F, Lupo C, Bacher C, Whittington RJ (2016) Infectious diseases in oyster aquaculture require a new
- 56 489 integrated approach. *Philos Trans R Soc B Biol Sci* 371:20150213 . doi: 10.1098/rstb.2015.0213
- 57
- 58
- 59
- 60
- 61
- 62
- 63
- 64
- 65

- Petton B, Pernet F, Robert R, Boudry P (2013) Temperature influence on pathogen transmission and subsequent mortalities in juvenile Pacific oysters *Crassostrea gigas*. *Aquac Environ Interact* 3:257–273 . doi: 10.3354/aei00070
- Petton, Boudry P, Alunno-Bruscia M, Pernet F (2015) Factors influencing disease-induced mortality of Pacific oysters *Crassostrea gigas*. *Aquac Environ Interact* 6:205–222 . doi: 10.3354/aei00125
- Poliseno L (2012) Pseudogenes: newly discovered players in human cancer. *Sci Signal* 5:5
- Puyraimond-Zemmour D, Vignot S (2013) Le métabolisme de la cellule tumorale : l'effet Warburg. *Oncologie* 15:435–440 . doi: 10.1007/s10269-013-2318-2
- Ramagli LS (1999) Quantifying protein in 2-D PAGE solubilization buffers. 2- *Proteome Anal Protoc* 99–103
- Renault T, Faury N, Barbosa-Solomieu V, Moreau K (2011) Suppression subtractive hybridisation (SSH) and real time PCR reveal differential gene expression in the Pacific cupped oyster, *Crassostrea gigas*, challenged with Ostreid herpesvirus 1. *Dev Comp Immunol* 35:725–735 . doi: 10.1016/j.dci.2011.02.004
- Rostovtseva TK, Komarov A, Bezrukov SM, Colombini M (2002) VDAC Channels Differentiate between Natural Metabolites and Synthetic Molecules. *J Membr Biol* 187:147–156 . doi: 10.1007/s00232-001-0159-1
- Schikorski D, Faury N, Pepin JF, et al (2011) Experimental ostreid herpesvirus 1 infection of the Pacific oyster *Crassostrea gigas*: Kinetics of virus DNA detection by q-PCR in seawater and in oyster samples. *Virus Res* 155:28–34 . doi: 10.1016/j.virusres.2010.07.031
- Segarra A, Baillon L, Tourbiez D, et al (2014) Ostreid herpesvirus type 1 replication and host response in adult Pacific oysters, *Crassostrea gigas*. *Vet Res* 45:103
- Segarra A, Pépin JF, Arzul I, et al (2010) Detection and description of a particular Ostreid herpesvirus 1 genotype associated with massive mortality outbreaks of Pacific oysters, *Crassostrea gigas*, in France in 2008. *Virus Res* 153:92–99 . doi: 10.1016/j.virusres.2010.07.011
- Shen X, Wang T, Xu D, Lu L (2014) Proteomic identification, characterization and expression analysis of *Ctenopharyngodon idella* VDAC1 upregulated by grass carp reovirus infection. *Fish Shellfish Immunol* 37:96–107 . doi: 10.1016/j.fsi.2014.01.009
- Su M-A, Huang Y-T, Chen I-T, et al (2014) An Invertebrate Warburg Effect: A Shrimp Virus Achieves Successful Replication by Altering the Host Metabolome via the PI3K-Akt-mTOR Pathway. *PLoS Pathog* 10:e1004196 . doi: 10.1371/journal.ppat.1004196
- Vander Heiden MG, Cantley LC, Thompson CB (2009) Understanding the Warburg Effect: The Metabolic Requirements of Cell Proliferation. *Science* 324:1029–1033 . doi: 10.1126/science.1160809
- Wang H-C, Kondo H, Hirono I, Aoki T (2010) The *Marsupenaeus japonicus* voltage-dependent anion channel (MjVDAC) protein is involved in white spot syndrome virus (WSSV) pathogenesis. *Fish Shellfish Immunol* 29:94–103 . doi: 10.1016/j.fsi.2010.02.020

- Wang H-C, Wang H-C, Leu J-H, et al (2007) Protein expression profiling of the shrimp cellular response to white spot syndrome virus infection. Dev Comp Immunol 31:672–686 . doi: 10.1016/j.dci.2006.11.001
- Warburg (1956) On the origin of cancer cells. Science 123:309–314
- Young T, Kesarcodi-Watson A, Alfaro AC, et al (2017) Differential expression of novel metabolic and immunological biomarkers in oysters challenged with a virulent strain of OsHV-1. Dev Comp Immunol 73:229–245 . doi: 10.1016/j.dci.2017.03.025

	% identity Cg VDAC	Epitope	Epitope identity (number aa)
<i>Cg</i> VDAC <i>Crassostrea gigas</i>	-	<u><b>QTGTKSGKIKTSYKM</b></u>	-
<i>Cr</i> VDAC <i>Crassostrea rhizophorae</i>	97 %	<u><b>QTGTKSGKIKSSYKM</b></u>	14/15
<i>Mg</i> VDAC <i>Mytilus galloprovincialis</i>	67 %	<u><b>QTGKKQGTIKTGYKQ</b></u>	10/15
<i>Me</i> VDAC <i>Mytilus edulis</i>	67 %	<u><b>QTGKKQGTIKTGYKQ</b></u>	10/15
<i>Pm</i> VDAC <i>Pecten maximus</i>	65 %	<u><b>QTGKKNGQIKTAYKM</b></u>	11/15
<i>Rp</i> VDAC <i>Ruditapes philippinarum</i>	64 %	<u><b>QTGKKSGKVKTGYKQ</b></u>	11/15
<i>Rd</i> VDAC <i>Ruditapes decussatus</i>	64 %	<u><b>QTGKKSGKVKTGFKQ</b></u>	10/15









Figure 3








Immunodetection  
Western Blot



Loading control  
Ponceau S



Well number	Sample
1	Molecular size markers
2	positive control
3	Site 37
4	Site 39
5	Site 38
6	Site 33
7	Site 36
8	Site 32
9	Molecular size markers
10	Molecular size markers