Università degli Studi di Padova

Padua Research Archive - Institutional Repository

The effect of experience and of dots' density and duration on the detection of coherent motion in dogs

Original Citation:

Availability:

This version is available at: 11577/3271645 since: 2019-10-05T16:06:14Z

Publisher:

Springer

Published version:

DOI: 10.1007/s10071-018-1200-4

Terms of use:

Open Access

This article is made available under terms and conditions applicable to Open Access Guidelines, as described at <http://www.unipd.it/download/file/fid/55401> (Italian only)

(Article begins on next page)

Animal Cognition

The effect of experience and of dots' density and duration on the detection of coherent motion in dogs --Manuscript Draft--

Manuscript Number:		
Full Title:	The effect of experience and of dots' density and duration on the detection of coherent motion in dogs	
Article Type:	Original Article	
Corresponding Author:	Paolo Mongillo, DVM, PhD Università degli Studi di Padova Dipartimento di Biomedicina Comparata e Alimentazione ITALY	
Corresponding Author Secondary Information:		
Corresponding Author's Institution:	Università degli Studi di Padova Dipartimento di Biomedicina Comparata e Alimentazione	
Corresponding Author's Secondary Institution:		
First Author:	Orsolya Kanizsár, PhD	
First Author Secondary Information:		
Order of Authors:	Orsolya Kanizsár, PhD	
	Paolo Mongillo, DVM, PhD	
	Luca Battaglini, PhD	
	Gianluca Campana, PhD	
	Miina Lööke	
	Lieta Marinelli, DVM, PhD	
Order of Authors Secondary Information:		
Funding Information:	Università degli Studi di Padova (DOR1673431)	Prof. Lieta Marinelli
	Università degli Studi di Padova (PhD scholarship)	Dr Orsolya Kanizsár
Abstract:	<p>Knowledge about the mechanisms underlying canine vision is far from being exhaustive, especially that concerning post-retinal elaboration. One aspect that has received little attention is motion perception, and in spite of the common belief that dogs are extremely apt at detecting moving stimuli, there is no scientific support to such assumption. In fact, we recently showed that dogs have higher thresholds than humans for coherent motion detection (Kanizsar et al. 2017). This term refers to the ability of the visual system to perceive several units moving in the same direction, as one, coherently moving global unit. Coherent motion perception is commonly investigated using random dot displays, containing variable proportions of coherently moving dots. Here, we investigated the relative contribution of local and global integration mechanisms to coherent motion perception, and changes in detection thresholds as a result of repeat-ed exposure to the experimental stimuli. Dogs who had been involved in the previous study were administered a conditioned discrimination task, in which we systematically manipulated dot density and duration and, eventually, we re-assessed our subjects' threshold after extensive exposure to the stimuli. Decreasing dot duration impacted on dogs' ac-curacy in detecting coherent motion only at very low duration values, revealing the efficacy of local integration mechanisms. Density impacted on dogs' accuracy in a linear fashion, indicating less efficient global integration. There was limited evidence of improvement in the re-assessment but, with an average threshold at re-assessment of 29%, dogs' ability to detect coherent motion</p>	

	remains much poorer than that of humans.
Suggested Reviewers:	Kun Guo, BSc PhD Reader, University of Lincoln kguo@lincoln.ac.uk Expertise in cognitive neuroscience of vision, including dog motion perception.
	Anna Kis, PhD Magyar Tudomanyos Akademia vargane.kis.anna@ttk.mta.hu The proposed reviewer has experience in dogs' cognition and visual perception
	George Mather, PhD Professor, University of Lincoln gmather@lincoln.ac.uk Professor of Vision Science
Opposed Reviewers:	

[Click here to view linked References](#)

The effect of experience and of dots' density and duration on the detection of coherent motion in dogs

Orsolya Kanizsár^a, Paolo Mongillo^{a*}, Luca Battaglini^b, Gianluca Campana^b, Miina Lööke^c, Lieta Marinelli^a

^aLaboratory of Applied Ethology, Department of Comparative Biomedicine and Food Science, University of Padua, Viale dell'Università 16, 35020, Legnaro (PD), Italy. ^bDepartment of General Psychology, University of Padua, Via Venezia 8, 35131, Padova, Italy ^cInstitute of Ecology and Earth Sciences, University of Tartu, Vanemuise 46, 51014, Tartu, Estonia

*Corresponding author's email: paolo.mongillo@unipd.it telephone number: +39 049 641862

ORCID:

Orsolya Kanizsar: 0000-0001-7351-7359

Paolo Mongillo: 0000-0001-8599-4005

Luca Battaglini: 0000-0002-5187-9225

Gianluca Campana: 0000-0003-4019-085X

Miina Lööke: 0000-0002-3981-1551

Lieta Marinelli: 0000-0003-0609-2172

Abstract

Knowledge about the mechanisms underlying canine vision is far from being exhaustive, especially that concerning post-retinal elaboration. One aspect that **has** received little attention is motion perception, and in spite of the common belief that dogs are extremely apt at detecting moving stimuli, there is no scientific support to such assumption. In fact, we recently showed that dogs have higher thresholds than humans for coherent motion detection (Kanizsar et al. 2017). This term refers to the ability of the visual system to perceive several units moving in the same direction, as one, coherently moving global unit. Coherent motion perception is commonly investigated using random dot displays, containing variable proportions of coherently moving dots. Here, we investigated the relative contribution of local and global integration mechanisms to coherent motion perception, and changes in detection thresholds as a result of repeated exposure to the experimental stimuli. Dogs who had been involved in the previous study were administered a conditioned discrimination task, in which we systematically manipulated dot density and duration and, eventually, we re-assessed our subjects' threshold after extensive exposure to the stimuli. Decreasing dot duration impacted on dogs' accuracy in detecting coherent motion only at very low duration values, revealing the efficacy of local integration mechanisms. Den-

32 sity impacted on dogs' accuracy in a linear fashion, indicating less efficient global integration. There was **limited** evi-
33 dence of improvement in the re-assessment but, with an average threshold at re-assessment of 29%, dogs' ability to de-
34 tect coherent motion remains much poorer than that of humans.

35

36 **keywords:** coherent motion, dot density, dot lifetime, perceptual learning, dog

37

38 **ACKNOWLEDGMENTS**

39 We are very grateful to the student Giulia Madumali Zotti for helping with experiments and to Dr. Carlo Poltronieri for
40 his technical assistance. The study was funded by the University of Padova (to LM, 2016 - prot. DOR1673431). Dr.
41 Orsolya Kanizár was supported by a PhD grant funded by the University of Padova.

42

INTRODUCTION

Dogs make extensive use of visual information, in tasks as simple as recognizing their owner (Mongillo et al. 2017b), to more cognitively complex activities, such as understanding human communicative signs (reviewed by Kaminsky and Nitzschner 2013), attentional states (Gácsi et al. 2004; Virányi et al. 2004) and emotions (Albuquerque et al. 2016; Nagasawa et al. 2011). Notwithstanding the demonstrated importance of such sensory modality, research on the mechanisms underlying canine vision is far from being exhaustive (Byosiere et al. 2017a). Most of the studies on the topic focused on the functional properties of eye structures, and their impact on basic characteristics of sight, like acuity, light sensitivity, and color discrimination. A much smaller number of studies looked at higher level (i.e. post-retinal) visual elaboration processes, like the perception and the discrimination of object shapes and sizes (e.g. Byosiere et al. 2017b; Milgram et al. 1994) or the spatial integration of local elements into a global percept (Mongillo et al. 2017a; Pitteri et al. 2014a, b).

One aspect that has received very little attention is dogs' perception of motion. However, the ability to detect and encode information about moving stimuli is likely to be involved in many aspects of the life of a dog. For instance, it could play a fundamental role in predatory behavior, with its desirable (e.g. for hunting dogs) or undesirable (for dogs expressing predation as a problematic behavior) consequences. In spite of common belief holding that dogs are extremely apt at detecting moving stimuli, there is no scientific data supporting such assumption. In fact, to date, the only study that dealt with dogs' ability to perceive motion, recently conducted by our research group, suggests that the opposite is the case (Kanizsar et al. 2017).

In such study, we specifically dealt with dogs' ability to detect coherent motion, that is the ability to perceive several local units moving with the same direction and speed as a single, coherently moving unit (Braddick 1993; Williams and Brannan 1994). According to a widely accepted theory, the perception of coherent motion represents the second of a two-stage motion processing mechanism, whereby higher order neurons integrate the local component of motion analyzed by neurons in the initial stage, providing a global percept (Rust et al. 2006). Such mechanism represents a crucial step to an organisms' ability to extrapolate complex information about relevant stimuli from motion cues (Berental and Pinto 1994; Blanke et al. 2007). The physiological boundaries of coherent motion perception are commonly investigated via discrimination tasks using random dot displays (Newsome and Pare, 1988), where a given number of dots moves in the same direction (signal dots), among a number of dots moving in random directions (noise). The smaller the proportion of signal dots that the subject needs in order to detect the coherent motion, the lower is the individual's threshold. Using this paradigm, we found that dogs have an average threshold of coherent motion detection of 42%, considerably higher than the 5% thresholds found in human subjects assessed in the same experimental condition (Kanizsar et al. 2017), and also higher than thresholds reported for other species, including monkeys (Newsome and

75 Pare 1988), seals (Weiffen et al. 2014) and cats (Huxlin and Pasternak 2004; Rudolph and Pasternak 1996). Thus, dogs’
 76 coherent motion perception abilities seem to be rather poor, not just compared to primates, but even to phylogenetically
 77 closer species.

78 One factor that could have contributed to our dog’s high thresholds, is their relatively little experience with the experi-
 79 mental stimuli. It has been widely demonstrated that the performance in many visual tasks improves after experience
 80 with the stimuli. Motion detection tasks make no exception, and improvement in detection thresholds through practice
 81 have been documented in humans, mice, monkeys and seals (Britten et al. 1992; Douglas et al. 2006; Watanabe et al.
 82 2001; Weiffen et al. 2014). The dogs who took part in our previous study were privately owned, and, although they had
 83 received extensive training, their overall exposure to the experimental stimuli was limited if compared to studies em-
 84 ploying animals housed in experimental facilities. It therefore remains a viable hypothesis that thresholds observed in
 85 our dogs did not represent their lower asymptote, and that such thresholds could be improved by giving dogs additional
 86 experience with the stimuli.

87 The thresholds of motion detection are also greatly influenced by the characteristics of the experimental stimuli. For
 88 instance, both a shorter duration (i.e. the amount of a dot remains visible before disappearing and being replaced by an-
 89 other dot elsewhere in the display)¹ and a lower density of dots in the display result in higher thresholds of coherent mo-
 90 tion detection in both human and non-human subjects (Snowden and Kavanagh 2006; Talcott et al. 2000; Weiffen et al.
 91 2014). It is unlikely that changing the duration or dot density used in our previous experiment (Kanizsar et al. 2017)
 92 would improve our dogs’ thresholds, since stimuli were designed in order to maximize the dogs’ performance and both
 93 parameters were set around the upper range end, in comparison with those used in other studies. Nonetheless, manipu-
 94 lating these parameters would still be useful in the attempt to clarify the relative contribution of different mechanisms to
 95 coherent motion perception (e.g. Bischof et al. 1999). Indeed, there are at least two basic processes through which co-
 96 herent motion detection can be attained, that is the integration of single motion units moving in the same direction
 97 across multiple time frames (local integration), which would be more greatly affected by shorter dots’ duration, and the
 98 integration of multiple motion units moving in the same direction, across as few as two subsequent frames (global inte-
 99 gration), which would be more greatly affected by lower dot densities. Although the role of local and global integration,
 100 and their neurophysiological substrates, have been largely addressed in the primate literature, there are substantial dif-
 101 ferences between primates and carnivoran (Aguirre et al. 2007; Djavadian and Harutiunian-Kozak 1983), to suggest that
 102 the mechanisms leading to te detection of coherent motion may differ between these taxa.

103 The experiments described in this paper represent an extension of our previous study (Kanizsar et al. 2017) and were
 104 aimed at investigating the relative contribution of local and global integration mechanisms and the role of experience in

¹In the vision literature this parameter is also called *lifetime*

determining dogs' coherent motion detection thresholds. To these aims, the same dogs who took part in our previous experiment underwent a two-alternative forced choice discrimination task using random dot displays, in which we systematically varied the dot density or duration. After the dogs completed these tasks we re-assessed the subjects' thresholds using the same testing procedure and parameters as described in Kanizsar et al. (2017), to investigate the effects of extensive stimulus exposure on dogs' coherent motion detection thresholds.

110

111 **MATERIALS AND METHODS**

112

113 **Subjects**

114

115 Our sample included five mesocephalic pet dogs, three females (1 Mongrel, 1 Mudi, 1 Siberian Husky) and two males
116 (1 Cocker Spaniel, 1 Labrador-Poodle Mix), between 3 and 11 years of age. All of these dogs had participated in the
117 previous study that investigated dogs' thresholds of coherent motion detection (Kanizsar et al. 2017). Dogs belonged to
118 private owners who participated in the experiments on a voluntary basis. The subjects underwent a veterinary examina-
119 tion before the enrollment in the experiments to exclude health conditions that would prevent them from participation.
120 Dogs were selected upon the requirement that they were highly motivated for food and willing to cooperate, while feel-
121 ing comfortable with being in the laboratory.

122

123 **Experimental setting**

124

125 All the experiments took place in the Laboratory of Applied Ethology of the Department of Biomedicine and Food Sci-
126 ence (University of Padova, Italy). A testing area of 2.5 x 3 m was established in a laboratory room. Stimuli were pre-
127 sented on two identical monitors (VG248QE, ASUSTeK Computer Inc., Taipei, Taiwan). Their refresh rate was set at
128 100 Hz, to prevent possible biases on motion detection, due to dogs' higher flicker fusion frequency (Miller and Mur-
129 phy 1995). Monitors were connected to a PC (Optiplex 960, Dell Inc., Round Rock, Texas, USA). They were placed
130 side by side at 25 cm from each other. During presentations, the dog viewed the monitors from a distance of 70 cm.
131 Monitors were attached to height-adjustable stands, so that their height could be set at eye level for each subject.
132 Presentations were controlled by the experimenter with a Bluetooth® keyboard (K400 Plus, Logitech International
133 S.A., Lausanne, Switzerland).

134

135 **Stimuli**

136

137 All experimental stimuli were created with MATLAB (MATLAB version 7.10.0. Natick, Massachusetts: The Math-
138 Works Inc., 2010), using features of the Psychtoolbox (Brainard 1997; Pelli 1997). The stimuli were displayed on a
139 black area of 31.1 cm x 31.1 cm (24.0 x 24.0 deg, from the viewing distance of 70 cm), where white dots with a diame-
140 ter of 0.16 cm moved **towards the left side of the monitor** at a speed of 19.4 cm/s (15.0 deg/s). For all the training trials
141 (including those used in the test phases), the positive stimulus was set at a coherence of 80%, i.e. 80% of the dots
142 moved in the same direction, whereas the remaining 20% moved in random directions. In all **test and training** trials, the
143 negative stimulus had a coherence level of 0%, that is all of the dots moved in random directions. In training trials, there
144 was a total of 5000 dots moving in the display, for a density of 5.9 dots/cm² (8.7 dots/deg²) and dots had a duration of
145 1000 ms. Dot density, dot duration and the percentage of coherence of the display were manipulated in the test trials of
146 the respective experiments, for which a detailed description of the stimuli is given below.

147

148 **Training phase and general test procedure**

149

150 Each dog underwent three tests, aimed at assessing respectively the effect of dot duration, dot density and experience on
151 their coherent motion detection threshold. The latter was defined as the percentage of coherently moving dots in the
152 positive stimulus at which dogs' accuracy in discriminating such stimulus reached an estimated value of 75%.

153 Prior to each test, dogs underwent a training phase, identical in all respects to the one they had already undergone in the
154 previous study (Kanizsar et al. 2017). Briefly, such phase consisted in a two-alternative forced choice discrimination
155 task, where dogs had to discriminate a random dot display with a coherence of 80% (**positive stimulus**), from one with a
156 coherence of 0%. Training sessions were composed of 20 trials, in each of which dogs were presented with the positive
157 and negative stimuli, and, after an inspection time of 4 s, they had to choose one of the two stimuli by approaching and
158 touching one of the two monitors with their snout. **The side of presentation of negative and positive stimuli were ran-**

159 **domly determined by the software, with the constraint that they were counterbalanced within a session. After touching**
160 **one of the two monitors, the stimuli disappeared, and dogs were reinforced with food (pieces of sausage of about 1cm³)**
161 **if they chose the positive stimulus; immediately after finishing eating, dogs were called to the starting position and the**
162 **next trial began. If dogs chose the negative stimulus, they received no reward and after an interval of approximately 7 s**
163 **they were called back to the starting position for the next trial.** Training was completed when dogs reached a criterion of

164 90% or better accuracy (i.e. at least 18 correct choices out of 20 trials), in 6 consecutive sessions. In this phase, as well
165 in the tests, dogs underwent a maximum of 4 sessions per day.

166 In the tests dogs underwent several sessions composed of a certain number of 'training' trials, in which the two stimuli
167 were identical to those presented in the training phase, and 'test' trials, where the characteristics of the stimuli were ma-

168 manipulated, according to the specifics of each test (described below in detail). The first 4 trials of every session were al-
169 ways of the training type, serving as a ‘warm-up’, whereas in rest of the session a set number of training and test trials
170 were intermingled in random order. The side of presentation of the positive stimulus was also randomized and counter-
171 balanced within each session. In test trials, dogs were never reinforced regardless of their choice. The inclusion of train-
172 ing trials in test sessions was meant to maintain dogs’ motivation throughout, and to allow an assessment of subjects’
173 discriminative performance during tests; if a dog failed to maintain a criterion of 85% correct responses in the training
174 trials of the test phase (e.g. more than 2 errors in the 14 training trials), it was sent back to the training phase.

175

176 **Dot density test**

177

178 This test was meant to investigate whether the manipulation of dots density in the presented displays affected individual
179 thresholds of coherent motion detection. The test comprised 10 test sessions, each including 14 training trials and 6 test
180 trials. In the latter, the density of dots in both the positive and negative stimuli was set to one of 3 different levels (8.7
181 dots/deg², 1.74 dots/deg² and 0.17 dots/deg²), so that every density level was shown twice per session, and 20 times in
182 the entire test. The choice of the density levels was based on the comparative literature and on pilot testing, with the aim
183 of maximizing the sensitivity of the assay. In test trials, the level of coherence of signal dots in the positive stimulus
184 was set for each dog to its individual threshold, as resulting from the previous study, where dot density was set to 8.7
185 dots/deg², (Kanisszar et al. 2017); the dots’ duration was the same as in the training stimuli (1000 ms).

186

187 **Dot duration test**

188

189 This test was meant to investigate whether the manipulation of dots’ duration affected individual thresholds of coherent
190 motion detection. The test comprised 10 test sessions, each including 14 training trials and 6 test trials. In the latter, the
191 duration of dots in both the positive and negative stimuli was set to one of 3 different levels (1000 ms, 200 ms, and 50
192 ms), so that every duration level was shown twice per session, and 20 times in the entire test. The choice of the duration
193 levels was based on the comparative literature and on pilot testing, with the aim of maximizing the sensitivity of the
194 assay. In test trials, the level of coherence of signal dots in the positive stimulus was set for each dog to its individual
195 threshold, as resulting from the previous study, where dot duration was set to 1 s (Kanisszar et al. 2017); the dots’ den-
196 sity was the same as in the training stimuli (8.7 dots/deg²).

197

198 **Re-assessment of coherent motion detection thresholds**

199

This test was meant to assess the effects of experience on the thresholds of coherent motion detection of dogs, by repeating the assessment procedure that the same dogs had previously undergone (Kanizsar et al. 2017), after having being exposed to the positive training stimulus in (at least) additional 640 trials since the first threshold assessment. The interval between the two assessments was on average 2 months.

The test consisted of 10 sessions, each composed of 14 training trials and 10 test trials. In the latter, the coherence of the positive stimulus was varied across 5 different levels (i.e. 60%, 50%, 40%, 30%, and 20% coherently moving signal dots); each of these levels was shown twice within the same session, and 20 times in the entire test.

Data collection and statistical Analysis

Linear regression analyses were run to find the best model for describing the relationships between the dots density and duration and the proportion of correct responses, respectively in the dot duration and in the dot density tests. Choice of the best fitting model was based on visual inspection of the data and on the comparison of the R^2 . After that, a two-tail one sample t-test was run to determine whether the mean slope of the functions was significantly different from zero, indicating better (> 0) or worse (< 0) performance when the independent variable increases.

For the re-assessment of coherent motion detection thresholds, data of each dog were fitted with a logistic function by using the routines provided by Palamedes (Prins and Kingdom 2009), which consider a proportion of correct response for the level of coherence given by as:

$$P(C; \alpha, \beta, \gamma, \lambda) = \gamma + \frac{1 - \gamma - \lambda}{1 + e^{-\beta(C - \alpha)}}$$

As the task was a 2-alternative forced-choice, the lower asymptote for guess (*Gamma*) was set to 0.5, while the upper asymptote (*Lambda*) was fixed by setting the lapse rate (probability of an incorrect response, independent of stimulus intensity) to 0.02. The parameters *Alpha* and *Beta* were left free. *Alpha* refers to the threshold, i.e. the value along the abscissa corresponding to the coherence level at which the function attains its steepest point. *Beta* is a discrimination parameter often referred to as the ‘slope’. Then, a one-tail paired t-test was run for threshold, slope and upper asymptote, regardless the small sample size, in order to investigate whether the parameters differed between the thresholds assessment performed in Kanizsar et al. (2107) and the re-assessment of the current study.

Finally, to determine whether improvements in thresholds after experience depended on the initial performance level, an analysis of correlations was run between the thresholds reported for each dog by Kanizsar et al. (2017), and the difference between such threshold and that observed for the same dog in the re-assessment test of the current study.

RESULTS

All dogs rapidly reached the learning criterion in the training phases preceding each test phase (median N of sessions needed to reach the learning criterion = 6; min = 6, max = 8). In test sessions, all dogs maintained the criterion of 85% accuracy in the training trials, thus no dog was retroceded to training once started any of the test phases.

Effect of dot density

Figure 1 shows the proportion of correct choices as a function of dot density. An optimum way to fit these data is a simple linear model (see Table 1 for individual *slopes*, *intercepts* and R^2). A two-tailed, one-sample t-test showed that the *slope* of the linear regression was significantly higher than zero ($t_4 = 3.58$, $P = 0.023$, Cohen's $d = 2.58$), indicating that dog's performance linearly improves with increasing dot density.

Effect of dot duration

Figure 2 shows the proportion of individual correct choices as a function of dot's duration. The dog's performance increases rapidly as the dot duration increases. In four out of five dogs, R^2 is higher than 0.7 (see Table 3). Furthermore, a two-tails one-sample t-test showed that the slope of the logarithmic regression was significantly higher than zero ($t_4 = 4.68$, $P = 0.008$, Cohen's $d = 3.3$). This indicates that dog's performance increases rapidly as the dot duration increases, but then it stabilizes for duration above 200 ms.

Effect of repeated exposition to the stimuli

Figure 3 shows the individual psychometric functions and the proportion of correct choices as a function of coherence, comparing the results reported in Kanizsar et al. (2017) with those of the re-assessment performed on the same dogs in the current study. Table 3 shows the *Alpha* (threshold) and *Beta* (slope) parameters for each dog. The mean threshold of coherent motion detection in dogs in the re-assessment was 29.3%, while the mean slope was 0.06; comparatively, the mean threshold assessed by Kanizsar et al. (2017) in the same dogs was 42.2%, whereas the mean slope was 0.08. The difference in thresholds between the two assessments approached significance ($t_4 = 2.08$, $P = 0.104$, Cohen's $d = 1.53$) as well as the difference in slope ($t_4 = 1.85$, $P = 0.12$, Cohen's $d = 0.97$). There was a significant correlation between the threshold observed in the first assessment, and the degree of improvement in the re-assessment ($r = 0.94$, $P = 0.014$), so that larger improvement was seen in dogs who had initially shown the higher thresholds.

DISCUSSION

263

264 In this study, we assessed dogs' threshold of coherent motion detection as a function of dot density, dot duration and of
265 the dogs' experience with the experimental stimuli. The results show that dogs' ability to detect coherent motion is neg-
266 atively affected by dots' duration only at very short duration values, while changes in density impact on dogs' thresh-
267 olds of motion detection in a linear fashion. Finally, re-assessment of the dogs' thresholds of motion detection after ex-
268 tensive practice with the experimental stimuli produced variable results within our sample, with indications of improve-
269 ment.

270 The dogs' ability to detect coherent motion as a function of dots' duration decreased according to a logarithmic curve:
271 dogs' performance showed only a trivial, if any, decrement when duration was lowered from 1000 to 200 ms, with four
272 out of five dogs still performing with an accuracy at or above 70%; however, when duration was further reduced to 50
273 ms, dogs' performance clearly dropped, with accuracy falling below 60% for all dogs. A short duration affects the pos-
274 sibility to rely on local integration, that is to detect the direction of movement of few, closely-spaced dots, which is a
275 prerequisite to detect coherent motion. The mechanism has been found to contribute to coherent motion detection in
276 humans as well as in other animal species, although to a different extent (e.g. Bischof et al. 1999; Talcott et al. 2000;
277 Weiffen et al. 2014). How do dogs compare to humans or other animal species? Talcott et al. (2000) found that humans'
278 motion detection improved when duration was increased from 200 ms up to about 900 ms, and remained unchanged
279 above such values. This would suggest a higher efficiency of local motion integration mechanism in dogs' than in hu-
280 mans, as our dogs' performance had already reached its asymptote with a dots' duration of 200 ms. However, when
281 stimuli are presented on a monitor, where the sensation of movement is provided by discrete image changes, the possi-
282 bility to determine a dot's displacement also depends on the number of frames displayed. In this respect, in Talcott et al.
283 (2000), the 200 ms and 900 ms conditions contained 4 and 18 frames, values that are respectively much more similar to
284 our 50 ms (5 frames) and 200 ms (20 frames) conditions than to the 200 and 1000 ms. Thus, if frame number rather
285 than time is considered, the improvement in performance of our dogs as a function of dots' duration is very similar to
286 that of humans. Regardless of which of the two parameters is considered, the results indicate that dogs' local integration
287 mechanisms are at least as efficient as they are in humans, and are therefore unlikely to play a major role in determining
288 dogs' higher threshold of coherent motion detection. In addition, the efficiency in local integration mechanisms is in
289 line with dogs' alleged skillfulness in detecting locally moving stimuli, such as a prey moving in the distance (Miller
290 and Murphy 1995).

291 Our results on local motion integration are less directly comparable to those obtained in other animal species, due to
292 relevant methodological differences. For instance, Weiffen et al. (2014) assessed a seal's motion detection sensitivity to
293 varying duration at 125, 250, and 500 ms (frame rates were not provided); across these values, the seals' decrement in

threshold appeared to be linear, with no indication of a stabilization. This may suggest that seal's local motion integration still had space for improvement (i.e. it is less efficient than that of our dogs) but without an assessment of the animals' sensitivity in a wider range, and considered other differences in the characteristics of the presented stimuli between the two studies, this remains only a speculative hypothesis.

Dogs' ability to detect coherent motion decreased in a linear fashion when dot density was reduced from 8.7 to 0.17 dots/deg², with four of the five dogs performing with < 60% accuracy at the lowest density level. The dependency of coherent motion detection from dot density is thought to reflect mechanisms of global integration: the higher total number of dots moving in a consistent direction determines the recruitment of a higher number of low-level motion detectors sensitive to that specific direction, while inhibiting those sensitive to other directions, thereby increasing the possibility to identify a set of coherently moving stimuli as a single entity. Our results indicate progressive improvement in dogs' reliance on such global integration mechanisms as density increases, at least within the range of densities that we investigated. By way of comparison, adult humans tested across a range of densities roughly spanning our two highest levels showed very little improvement in their detection threshold (Talcott et al. 2000) and the seal in the study of Weiffen et al. (2014) only showed improvement when density was increased up to 0.77 dots/deg², but no further improvement with higher density values. Although proper comparisons are hindered by methodological differences, the results suggest that dogs' global integration mechanisms for motion detection are less efficient than in humans and also other species; in turn, such lower efficiency in global motion detection may be responsible for the higher thresholds of coherent motion detection found in dogs.

As far as nervous structures are concerned, local integration is thought to rely on low-level direction-sensitive detectors, found in the primary visual cortex. The sensitivity to a specific direction of local motion units emerges at this level in both primates (Wurtz and Kandel 2000) and cats (Humphrey and Saul 2002). By contrast, global integration occurs at a higher level, in specialized areas of the extra-striate cortex. As opposed to primary visual cortex, there is substantial difference in the neuro-functional organization of these areas between primates, where global motion detection occurs in the middle-temporal area (Newsome and Pare 1988), and other mammals, such as the cat, where the same processes occur in the lateral suprasylvian area (Gizzi et al. 1990; Rudolph and Pasternak 1996); although dogs' visual cortex has not been studied as thoroughly as that of the cat, and no data directly comparable to our results exists in the cat literature, there seems to be a good degree of correspondence in the neuro-functional organization of these areas between the two species (Aguirre et al. 2007). Thus, the difference in the functions and organization of these areas between dogs and primates could be responsible for the different abilities of global integration between dogs and humans. The larger variability observed in the effect of dot density than duration is also in agreement with the higher-level origin (i.e. further from sensory origin) of processes that affect dogs' performance.

325 After the extensive exposure to random dots motion displays, dogs' thresholds of coherent motion perception were re-
326 assessed using the same procedure they underwent earlier (Kanizsar et al. 2017) and dogs' average thresholds passed
327 from 42% of the first study, to about 29%. The analysis did not result in a significant difference; however, for three out
328 of the five dogs the improvement seemed to be very clear ($>10\%$); the remaining two dogs showed little change from
329 the first assessment.

330 This suggests that some of the dogs had not reached their lower asymptote in threshold in the course of the first experi-
331 ment, but needed further exposition to the stimuli in order to optimize their performance. Similar effects of experience
332 in improving coherent motion detection have been reported in many other species, including rodents (Douglas et al.
333 2006), seals (Weiffen et al. 2014), monkeys (Britten et al. 1992) and humans (Sagi 2011). Most likely, such improve-
334 ment is the result of perceptual learning, a well-known effect often observed in visual tasks, which is believed to reflect
335 improvement in low-level cortical processes, consequent to extensive exposure to specific stimuli. In the context of
336 global motion detection, the neurobiological substrate of such improvement has been localized in the middle-temporal
337 area of primates' brain (Thomson and Liu 2006). The improvement in threshold in our dogs showed some inter-individ-
338 ual variability, and their degree of improvement was correlated to the threshold observed in the first test, so that dogs
339 with the highest detection thresholds, also showed the largest improvement in the present study. Both variability in the
340 degree of perceptual learning and its dependency on the initial performance have been observed before in visual tasks in
341 humans and are consistent with perceptual learning processes (Doshier and Lu 2005; Fahle and Henke-Fahle 1996).

342 The difference in average threshold between the first assessment and that done in the present did not reach statistical
343 significance; however, considered the small number of subjects and the extent of the improvement observed in some of
344 them, it seems sensible to conclude that experience had an effect in improving detection thresholds. In spite of such im-
345 provement, final thresholds for all dogs were still markedly higher than those observed in humans in our first experi-
346 ment.

347

348 **Conclusions**

349

350 This study confirms our previous findings that dogs coherent motion perception is less efficient than that of humans, and
351 suggests that the source of such differences is to be found in dogs' less efficient global integration mechanisms. By
352 contrast, dogs' local motion integration seems to be highly efficient, possibly more than the corresponding mechanism in
353 humans. These findings bear on some relevant practical aspects. For instance, it would be important to determine how
354 the mechanisms of motion detection contribute to prey drive, or whether the efficiency of dogs' local motion integration
355 mechanisms allows them to be particularly good at spotting movements in the distance, as it was earlier suggested.

356 The study also shows that the performance of dogs in motion detection tasks can be improved through perceptual learning,
357 although the presence and the extent of such improvement are variable, with larger improvements seen in subjects with
358 the worst initial performance. In any case, such improvement is not sufficient to bring dogs' thresholds near the level of
359 human ones. Whether dogs' poor ability to detect coherent motion extends to other aspects of motion detection (for
360 instance, the ability to detect minimum amounts of global motion) remains an aspect to be addressed by future studies.

361

362 COMPLIANCE WITH ETHICAL STANDARDS

363 None of the authors of this paper has any financial or personal relationship with other people or organizations which
364 might inappropriately influence or bias its content.

365 All applicable international, national, and/or institutional guidelines for the care and use of animals were followed.

366

367

368 REFERENCES

369

- 370 1. Albuquerque N, Guo K, Wilkinson A, Savalli C, Otta E, Mills D (2016) Dogs recognize dog and human emo-
371 tions. *Biol Lett* 12: 20150883. [https://doi.org/ 10.1098/rsbl.2015.0883](https://doi.org/10.1098/rsbl.2015.0883)
- 372 2. Aguirre GK, Komáromy AM, Cideciyan AV, Brainard DH, Aleman TS, Roman AJ, Avants BB, Gee JC, Kor-
373 czykowski M, Hauswirth WW, Acland GM (2007) Canine and human visual cortex intact and responsive de-
374 spite early retinal blindness from RPE65 mutation. *Plos Med* 4:e230. [https://doi.org/10.1371/jour-](https://doi.org/10.1371/journal.pmed.0040230)
375 [nal.pmed.0040230](https://doi.org/10.1371/journal.pmed.0040230)
- 376 3. Bertenthal BI, Pinto J. (1994). Global processing of biological motions. *Psychol Sci* 5:221–225. doi:
377 [10.1111/j.1467-9280.1994.tb00504.x](https://doi.org/10.1111/j.1467-9280.1994.tb00504.x)
- 378 4. Bischof WF, Reid SL, Wylie DR, Spetch ML (1999) Perception of coherent motion in random dot displays by
379 pigeons and humans. *Percept Psychophys* 61:1089-1101. <https://doi.org/10.3758/BF03207616>
- 380 5. Blanke O, Brooks A, Mercier M, Spinelli L, Adriani M, Lavanchy L, Safran AB, Landis T (2007) Distinct
381 mechanisms of form-from-motion perception in human extrastriate cortex. *Neuropsychol* 45:644-653.
382 [https://doi.org/10.1016/j.neuropsychologia.2006.07.019.](https://doi.org/10.1016/j.neuropsychologia.2006.07.019)
- 383 6. Braddick O (1993) Segmentation vs integration in visual motion processing. *Trends Neorosci* 16:263-268.
384 [https://doi.org/10.1016/0166-2236\(93\)90179-P](https://doi.org/10.1016/0166-2236(93)90179-P)
- 385 7. Brainard DH (1997) The psychophysics toolbox. *Spat Vision* 10:433-436.
- 386 8. Britten KH, Shadlen MN, Newsome WT, Movshon J (1992) The analysis of visual motion: a comparison of
387 neuronal and psychophysical performance. *J Neurosci* 12:4745-4765.

9. Byosiere SE, Chouinard PA, Howell TJ, Bennett PC (2017a) What do dogs (*Canis familiaris*) see? A review of vision in dogs and implications for cognition research. *Psychon Bull Rev* 1-16. <https://doi.org/10.3758/s13423-017-1404-7>
10. Byosiere SE, Feng LC, Chouinard PA, Howell TJ, Bennett PC (2017b) Relational concept learning in domestic dogs: Performance on a two-choice size discrimination task generalises to novel stimuli. *Behav Process* 145:93-101. <https://doi.org/10.1016/j.beproc.2017.10.009>
11. Djavadian RL, Harutiunian-Kozak BA (1987) Retinotopic organization of the lateral suprasylvian area of the cat. *Acta Neurobiol Exp* 43:251-262.
12. Doshier BA, Lu ZL (2005) Perceptual learning in clear displays optimizes perceptual expertise: Learning the limiting process. *Proc Natl Acad Sci USA* 102:5286-5290. <https://doi.org/10.1073/pnas.0500492102>
13. Douglas RM, Neve A, Quittenbaum JP, Alam NM, Prusky GT (2006) Perception of visual motion coherence by rats and mice. *Vision Res* 46:2842-2847. <https://doi.org/10.1016/j.visres.2006.02.025>
14. Fahle M, Henkle-Fahle S (1996) Interobserver variance in perceptual performance and learning. *Invest Ophthalmol Vis Sci* 37:869-877.
15. Gácsi M, Miklósi Á, Varga O, Topál J, Csányi V (2004) Are readers of our face readers of our minds? Dogs (*Canis familiaris*) show situation-dependent recognition of human's attention. *Anim Cogn* 7:144-153. <https://doi.org/10.1007/s10071-003-0205-8>
16. Gizzi MS, Katz E, Movshon JA (1990) Spatial and temporal analysis by neurons in the representation of the central visual field in the cat's lateral suprasylvian visual cortex. *Visual Neurosci* 5:463-468. <https://doi.org/10.1017/S0952523800000584>
17. Humphrey AL, Saul AB (2002) The emergence of direction selectivity in cat primary visual cortex. In: Payne B, Peters A (ed) *The Cat Primary Visual Cortex*. Academic Press, London, pp 343-386
18. Huxlin KR, Pasternak T (2004) Training-induced recovery of visual motion perception after extrastriate cortical damage in the adult cat. *Cereb Cortex* 14:81-90. <https://doi.org/10.1093/cercor/bhg106>
19. Kaminski J, Nitzschner M (2013) Do dogs get the point? A review of dog-human communication ability. *Learn Motiv* 44:294-302. <https://doi.org/10.1016/j.lmot.2013.05.001>
20. Kanizsar O, Mongillo P, Battaglini L, Campagna G, Marinelli L (2017) Dogs are not better than humans at detecting coherent motion. *Sci Rep-UK* 7:11259. <https://doi.org/10.1038/s41598-017-11864-z>
21. Milgram NW, Head E, Weiner E, Thomas E (1994) Cognitive functions and aging in the dog: acquisition of nonspatial visual tasks. *Behav Neurosci* 108:57. <http://dx.doi.org/10.1037/0735-7044.108.1.57>
22. Miller PE, Murphy CJ (1995) Vision in dogs. *J Am Vet Med Assoc* 15:1623-1634.

23. Mongillo P, Pitteri E, Sambugaro P, Carnier P, Marinelli L (2017a) Global bias reliability in dogs (*Canis familiaris*). *Anim Cogn* 20:257-265. <https://doi.org/10.1007/s10071-016-1044-8>
24. Mongillo P, Scandurra A, Kramer RSS, Marinelli L (2017b) Recognition of human faces by dogs (*Canis familiaris*) requires visibility of head contour. *Anim Cogn* 20:881-890. <https://doi.org/10.1007/s10071-017-1108-4>
25. Nagasawa M, Murai K, Mogi K, Kikusui T (2011) Dogs can discriminate human smiling faces from blank expressions. *Anim Cogn* 14:525-533. <https://doi.org/10.1007/s10071-011-0386-5>
26. Newsome WT, Pare BT (1988) A selective impairment of motion perception following lesions of the middle temporal visual area (MT). *J Neurosci* 8:2201-2211.
27. Pelli DG (1997) The VideoToolbox software for visual psychophysics: Transforming numbers into movies. *Spat Vis* 10:437-442. <https://doi.org/10.1163/156856897X00366>
28. Pitteri E, Mongillo P, Carnier P, Marinelli L (2014a) Hierarchical stimulus processing by dogs (*Canis familiaris*). *Anim Cogn* 17:869-877. <https://doi.org/10.1007/s10071-013-0720-1>
29. Pitteri E, Mongillo P, Carnier P, Marinelli L, Huber L (2014b) Part-based and configural processing of owner's face in dogs. *PlosONE* 9:e108176. <https://doi.org/10.1371/journal.pone.0108176>
30. Prins N, Kingdom FAA (2009) Palamedes: Matlab routines for analyzing psychophysical data. <http://www.palamedestoolbox.org>.
31. Rudolph KK, Pasternak T (1996) Lesions in cat lateral suprasylvian cortex affect the perception of complex motion. *Cereb Cortex* 6:814-822. <https://doi.org/10.1093/cercor/6.6.814>
32. Rust NC, Mante V, Simoncelli EP, Movshon JA (2006). How MT cells analyze the motion of visual patterns. *Nat Neurosci* 9:1421.
33. Sagi D (2011) Perceptual learning in vision research. *Vision Res* 51:1552-1566. <https://doi.org/10.1016/j.visres.2010.10.019>
34. Snowden RJ, Kavanagh E (2006) Motion perception in the ageing visual system: minimum motion, motion coherence, and speed discrimination thresholds. *Perception* 35:9-24. <https://doi.org/10.1068/p5399>
35. Talcott JB, Hansen PC, Assoku EL, Stein JF (2000) Visual motion sensitivity in dyslexia: evidence for temporal and energy integration deficits. *Neuropsychologia* 38:935-943. [https://doi.org/10.1016/S0028-3932\(00\)00020-8](https://doi.org/10.1016/S0028-3932(00)00020-8)
36. Thompson B, Liu Z (2006) Learning motion discrimination with suppressed and un-suppressed MT. *Vision Res* 46:2110-2121. <https://doi.org/10.1016/j.visres.2006.01.005>
37. Virányi Z, Topál J, Gácsi M, Miklósi Á, Csányi V (2004) Dogs respond appropriately to cues of humans' attentional focus. *Behav Process* 66:161-172. <https://doi.org/10.1016/j.beproc.2004.01.012>

38. Watanabe T, Nanez JE, Sasaki Y (2001) Perceptual learning without perception. *Nature* 413:844-847.
<https://doi.org/10.1038/35101601>
39. Weiffen M, Mauck B, Dehnhardt G, Hanke FD (2014) Sensitivity of a harbor seal (*Phoca vitulina*) to coherent visual motion in random dot displays. *Springerplus*. 3:688. <https://doi.org/10.1186/2193-1801-3-688>
40. Williams D, Brannan, J (1994) Spatial integration of local motion signals. In: Smith A, Snowden R (ed) *Visual Detection of Motion*. Academic Press, London, pp 291-303
41. Wurtz RH, Kandel ER (2000) Perception of motion, depth and form. In: Kandel ER, Schwartz JH, Jessel TM (ed) *Principles of Neural Science*, 4th edn. McGraw-Hill, New York, pp 548-571.

462 **Figure 1.** Dot density test: proportion of correct choices performed by each dog as a function of dot density. Symbols
463 indicate the proportion of correct choices; lines represent the linear regression of the data of each subject.

464
465 **Figure 2.** Dot duration test: proportion of correct choices performed by each dog as a function of dot's duration. Sym-
466 bols are indicating the proportion of correct choices while lines represent the linear regression of the data of each sub-
467 ject.

468
469 **Figure 3.** Psychometric curves and proportion of correct choices as a function of coherence, assessed in Kanizsar et al.
470 (2017) (dotted line, empty circles) and in present study (black line, filled circles) on the same five dogs.

1 **Table 1.** Dot density test: values of *slope*, *intercept* and R^2 of the linear regression of the data of each of the
2 five dogs.

	<i>Slope</i>	<i>Intercept</i>	R^2
Dog 1	0.046	0.46	0.92
Dog 2	0.023	0.57	0.92
Dog 3	0.023	0.56	0.83
Dog 4	0.006	0.69	0.97
Dog 5	0.018	0.54	0.81

3

Table 2. Dot lifetime test: values of *slope*, *intercept* and R^2 of the linear regression of the data of each of the five dogs.

	<i>Slope</i>	<i>intercept</i>	R^2
Dog 1	0.14	0.91	0.75
Dog 2	0.08	0.76	0.97
Dog 3	0.09	0.88	0.84
Dog 4	0.06	0.81	0.71
Dog 5	0.03	0.66	0.28

1 **Table 3.** Values of the *Alpha* (threshold) and *Beta* (slope) parameters for each of the five dogs in the first
2 assessment (Kanizsar et al., 2017) and in the current study's re-assessment

	<i>Alpha</i>	<i>Beta</i>	<i>Alpha</i>	<i>Beta</i>
	<i>1st assessment</i>	<i>1st assessment</i>	<i>re-assessment</i>	<i>re-assessment</i>
Dog 1	37.6	0.095	33.72	0.058
Dog 2	37.4	0.048	39.42	0.048
Dog 3	40.5	0.068	24.19	0.069
Dog 4	41.8	0.086	29.42	0.075
Dog 5	53.9	0.104	19.63	0.05

Click here to access/download
Supplementary Material
Response to reviewers.docx

Click here to access/download
Supplementary Material
Cover Letter_200518.docx