Università degli Studi di Padova

Padua Research Archive - Institutional Repository

Does aquatic thermal therapy improve quality of life after total hip replacement? A retrospective preliminary pilot study

Original Citation:

Availability:

This version is available at: 11577/3327774 since: 2020-02-27T19:00:41Z

Publisher:

Published version:

DOI: 10.1007/s00484-019-01846-3

Terms of use:

Open Access

This article is made available under terms and conditions applicable to Open Access Guidelines, as described at <http://www.unipd.it/download/file/fid/55401> (Italian only)

(Article begins on next page)

International Journal of Biometeorology

Does aquatic thermal therapy improve quality of life in subjects who underwent total hip replacement? A retrospective pilot study.

--Manuscript Draft--

Manuscript Number:	
Full Title:	Does aquatic thermal therapy improve quality of life in subjects who underwent total hip replacement? A retrospective pilot study.
Article Type:	Special Issue: Balneology (invited only)
Keywords:	Hydrokinesitherapy; Balneotherapy; Rehabilitation; Quality of life; Total hip replacement; Osteoarthritis.
Corresponding Author:	Stefano Masiero Universita degli Studi di Padova ITALY
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	Universita degli Studi di Padova
Corresponding Author's Secondary Institution:	
First Author:	Stefano Masiero
First Author Secondary Information:	
Order of Authors:	Stefano Masiero
	Giulia Pranovi
	Martina Di Pumpo
	Simone Bernardini
	Alberto Dattilo
	Alessandra Del Felice, MD
	Patrizio Sale
Order of Authors Secondary Information:	
Funding Information:	
Abstract:	<p>Aquatic thermal therapy is prescribed for its antalgic, muscle relaxant, and trophic properties, as well as anti-edemigenous and anti-inflammatory effects. Currently, the management of persons with total hip replacement (THR) consists of multiple pharmacologic and non-pharmacologic therapies, which include therapeutic exercise (joint-specific exercises for strength and range of motion, stretching, proprioceptive exercises and functional training), physical therapies and education. Our hypothesis is that therapeutic exercise in thermal water would improve functional outcome. The aim of the study was to assess thermal water rehabilitation efficacy on pain, mood and quality of life. We enrolled 11 subjects (mean age of 70.55±14 years) with Total Hip Replacement (THR) after early hospital discharge at a residential rehabilitation facility. All participants underwent a standardized combined land-based and hydrokinesitherapy rehabilitation program for 2 consecutive weeks. Clinical data were collected at admission and discharge from the rehabilitation facility; scoring from scales on pain, anxiety and quality of life were collected at admission, discharge, 3 and 6 months from surgery. Quality of life, anxiety, mood and pain improved after treatment and persisted at 6 months' follow-up. These preliminary data demonstrate that thermal hydrokinesitherapy can positively influence functional recovery in orthopedic post-surgical patients. Further randomized controlled trials are mandatory to consolidate this finding.</p>
Suggested Reviewers:	Filippo Vittadini

	filippo.vittadini@gmail.com
	Massimiliano Mangone massimiliano.mangone@uniroma1.it

[Click here to view linked References](#)

1 **Abstract**

2 Aquatic thermal therapy is prescribed for its antalgic, muscle relaxant, and trophic properties, as well as anti-edemigenous and anti-inflammatory effects.
3 Currently, the management of persons with total hip replacement (THR) consists of multiple pharmacologic and non-pharmacologic therapies, which
4 include therapeutic exercise (joint-specific exercises for strength and range of motion, stretching, proprioceptive exercises and functional training),
5 physical therapies and education. Our hypothesis is that therapeutic exercise in thermal water would improve functional outcome. The aim of the study
6 was to assess thermal water rehabilitation efficacy on pain, mood and quality of life. We enrolled 11 subjects (mean age of 70.55 ± 14 years) with Total
7 Hip Replacement (THR) after early hospital discharge at a residential rehabilitation facility. All participants underwent a standardized combined land-
8 based and hydrokinesitherapy rehabilitation program for 2 consecutive weeks. Clinical data were collected at admission and discharge from the
9 rehabilitation facility; scoring from scales on pain, anxiety and quality of life were collected at admission, discharge, 3 and 6 months from surgery.
10 Quality of life, anxiety, mood and pain improved after treatment and persisted at 6 months' follow-up. These preliminary data demonstrate that thermal
11 hydrokinesitherapy can positively influence functional recovery in orthopedic post-surgical patients. Further randomized controlled trials are mandatory
12 to consolidate this finding.

13

14

15

16

17

18

19

20

21

22

Introduction

Thermal water physiotherapy reduces postural instability and risk of falling in elderly subjects, with a positive impact on quality of life (Vivas et al. 2011), and maximizes rehabilitative treatment efficacy in different musculoskeletal disorders (Devereux et al. 2005; Masiero 2008; Arnold et al. 2008; Paoloni et al. 2017; Mazzoli et al. 2017; Masiero et al. 2018). Thermal rehabilitative therapies provide also a favorable social environment (Miller and Kaptchuk 2008). Previous studies demonstrated anti-edemigenous, anti-inflammatory, muscle relaxant, muscle trophic and antalgic physical properties of thermal waters (Masiero 2008; Masiero et al. 2011, 2018; Paoloni et al. 2017; Mazzoli et al. 2017). Current evidence indicates that aquatic thermal therapy may have the potential to decrease cortisol levels in the short term and to improve cortisol awakening response in healthy subjects (Toda et al. 2006).

Our hypothesis is that intensive thermal water rehabilitation contributes to recovery after total hip replacement arthroplasty (THR). Main outcome was improvement of quality of life; secondary outcomes were pain control, anxiety and mood reduction.

Materials and methods

Eleven subjects with total hip replacement (THR) were subsequently recruited and enrolled among participants to a thermal water standardized rehabilitation program at a residential rehabilitation facility [Thermal Rehabilitation Center in Abano Terme (Italy)].

Inclusion criteria were total hip replacement for primary hip osteoarthritis 15 to 30 days before enrollment and age between 45 and 90 years. Exclusion criteria were surgical open wounds, cardiovascular (included phlebitis or deep vein thrombosis), pulmonary, metabolic (e.g. complicated diabetes), cutaneous, neoplastic and mental disorders. Different surgical operators were not an exclusion criterion.

Rehabilitation program was performed in hyper-thermal “salso-bromo-iodic” water pools.

All participants underwent a thorough medical examination before and at the end of the rehabilitation program.

Treatment consisted of six sessions/week for 2 consecutive weeks of two and half hours of land-based (1 hour and a half) and hydrokinesitherapy (1 hour) with active and passive joint mobilization, respiratory and functional re-education exercises, gait and balance training, resistance exercise, and power training associated with physical therapy (electrotherapy and low-level-laser therapy). An educational program was provided to participants and

families (Musumeci et al. 2018). All subjects were evaluated the day of admission (T0) and at the end of treatment (T1); after 3 months (T2) and 6 months after the end of the treatment (T3) scales scores were collected by telephone interview.

At T0 and T1 collected parameters were: Range of Motion (ROM) of hip flexion/abduction and Harris Hip Score (HHS), a clinician-based outcome measure for the assessment of the results of the hip replacement (Nilsson and Bremander 2011). At T0, T1, T2 and T3 self-administered scale scores were collected: Numeric Pain Rating Scale (NRS_p) measurement for pain assessment (where zero describes no pain and 10 severe pain) (Huskisson 1974); Numeric Mood Rating Scale (NRS_m) measurement for mood assessment (where zero describes severe depression and 10 euphoria); Numeric Anxiety Rating Scale (NRS_a) measurement for anxiety assessment (where zero describes no anxiety and 10 severe anxiety); EQ5D Euroqol Quality of Life Scale (Eq-5D) (Balestroni and Bertolotti 2015).

A preliminary descriptive analysis was performed to check the normal distribution of participants' data using Shapiro-Wilk test. The Mann-Whitney test (T0 vs T1) for clinical tests and Friedman test (T0 vs T1 vs T2 vs T3) for self-administered scales was used to determine any significant differences between time points. Significance was set at $p < 0.05$.

Results

Average age of participants was 70.55 ± 14.03 years (5 males). The ANOVA analysis showed a significant improvement, more evident at T1 compared with T0, which persisted at T2 and T3 for NRS_a (T1>T0: $p=0.0305$; T2>T0: $p=0.0453$; T3>T0: $p=0.0447$), NRS_m (T1>T0: $p=0.0435$; T2>T0: $p=0.037$; T3>T0: $p=0.0501$) and EQ5D (T1>T0: $p=0.0027$; T2>T0: $p=0.013$; T3>T0: $p=0.024$). The HHS score before (T0) and after treatment (T1) showed a statistical improvement ($p=0.0374$). There was a non-statistically significant improvement in hip flexion and abduction between T0 and T1. There was no adverse event during spa treatment. Results are reported in table 1 and figure 1.

Discussion

This study provides evidence of long-term improvement of quality of life of subjects with THR after aquatic thermal therapy. Currently, the management of persons with THR consists of multiple pharmacologic and non-pharmacologic therapies, which include therapeutic exercise (joint-specific exercises

for improving strength and range of motion, stretching, proprioceptive exercises and functional training) physical therapies and education (Wijnen et al. 2018).

Our results support the role of aquatic thermal therapy in the rehabilitation process of this population. Thermal hydrokinesitherapy may improve mood, anxiety and stress management, reduce fatigue and pain and thus facilitate rehabilitation (Oláh et al. 2011; Musumeci et al. 2018) without no reported side effects. Stress has a direct association with fatigue, which relates to pain (Rapolienė et al. 2016). In fact, mineral bathing appears to be more efficient to relieve pain than non-mineral baths or muds, likely due to a specific effect of the mineral components of thermal water (Morer et al. 2017). Mineral water has also a crucial role in the rehabilitation intervention due to the potential to stimulate the healing process through minerals and gaseous compounds (Masiero 2008). Clinical experience suggests that density, hydrostatic pressure, buoyancy, viscosity and thermodynamics of aquatic thermal therapy contribute to the rehabilitation process. The long-term improvement in quality of life scores (Rahmann et al. 2009; Villalta and Peiris 2013) further supports the integration of thermal hydrokinesitherapy in THR rehabilitation.

The main limitation of the study was the lack of a control group and the selection bias. Indeed, we are confident of the impact of thermal water therapy in light of the clear-cut improvements and their long-term persistence, which calls for randomized controlled trials to demonstrate our preliminary results.

Conclusions

The lack of side effects and the easiness of providing intervention support the develop of large randomized clinical trial, which are mandatory to confirm that thermal aquatic rehabilitation might become a useful strategy in the rehabilitation program of total hip replacement. In conclusion, thermal settings appear promising rehabilitation environments for people with orthopedic musculoskeletal disability.

89 **Acknowledgements**

90 We acknowledge the Medical Hotel Hermitage Bel Air for providing the thermal setting.

91

92

93

94

95

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111 **Competing interests**

112 Authors have no conflicts of interest or any financial interest. All authors confirm that the material has not been and will not be submitted for publication
113 elsewhere.

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

References

- Arnold CM, Busch AJ, Schachter CL, et al (2008) A Randomized Clinical Trial of Aquatic versus Land Exercise to Improve Balance, Function, and Quality of Life in Older Women with Osteoporosis. *Physiother Canada* 60:296–306. doi: 10.3138/physio.60.4.296
- Balestroni G, Bertolotti G (2015) EuroQol-5D (EQ-5D): an instrument for measuring quality of life. *Monaldi Arch Chest Dis* 78:155–9. doi: 10.4081/monaldi.2012.121
- Devereux K, Robertson D, Briffa NK (2005) Effects of a water-based program on women 65 years and over: a randomised controlled trial. *Aust J Physiother* 51:102–8
- Huskisson EC (1974) Measurement of pain. *Lancet (London, England)* 2:1127–31
- Masiero S (2008) Thermal rehabilitation and osteoarticular diseases of the elderly. *Aging Clin Exp Res* 20:189–194. doi: 10.1007/BF03324772
- Masiero S, Bonaldo L, Pigatto M, et al (2011) Rehabilitation Treatment in Patients with Ankylosing Spondylitis Stabilized with Tumor Necrosis Factor Inhibitor Therapy. A Randomized Controlled Trial. *J Rheumatol* 38:1335–1342. doi: 10.3899/jrheum.100987
- Masiero S, Vittadini F, Ferroni C, et al (2018). The role of thermal balneotherapy in the treatment of obese patient with knee osteoarthritis. *Int J Biometeorol* 62:243–252. doi: 10.1007/s00484-017-1445-7
- Mazzoli D, Giannotti E, Longhi M, et al (2017) Age explains limited hip extension recovery at one year from total hip arthroplasty. *Clin Biomech* 48:35–41. doi: 10.1016/j.clinbiomech.2017.07.003
- Miller FG, Kaptchuk TJ (2008). The power of context: reconceptualizing the placebo effect. *J R Soc Med* 101:222–225. doi: 10.1258/jrsm.2008.070466
- Morer C, Roques CF, Françon A, Forestier R, Maraver F. The role of mineral elements and other chemical compounds used in balneology: data from double blind randomized clinical trials. *Int J Biometeorol* (2017) 61:2159–2173. DOI 10.1007/s00484-017-1421-2
- Musumeci A, Pranovi G, Masiero S (2018) Patient education and rehabilitation after hip arthroplasty in an Italian spa center: a pilot study on its feasibility. *Int J Biometeorol*. doi: 10.1007/s00484-018-1548-9

155 Nilsdotter A, Bremander A (2011) Measures of hip function and symptoms: Harris Hip Score (HHS), Hip Disability and Osteoarthritis Outcome Score
 156 (HOOS), Oxford Hip Score (OHS), Lequesne Index of Severity for Osteoarthritis of the Hip (LISOH), and American Academy of Orthopedic
 157 Surgeons (A. Arthritis Care Res (Hoboken) 63:S200–S207. doi: 10.1002/acr.20549

158 Oláh M, Koncz A, Fehér J, Kálmánczhey J, Oláh C, Balogh S, Nagy G, Bender T (2011). The effect of balneotherapy on C-reactive protein, serum
 159 cholesterol, triglyceride, total antioxidant status and HSP-60 levels,” International Journal of Biometeorology, vol. 54, no. 3, pp. 249–254.

160 Paoloni M, Bernetti A, Brignoli O, et al (2017) Appropriateness and efficacy of Spa therapy for musculoskeletal disorders. A Delphi method consensus
 161 initiative among experts in Italy. Ann Ist Super Sanita 53:70–76. doi: http://old.iss.it/binary/publ/cont/ANN_17_01_13.pdf

162 Rahmann AE, Brauer SG, Nitz JC (2009) A Specific Inpatient Aquatic Physiotherapy Program Improves Strength After Total Hip or Knee
 163 Replacement Surgery: A Randomized Controlled Trial. Arch Phys Med Rehabil 90:745–755. doi: 10.1016/j.apmr.2008.12.011

164 Rapolienė L, Razbadauskas A, Sąlyga J, Martinkėnas A. Stress and Fatigue Management Using Balneotherapy in a Short-Time Randomized
 165 Controlled Trial. Evid Based Complement Alternat Med. 2016;2016:9631684. doi: 10.1155/2016/9631684. Epub 2016 Mar 9.

166 Toda M, Morimoto K, Nagasawa S, Kitamura K (2006) Change in salivary physiological stress markers by spa bathing. Biomed Res 27:11–14. doi:
 167 10.2220/biomedres.27.11

168 Villalta EM, Peiris CL (2013) Early Aquatic Physical Therapy Improves Function and Does Not Increase Risk of Wound-Related Adverse Events for
 169 Adults After Orthopedic Surgery: A Systematic Review and Meta-Analysis. Arch Phys Med Rehabil 94:138–148. doi:
 170 10.1016/j.apmr.2012.07.020

171 Vivas J, Arias P, Cudeiro J (2011) Aquatic Therapy Versus Conventional Land-Based Therapy for Parkinson’s Disease: An Open-Label Pilot Study.
 172 Arch Phys Med Rehabil 92:1202–1210. doi: 10.1016/j.apmr.2011.03.017

173 Wijnen A, Bouma SE, Seeber GH, van der Woude LHV, Bulstra SK, Lazovic D, Stevens M, van den Akker-Scheek I (2018). The therapeutic validity
 174 and effectiveness of physiotherapeutic exercise following total hip arthroplasty for osteoarthritis: A systematic review. PLoS One. 2018 Mar
 175 16;13(3):e0194517. doi: 10.1371/journal.pone.0194517.

176 **Figure Caption**

177

178 **Figure 1.** Mean value of self-administered scales score pre and post treatment T0 vs T1 and 3 (T2) and 6 (T3) months follow-up. Numeric Pain
179 Rating Scale (NRSp), Numeric Mood Rating Scale (NRSm), Numeric Anxiety Rating Scale (NRSa), EQ5D Euroqool Quality of Life Scale (Eq-5D).

1

	NRSp			NRSa			NRSm			EQ5D		
	Mean	SD	p	Mean	SD	p	Mean	SD	p	Mean	SD	p
T0	2.81	2.52	ns	2.36	2.97	> T0	4.54	3.11	> T0	0.53	0.20	> T0
T1	2.01	2.09	ns	0.818	1.53	0.030	6.72	2.72	0.043	0.67	0.11	0.002
T2	2.45	2.54	ns	2.45	2.80	0.045	7.36	1.62	0.037	0.74	0.27	0.01
T3	2.36	2.33	ns	2.36	2.69	0.045	7.45	1.63	0.050	0.77	0.27	0.02

2

3

4 **Table 1.** Friedman test for sSelf-administered scales scores for quality of life and pain pre- and post-treatment T0 vs T1 and at 3 (T2) and 6 (T3) months follow-up: Numeric Pain
5 Rating Scale (NRSp), Numeric Mood Rating Scale (NRSm), Numeric Anxiety Rating Scale (NRSa), and EQ5D Euroqool Quality of Life Scale (Eq-5D).

6

Figure 1

[Click here to access/download](#)

Electronic Supplementary Material
Title Page.docx

[Click here to access/download](#)

Electronic Supplementary Material
Manuscript pdf.pdf