

Prognostic significance of the sum of the diameters of single foci in multifocal papillary thyroid cancer: the concept of new-old tumor burden

Jacopo Manso , Simona Censi, Amir Roberti, Maurizio Iacobone, Susi Barollo, Loris Bertazza, Francesca Galuppini, Federica Vianello, Nora Albinger, Carla Scaroni, Gianmaria Pennelli and Caterina Mian

Abstract

Aim: The prognostic value of multifocality (Mu) in papillary thyroid cancer (PTC) remains controversial. The present study aimed to investigate this issue and test the possible prognostic significance of the sum of the diameters of single foci (SDSF), the total number of foci (TNF), and primary tumor size (PTS) in multifocal PTC.

Methods: We retrospectively analyzed a single-center consecutive series of 370 PTCs. For multifocal cases we analyzed bilaterality occurrence, SDSF, TNF, and PTS.

Results: Mu was observed in 41.1% PTCs, and bilaterality in 30%. Mu was associated with an advanced T-category. In bilateral multifocal PTC, the PTS was larger, and microPTC was less frequent, while T-categories were higher. Mu and bilaterality *per se* had no impact on prognosis. At univariate analysis, PTS, SDSF, vascular invasion, lymph node metastases, distant metastases, T-categories, Initial Risk Stratification System score, second treatment and TERT promoter mutation correlated with persistence/recurrence or death in the multifocal PTC group. On multivariate Cox proportional hazards regression analyses, SDSF again independently predicted persistence/recurrence or death in multifocal PTCs. We found that a cut-off for SDSF less than 40 mm was able to identify multifocal PTC patients with a very low risk of persistence/recurrence (negative predictive value 96.9%). Disease-free survival was significantly shorter in patients with multifocal PTCs and SDSF ≥ 40 mm.

Conclusions: Mu and bilaterality *per se* were not prognostically significant. SDSF emerged as a new independent prognostic factor for persistence/recurrence of multifocal PTC. SDSF might better represent the tumor burden in multifocal PTC, with SDSF < 40 mm identifying multifocal PTC patients with a good prognosis.

Keywords: bilaterality, multifocality, papillary thyroid cancer, sum of diameters of single foci, tumor burden

Received: 10 June 2020; revised manuscript accepted: 11 September 2020.

Introduction

Well-differentiated thyroid cancer includes papillary and follicular histotypes, and accounts for the vast majority ($>90\%$) of all thyroid cancers (TCs).¹ In the United States (US), the incidence of TC has tripled in recent times, from 4.9 per 100,000 in 1975 to 14.3 per 100,000 in 2009.²

Papillary thyroid cancer (PTC), the main culprit responsible for the constantly rising incidence of TC, has an excellent prognosis, with survival rates exceeding 97% at 5 years, and $>90\%$ at 10 years.^{3,4} Given this low mortality rate for PTC, clinical and pathological factors affecting disease-free survival – in other words, disease persistence

Ther Adv Endocrinol Metab

2020, Vol. 11: 1–15

DOI: 10.1177/
2042018820964326

© The Author(s), 2020.
Article reuse guidelines:
sagepub.com/journals-
permissions

Correspondence to:

Jacopo Manso
Department of Medicine
(DIMED), Endocrine Unit,
Università di Padova,
Via Ospedale Civile 105,
Padova, 35128, Italy
jacopo.manso@gmail.com

Simona Censi
Amir Roberti
Susi Barollo
Loris Bertazza
Carla Scaroni
Caterina Mian
Department of Medicine
(DIMED), Endocrinology
Unit, Padua University,
Padua, Italy

Maurizio Iacobone
Department of Surgical,
Oncological and
Gastroenterological
Sciences (DiSCOG),
Endocrine Surgery Unit,
Padua University Hospital,
Padua, Italy

Department of Cardiac,
Thoracic and Vascular
Sciences (DCTV),
Biostatistics Epidemiology
and Public Health Unit,
Padua University Hospital,
Padua, Italy

Francesca Galuppini
Gianmaria Pennelli
Department of
Medicine (DIMED),
Surgical Pathology and
Cytopathology Unit, Padua
University, Padua, Italy

Federica Vianello
Nora Albinger
Department of
Radiotherapy, Istituto
Oncologico del Veneto,
IOV-IRCCS, Padova, Italy

and recurrence – rather than disease-related death are the key elements to consider at diagnosis and during patient follow up in every day clinical practice.

So, while the AJCC/UICC staging system (8th edition) is the most powerful tool for predicting the mortality risk,^{5–8} a combination of tumor size and gross extrathyroidal extension, more or less aggressive histologies, the burden of metastatic lymph nodes, incomplete tumor resections, and particular molecular profiles have all emerged as significant factors influencing the risk of recurrent structural disease.

On the other hand, vascular invasion, capsular invasion, and multifocality (Mu) are less clear prognostic determinants, and any role they may have is still being debated.⁹

PTC can occur as single or multifocal tumors (involving two or more anatomically separate foci). The prevalence of Mu in PTC ranges from 32% to 39%.^{10,11} Mu most often presents as multiple microPTC (maximum tumor size <10 mm), and only occasionally with lesions visible on ultrasound.¹² The latest guidelines issued by the American and European Thyroid Associations (ATA and ETA) place patients with multifocal PTC in the category at low risk of persistence/recurrence.^{13,14} Recent studies have found, however, that Mu is associated with lymph node metastases at diagnosis, and with persistent/recurrent disease during follow-up.^{10,11,15,16} Mu in PTC is also often empirically interpreted by clinicians as a high-risk factor, and this prompts a more aggressive treatment.¹⁷

In the literature, Mu in PTC is associated with older age,^{15,18} male sex,^{18,19} and extrathyroidal extension.^{11,15,17} It is only in some studies that Mu seemed to be associated with vascular invasion, lymph node metastases, more advanced disease and higher Initial Risk Stratification System (IRSS) scores.^{16,18} Nevertheless, the real prognostic significance of Mu in contributing to PTC persistence/recurrence, or even mortality, remains controversial. As an example, two recent meta-analyses reported opposite findings. The one conducted by Guo *et al.* included 7048 patients from 13 studies, and found Mu *per se* of no value in predicting recurrent disease.¹⁹ The other meta-analysis, by Joseph *et al.*, concerned 178,550 patients with PTC from five predominantly

registry-based studies, found that Mu had an impact on disease recurrence with a hazard ratio (HR) of 2.81; no data on mortality were considered.¹⁵ An interesting paper by Qu *et al.* analyzed whether the number of single tumor foci affected the prognosis: Mu emerged as a risk factor for recurrence, with a strong linear effect: the more numerous the foci, the higher the risk of recurrence.²⁰

In a very recent study by Geron *et al.*, on 1039 consecutive PTC patients, Mu confirmed its association with a more aggressive disease in terms of baseline characteristics, intensity of treatment, persistence/recurrence rates, and mortality. That said, after adjusting for confounding variables using a propensity score matching, Mu was no longer significantly associated with recurrence, long-term outcome, and mortality rates. The authors thus concluded that Mu in PTC is a marker of more extensive disease on presentation, but not an independent prognostic factor of disease outcome.¹⁸

As far as bilaterality is concerned, multifocal PTC may be bilateral in 13–71% of cases,¹¹ and the prognostic significance of this characteristic is still debated. A study by Kim *et al.* found Mu, but not bilaterality, associated with disease persistence/recurrence, and the authors judged that the number of tumor foci was of greater prognostic value than their location.¹¹ Another retrospective study analyzing 496 PTC patients found instead that bilaterality, as opposed to unilateral Mu, was an independent risk factor for neck recurrence (HR=4.052), distant metastasis (HR=3.860), and cancer-related death (HR=7.252).²¹

In the present study, we investigated the prognostic value of Mu and bilaterality in PTC in a large consecutive series of patients treated at a single center. We also made an effort to fully characterize multifocal PTC from a clinical, histopathological and molecular point of view. Our aim was ultimately to establish whether the sum of the diameters of single foci (SDSF), the total number of foci (TNF), and primary tumor size (PTS) are of prognostic significance in multifocal PTC.

Materials and methods

Patients

We conducted a retrospective analysis on a consecutive series of 370 adult patients at a single

center with thyroid nodules found malignant or suspect for TC (TIR 4–5 according to the SIAPEC 2014 consensus statement)²² on fine-needle aspiration cytology (FNAC). Between 2007 and 2017, all patients underwent total thyroidectomy and were histologically confirmed as cases of PTC.²³ Decisions regarding the extent of initial surgery in these patients (total thyroidectomy with or without prophylactic neck compartment dissection), ¹³¹I treatment and follow-up modalities have been reported elsewhere.²³

The histological variants of these cases of PTC were identified in accordance with World Health Organization (WHO) criteria.²⁴ Histological diagnostics and staging were managed according to the AJCC/UICC system (8th edition), and on the grounds of the first whole-body scan after ¹³¹I remnant ablation.^{5–8} The maximum diameter of the largest tumor focus was used to define primary tumor size (PTS) in each case. MicroPTC was diagnosed when the largest tumor was ≤ 1.0 cm in diameter. The methodology adopted for the molecular analysis to identify somatic TERT promoter and BRAF mutations has been described elsewhere.²³

Patient outcome was defined as an excellent response (ER), indeterminate response (IR), biochemically persistent disease (BPD), or structurally-persistent disease (SPD), according to the ATA guidelines for patients given ¹³¹I therapy,¹³ and the criteria proposed by Momesso *et al.* for non-radioiodine-treated patients.²⁵

Considering the low mortality rate in our series, patients were divided for the purposes of our study into three possible outcome groups: “BPD + SPD + TC-related death (TCD)” or “IR” or “ER”. The median follow-up period was 69 months (IQR: 42–92 months).

All studies were conducted in accordance with the guidelines of the Declaration of Helsinki. The present study was approved by the ethics committee of Azienda Ospedaliera di Padova (code N° AOP1303). All participants gave their written informed consent before enrolling for the study.

Unifocality, multifocality, bilaterality, and Mu-related variables: definitions

Two experienced pathologists (F.G. and G.P.) reviewed all pathology specimens to confirm all

diagnoses of PTC, their anatomopathological features, cases of Mu, the number of tumor foci, and the diameter of each one. Unifocality was defined as a solitary focus of PTC, and Mu as the presence of two or more tumor foci in the pathological specimen of thyroid. Bilaterality was defined as the presence of tumor foci in both thyroid lobes.

In cases of multifocal PTC, the Mu-related variables of interest were defined as follows: (1) the SDSF was the sum of the largest diameters of all tumor foci; (2) the TNF was the total number of tumor foci; and (3) the PTS corresponded to the size of the PTC focus with the largest diameter.

Statistical analysis

The Kolmogorov–Smirnov test was used to assess the normal distribution of each variable. All data were expressed as mean \pm standard deviation for normally distributed variables, and as median with interquartile range (IQR) for those not normally distributed. Mann–Whitney, and chi-square tests were used to compare clinical and pathological features, molecular mutational status, and oncological outcomes between the unifocal and multifocal PTC groups, as appropriate. When a dichotomized oncological outcome was needed for statistical purposes [multivariate Cox proportional hazards regression analyses, receiver operating characteristic (ROC) curves and Kaplan–Meier analysis], we pooled ER and IR in a “good outcome” group, and BPD, SPD, TCD in a “worse outcome” group. The same statistical analyses were used to compare the unilateral multifocal PTC, and bilateral-multifocal PTC groups.

To examine the prognostic factors on univariate analysis in the subset of multifocal PTCs, the Mann–Whitney and Kruskal–Wallis tests (for non-parametric variables), and Student’s *t*-test and one-way analysis of variance (for parametric variables) were used to correlate continuous variables with final oncological outcomes, as appropriate; and categorical variables were compared with outcome using the Chi-square test. Multivariate Cox proportional hazards regression analyses was used to identify the independent prognostic factors associated with outcome in the subset of multifocal PTCs, using a backward stepwise selection procedure with all clinically relevant variable. Disease-free survival (DFS) data were also analyzed using the Kaplan–Meier method. A *p* value < 0.05 was considered statistically significant.

Table 1. Descriptive characteristics of multifocality in patients with PTC.

	<i>n</i>	%
TNF		
2	80/152	52.6%
3	26/152	17.1%
4	14/152	9.2%
5	12/152	7.9%
6–10	10/152	6.6%
11–20	3/152	2%
21–30	6/152	4%
31–50	1/152	0.7%
SDSF (mm)		
5–10	12/152	7.9%
11–20	54/152	35.5%
21–30	44/152	29%
31–40	19/152	12.5%
41–50	8/152	5.2%
51–70	10/152	6.6%
71–116	5/152	3.3%
PTC, papillary thyroid cancer; SDSF, sum of diameters of single foci, TNF, total number of foci.		

Results

Patient characteristics

This study included 370 patients with histologically confirmed PTC who underwent total thyroidectomy, 282 (76.2%) females and 88 (23.8%) males, with a median age of 47 years (IQR 38–57 years). The median follow-up period was 69 months (IQR: 42–92 months).

In detail, the PTCs were histologically classified as follows: 248/370 (67%) classical variant; 24/370 (6.5%) follicular variant; 31/370 (8.4%) oxyphilic variant; and 67/370 (18.1%) were aggressive variants of PTC. Mu was seen in 152/370 PTCs (41.1%), and bilaterality in 111/370 (30%). Of the 152 multifocal PTCs, 41 (27%) were unilateral, and 111/152 (73%) were bilateral. SDSF and TNF descriptive characteristics of our population are shown in Table 1.

The median PTS in the study population as a whole was 14 mm (IQR 11–20 mm); 89/370 patients (24%) had microPTCs. Histology showed vascular invasion in 123/196 PTCs (62.2%). Cervical lymph node metastases (globally N1a and N1b) were identified in 160/341 patients (46.9%). In particular, level VI lymph node metastases (N1a) were confirmed histologically in 99/341 patients (29%), and lateral neck lymph node metastases were present in 61/341 patients (17.9%). Distant metastases were identified in 8/370 patients (2.2%).

According to the 8th edition of the TNM classification system (AJCC/UICC^{5–8}), 314/370 patients (84.9%) with PTC were classified as stage I, 50/370 (13.5%) in stage II, 1/370 (0.3%) in stage III, and 5/370 (1.4%) in stage IV at diagnosis. As concerns T-categories, 90/370 patients (24.3%) were in T1a, 181/370 (48.9%) in T1b, 69/370 (18.6%) in T2, 17/370 (4.6%) in T3a, 6/370 (1.6%) in T3b, 7/370 (1.9%) in T4a, and none in T4b.

BRAF mutations were identified in 232/368 (63%) PTCs, and TERT promoter mutations in 18/370 (4.9%).

According to the ATA guidelines, patients were IRSS scored as follows: 116/370 (31.4%) were at low risk; 234/370 (63.2%) were at intermediate risk; and 20/370 (5.4%) were at high risk.

At the end of follow up, 310/370 patients (83.8%) had an ER, 37/370 (10%) had an IR, 11/370 (3.0%) had a BPD, 9/370 (2.4%) had a SPD, and 3/370 (0.8%) had died of their disease (TCD).

For the purposes of our study, patients' final oncological outcomes were grouped as follows: 310/370 (83.8%) patients had an ER, 37/370 (10%) patients an IR and 23/370 (6.2%) a BPD or SPD or TCD.

Comparing unifocal with multifocal PTC

The unifocal and multifocal PTC groups did not differ significantly in terms of age at diagnosis, sex, histological variants, PTS, microPTC, vascular invasion, cervical lymph node involvement (central and lateral neck), distant metastases, TNM stage, IRSS score, postoperative radioactive iodine (RAI) therapy, administered activity of RAI, second treatment, follow-up period or BRAF and TERT mutation status (Table 2). When the two groups were compared by T-categories, on the other hand,

Table 2. Comparison of characteristics of patients with unifocal *versus* multifocal PTC patients.

	Unifocal PTC <i>n</i> = 218 (%)	Multifocal PTC <i>n</i> = 152 (%)	<i>p</i> value
Age at diagnosis [median; (IQR)]	47 years old (38–56)	46 years old (36.5–58)	NS
Gender			NS
Male	46 (21.1%)	42 (27.6%)	
Female	172 (78.9%)	110 (72.4%)	
PTC histological classification			NS
Classical variant	146 (66.9%)	102 (67.2%)	
Follicular variant	15 (6.9%)	9 (5.9%)	
Oxyphilic variant	20 (9.2%)	11 (7.2%)	
Aggressive Variants	37 (17.0%)	30 (19.7%)	
Primary tumor size [median; (IQR)]	15 mm (11–20.25)	13 mm (10–20)	NS
Primary tumor size			NS
≤10 mm (microPTC)	45 (20.6%)	44 (28.9%)	
>10 mm	172 (79.4%)	108 (71.1%)	
Vascular invasion	55 (57.3%)	68 (68%)	NS
Extrathyroidal extension	115 (57.2%)	85 (58.6%)	NS
Cervical lymph node involvement	88 (43.8%)	72 (51.4%)	NS
Cervical lymph node involvement			NS
Central, N1a	57 (28.4%)	42 (30%)	
Lateral, N1b	31 (15.4%)	30 (21.4%)	
Distant metastases	6 (2.8%)	2 (1.3%)	NS
TNM stage (8th edition)			NS
Stage I	184 (84.4%)	130 (85.5)	
Stage II	29 (13.3%)	21 (13.8%)	
Stage III	0	1 (0.7%)	
Stage IV	5 (2.3%)	0	
T-categories (TNM, 8th edition)			0.04
T1a	47 (21.6%)	43 (28.3%)	
T1b	113 (51.8%)	68 (44.7%)	
T2	47 (21.6%)	22 (14.5%)	
T3a	6 (2.8%)	11 (7.2%)	
T3b	2 (0.9%)	4 (2.6%)	
T4a	3 (1.4%)	4 (2.6%)	
T4b	0	0	

(Continued)

Table 2. (Continued)

	Unifocal PTC <i>n</i> = 218 (%)	Multifocal PTC <i>n</i> = 152 (%)	<i>p</i> value
Postoperative RAI therapy			NS
Yes	208 (95.4%)	139 (91.4%)	
No	10 (4.6%)	13 (8.6%)	
Administered activity of RAI [median; (IQR)]	100 mCi (100–150)	100 mCi (70–150)	NS
Second treatment	20 (9.2%)	22 (14.5%)	NS
Median follow-up period [median; (IQR)]	70.5 months (40–94)	65 months (44.5–87)	NS
BRAF mutation (FNAC)	132 (60.8%)	100 (66.2%)	NS
TERT promoter mutation (FNAC)	12 (5.5%)	6 (3.9%)	NS
Initial risk stratification system scores			NS
Low	74 (33.9%)	41 (27%)	
Intermediate	135 (61.9%)	100 (65.8%)	
High	9 (4.1%)	11 (7.2%)	
Disease status at latest follow-up (ongoing risk stratification)			NS
ER	190 (87.2%)	120 (78.9%)	
IR	16 (7.3%)	21 (13.8%)	
BPD	5 (2.3%)	6 (3.9%)	
SPD	5 (2.3%)	4 (2.6%)	
TCD	2 (0.9%)	1 (0.7%)	
Final oncological outcome			NS
ER	190 (87.2%)	120 (79.8%)	
IR	16 (7.3%)	21 (13.8%)	
BPD + SPD + TCD	12 (5.5%)	11 (7.2%)	

BPD, biochemically persistent disease; ER, excellent response; FNAC, fine needle aspiration cytology; IQR, interquartile range; IR, indeterminate response; NS, not significant; PTC, papillary thyroid cancer; RAI, radioactive immunotherapy; SPD, structurally persistent disease; TCD, death due to thyroid cancer.

there was a statistically significant difference between the multifocal and unifocal PTC groups, with a moderately higher frequency of cases in T3a, T3b, and T4a in the multifocal PTC group ($p=0.04$). At the end of the follow up, the multifocal and unifocal PTC groups did not differ in terms of final oncological outcome, however.

Comparing unilateral multifocal with bilateral multifocal PTC

The multifocal PTC groups with unilateral as opposed to bilateral disease did not differ significantly in terms of age at diagnosis, sex, histological

variants, vascular invasion, cervical lymph node involvement (central and lateral neck), distant metastases, TNM stage, IRSS score, postoperative RAI therapy, administered activity of RAI, second treatment, follow-up period or BRAF and TERT mutation status (Table 3). Patients with bilateral multifocal PTCs had a larger PTS ($p=0.01$), and microPTCs were less frequent ($p=0.02$) than in the group with unilateral multifocal PTCs. Comparing the two groups by T-categories, a significant difference emerged, with higher T-categories in the bilateral multifocal PTC group ($p=0.04$). Here again, at the end of the follow up, the groups with unilateral *versus* bilateral-multifocal

Table 3. Comparison between unilateral multifocal and bilateral multifocal PTC.

	Unilateral multifocal PTC <i>n</i> =41 (%)	Bilateral multifocal PTC <i>n</i> =111 (%)	<i>p</i> value
Age at diagnosis (mean \pm DS)	46years old \pm 14	48years old \pm 15	NS
Gender			NS
Male	12 (29.3%)	30 (27%)	
Female	29 (70.7%)	81 (73%)	
PTC histological classification			NS
Classical variant	31 (75.6%)	71 (64%)	
Follicular variant	2 (4.9%)	7 (6.3%)	
Oxyphilic variant	1 (2.4%)	10 (9%)	
Aggressive Variants	7 (17.1%)	23 (20.7%)	
Primary tumor size [median; (IQR)]	11 mm (8–16)	14 mm (11–21.75)	0.01
Primary tumor size			0.02
≤ 10 mm (microPTC)	18 (43.9%)	26 (23.4%)	
> 10 mm	23 (56.1%)	85 (76.6%)	
Vascular invasion	21 (84%)	47 (62.7%)	NS
Cervical lymph node involvement	17 (44.7%)	55 (53.9%)	NS
Cervical lymph node involvement			NS
Central, N1a	9 (23.7%)	33 (32.4%)	
Lateral, N1b	8 (21.1%)	22 (21.6%)	
Distant metastases	0	2 (1.8%)	NS
TNM stage (8th edition)			NS
Stage I	37 (90.2%)	93 (83.3%)	
Stage II	3 (7.3%)	18 (16.2%)	
Stage III	1 (2.4%)	0	
Stage IV	0	0	
T-categories (TNM, 8th edition)			0.04
T1a	19 (46.3%)	24 (21.6%)	
T1b	15 (36.6%)	53 (47.7%)	
T2	5 (12.2%)	17 (15.3%)	
T3a	0	11 (9.9%)	
T3b	1 (2.4%)	3 (2.7%)	

(Continued)

Table 3. (Continued)

	Unilateral multifocal PTC <i>n</i> =41 (%)	Bilateral multifocal PTC <i>n</i> =111 (%)	<i>p</i> value
T4a	1 (2.4%)	3 (2.7%)	
T4b	0	0	
Postoperative RAI therapy	35 (85.4%)	103 (93.6%)	NS
Administered activity of RAI [median; (IQR)]	100 mCi (77.5–150)	100 mCi (70–150)	NS
Second treatment	5 (12.2%)	17 (15.3%)	NS
Follow-up period (mean ± DS)	71.5 months ± 35	65.5 months ± 30	NS
BRAF mutation (FNAC)	27 (65.9%)	73 (66.4%)	NS
TERT promoter mutation (FNAC)	1 (2.4%)	5 (4.5%)	NS
Initial risk stratification system scores			NS
Low	13 (31.3%)	28 (25.2%)	
Intermediate	25 (61%)	75 (67.6%)	
High	3 (7.3%)	8 (7.2%)	
Disease status at latest follow-up (ongoing risk stratification)			NS
ER	34 (82.9%)	86 (77.5%)	
IR	6 (14.6%)	15 (13.5%)	
BPD	1 (2.4%)	5 (4.5%)	
SPD	0	4 (3.6%)	
TCD	0	1 (0.9%)	
Final oncological outcome			NS
ER	34 (82.9%)	86 (77.5%)	
IR	6 (14.6%)	15 (13.5%)	
BPD + SPD + TCD	1 (2.4%)	10 (9%)	
BPD, biochemically persistent disease; ER, excellent response; FNAC, fine needle aspiration cytology; IQR, interquartile range; IR, indeterminate response; NS, not significant; PTC, papillary thyroid cancer; RAI, radioactive immunotherapy; SPD, structurally persistent disease; TCD, death due to thyroid cancer.			

PTCs did not differ in terms of final oncological outcome.

Risk factors for persistent/recurrent disease or disease-related death in multifocal PTC

We analyzed the factors that could predict persistent/recurrent disease or disease-related death

in the group of patients with multifocal PTC (Table 4).

On univariate analysis, PTS ($p=0.004$), SDSF ($p=0.004$), vascular invasion ($p=0.03$), lymph node involvement ($p=0.03$), with N1b carrying a higher risk than N1a ($p=0.001$), distant metastases ($p<0.0001$), stage at diagnosis ($p=0.03$),

Table 4. Univariate analysis of prognostic factors for persistent/recurrent disease or disease-related death in patients with multifocal PTC.

	Excellent response n = 120 (%)	Indeterminate response n = 21 (%)	BPD + SPD + TCD n = 11 (%)	p value
Age at diagnosis (mean ± SD)	48 years ± 14	47 years ± 15	46 years ± 23	NS
Gender				NS
Male	33 (27.5%)	4 (19%)	5 (45.5%)	
Female	87 (72.5%)	17 (81%)	6 (54.5%)	
PTC histological classification				NS
Classical variant	81 (67.5%)	15 (71.4%)	6 (54.5%)	
Follicular variant	7 (5.8%)	1 (4.8%)	1 (9.1%)	
Oxyphilic variant	9 (7.5%)	1 (4.8%)	1 (9.1%)	
Aggressive variants	23 (19.2%)	4 (19%)	3 (27.3%)	
PTS [median; (IQR)]	13 mm (9–20)	12 mm (10.5–15)	25 mm (16.25–41.5)	0.004
Primary tumor size				NS
≤10 mm (microPTC)	39 (32.5%)	5 (23.8%)	0	
>10 mm	81 (67.5%)	16 (76.2%)	11 (100%)	
TNF [median; (IQR)]	2 (2–4)	3 (2–6.25)	3 (2–3)	NS
SDSF [median; (IQR)]	21 mm (14–30)	23 mm (15.5–32.75)	62 mm (24–67.5)	0.004
Vascular invasion	46 (61.3%)	15 (83.3%)	7 (100%)	0.03
Cervical lymph node involvement	51 (45.9%)	13 (68.4%)	8 (80%)	0.03
Cervical lymph node involvement				0.001
Central, N1a	33 (29.7%)	8 (42.1%)	1 (10%)	
Lateral, N1b	18 (16.2%)	5 (26.3%)	7 (70%)	
Distant metastases	0	0	2 (18.2%)	<0.0001
Bilaterality	86 (71.7%)	15 (71.4%)	10 (90.9%)	NS
TNM stage (8th edition)				0.03
Stage I	105 (87.5%)	19 (90.5%)	6 (54.5%)	
Stage II	14 (11.7%)	2 (9.5%)	5 (45.5%)	
Stage III	1 (0.8%)	0	0	
Stage IV	0	0	0	
T-categories based on PTS (TNM, 8th edition)				0.004
T1a	37 (30.8%)	5 (23.8%)	0	
T1b	53 (44.5%)	14 (66.7%)	4 (36.4%)	

(Continued)

Table 4. (Continued)

	Excellent response <i>n</i> = 120 (%)	Indeterminate response <i>n</i> = 21 (%)	BPD + SPD + TCD <i>n</i> = 11 (%)	<i>p</i> value
T2	17 (14.2%)	1 (4.8%)	2 (18.2%)	
T3a	7 (5.8%)	1 (4.8%)	3 (27.3%)	
T3b	4 (3.3%)	0	0	
T4a	2 (1.7%)	0	2 (18.2%)	
T4b	0	0	0	
T-categories based on SDSF (TNM, 8th edition)				0.0003
T1a	12 (10%)	2 (9.5%)	0	
T1b	42 (35%)	7 (33.3%)	1 (9.1%)	
T2	48 (40%)	10 (47.6%)	2 (18.2%)	
T3a	12 (10%)	2 (9.5%)	6 (54.5%)	
T3b	4 (3.3%)	0	0	
T4a	2 (1.7%)	0	2 (18.2%)	
T4b	0	0	0	
Postoperative RAI therapy	109 (91.6%)	19 (90.5%)	10 (90.9%)	NS
Administered activity of RAI [median; (IQR)]	100 mCi (70–150)	125 mCi (50–150)	150 mCi (150–150)	0.03
Follow-up period (mean ± DS)	69 months ± 30	68 months ± 31	46 months ± 40	NS
Second treatment	6 (5%)	8 (38.1%)	8 (72.7%)	<0.0001
BRAF mutation (FNAC)	78 (65.5%)	13 (61.9%)	9 (81.8%)	NS
TERT promoter mutation (FNAC)	4 (3.3%)	0	2 (18.2%)	0.03
Initial risk stratification system scores				0.0003
Low	37 (31.1%)	2 (9.5%)	2 (18.2%)	
Intermediate	78 (65.5%)	16 (76.2%)	5 (45.5%)	
High	4 (3.4%)	3 (14.3%)	4 (36.4%)	

BPD, biochemical persistent disease; FNAC, fine needle aspiration cytology; IQR, interquartile range; NS, not significant; PTS, primary tumor size; RAI, radioactive immunotherapy; SDSF, sum of diameters of single foci; SPD, structural persistent disease; TCD, death due to thyroid cancer; TNF, total number of foci.

T-categories ($p = 0.004$), IRSS score ($p = 0.0003$), administered activity of RAI ($p = 0.03$), second treatment ($p < 0.0001$) and TERT promoter mutation ($p = 0.03$) all correlated significantly with the risk of persistent/recurrent disease or disease-related death in the multifocal PTC group. When we reclassified patients'

T-categories based on SDSF instead of PTS, T-categories was even more strongly associated with PTC persistence/recurrence or death ($p = 0.0003$).

On multivariate Cox proportional hazards regression analyses, only distant metastases




Figure 1. ROC curve analysis on the sum of the diameters of single foci and persistent/recurrent disease or death in multifocal PTC.

AUC, area under the curve; NPV, negative predictive value; PPV, positive predictive value; PTC, papillary thyroid cancer; ROC, receiver operating characteristic.

(HR 39.4761, 95% CI 6.4794–240.5100, $p=0.0001$), and SDSF (HR 1.0734, 95% CI 1.0212–1.128, $p=0.0056$) independently predicted persistent/recurrent disease or disease-related death in multifocal PTC patients.

Considering the prognostic impact of SDSF in multifocal PTCs, we were able to identify a cut-off of 40 mm using ROC curve analysis ($p=0.0002$, area under the curve 0.792; Figure 1). In other words, SDSF >40 mm in cases of multifocal PTC can predict a worse prognosis with a sensitivity of 63.6% and specificity of 88.7%. The cut-off also showed a very high negative predictive value of 96.9%. Adopting this SDSF cut-off of 40 mm, we analyzed DFS using the Kaplan–Meier method, and dividing multifocal PTC patients by SDSF ≥ 40 mm or <40 mm (Figure 2). The 5-year DFS rates were 71.1% in multifocal PTC patients with SDSF ≥ 40 mm and 96.8% in those with SDSF <40 mm. A log-rank test showed that the DFS rate was significantly lower ($p < 0.0001$) in patients with SDSF ≥ 40 mm.




Figure 2. Kaplan–Meier curves estimating disease-free survival for patients with multifocal PTC based on a SDSF ≥ 40 mm or <40 mm.

PTC, papillary thyroid cancer; SDSF, sum of the diameters of single foci.

Discussion

Proper risk stratification is the mainstay of appropriate clinical management for patients with PTC. In this scenario, Mu is often empirically considered by clinicians as a factor suggesting a poor prognosis, prompting them to opt for more aggressive treatments and follow up, although the ATA's IRSS includes Mu as a low-risk feature with an estimated risk of 4–6% of persistent/recurrent disease. The recent ATA guidelines only recommend considering Mu in cases of microPTC, however, and offer no specific recommendations on Mu in patients with PTCs larger than 10 mm.¹³ In short, the prognostic significance of Mu in PTC has often been investigated, but remains an open question.

In our study, the prevalence of Mu was 41.1% – a figure consistent with the literature.^{10,11} In previous studies, Mu in PTC was associated with older age,^{15,18} male sex,^{18,19} and extrathyroidal extension.^{11,15,17} It is only in some studies that it was also associated with macroPTC, vascular invasion, lymph node involvement, more advanced disease and higher IRSS scores on presentation.^{16,18,26} In our series, patients with multifocal as opposed to unifocal PTCs were not dissimilar in terms of age at diagnosis, sex, histological

variants, PTS, presence of microPTC, vascular invasion, cervical lymph node involvement (central and lateral neck), distant metastases, TNM stage, IRSS score, or BRAF and TERT mutational status. On the other hand, more advanced T-categories (on TNM staging) were more frequently associated with multifocality, as reported by other groups.^{10,16,18}

Bilaterality was seen in 30% of our whole population of PTC patients, and in 73% of those with multifocal PTC; these proportions are similar to those found in other series.^{11,21} Only a few studies have investigated bilateral multifocal PTC characteristics.^{11,21} In our population, bilaterality was associated with more aggressive features, such as larger PTS, fewer cases of microPTC and more advanced T-categories on TNM staging. This is consistent with previous studies by Qu *et al.*, and Hwang *et al.*, who found that bilateral multifocality tended to be associated with more aggressive tumor characteristics.^{21,27}

How Mu and bilaterality correlate with oncological outcome in PTC is a more complicated issue, however.

A powerful multicenter study by Wang *et al.* investigated the prognostic value of Mu in 2638 patients from 11 centers, whose data were fully replicated and validated in a complementary database of 89,680 patients. As found in previous studies, univariate analyses confirmed an association between Mu and lymph node involvement, extrathyroidal extension, T-categories, and recurrence. On multivariate analysis, however, after adjusting for classical clinicopathological risk factors, Mu was no longer associated with disease recurrence. The authors concluded that Mu had no independent role in predicting PTC recurrence, nor any prognostic impact on patient mortality, on univariate or multivariate analysis.¹⁷ In contrast, another meta-analysis involving 178,550 patients with PTC from five studies concluded that Mu could predict disease recurrence with an HR of 2.81.¹⁵ Two separate Korean groups also found in retrospective series that Mu was an independent risk factor for disease persistence/recurrence of PTCs larger than 10 mm.^{10,28} As far as Mu-related variables are concerned, an interesting paper by Qu *et al.* analyzed the possible prognostic significance of TNF, finding that having more numerous tumor foci had a strong linear correlation with the probability of recurrent disease.²⁰

In our study population, neither Mu nor bilaterality *per se* had any prognostic impact on PTC persistence/recurrence or mortality. We therefore wondered whether we could identify any prognostic factors in the setting of multifocal PTC that would enable us to customize patient follow-up based on the risk of persistence/recurrence or death.

In multifocal PTC, the classical prognostic factors such as vascular invasion, cervical lymph node involvement, distant metastases, TERT mutation, IRSS score, TNM stage and T-categories all confirmed their significance in predicting persistent/recurrent disease.^{9,10,25,28} Interestingly, among the Mu-related variables, only SDSF and PTS correlated with the risk of persistence/recurrence or TCD, while TNF failed to predict recurrent/persistent disease.

Only few studies have investigated the role of SDSF in multifocal PTC. In particular, a recent paper by Feng *et al.* found that multifocal microPTC with SDSF > 10 mm tend to behave like the multifocal macroPTC in terms of clinicopathological features and prognosis.²⁹ Another paper by Liu *et al.* reached similar conclusions observing that multifocal PTC with SDSF > 10 mm showed a lower DFS than that with SDSF ≤ 10 mm.³⁰ However, authors analyzed SDSF only as a categorical variable using 10 mm as a cut-off.

Another interesting work by Tam *et al.* tackles the topic from a different point of view.³¹ They investigated the role of the tumor diameter ratio (ratio of primary tumor diameter to total tumor diameter) in identifying multifocal PTCs with more aggressive characteristics. In their study a decreased tumor diameter ratio was associated with capsular invasion, extrathyroidal extension and lymph node metastasis in patients with multifocal microPTC and PTC, but they did not analyze its impact on final prognosis.

Instead, we investigated the impact of SDSF on prognosis as a continuous variable for the first time in literature. In our opinion, the prognostic importance of SDSF is particularly noteworthy because this novel anatomopathological factor emerged as an independent predictor of persistence/recurrence or death in multifocal PTC. SDSF might represent what we could call the total tumor burden of multifocal PTC. There is

certainly plenty of evidence to suggest that tumor burden has prognostic value in numerous human tumors.^{32–37} In the particular case of multifocal PTC, SDSF may reflect the tumor burden better than the size of the largest focus, that is, PTS. This impression is supported by the SDSF-revised T-categories showing an even stronger association with persistence/recurrence ($p = 0.0003$ *versus* $p = 0.01$) in our series than the T-categories based on PTS.

Using ROC curve analysis, we identified a cut-off of 40 mm for SDSF as capable of distinguishing between multifocal PTC patients at higher or lower risk of persistence/recurrence or TCD. This is remarkably consistent with the well-known prognostic significance of tumor size larger than 4 cm, even in unifocal PTC.^{9,25} The strength of such a cut-off is confirmed by its high negative predictive value (96.9%). In short, SDSF can be useful as a rule-out test: multifocal PTC patients with SDSF < 40 mm are at very low risk of persistent/recurrent disease or TCD. This new parameter could be useful for proper risk stratification and for customizing the frequency of follow-up for multifocal PTC patients.

As further confirmation, the DFS rate in multifocal PTCs with SDSF ≥ 40 mm was also significantly lower.

Our study has some limitations to bear in mind, including: its retrospective nature and the relatively short follow-up period; the relatively low rate of persistent/recurrent disease or cancer-related death in our series; and the small number of patients, which could have affected the chances of finding substantial differences between multifocal and unifocal PTC (although the small size of our series of multifocal PTCs enabled us to conduct a careful and extensive pathological review to define the tumor burden accurately).

In conclusion, multifocality and bilaterality *per se* seemed to have no prognostic impact on PTC persistence/recurrence or cancer-related death in the present study. To our knowledge, this is the first study to investigate the prognostic role of SDSF as a continuous variable in multifocal PTC, which emerged as a novel independent predictor of tumor persistence/recurrence or cancer-related death. SDSF might better represent the tumor burden in cases of multifocal

PTC. A cut-off of 40 mm enabled us to identify multifocal PTC patients with a good prognosis, making SDSF a useful tool for risk stratification in patients with multifocal PTC.

Acknowledgements

We thank Frances Coburn for text editing.

Author contribution(s)

Jacopo Manso: Conceptualization; Data curation; Formal analysis; Investigation; Methodology; Writing-original draft.

Simona Censi: Data curation; Investigation; Writing-review & editing.

Amir Roberti: Data curation; Investigation; Writing-original draft.

Maurizio Iacobone: Data curation; Investigation; Writing-review & editing.

Susi Barollo: Data curation; Investigation; Methodology; Writing-review & editing.

Loris Bertazza: Data curation; Formal analysis; Writing-original draft.

Francesca Galuppini: Data curation; Investigation; Writing-review & editing.

Federica Vianello: Data curation; Investigation; Writing-review & editing.

Nora Albiger: Data curation; Investigation; Writing-review & editing.

Carla Scaroni: Conceptualization; Methodology; Writing-review & editing.

Gianmaria Pennelli: Data curation; Investigation; Writing-review & editing.

Caterina Mian: Conceptualization; Data curation; Formal analysis; Investigation; Methodology; Writing-original draft; Writing-review & editing.

Consent for publication

All participants gave their written informed consent for publication of the results.

Consent to participate

All participants gave their written informed consent before enrolling for the study.

Conflict of interest statement

The authors declare that there is no conflict of interest.

Ethical approval

We declare under our responsibility that the present study was approved by our local ethical committee (Azienda Ospedaliera di Padova, approval code number: AOP1303), and all patients gave their written informed consent to the use of their thyroid cytology and histology findings for research purpose.

Funding

The authors received no financial support for the research, authorship, and/or publication of this article.

ORCID iD

Jacopo Manso  <https://orcid.org/0000-0003-2403-4339>

References

1. Siegel R, Ma J, Zou Z, *et al.* Cancer statistics, 2014. *CA Cancer J Clin* 2014; 64: 9–29.
2. Davies L and Welch HG. Current thyroid cancer trends in the United States. *JAMA Otolaryngol Head Neck Surg* 2014; 140: 317–322.
3. Sherman SI. Thyroid carcinoma. *Lancet* 2003; 361: 501–511.
4. Eustatia-Rutten CFA, Corssmit EPM, Biermasz NR, *et al.* Survival and death causes in differentiated thyroid carcinoma. *J Clin Endocrinol Metab* 2006; 91: 313–319.
5. Brierley JD, Panzarella T, Tsang RW, *et al.* A comparison of different staging systems predictability of patient outcome: thyroid carcinoma as an example. *Cancer* 1997; 79: 2414–2423.
6. Lo CY, Chan WF, Lam KY, *et al.* Follicular thyroid carcinoma: the role of histology and staging systems in predicting survival. *Ann Surg* 2005; 242: 708–715.
7. Huang SH and O'Sullivan B. Overview of the 8th edition TNM classification for head and neck cancer. *Curr Treat Options Oncol* 2017; 18: 40.
8. Amin MB, Edge S, Greene FL, *et al.* (eds). *AJCC cancer staging manual*. 11th ed. New York: Springer, 2017.
9. Kim TH, Kim YN, Kim HI, *et al.* Prognostic value of the eighth edition AJCC TNM classification for differentiated thyroid carcinoma. *Oral Oncol* 2017; 71: 81–86.
10. Kim K-J, Kim S-M, Lee YS, *et al.* Prognostic significance of tumor multifocality in papillary thyroid carcinoma and its relationship with primary tumor size: a retrospective study of 2,309 consecutive patients. *Ann Surg Oncol* 2015; 22: 125–131.
11. Kim HJ, Sohn SY, Jang HW, *et al.* Multifocality, but not bilaterality, is a predictor of disease recurrence/persistence of papillary thyroid carcinoma. *World J Surg* 2013; 37: 376–384.
12. Kiriakopoulos A, Petralias A and Linos D. Multifocal versus solitary papillary thyroid carcinoma. *World J Surg* 2016; 40: 2139–2143.
13. Haugen BR, Alexander EK, Bible KC, *et al.* 2015 American thyroid association management guidelines for adult patients with thyroid nodules and differentiated thyroid cancer: the American thyroid association guidelines task force on thyroid nodules and differentiated thyroid cancer. *Thyroid* 2016; 26: 1–133.
14. Pacini F, Schlumberger M, Dralle H, *et al.* European consensus for the management of patients with differentiated thyroid carcinoma of the follicular epithelium. *Eur J Endocrinol* 2006; 154: 787–803.
15. Joseph KR, Edirimanne S and Eslick GD. Multifocality as a prognostic factor in thyroid cancer: a meta-analysis. *Int J Surg* 2018; 50: 121–125.
16. Genpeng L, Jianyong L, Jiaying Y, *et al.* Independent predictors and lymph node metastasis characteristics of multifocal papillary thyroid cancer. *Medicine (Baltimore)* 2018; 97: e9619.
17. Wang F, Yu X, Shen X, *et al.* The prognostic value of tumor multifocality in clinical outcomes of papillary thyroid cancer. *J Clin Endocrinol Metab* 2017; 102: 3241–3250.
18. Geron Y, Benbassat C, Shteinshneider M, *et al.* Multifocality is not an independent prognostic factor in papillary thyroid cancer: a propensity score-matching analysis. *Thyroid* 2019; 29: 513–522.
19. Guo K and Wang Z. Risk factors influencing the recurrence of papillary thyroid carcinoma: a systematic review and meta-analysis. *Int J Clin Exp Pathol* 2014; 7: 5393–5403.
20. Qu N, Zhang L, Ji Q, *et al.* Number of tumor foci predicts prognosis in papillary thyroid cancer. *BMC Cancer* 2014; 14: 914.
21. Qu N, Zhang L, Wu W-L, *et al.* Bilaterality weighs more than unilateral multifocality in predicting prognosis in papillary thyroid cancer. *Tumour Biol* 2016; 37: 8783–8789.
22. Nardi F, Basolo F, Crescenzi A, *et al.* Italian consensus for the classification and reporting of

- thyroid cytology. *J Endocrinol Invest* 2014; 37: 593–599.
23. Censi S, Barollo S, Grespan E, *et al.* Prognostic significance of TERT promoter and BRAF mutations in TIR-4 and TIR-5 thyroid cytology. *Eur J Endocrinol* 2019; 181: 1–11.
 24. DeLellis RA, Lioyd RV, Heitz PU, *et al.* (eds). *WHO classification of tumours: pathology and genetics of tumours of endocrine organs*. Lyon: IARC Press, 2004.
 25. Nam SH, Bae MR, Roh JL, *et al.* A comparison of the 7th and 8th editions of the AJCC staging system in terms of predicting recurrence and survival in patients with papillary thyroid carcinoma. *Oral Oncol* 2018; 87: 158–164.
 26. Gur EO, Karaisli S, Hacıyanli S, *et al.* Multifocality related factors in papillary thyroid carcinoma. *Asian J Surg* 2019; 42: 297–302.
 27. Hwang E, Pakdaman MN, Tamilia M, *et al.* Bilateral papillary thyroid cancer and associated histopathologic findings. *J Otolaryngol Head Neck Surg* 2010; 39: 284–287.
 28. Choi WR, Roh JL, Gong G, *et al.* Multifocality of papillary thyroid carcinoma as a risk factor for disease recurrence. *Oral Oncol* 2019; 94: 106–110.
 29. Feng JW, Pan H, Wang L, *et al.* Total tumor diameter: the neglected value in papillary thyroid microcarcinoma. *J Endocrinol Invest* 2020; 43: 601–613.
 30. Liu C, Wang S, Zeng W, *et al.* Total tumour diameter is superior to unifocal diameter as a predictor of papillary thyroid microcarcinoma prognosis. *Sci Rep*. Epub ahead of print 1 December 2017. DOI: 10.1038/s41598-017-02165-6.
 31. Tam AA, Özdemir D, Çuhacı N, *et al.* Can ratio of the biggest tumor diameter to total tumor diameter be a new parameter in the differential diagnosis of aggressive and favorable multifocal papillary thyroid microcarcinoma? *Oral Oncol* 2017; 65: 1–7.
 32. Hsu CY, Liu PH, Ho SY, *et al.* Impact of tumor burden on prognostic prediction for patients with terminal stage hepatocellular carcinoma: a nomogram study. *PLoS One*. Epub ahead of print 1 November 2017. DOI: 10.1371/journal.pone.0188031.
 33. Chen C, Fei Z, Huang C, *et al.* Prognostic value of tumor burden in nasopharyngeal carcinoma. *Cancer Manag Res* 2018; 10: 3169–3175.
 34. Kulyova SA and Karitsky AP. Predictive value of Hodgkin's lymphoma tumor burden in present. *Vestn Ross Akad meditsinskikh Nauk* 2014; 69: 67–71.
 35. Poklepovic AS and Carvajal RD. Prognostic value of low tumor burden in patients with melanoma. *Oncology (Williston Park)* 2018; 32: e90–e96.
 36. Sasaki K, Morioka D, Conci S, *et al.* The tumor burden score: a new 'metro-ticket' prognostic tool for colorectal liver metastases based on tumor size and number of tumors. *Ann Surg* 2018; 267: 132–141.
 37. Gobbi PG. Tumor burden in Hodgkin's lymphoma: much more than the best prognostic factor. *Crit Rev Oncol Hematol* 2014; 90: 17–23.

Visit SAGE journals online
journals.sagepub.com/
home/tae

 SAGE journals