

## Supplementary information

### Enhancing biological activities and phenolic content of wild grapevine roots by severe drought stress

Samia Daldoul<sup>1\*</sup>, Faouzia Hanzouli<sup>1,2</sup>, Nadia fares<sup>3</sup>, Olfa Tabbene<sup>3</sup>, Hassène Zemni<sup>1</sup>, Simone Vincenzi<sup>4</sup>, Ahmed Mliki<sup>1</sup>, Mahmoud Gargouri<sup>1</sup>


<sup>1</sup>Laboratory of Plant Molecular Physiology, Center of Biotechnology of Borj-Cedria, Hammam-Lif, BP 901, 2050 Tunisia.

<sup>2</sup>Faculty of Sciences of Tunis, University Tunis El-Manar, 2092, El Manar II, Tunis, Tunisia.

<sup>3</sup>Laboratory of Bioactive Substances, Center of Biotechnology of Borj Cedria, BP-901, 2050 Hammam-lif, Tunisia.

<sup>4</sup>University of Padova, Department of Agronomy, Food, Natural resources, Animals and Environment (DAFNAE), Viale dell'Università, 16, 35020 Legnaro (PD). Italy.

**Corresponding author:** Samia Daldoul, email: Samia.daldoul@cbbc.rnrt.tn



60

61 **Fig. S1** Representative HPLC-UV (310 nm) profile of root ethanolic extracts of "Tebaba" genotype  
62 from control (C) and drought stress (DS) treatments of 15 days (a, b) and 20 days (c,d). comparing  
63 to the standard stilbene solution (e). Numbers on the arrow refers to (1) *t*-piceid, (2) *t*-piceatannol,  
64 (3) *t*-resveratrol, (4) *t*- $\epsilon$ -viniferin, (5) *t*-pterostilbene and (6) *t*-hydroxystilbene

65