

Kinetics of lymphocytosis in naïve chronic lymphocytic leukemia patients treated with covalent Bruton's tyrosine kinase inhibitors: An Italian multicenter real-life experience

Idanna Innocenti¹ | Antonio Mosca² | Annamaria Tomasso²  | Andrea Galitzia³ | Lydia Scarfò^{4,5} | Francesca Morelli⁶ | Eugenio Galli^{1,2} | Francesca Martini⁴ | Eugenio Sangiorgi⁷ | Roberta Laureana⁸ | Giulia Benintende⁹ | Veronica Mattiello¹⁰ | Sabrina Chiriu¹¹ | Maria I. Del Principe⁸ | Giulia Zamprognà¹² | Massimo Gentile¹³ | Enrica A. Martino¹³ | Emilia Cappello¹⁴ | Maria C. Montalbano¹⁵ | Giuliana Farina¹⁶ | Vanessa Innao¹⁷ | Luca Stirparo² | Caterina Patti¹⁸ | Paolo Sportoletti¹⁹ | Alberto Fresa^{1,2} | Gioacchino Catania²⁰ | Marta Coscia¹⁵ | Silvia Bellesi¹ | Alessandra Tedeschi²¹ | Alessandro Sanna²² | Andrea Visentin²³ | Francesco Autore¹ | Raffaella Pasquale²⁴ | Livio Trentin²³ | Marzia Varettoni²⁵ | Paolo Ghia^{4,5}  | Roberta Murru²⁶ | Luca Laurenti^{1,2}

Correspondence: Luca Laurenti (luca.laurenti@unicatt.it)

¹Dipartimento di Scienze di Laboratorio ed Ematologiche, Area Ematologia, Fondazione Policlinico Universitario A. Gemelli IRCCS, Università Cattolica del Sacro Cuore, Rome, Italy

²Sezione di Ematologia, Dipartimento di Scienze Radiologiche ed Ematologiche, Università Cattolica del Sacro Cuore, Rome, Italy

³Hematology and Stem Cell Transplantation Unit, Ospedale San Francesco, Nuoro, Italy

⁴Strategic Research Program on CLL, IRCCS Ospedale San Raffaele, Milano, Italy

⁵Medical School Università Vita Salute San Raffaele, Milano, Italy

⁶Hematology, University of Florence, Florence, Italy

⁷Dipartimento di Scienze della Vita e Sanità Pubblica, Sezione di Medicina Genomica, Università Cattolica del Sacro Cuore, Rome, Italy

⁸Division of Haematology, University of Tor Vergata, Rome, Italy

⁹Department of Medicine 5, Haematology and Oncology, Friedrich-Alexander-Universität Erlangen-Nürnberg (FAU), Universitätsklinikum, Erlangen, Germany

¹⁰Hematology Unit, Fondazione IRCCS Ca' Granda Policlinico, Milano, Italy

¹¹Department of Medical Sciences and Public Health, University of Cagliari, Cagliari, Italy

¹²Department of Hematology, Niguarda Cancer Center, ASST Grande Ospedale Metropolitano Niguarda, Milano, Italy

¹³Hematology Section, Cosenza Hospital, Cosenza, Italy

¹⁴Dipartimento di Medicina Molecolare, Università di Pavia, Pavia, Italy

¹⁵Department of Molecular Biotechnology and Health Sciences, University of Torino and Division of Hematology, University Hospital (A.O.U.) Città della Salute e della Scienza di Torino, Torino, Italy

¹⁶Hematology and Medical Oncology, AORN Sant'Anna e San Sebastiano, Caserta, Italy

¹⁷UOC di Ematologia, Azienda Ospedaliera di rilievo nazionale di alta specializzazione, ARNAS - Garibaldi di Catania, Catania, Italy

¹⁸Divisione di Oncematologia, Azienda Villa Sofia Cervello, Palermo, Italy

¹⁹Department of Medicine and Surgery, Institute of Hematology, Centro di Ricerca Emato Oncologica (CREO), University of Perugia, Perugia, Italy

²⁰Division of Hematology, Hospital Saints (A. O. SS) Antonio e Biagio and Cesare Arrigo, Alessandria, Italy

²¹Department of Hematology, Azienda Socio Sanitaria Territoriale Grande Ospedale Metropolitano Niguarda, Milano, Italy

²²Hematology, Department of Oncology, AOU Careggi, Firenze, Italy

²³Hematology and Clinical Immunology Unit, Department of Medicine - DIMED, University of Padua, School of Medicine, Padova, Italy

²⁴SOC Clinica Ematologia, Azienda Sanitaria Universitaria Friuli Centrale (ASU FC), Udine, Italy

²⁵Divisione di Ematologia, Fondazione IRCCS Policlinico San Matteo, Pavia, Italy

²⁶Hematology and Stem Cell Transplantation Unit, Ospedale Oncologico Armando Businco, ARNAS G. Brotzu, Cagliari, Italy

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2024 The Author(s). *HemaSphere* published by John Wiley & Sons Ltd on behalf of European Hematology Association.

Chronic lymphocytic leukemia (CLL) therapy has recently undergone a revolution with the introduction of a new class of drugs: covalent Bruton's tyrosine kinase inhibitors (cBTKi), paving the way for a chemotherapy-free approach.¹⁻⁴ Presently, cBTKi can be utilized in the first line of CLL management, thanks to the results of phase III clinical trials such as RESONATE-2 and ELEVATE-TN, which demonstrated the superiority of Ibrutinib over chemotherapy with chlorambucil⁵ and acalabrutinib over chemoimmunotherapy with chlorambucil + obinutuzumab,⁶ in terms of progression-free survival (PFS) in both cases. Ibrutinib exhibited better PFS, overall survival (OS), and overall response rate than the monoclonal anti-CD20 antibody ofatumumab in previously treated patients with CLL.⁷ Additionally, the ASCEND study, another phase III randomized clinical trial, demonstrated that acalabrutinib significantly improved PFS compared to a physician's choice of Idelalisib + rituximab or bendamustine + rituximab, in patients with relapsed/refractory CLL.⁸

BTK plays a pivotal role in B-cell receptor (BCR) signal transduction,⁹ stimulating important pathways such as NFKB^{10,11} and CXCR4.¹² Consequently, BTK is involved in B-cell survival, proliferation, and adhesion, while its activation promotes B-cell proliferation.¹³ Paradoxically, ibrutinib has shown to increase absolute lymphocyte count (ALC) in the initial phase of treatment, regardless of previous lines of therapy. Ibrutinib-induced lymphocytosis may be explained by the redistribution of lymphocytes from neoplastic nodal compartments into the peripheral blood.¹⁴ Furthermore, it was noted that Ibrutinib-induced lymphocytosis is transient in most patients, resolving within 8 months, but may rarely persist for over 12 months without impacting survival.¹⁴ This evidence led to the introduction of a new criterion in the assessment of CLL therapy response: partial response with lymphocytosis (PR-L).¹⁵ Subsequently, the kinetics of lymphocytosis in CLL treated with ibrutinib monotherapy showed that lymphocytosis occurred in the majority of patients treated in first line was higher in immunoglobulin variable heavy chain (IGHV) mutated settings and resolved in 95% of patients after a median of 18.4 months.¹⁶

Little is known about frequency and duration of lymphocytosis in patients treated with the second-generation cBTKi acalabrutinib. Therefore, the aim of this study is to outline the kinetics of lymphocytosis in CLL patients treated with acalabrutinib compared to ibrutinib.

We conducted a multicenter retrospective real-life study involving 17 Italian centers. The study was carried out according to the Helsinki Declaration, Good Clinical Practice, and the applicable national regulations and approved by the local ethical committee. All patients provided written informed consent. The primary endpoint was to define the kinetics of lymphocytosis in naïve patients treated with acalabrutinib monotherapy compared to those treated with ibrutinib, over a 12-month observational period. We included patients receiving therapy with ibrutinib or acalabrutinib, in the first line at the target dose of 420 mg/day for ibrutinib and 200 mg/day for acalabrutinib.

We enrolled 204 patients divided into two arms: the ibrutinib arm ($n = 136$) and the acalabrutinib arm ($n = 68$). The median age was 73 years for the ibrutinib arm and 71 for the acalabrutinib arm. For each patient, we defined the clinical and biological features of disease at baseline, including IGHV mutational status, chromosomal abnormalities by FISH, and molecular biology mutations. At baseline, we considered stage, lymph node involvement, and the presence of splenomegaly. Clinical characteristics and molecular features are reported in Table 1.

Subsequently, we assessed the ALC through serial blood count tests at baseline and at different time points: 2 weeks, 1 month, 2 months, 3 months, 6 months, 9 months, and 12 months after the start of treatment. We calculated the median ALC, expressed both in

terms of cells/mm³ and as a percentage compared to the baseline, at each time point. For statistical analysis, we used the Mann-Whitney test to compare median ALC values, considering only those with a p -value < 0.05 as statistically significant.

We observed that in the ibrutinib group (IBR), the median ALC at baseline was 63,270/mm³, while in the acalabrutinib group (ACALA), it was 82,905/mm³. Median ALC peaked at two weeks in both arms and then immediately began to decline, reaching the baseline level at 1 month. From Month 1 to Month 12 ALC steadily declined, reaching a normal lymphocyte count (<4000/mm³, according to iwCLL guidelines) at Month 12 for ACALA. ACALA exhibited lower ALC from Month 6 to Month 12 compared to IBR. At Month 12, IBR did not

TABLE 1 Clinical and molecular features of patients treated with ibrutinib (IBR) or acalabrutinib (ACALA).

Clinical and molecular features		IBR (136) N (%)	ACALA (68) N (%)	p-Value
Age	Median years	73	71	0.593
Gender	Female	54 (40)	23 (34)	0.413
	Male	82 (60)	45 (66)	
Stage	Low	79 (58)	32 (47)	0.136
	High (III-IV/C)	57 (42)	36 (53)	
Binet	A	20 (15)	3 (4)	0.040
	B	63 (46)	29 (43)	
	C	53 (39)	36 (53)	
Lymphonode burden	No nodes	11 (8)	3 (4)	0.340
	<5	80 (59)	38 (56)	
	5-10	27 (20)	19 (28)	
	>10	18 (13)	8 (12)	
Lymphonodes	Absent	11 (8)	3 (4)	0.327
	Present	125 (92)	65 (96)	
Lymphonodes	Up to 5 cm	91 (67)	41 (60)	0.351
	Over 5 cm	45 (33)	27 (40)	
Lymphonodes	Up to 10 cm	118 (87)	60 (88)	0.766
	Over 10 cm	18 (13)	8 (12)	
Splenomegaly	No	47 (35)	19 (28)	0.340
	Yes	89 (65)	49 (72)	
Del17p	No	92 (69)	58 (91)	0.0007
	Yes	42 (31)	6 (9)	
Del11q	No	128 (96)	52 (80)	0.0005
	Yes	6 (4)	13 (20)	
Del13q	No	88 (66)	47 (73)	0.272
	Yes	46 (34)	17 (27)	
Trisomy 12	No	111 (83)	51 (80)	0.591
	Yes	23 (17)	13 (20)	
TP53	Unmutated	73 (55)	54 (89)	<0.001
	Mutated	59 (45)	7 (11)	
NOTCH1	Unmutated	81 (93)	32 (80)	0.028
	Mutated	6 (7)	8 (20)	
IGHV	Mutated	44 (41)	21 (35)	0.464
	Unmutated	64 (59)	39 (65)	

reach the median normal lymphocyte count. For each data point, we calculated the percentage compared to the baseline, which provides a more representative view of lymphocyte count changes at each time point of the study (Supporting Information S1: Table S1).

Subsequently, we examined the decline in lymphocytosis to understand if well-defined clinical or biological features could independently impact the kinetics of lymphocyte count in both arms. ACALA had less patients with a molecularly unfavorable prognostic profile; despite this, it seems to have no impact on the kinetics of lymphocytosis in both arms according to our subanalysis (data not shown). IGHV mutational status did not differ between the two arms, but IGHV mutational status had an impact on the lymphocytosis decline. We considered separately cases with unmutated and mutated IGHV genes for IBR and ACALA. They exhibited a similar increment at Day 14 followed by a steady decline. Unmutated IGHV curves of IBR and ACALA overlapped until Month 12. Mutated IGHV for IBR and ACALA followed a similar pattern until Month 6, after which ACALA exhibited a stronger reduction, albeit only as a percentage of baseline, due to ACALA starting from a higher baseline level (Figure 1, Supporting Information S1: Tables S2 and S3). Because there were more cases with TP53 mutations (del17p or TP53 mutated), we evaluated whether the TP53 mutational status had an impact on ALC kinetics within the subgroups. As shown in Supporting Information S1: Table S4, there was no difference in ALC kinetics

according to TP53 mutational status among the IGHV mutated and unmutated cases in the IBR arm.

Regarding clinical features and the burden of disease at baseline, the two arms were homogeneous with only one exception: more patients treated with acalabrutinib (53%) had Binet staging C before therapy, while only (39%) of patients treated with ibrutinib had stage C. Despite this difference being statistically significant ($p = 0.04$), our analysis demonstrated that the stage had no impact on the kinetics of lymphocytosis (data not shown). Even with different grades of lymphadenopathy and splenomegaly, including the presence of bulky masses, the trend of lymphocytosis during treatment with both cBTKi remained unaltered (data not shown).

We have provided the first description of the kinetics of lymphocytosis in patients treated with acalabrutinib and conducted the first comparative study of lymphocyte counts during 12 months of treatment with two cBTK inhibitors. We observed that, similar to ibrutinib, acalabrutinib leads to an increase in lymphocyte count immediately after starting therapy. This observation confirms previous works identifying lymphocytosis as a "class effect" of cBTKi.¹⁴

We observed that while there is an appreciable reduction in lymphocyte count starting from the second month of treatment in both groups, the decrease in lymphocytosis appears to be faster and more profound in patients treated with acalabrutinib from the 6th month onward.




FIGURE 1 Description of lymphocytosis in IGHV-mutated and unmutated patients in the ibrutinib group (IBR) and acalabrutinib (ACALA) treatment arms. In both arms, the two curves followed a similar trend up to Month 6. However, the ACALA arm exhibited a sharper and deeper decline compared to IBR in the mutated group. This difference was observed solely in the median percentage of baseline, attributable to ACALA starting from higher baseline values.

We also investigated whether certain disease's features could explain the differences in the kinetics of lymphocytosis. There was a significant difference in the distribution of patients with a poor molecular prognosis (del17p/TP53) between the two groups, with a marked prevalence in the IBR arm. This difference can be attributed to the evolution of cBTKi use in clinical practice: when ibrutinib was introduced, it was only allowed for patients with a poor prognosis. Conversely, acalabrutinib was used in clinical practice when the administration of cBTKi was already well established in all untreated patients. However, according to our data, having a del17p/TP53 mutation or NOTCH1 mutation does not interfere with the kinetics of lymphocytosis during treatment with the two different cBTKi. NOTCH1 mutations and higher CD49D expression have been associated with reduced ibrutinib-induced lymphocytosis,^{17,18} but in this cohort, we did not observe any effect. The IGHV mutational status did not differ between the two groups; however, when examining lymphocyte count curves in mutated IGHV, starting from the 6th month to the end of the study period, the median percentage of baseline declined more in the ACALA arm reaching a statistical difference. Overall, IGHV mutated/unmutated patients treated with acalabrutinib had a similar increase in lymphocyte count after 14 days, followed by a sharper decline, achieving median normal lymphocyte count earlier than the IBR arm. These data suggest that the main differences of lymphocytes count between the two cBTKi is due to IGHV mutated status. The clinical burden of disease at baseline had no impact on the kinetics of lymphocytosis between the two arms.

Currently, there is lack of data in the literature comparing the differences between ACALA and IBR concerning lymphocytosis, IGHV mutational status, and the extent of lymphocyte mobilization in peripheral blood. Particularly, there is scarcity of mechanistic explanations for the divergent behavior of the two cBTK inhibitors. In order to verify this hypothesis, our group is conducting research into chemokine receptors and adhesion molecules.

In conclusion, we describe the kinetics of peripheral blood lymphocytosis after ibrutinib or acalabrutinib in patients with CLL treated in front line. Both treatments exhibited a similar peak at 2 weeks after initiation, followed by a more pronounced and rapid decrease in the acalabrutinib treatment group, particularly in IGHV mutated cases.

ACKNOWLEDGMENTS

MoH ricerca corrente 2024.

AUTHOR CONTRIBUTIONS

Antonio Mosca, Annamaria Tomasso, Andrea Galitzia, Luca Stirparo, Francesca Martini, Francesca Morelli, Roberta Laureana, Giulia Benintende, Veronica Mattiello, Sabrina Chiriu, Maria I. Del Principe, Giulia Zamproga, Massimo Gentile, Enrica A. Martino, Emilia Cappello, Maria C. Montalbano, Giuliana Farina, Vanessa Innao, Lydia Scarfò, Caterina Patti, Paolo Sportoletti, Alberto Fresa, Gioacchino Catania, Marta Coscia, Silvia Bellesi, Alessandra Tedeschi, Alessandro Sanna, Andrea Visentin, Francesco Autore, Raffaella Pasquale, Livio Trentin, Marzia Varettoni, Paolo Ghia, and Roberta Murru collected the data. Eugenio Galli performed the statistical analysis. Idanna Innocenti and Antonio Mosca wrote the manuscript and Eugenio Sangiorgi and Luca Laurenti supervised the project.

CONFLICT OF INTEREST STATEMENT

Paolo Ghia received honoraria from AbbVie, AstraZeneca, BeiGene, BMS, Galapagos, Janssen, Lilly/LoxoOncology, MSD, and Roche, and research funding from AbbVie, AstraZeneca, BMS, and Janssen, and is an Editor of HemaSphere. Marzia Varettoni received honoraria from AbbVie, AstraZeneca, BeiGene, and Janssen. The remaining authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

FUNDING

This research received no external funding.

ETHICS STATEMENT

This study was conducted in accordance with the Declaration of Helsinki and approved by the Institutional Review Board of Fondazione Policlinico Universitario Agostino Gemelli IRCCS (Prot. ID 5765 13/10/2023). Written informed consent was obtained from the patients to publish this study.

ORCID

Annamaria Tomasso  <http://orcid.org/0009-0005-5361-7463>

Paolo Ghia  <http://orcid.org/0000-0003-3750-7342>

SUPPORTING INFORMATION

Additional supporting information can be found in the online version of this article.

REFERENCES

- Ahn IE, Brown JR. Targeting Bruton's Tyrosine Kinase in CLL. *Front Immunol.* 2021;12:12687458.
- Hutchinson CV, Dyer MJS. Breaking good: the inexorable rise of BTK inhibitors in the treatment of chronic lymphocytic leukaemia. *Br J Haematol.* 2014;166(1):12-22.
- Innocenti I, Fresa A, Tomasso A, et al. Treatment sequencing and outcome of chronic lymphocytic leukemia patients treated at Fondazione Policlinico Universitario Agostino Gemelli IRCCS: a thirty-year single-center experience. *Cancers.* 2023;15(23):5592.
- Martino EA, Mauro FR, Reda G, et al. Ibrutinib as first line therapy in chronic lymphocytic leukemia patients over 80 years old: a retrospective real-life multicenter Italian cohort. *Hematol Oncol.* 2024;42(1):e3249.
- Burger JA, Tedeschi A, Barr PM, et al. Ibrutinib as initial therapy for patients with chronic lymphocytic leukemia. *N Engl J Med.* 2015;373(25):2425-2437.
- Sharman JP, Egyed M, Jurczak W, et al. Acalabrutinib with or without obinutuzumab versus chlorambucil and obinutuzumab for treatment-naive chronic lymphocytic leukaemia (ELEVATE-TN): a randomised, controlled, phase 3 trial. *Lancet.* 2020;395(10232):1278-1291.
- Byrd JC, Brown JR, O'Brien S, et al. Ibrutinib versus ofatumumab in previously treated chronic lymphoid leukemia. *N Engl J Med.* 2014;371(3):213-223.
- Ghia P, Pluta A, Wach M, et al. ASCEND: phase III, randomized trial of acalabrutinib versus idelalisib plus rituximab or bendamustine plus rituximab in relapsed or refractory chronic lymphocytic leukemia. *J Clin Oncol.* 2020;38(25):2849-2861.
- Efremov DG, Turkalj S, Laurenti L. Mechanisms of B cell receptor activation and responses to B cell receptor inhibitors in B cell malignancies. *Cancers.* 2020;12(6):1396.
- Zanesi N, Balatti V, Riordan J, et al. A sleeping beauty screen reveals NF- κ B activation in CLL mouse model. *Blood.* 2013;121(21):4355-4358.
- Doménech E, Gómez-López G, Gzlez-Peña D, et al. New mutations in chronic lymphocytic leukemia identified by target enrichment and deep sequencing. *PLoS One.* 2012;7(6):e38158.
- Vlad A, Deglesne P-A, Letestu R, et al. Down-regulation of CXCR4 and CD62L in chronic lymphocytic leukemia cells is triggered by B-cell receptor ligation and associated with progressive disease. *Cancer Res.* 2009;69(16):6387-6395.

13. Deglesne P-A, Chevallier N, Letestu R, et al. Survival response to B-cell receptor ligation is restricted to progressive chronic lymphocytic leukemia cells irrespective of Zap70 expression. *Cancer Res.* 2006;66(14):7158-7166.
14. Woyach JA, Smucker K, Smith LL, et al. Prolonged lymphocytosis during ibrutinib therapy is associated with distinct molecular characteristics and does not indicate a suboptimal response to therapy. *Blood.* 2014;123(12):1810-1817.
15. Hallek M, Cheson BD, Catovsky D, et al. iwCLL guidelines for diagnosis, indications for treatment, response assessment, and supportive management of CLL. *Blood.* 2018;131(25):2745-2760.
16. Barrientos JC, Burger JA, Byrd JC, et al. Characterizing the kinetics of lymphocytosis in patients with chronic lymphocytic leukemia treated with single-agent ibrutinib. *Leuk Lymphoma.* 2019;60(4):1000-1005.
17. Giovanni Del Poeta P, Annalisa Biagi B, Luca Laurenti L, et al. Impaired nodal shrinkage and apoptosis define the independent adverse outcome of NOTCH1 mutated patients under ibrutinib therapy in chronic lymphocytic leukaemia. *Haematologica.* 2021;106(9):2345-2353.
18. Tissino E, Benedetti D, Herman SEM, et al. Functional and clinical relevance of VLA-4 (CD49d/CD29) in ibrutinib-treated chronic lymphocytic leukemia. *J Exp Med.* 2018;215(2):681-697.