



Review Article

Prunus spp. fruits as practical natural colouring agents in foods

Dasha Mihaylova¹✉, Aneta Popova², Sashka Savchovska³, Anna Lante⁴, Maria Dimitrova-Dimova²

¹Department of Biotechnology, Technological Faculty, University of Food Technologies, Plovdiv, Bulgaria

²Department of Catering and nutrition, Economics Faculty, University of food technology, Plovdiv, Bulgaria

³Department of Breeding and Genetic Resources, Fruit Growing Institute, Agricultural Academy, Plovdiv, Bulgaria

⁴Department of Agronomy Food Natural resources Animal and Environment (DAFNAE). University of Padova. Legnaro (Padova). Italy

Abstract

Fruits and vegetables are among the most important phytochemical sources worldwide, which can also provide an anticipated food colour. The sensory perception of the food is directly linked to its colour. Consumers have a specific expectation for the colour of food i.e. desired and vivid ones. Thus, colour additives, both natural and synthetic, are so important to the food industry. Colour additives are able to enhance the sensory attributes lost during processing, and to expand product variety. However, studies report a link between synthetic colours and some health disorders. That is why serious attention is paid to natural dyes. In this context, Prunus fruits present an excellent alternative source of natural compounds that enable the production of a wide range of colouring molecules, such as anthocyanins, carotenoids and chlorophylls. Moreover, in addition to their ability to colour, they also contribute by increasing food's bioactive qualities. Carotenoids are the pigments that give the yellow and orange colour to the pulp and skin of apricots and other Prunus fruits. Along with them, flavonoids, anthocyanins (purple, blue colours), etc. are also found in prunes, for example. Still, whether the fruit can be used as a reliable source of food colouring is an open question worth answering. Therefore, more research is currently needed to better understand the behaviour of natural compounds during the extraction processes and further incorporation into food matrices. In this regard, scientists add fruits directly in the food in order to avoid extraction and potential loss of bioactive compounds, i.e. natural dyes. Regulation approvals should always be considered as well and it has to be noted that different legislation applies to different countries. In view of the sustainable food cycle provision and the seek for more value added ingredients, natural dyes ought to be taken into account with priority.

✉Corresponding author: Assoc. Prof. Dasha Mihaylova, PhD, Department of Biotechnology, Technological Faculty, University of Food Technologies, 26 Maritza Blvd., 4002 Plovdiv, Bulgaria; E-mail: dashamihaylova@yahoo.com

Article history:

Received

Reviewed

Accepted

Available on-line

<https://doi.org/10.30721/fsab202x.xx.xx.xx>

© 202x The Authors. UFT Academic publishing house, Plovdiv



Food Science and Applied Biotechnology

e-ISSN: 2603-3380

Journal home page: www.ijfsab.com
<https://doi.org/10.30721/fsab202x.xx.xx>



Keywords

value added ingredients, natural sources, bioactive compounds

Abbreviations

EFSA - European Food and Safety Authority; FDA - Food and Drug Administration

✉ Corresponding author: Assoc. Prof. Dasha Mihaylova, PhD, Department of Biotechnology, Technological Faculty, University of Food Technologies, 26 Maritza Blvd., 4002 Plovdiv, Bulgaria; E-mail: dashamihaylova@yahoo.com

Article history:

Received

Reviewed

Accepted

Available on-line

<https://doi.org/10.30721/fsab202x.xx.xx.xx>

© 202x The Authors. UFT Academic publishing house, Plovdiv

Introduction


Consumers use colour to assess the quality of products. The colour is associated with not only freshness, maturity state, and safety but also expected taste, especially when food colorants are being used (Figueiredo et al., 2023). For example, the orange colour is constantly linked to the taste of the orange fruit.




Synthetic colours have been long used in various industries but their harmful effects (inability to degrade, health issues, among others) exceed their practical use (Miller et al., 2022). Natural colorants, on the other hand, can not only contribute to the physical appearance of products but also carry important health benefits due to their chemical structure (Shakoor et al., 2022). Colour molecules can be extracted from roots, flowers,

leaves, or the whole raw material. Natural colorants belong to chemical structures like anthocyanins, chlorophylls, carotenoids, betalains, phenolic compounds, among others (Singh et al., 2023). Countries with better biodiversity have greater possibilities to produce dyes. Food safety authorities like the EFSA or the FDA are the ones that monitor, regulate, allow or suspend the use of both synthetic and natural dyes.

The genus *Prunus* is known worldwide and can be evaluated as a potential source for natural dyes (Table 1). The genus comprises of some of the well accepted and cherished fruits like cherries, peaches, sour cherries, plums, apricots, and almonds (Popova et al., 2022).

Table 1. *Prunus* spp. representatives and their dyeing compounds

Representative	Chemical class	Compound	Extraction solvents and techniques	Potential beneficial properties	References
 <i>Prunus persica</i> L.	Carotenoids flavonoids anthocyanidins	α -carotene, β -carotene, β -cryptoxanthin, lycopene, lutein, and zeaxanthin	hexane, petroleum ether, acetone, ethanol, methanol; liquid-liquid extraction, solid-liquid extraction, Soxhlet extraction, ultrasonic extraction	antioxidant propensity, various anti-inflammatory actions, cardiovascular health, cancer and neurodegenerative protection	Cheng et al., 2020 Molina et al., 2023

	<p>carotenoids flavonoids chlorophylls</p>	<p>Lutein; α-carotene, β-carotene, γ-carotene, lycopene, and cryptoxanthin; β-cryptoxanthin, phytoene, phytofluene and zeaxanthin quercetin, kaempferol and rutin Chlorophyll a and b</p>	<p>acetone; n-hexane: acetone; heat reflux extraction</p>	<p>antioxidant activity; degenerative diseases protection; cardiovascular health</p>	<p>Ali et al., 2015 Makrygiann et al, 2022</p>	
	<p>anthocyanin carotenoids</p>	<p>cyanidin-3-rutinoside, cyanidin-3-glucoside and peonidin-3-rutinoside</p>	<p>methanol, ethanol, acidified water; thermo stated cell extraction</p>	<p>preventing inflammation, hypertension control, thrombotic risk management, and cancer prevention</p>	<p>Zbrzeźniak et al., 2015 Hn et al., 2020 Liao et al., 2023</p>	
	<p>carotenoids flavonoids anthocyanins</p>	<p>Lutein; Zeaxanthin; α-Carotene; β-Carotene; quercetin-3-O-rutinoside; kaempferol; cyanidin-3-O-rutinoside</p>	<p>acetone; ethanol; methanol; UAE</p>	<p>modulation of the antioxidant response; protect against oxidative damage</p>	<p>Średnicka-Tober et al., 2019</p>	
<p>Images</p>	<p>are</p>	<p>from</p>	<p>freepik.com</p>	<p>-</p>	<p>public</p>	<p>domain</p>

These representatives have been studied for their potential bioactivity and health-promoting properties. Some of the reported compounds include polyphenols, carotenoids, phenolic acids, carbohydrates, among others (Nowicka et al., 2023).

The object of this review is to synthesize the availability of natural colouring compounds as well as to make reference to the potential health benefits as well as to present *Prunus* spp. as possible natural dyes. In addition, some future perspectives in view of the trending research topics will be provided.

Natural Colouring Compounds

The colouring compounds present in *Prunus* spp. fruits are not exclusive to the genus. Some of the most common natural colorants in fruits, in general, are carotenoids, flavonoids, anthocyanidins (Lu et al. 2021).

Flavonoids can be divided into flavanols, flavanols, flavanones, flavones, isoflavones, and anthocyanins (Akhlaghi & Foshati, 2017). Flavonoids possess various biological activities (Liga et al., 2023). The colour of flavonoid pigments varies from white-cream to orange-red

and purple-blue (Miao et al., 2022). Anthocyanins are glycosylated forms of aglycones, and a class of water-soluble flavonoids (Mattioli et al., 2020). They can exhibit red, blue and purple colours. Their colour is pH dependent, meaning that in an acidic pH ($\text{pH} < 3$) cyanidin will be red to orange, while in $\text{pH} > 11$ it will appear blue-green. Appropriate extraction techniques should be used in order to enhance their stability and application (Guo et al., 2022). Different approaches have been used in order to recover anthocyanin pigments (Tan et al., 2022). Some of the frequently used extraction methods include ultrasonic assisted extraction, microwave assisted extraction, supercritical carbon dioxide extraction and combined methods (Tan et al., 2022). Alcohols like methanol and ethanol, along with acidified water or ethanol are some of the most common solvents for the anthocyanin recovery (Muangrat et al., 2017). Anthocyanins are vacuolar pigments and are also reported for their antioxidant properties (Tena & Asuero, 2020). They can be found in various representatives of the genus *Prunus* (Mollaamin et al., 2022; Liao et al., 2023). Pelargonidin and cyanidin 3-glucoside have been reported in cherry jams (Swami et al., 2020). The brownish coloration of prune juice is a result of the reaction between phenolic compounds and anthocyanins (Swami et al., 2020). Carotenoids can be responsible for yellow, orange, and red colours (González-Peña et al., 2023). They are classified into carotenes and xanthophylls (Schawartz et al., 2017). Research showed that apricots accumulate more carotenoids compared to other *Prunus* representatives (Han et al., 2020). A total of twelve carotenoids were detected in *Prunus* species, *i.e.* violaxanthin, luteoxanthin, antheroxanthin, lutein, zeaxanthin, β -cryptoxanthin, β -carotene, among others (Han et al., 2020). Carotenoids extract well in polar solvents (Saini et al., 2022). Oxygen concentration, pH-values, light and temperature are some of the most important parameters for the effective carotenoid pigment extraction (Flieger et al., 2018). Additionally mechanical treatment for the disruption of the thick cell walls and appropriate solvents are recommended for better pigment recovery (Gu et al., 2008). Oxidation is the primary cause for the loss of carotenoids (Ribeiro et al., 2018). Chlorophylls are green oil-soluble pigments (Ebrahimi et al., 2023). Their stability is

dependent on pH fluctuation, temperature, and storage duration (Kwartiningsih et al., 2021). Some of the reported effective extraction techniques include enzyme-assisted extraction, supercritical fluid CO_2 , methanol extraction (Lu et al., 2014). The bioactivity of chlorophylls has been confirmed by researchers (Martins et al., 2023).

Applications

Many branches of the food industry benefit from both natural and synthetic colorants. The colour of ice cream and yogurt is due to colorants. If natural ones are considered research shows that betalains can be used (Lis & Bartuzi, 2023). Anthocyanins can be found as pigments in beverages, desserts, ice cream, and dairy products (de Mejia et al., 2020). Beverages are an example of food products that most other need colorants. The popular “vitamin waters” are most often coloured. Bakery products are other examples of the existence of food dyes. The colour of bakery products is linked to the Maillard reaction but the presence of another colour can contribute to the sensory profile in terms of not only colour and appearance but also aroma (Luzardo-Ocampo et al., 2021). There are examples of the microencapsulation techniques for the colouring of jelly candies (Dewi et al., 2018). Confectionary is a sector of the food industry with high demand for colouring agents (Suryanarayana et al., 2017). Yogurt and flavoured milks are coloured due to the expectations of the consumer. For example, a strawberry yogurt should be pink, and chocolate milk should be brown. Cheeses also take advantage of natural colorants (Sharma et al., 2020). Furthermore, meat products are a niche for natural colorants not only because of potential colour loss, during processing, but also because of the antioxidant properties of natural dyes like carotenoids (Hamdi et al., 2018). Lastly, coloured pasta has recently gained popularity (Porto Dalla Costa et al., 2016). It is often green, orange and red with reference to tomato, carrot and spinach.

Conclusions and Future Perspectives

Natural pigments in fruit are not only powerful antioxidants but also bear potential health benefits. Living in the era of synthetic pollution, the seek for natural ingredients has become even more valuable. Natural pigments are renewable, non-toxic but their bio accessibility and availability is

still to be thoroughly researched. Colour will always be one of the most important attributes of food. Contemporary research design is turning to replacing synthetic dyes with natural as much as possible, because of their harmful nature. The spread of bio dyes can potentially expand biodiversity and thriving availability of native cultivars. The use of fruit waste for colouring can contribute to the sustainability cycle and zero-waste policy. However, research is looking into the possibility of allergies due to natural dyes interactions in the body (Lis & Bartuzi, 2023). Additionally, the stability of the colour is sometimes challenging.

Natural pigments should be carefully regulated with safe dosages so that they can act as an

References

- Akhlaghi M., Foshati S. Bioavailability and metabolism of flavonoids: A Review. *International Journal of Nutrition Sciences*. 2017, 2(4):180-184.
- Ali S., Masud T., Abbasi K., Mahmood T., Hussain A. Apricot: nutritional potentials and health benefits-A review. *Annals. Food Science and Technology*. 2015, 16(1): 175-189.
- Cheng S.-H., Khoo H., Kong K., Prasad K., Galanaki, C. Extraction of carotenoids and applications. In *Carotenoids: Properties, Processing and Applications*; Elsevier: Amsterdam, The Netherlands, 2020; pp. 259-288. ISBN 9780128170670
- De Mejia E., Zhang Q., Penta K., Eroglu A., Lila M. The colors of health: Chemistry, bioactivity, and market demand for colorful foods and natural food sources of colorants. *Annual review of food science and technology*. 2020, 11: 145-182. <https://doi.org/10.1146/annurev-food-032519-051729>
- Dewi E., Kurniasih R., Purnamayati L. The application of microencapsulated phycocyanin as a blue natural colorant to the quality of jelly candy. *IOP Conference Series: Earth and Environmental Sciences*. 2018, 116: 012047. <https://doi.org/10.1088/1755-1315/116/1/012047>
- Ebrahimi P., Shokramraji Z., Tavakkoli S., Mihaylova D., Lante A. Chlorophylls as natural bioactive compounds existing in food by-products: A critical review. *Plants (Basel, Switzerland)*. 2023, 12: 1533. <https://doi.org/10.3390/plants12071533>
- Figueiredo Muniz V., Ribeiro I., Beckmam K., Godoy R. The impact of color on food choice. *Brazilian Journal of Food Technology*. 2023, 26: e2022088. <https://doi.org/10.1590/1981-6723.08822>
- Flieger K., Knabe N., Toepel J. Development of an Improved Carotenoid Extraction Method to Characterize the Carotenoid Composition under Oxidative Stress and Cold Temperature in the Rock Inhabiting Fungus *Knufia petricola* A95. *J Fungi (Basel)*. 2018, 4(4): 124. <https://doi.org/10.3390/jof4040124>
- González-Peña M., Ortega-Regules A., Anaya de Parrodi C., Lozada-Ramírez J. Chemistry, occurrence, properties, applications, and encapsulation of carotenoids - A Review. *Plants (Basel, Switzerland)*. 2023, 12: 313. <https://doi.org/10.3390/plants12020313>
- Gu Z., Deming C., Yongbin H., Zhigang C., Feirong G. Optimization of carotenoids extraction from *Rhodobacter sphaeroides*. *LWT-Food Science and Technology*. 2008, 41: 1082-1088. <https://doi.org/10.1016/j.lwt.2007.07.005>
- Guo Y., Zhang H., Shao S., Sun S., Yang D., Lv S. Anthocyanin: a review of plant sources, extraction, stability, content determination and modifications. *International Journal of Food Science & Technology*. 2022, 57: 7573-7591. <https://doi.org/10.1111/ijfs.16132>
- Hamdi M., Nasri R., Dridi N., Moussa H., Ashour L., Nasri M. Improvement of the quality and the shelf life of reduced-nitrites turkey meat sausages incorporated with carotenoproteins from blue crabs shells. *Food Control*. 2018, 91: 148-159. <https://doi.org/10.1016/j.foodcont.2018.03.048>
- Han Y., Wang P., Hyden B., Qu P., Liu B., Zhang F., Cao H., Chen H. Diversity of carotenoid composition, sequestering structures and gene transcription in mature fruits of four *Prunus* species. *Plant Physiology and Biochemistry*. 2020, 151: 113-123. <https://doi.org/10.1016/j.plaphy.2020.03.015>

- Kwartiningsih E., Ramadhani A., Damara N. Chlorophyll extraction methods review and chlorophyll stability of Katuk leaves (*Sauropus androgynous*). *Journal of Physics: Conference Series*. 2021, 1858: 012015. <https://doi.org/10.1088/1742-6596/1858/1/012015>
- Liao L, Li Y, Lan X, Yang Y, Wei W, Ai J, Feng X, Chen H, Tang Y, Xi L, Wang Z. Integrative analysis of fruit quality and anthocyanin accumulation of Plum cv. 'Cuihongli' (*Prunus salicina* Lindl.) and its bud mutation. *Plants (Basel, Switzerland)*. 2023, 12(6): 1357. <https://doi.org/10.3390/plants12061357>
- Liga S., Paul C., Péter F. Flavonoids: overview of biosynthesis, biological activity, and current extraction techniques. *Plants (Basel, Switzerland)*. 2023, 12: 2732. <https://doi.org/10.3390/plants12142732>
- Lis K., Bartuzi Z. Plant food dyes with antioxidant properties and allergies - Friend or enemy? *Antioxidants (Basel, Switzerland)*. 2023, 12(7): 1357. <https://doi.org/10.3390/antiox12071357>
- Lu J., Feng X., Han Y., Xue C. Optimization of subcritical fluid extraction of carotenoids and chlorophyll a from *Laminaria japonica* Aresch by response surface methodology. *Journal of the science of food and agriculture*. 2014, 94(1): 139–145. <https://doi.org/10.1002/jsfa.6224>
- Lu W., Shi Y., Wang R., Su D., Tang M., Liu Y., Li Z. Antioxidant activity and healthy benefits of natural pigments in fruits: A review. *International Journal of Molecular Sciences*, 2021, 22(9): 4945. <https://doi.org/10.3390/ijms22094945>
- Luzardo-Ocampo I., Ramírez-Jiménez A., Yañez J., Mojica L., Luna-Vital D. Technological applications of natural colorants in food systems: A Review. *Foods (Basel, Switzerland)*. 2021, 10(3): 634. <https://doi.org/10.3390/foods10030634>
- Makrygiannis I., Athanasiadis V., Bozinou E., Chatzimitakos T., Makris D., Lalas SI. An investigation into apricot pulp waste as a source of antioxidant polyphenols and carotenoid pigments. *Biomass*. 2022, 2(4): 334-347. <https://doi.org/10.3390/biomass2040022>
- Martins T., Barros A., Rosa E., Antunes L. Enhancing health benefits through chlorophylls and chlorophyll-rich agro-food: A Comprehensive review. *Molecules (Basel, Switzerland)*, 2023, 28: 5344. <https://doi.org/10.3390/molecules28145344>
- Mattioli R., Francioso A., Mosca L., Silva P. Anthocyanins: A comprehensive review of their chemical properties and health effects on cardiovascular and neurodegenerative diseases. *Molecules (Basel, Switzerland)*, 2020 25(17): 3809. <https://doi.org/10.3390/molecules25173809>
- Miao L., Zhang H., Yang L., Chen L., Xie Y., Xiao J., Part 4 - Natural occurring antioxidants: bright and the dark side, Editor(s): Seyed Mohammad Nabavi, Ana Sanches Silva, Antioxidants Effects in Health, Elsevier, 2022. p.235-236. ISBN 9780128190968
- Miller M., Steinmaus C., Golub M., Castorina R., Thilakartne R., Bradman A., Marty M. Potential impacts of synthetic food dyes on activity and attention in children: a review of the human and animal evidence. *Environmental health : a global access science source*, 2022, 21(1): 45. <https://doi.org/10.1186/s12940-022-00849-9>
- Molina A., Corrêa R., Prieto M., Pereira C., Barros L. Bioactive Natural Pigments' Extraction, Isolation, and Stability in Food Applications. *Molecules (Basel, Switzerland)*. 2023, 28: 1200. <https://doi.org/10.3390/molecules28031200>
- Mollaamin F., Kandemirli F., Monajjemi M. Anthocyanin compounds of the Persian sour cherry (*Prunus cerasus* L) as the fruit with high antioxidant properties. *Biointerface research and applied chemistry*. 2022, 12,(3): 3696-3709. <https://doi.org/10.33263/BRIAC123.36963709>
- Muangrat R., Williams P., Saengcharoenrat P. Subcritical solvent extraction of total anthocyanins from dried purple waxy corn: influence of process conditions *Journal of Food Processing and Preservation*. 2017, 41: 46-50. <https://doi.org/10.1111/jfpp.13252>
- Nowicka P., Wojdyło A., Tkacz K., Turkiewicz I. Quantitative and qualitative determination of carotenoids and polyphenolics compounds in selected cultivars of *Prunus persica* L. and their ability to in vitro inhibit lipoxygenase, holinesterase, α -amylase, α -glucosidase and pancreatic lipase. *Food Chemistry*. 2023, 17: 100619. <https://doi.org/10.1016/j.fochx.2023.100619>
- Popova A., Mihaylova D., Alexieva I., Doykina P. Ethnopharmacology and phytochemistry of some representatives of the genus *Prunus*. *Journal of Central European Agriculture*, 2022, 23(3):665-678. <https://doi.org/10.5513/JCEA01/22.3.3629>
- Porto Dalla Costa A., Cruz Silveira S. Thays R., De Oliveira Rios A., Hickmann Flôres S. Carrot flour from minimally processed residue as substitute of β -carotene commercial in dry pasta prepared with common wheat (*Triticum aestivum*). *Journal of Food Quality*. 2016, 39: 590-598. <https://doi.org/10.1111/jfq.12253>
- Ribeiro D., Freitas M., Silva A., Carvalho F., Fernandes E. Antioxidant and pro-oxidant activities of carotenoids and their oxidation products. *Food and chemical toxicology: an international journal published for the British Industrial Biological Research Association*, 2018, 120, 681–699. <https://doi.org/10.1016/j.fct.2018.07.060>
- Saini R., Prasad P., Lokesh V., Shang X., Shin J., Keum Y.-S., Lee J.-H. Carotenoids: dietary sources,

- extraction, encapsulation, bioavailability, and health benefits - A review of recent advancements. *Antioxidants (Basel, Switzerland)*, 2022, 11: 795. <https://doi.org/10.3390/antiox11040795>
- Schawartz S., Cooperstone J., Cichon M., von Elbe J., Giusti M. 2017. Colorants. In Fennema's Food Chemistry; Damodaran, S., Parkin, K.L., Eds.; CRC Press Taylor & Francis: Boca Raton, FL, USA, p. 681. ISBN 978-1-4822-4361-1
- Shakoor S., Ismail A., Sabran M., Mohtarrudin N., Kaka U., Nadeem M. In-vivo study of synthetic and natural food colors effect on biochemical and immunity parameters. *Food Science and Technology (Campinas)*. 2022, 42: e41420. <https://doi.org/10.1590/fst.41420>
- Sharma P., Segat A., Kelly A., Sheehan J. Colorants in cheese manufacture: Production, chemistry, interactions, and regulation. *Comprehensive Reviews in Food Science and Food Safety*. 2020, 19: 1220-1242. <https://doi.org/10.1111/1541-4337.12519>
- Singh T., Pandey V., Dash K., Zanwar S., Singh R. Natural bio-colorant and pigments: Sources and applications in food processing. *Journal of Agriculture and Food Research*. 2023, 12: 100628. <https://doi.org/10.1016/j.jafr.2023.100628>
- Średnicka-Tober D, Ponder A, Hallmann E, Głowacka A, Rozpara E. The profile and content of polyphenols and carotenoids in local and commercial sweet cherry fruits (*Prunus avium* L.) and their antioxidant activity in vitro. *Antioxidants (Basel, Switzerland)*. 2019, 8(11): 534. <https://doi.org/10.3390/antiox8110534>
- Suryanarayana R., Chandrappa M., Santhosh R. Awareness of use of artificial colourants in sweets preparation and their harmful effects. *International Journal of Community Medicine and Public Health*. 2017, 4(10): 3893–3898. <https://doi.org/10.18203/2394-6040.ijcmph20174270>
- Swami S., Ghgare S., Swami S.S., Shinde K., Kalse S., Pardeshi I. Natural pigments from plant sources: A review. *The Pharma Innovation Journal*. 2020, 9(10): 566-574.
- Tan J., Han Y., Han B., Qi X., Cai X., Ge S., Xue H. Extraction and purification of anthocyanins: A review. *Journal of Agriculture and Food Research*. 2022, 8: 100306. <https://doi.org/10.1016/j.jafr.2022.100306>
- Tena N., Martín J., Asuero A. State of the art of anthocyanins: antioxidant activity, sources, bioavailability, and therapeutic effect in human health. *Antioxidants (Basel, Switzerland)*, 2020, 9(5): 451. <https://doi.org/10.3390/antiox9050451>
- Zbrzeźniak M., Nordlund E., Mieszczakowska-Frać M., Płocharski W., Konopacka D. Quality of cloudy plum juice produced from fresh fruit of *Prunus domestica* L. - the effect of cultivar and enzyme treatment. *Journal of Horticultural Research*. 2015, 23(2): 83-94. <https://doi.org/10.2478/johr-2015-0018>