

Earth's Future



RESEARCH ARTICLE

10.1029/2021EF002397

Key Points:

- End of 21st century heavy precipitation events in the eastern Mediterranean are projected to have substantially reduced rainfall yield
- This reduction results mainly from a major decrease in rain area during future events, despite increased conditional rain rate
- Changes in rain yield, rate, and area are consistent across many events, suggesting significant hydrological implications for the region

Supporting Information:

Supporting Information may be found in the online version of this article.

Correspondence to:

M. Armon,
moshe.armon@mail.huji.ac.il

Citation:

Armon, M., Marra, F., Enzel, Y., Rostkier-Edelstein, D., Garfinkel, C. I., Adam, O., et al. (2022). Reduced rainfall in future heavy precipitation events related to contracted rain area despite increased rain rate. *Earth's Future*, 10, e2021EF002397. <https://doi.org/10.1029/2021EF002397>

Received 1 SEP 2021
Accepted 25 NOV 2021

Reduced Rainfall in Future Heavy Precipitation Events Related to Contracted Rain Area Despite Increased Rain Rate

Moshe Armon¹ , Francesco Marra² , Yehouda Enzel¹ , Dorita Rostkier-Edelstein^{1,3}, Chaim I. Garfinkel¹ , Ori Adam¹ , Uri Dayan⁴, and Efrat Morin¹ 

¹Fredy and Nadine Herrmann Institute of Earth Sciences, The Hebrew University of Jerusalem, Jerusalem, Israel, ²National Research Council of Italy, Institute of Atmospheric Sciences and Climate, CNR-ISAC, Bologna, Italy, ³Department of Environmental Physics, Environmental Sciences Division, IIBR, Ness-Ziona, Israel, ⁴Department of Geography, The Hebrew University of Jerusalem, Jerusalem, Israel

Abstract Heavy precipitation events (HPEs) can lead to deadly and costly natural disasters and are critical to the hydrological budget in regions where rainfall variability is high and water resources depend on individual storms. Thus, reliable projections of such events in the future are needed. To provide high-resolution projections under the RCP8.5 scenario for HPEs at the end of the 21st century, and to understand the changes in sub-hourly to daily rainfall patterns, weather research and forecasting (WRF) model simulations of 41 historic HPEs in the eastern Mediterranean are compared with “pseudo global warming” simulations of the same events. This paper presents the changes in rainfall patterns in future storms, decomposed into storms' mean conditional rain rate, duration, and area. A major decrease in rainfall accumulation (−30% averaged across events) is found throughout future HPEs. This decrease results from a substantial reduction of the rain area of storms (−40%) and occurs despite an increase in the mean conditional rain intensity (+15%). The duration of the HPEs decreases (−9%) in future simulations. Regionally maximal 10-min rain rates increase (+22%), whereas over most of the region, long-duration rain rates decrease. The consistency of results across events, driven by varying synoptic conditions, suggests that these changes have low sensitivity to the specific synoptic evolution during the events. Future HPEs in the eastern Mediterranean will therefore likely be drier and more spatiotemporally concentrated, with substantial implications on hydrological outcomes of storms.

Plain Language Summary Heavy precipitation events are large storms that can recharge freshwater reservoirs, but can also lead to hazardous outcomes such as flash floods. Therefore, understanding the impacts of climate change on such storms is critical. Here, a weather model similar to those used in weather forecasts is used to simulate heavy precipitation events in the eastern Mediterranean. A large collection of storms is simulated in pairs: (a) historic storms, selected for their high impact, and (b) the same storms placed in a global warming scenario projected for the end of the 21st century. Using these simulations we ask how present-day storms would look like were they to occur at the warmer end of the 21st century. The future storms are found to produce much less rainfall compared to their historic counterparts. This decrease in rainfall is attributed mainly to the reduction in the area covered by storms' rainfall, and happens despite increasing rainfall intensities. These results suggest that the region will be drier in the future with larger dry areas during storms; however, over short durations, it would rain more intensely over contracted areas—increasing local hazards associated with heavy precipitation events.

1. Introduction

Expected impacts of climate change on rainfall during heavy precipitation events (HPEs) have the potential to significantly alter their influence on future populations. Where precipitation variability is high, such as in Mediterranean and arid climates, the impact of individual HPEs in terms of both peril (e.g., Borga et al., 2014; Dayan et al., 2021; De Vries et al., 2013; Raveh-Rubin & Wernli, 2016; Rinat et al., 2021) and water resources (Flaounas et al., 2021; Nasta et al., 2018; Samuels et al., 2009; R. G. Taylor et al., 2013) is great, and reliable projections of HPEs are needed (e.g., Sillmann et al., 2021).

Individual HPEs are controlled by specific large scale and synoptic circulation patterns. However, projected changes in the atmospheric circulation are highly uncertain across global climate models (GCMs) due to the wide variety of factors at play (Shepherd, 2014). Furthermore, climate change impact on HPEs can be quite

© 2021 The Authors. Earth's Future published by Wiley Periodicals LLC on behalf of American Geophysical Union. This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs License](https://creativecommons.org/licenses/by-nc-nd/4.0/), which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

different from the well-studied impact on the mean rainfall or even on high precipitation percentiles (e.g., Donat et al., 2016; Kendon et al., 2018; Moustakis et al., 2021; O’Gorman, 2015; Pfahl et al., 2017; Trenberth et al., 2015).

Detailed projections of the regional rainfall during a specific event can only be provided by models that can explicitly resolve the convective processes governing precipitation during HPEs (e.g., Fosser et al., 2015). Indeed, convection-permitting models (CPMs) are more reliable than GCMs in simulating spatiotemporal precipitation patterns (Ban et al., 2014; Cannon & Innocenti, 2019; Crook et al., 2019; Kendon et al., 2014; Meredith et al., 2020; Poujol et al., 2020; Prein et al., 2015, 2017; Westra et al., 2014). Recent methodological and computing advances enable “climate” CPM simulations with long-term (~10 years), large-scale (continental), and high resolution (a few kilometers) outputs with some groups already running ensemble simulations over specific regions (Chan et al., 2020; Coppola et al., 2020; Pichelli et al., 2021). These give probabilistic projections of changes in precipitation extremes with expectations to achieve better quantification of future HPEs (e.g., Kendon et al., 2014; Poujol et al., 2020). However, rare extremes or HPEs are, by definition, hard to assess even with such simulations (e.g., Fatichi et al., 2016; Kendon et al., 2021). Moreover, a resolution of a few kilometers may still not be sufficient to represent the local nature of convective clouds, especially when shallow convection is present (Kendon et al., 2021; Prein et al., 2015). Therefore, trying to provide reliable projections of the changes in rainfall patterns during HPEs will probably take many more years of improvement in climate modeling. A complementing approach, aimed at resolving extreme events and intra-event characteristics (Fowler, Ali, et al., 2021; O’Gorman, 2015), is to provide projections of specific high impact events either by identifying interesting events such as hurricanes over long-term simulations (Gutmann et al., 2018), or through the simulation of individual events known for their high-impact, such as snowstorms (G. Chen, Wang, Cheng, & Hsu, 2021), tropical cyclones (J. Chen, Wang, Tam, et al., 2020), or HPEs (Ferreira, 2021).

Pseudo global warming (PGW) is an emerging methodology for event-based projections, enabling assessment of the impacts of one or more meteorological parameters over local-scale weather events (Brogli et al., 2019; Fowler, Lenderink, et al., 2021; Moustakis et al., 2021; Prein et al., 2017; Sato et al., 2007; Schär et al., 1996). The PGW methodology imposes a certain climate change, for example, temperature rise, over the initial and boundary conditions of a regional model, by prescribing the synoptic and larger-scale changes from GCMs, while allowing smaller scale features to develop freely within a downscaled modeled domain in a physically consistent manner. Further, projections of precipitation extremes under global warming scenarios commonly focus on daily resolutions (Donat et al., 2016; O’Gorman, 2015; Pfahl et al., 2017), which hinders the possible impact of short-duration extremes; only recently more studies have directed attention to changes expected over subdaily or even subhourly extremes (Fowler, Ali, et al., 2021; Fowler, Wasko, & Prein, 2021; Morrison et al., 2019). However, to understand the potential effects of changes in precipitation extremes, not only their changing intensity and frequency are important, but also high-resolution changes in intra-event characteristics, such as the spatiotemporal organization of the storms (Li et al., 2018). This requires high-resolution analysis of many high-impact storms of different synoptic-scale circulations, as there is no guarantee that different HPEs behave the same way (Fowler, Ali, et al., 2021).

The goal of this study is to identify and quantify changes in rainfall patterns during HPEs induced by global warming, and to examine whether a common change emerges over a variety of HPEs. To do so, we exploit the case of the eastern Mediterranean (Section 2.1) to simulate a large number of HPEs using the PGW methodology with a very high spatiotemporal resolution, and explicitly consider space-time patterns of rainfall during the events over durations of 10-min to 24-hr.

The paper is organized as follows: Section 2 describes the study region and outlines the modeling strategy and the analyses of rainfall patterns. We first demonstrate the expected changes for a specific HPE case (Section 3.1), and then examine changes in rainfall accumulation over a large set of HPEs (Section 3.2). Changes in specific rainfall properties are outlined in Sections 3.2–3.4, with the unique role of the rain area shown in Section 3.3.1. Section 4 begins with a discussion of the event-based approach (Section 4.1) and continues with an examination of the change in rainfall patterns in future HPEs (Section 4.2). Our conclusions are presented in Section 5.




Figure 1. Map of the study area. (a) Köppen-Geiger climate classification of the eastern Mediterranean (Atlas of Israel, 2011), weather research and forecasting (WRF) model domains (D1–D3; Section 2.3), and the range of the weather radar used for the identification of events (Section 2.2). (b) Mean annual rainfall based on 1960–1990 interpolated rain gauge data (Enzel et al., 2003), the innermost model domain, and the weather radar range. Green and yellow colors, corresponding to drier and wetter than 200 mm yr⁻¹, respectively, roughly mark the extent of the desert and Mediterranean climate regions. SND = Sinai-Negev Desert, LM = Lebanon Mountains.

2. Study Region, Data, and Methods

2.1. Study Region

The focus here is on the eastern Mediterranean (Figure 1), which (a) is expected to suffer from a large future decrease in total rainfall (Garfinkel et al., 2020; Giorgi & Lionello, 2008; Zappa et al., 2015), (b) may experience an increase in extreme precipitation occurrence (Alpert et al., 2002; Marra et al., 2021; Samuels et al., 2017), (c) is characterized by the least precipitation per capita in the world (Dirmeyer et al., 2009), and (d) is exposed to large rainfall variability (Morin, 2011). These characteristics result in a large dependency on HPEs, in terms of water resources and vulnerability to natural hazards; therefore, we explore here possible future changes in HPEs in the region, and disassemble them to their distinct hydrometeorological constituents. It is important to note there is currently no CPM with future projections available for the study region.

In the eastern Mediterranean (Figure 1a), the Mediterranean climate abuts the semiarid to hyperarid climates characterizing the region to the south and east of the Mediterranean Sea. Yearly rainfall amounts drop from >1000 mm in the northern mountains, to <<100 mm at the southeast regions (Figure 1b). Summers are dry, and the rainy season is from October to May, with a few rare exceptions in September and June (Goldreich, 2012; Kushnir et al., 2017). The core of the rainy season is December–February (>65% of precipitation). However, the rainy season's midpoint changes from the beginning of January near the Mediterranean Sea to the end of January farther inland (Goldreich, 1994, 1995). This reflects the important contribution of the warm Mediterranean Sea water to building up of Mediterranean Cyclones (MCs), the favorable synoptic condition prevailing during rainy days, generating >90% of all rainfall in the northern, wetter part of the region (Alpert & Shay-EL, 1994; El-Fandy, 1946; Ziv et al., 2006). Other synoptic systems contribute relatively large rain amounts to the interior-desert

area mainly during the transitional seasons (Armon et al., 2019; Dayan & Morin, 2006; Kahana et al., 2002), including the more frequent (a) active Red Sea troughs (ARSTs) (Ashbel, 1938; De Vries et al., 2013), occurring mainly in fall, and (b) less frequent disturbances in the Subtropical Jet sometimes termed Tropical Plumes or Active Subtropical Jet (Armon et al., 2018; Dayan & Abramski, 1983; Rubin et al., 2007; Tubi et al., 2017).

2.2. HPEs Identification

A collection of carefully selected HPEs was used in this study (Table S1 in the Supporting Information S1) following Armon et al. (2020) and described here briefly. It consists of 41 HPEs identified based on their magnitude compared to a 24-year (1990/1991–2013/2014) rainfall climatology from physically-corrected and gauge-adjusted weather radar rainfall data (Marra & Morin, 2015, Figure 1b). A HPE was identified when at least a thousand 1-km² radar pixels exhibited a rain rate greater than the local 99.5th quantile of the conditional rain rate (i.e., where the rate is ≥ 0.1 mm hr⁻¹) for at least one of the durations analyzed, as detailed below and in Table S1 in Supporting Information S1, thus revealing events which can be considered locally intense. To have a good representation of both short- and long-duration HPEs, this process was repeated for durations of 1–72 hr. Events identified as a HPE for more than one duration were merged. Return levels of the 99.5th quantile thresholds are roughly 2–10 years. Events were separated by at least 24 hr with less than 100 pixels displaying rainfall of more than 0.1 mm, and they span 3.4 ± 1.6 days (mean and standard deviation). This collection of HPEs represents a large variance of synoptic conditions (Alpert et al., 2004), associated with both MCs (35 events) and ARSTs (six events). A detailed description of the collection of events and their rainfall characteristics is given in Armon et al. (2020).

2.3. Weather Research and Forecasting Simulations

Each of the 41 HPEs was simulated twice, using version 3.9.1.1 of the weather research and forecasting (WRF) atmospheric model, at a convection permitting resolution. The first simulation of each event represents the historic conditions during the storm (near the end of the 20th century; Section 2.3.1). Results of these simulations were previously published (Armon et al., 2020). The second simulation represents a hypothetical storyline in which the same HPE hits the area by the end of the 21st century, when global warming conditions at a representative concentration pathway (RCP) 8.5 prevail in the region (Section 2.3.2).

2.3.1. Simulation of Historic HPEs

Simulation of historic HPEs was conducted at a configuration suitable for a skillful representation of rainfall patterns in the eastern Mediterranean (Armon et al., 2020; Romine et al., 2013; Rostkier-Edelstein et al., 2014; Schwartz et al., 2015). This configuration includes three two-way nested domains (Figure 1a; Table S2 in Supporting Information S1) in which the innermost domain is simulated at a very high spatial and temporal resolution (1 km², 4–8 s). Convective parametrization (Tiedtke, 1989) was used only in the two outer nests, while in the inner nest the resolution is high enough to explicitly represent convection (e.g., Prein et al., 2015). Further details, including parametrization schemes (Iacono et al., 2008; Janjić, 1994; Tewari et al., 2004; Thompson et al., 2008), are described in Table S2 in Supporting Information S1 and in Armon et al. (2020).

Initial and boundary conditions for the historic simulations are 6-hourly ERA-Interim reanalysis data, at 60 vertical layers with a T255 spectral spatial resolution (~ 80 km) (Dee et al., 2011). HPEs were simulated starting 24 hr before the beginning of the observed rainfall (rounded down to the previous 6 hr) and lasted until the end of the HPE (rounded to the following 6 hr). The 24 hr period before the event is considered a spin-up phase, for which we discard the rain fields. This duration is considered long enough to correct spatial heterogeneities arising from the initial conditions (Gómez-Navarro et al., 2019; Picard & Mass, 2017; Warner, 2011: pp 215–216), which is crucial in correcting non-physical properties of the atmosphere, expected to be present in the PGW simulations because of the usage of ensemble mean fields (Shepherd, 2019; Tebaldi & Knutti, 2007) (Section 2.3.2). Precipitation outputs for the innermost domain were saved at 10 min intervals.

As was shown by Armon et al. (2020), the total precipitation during most of the HPEs reproduces the structure, location, and the seasonal change of the precipitation's center-of-mass of radar-observed precipitation, albeit with a positive bias. Given that HPEs in the region are characterized by small spatiotemporal scale rain-cells, it

is important to note the model's skill is particularly good at its “raw” (1 km²) resolution for total rain amounts of <25 mm, however, for larger amounts, a skillful representation must include a spatial averaging of at least a few tens of square kilometers. The model also well represents areal mean rainfall amounts, for various durations, which are crucial drivers of the hydrological response to precipitation. MC-type HPEs are better simulated compared to ARSTs, which are in general shorter and more local in nature.

2.3.2. Simulation of “Future” HPEs

To simulate the occurrence of the same HPEs in the future we used the PGW methodology (Kawase et al., 2009; Rasmussen et al., 2011; Schär et al., 1996). Each of the simulated HPEs was forced with the same input data as the historic events (Section 2.3.1) after adding the signal of climate change to the following input variables: surface pressure, skin temperature (including sea surface temperature), and 3D fields of temperature, wind, and specific humidity. In contrast to homogeneous changes common in the surrogate climate-change methodology (Keller et al., 2018; Schär et al., 1996), 3D spatial heterogeneity in the altered fields used in PGW experiments allows for representation of non-uniform spatial response to global warming (e.g., Rasmussen et al., 2011).

The changes applied over the initial and boundary conditions, for each pixel and time step (denoted hereon as Δ), were derived from the monthly values (October–April) of the ensemble mean of 29 models of the Coupled Model Intercomparison Project Phase 5 (CMIP5; Table S3 in Supporting Information S1) (K. E. Taylor et al., 2012). They were based on the difference in the corresponding parameter values for the end of the 21st century and the end of the 20th century under an RCP 8.5 scenario, as follows:

$$\Delta X_j = \overline{\overline{X_j}}_{2074-2099}^{29 \text{ models}} - \overline{\overline{X_j}}_{1979-2004}^{29 \text{ models}} \quad (1)$$

where X is a specified meteorological variable, defined for the particular month (j) of the HPE occurrence. The double overbar represents the mean of this parameter over the future (2074–2099) or historic (1979–2004) periods, averaged among the 29 CMIP5 models. Δ fields were linearly interpolated into a common grid, similar to the ERA-Interim horizontal grid (T255) and consisting of 42 levels in the vertical (model top = 10 hPa) for the 3D fields, over the entirety of the outermost domain. It should be noted that the WRF simulations are atmosphere-only simulations forced with the ensemble mean sea surface temperature warming computed from the “coupled” CMIP5 models; air-sea coupling processes not represented by the CMIP5 ensemble mean are necessarily missed in such a setup.

The changes applied represent a major warming of the region over the whole troposphere, but specifically over its upper levels, in concert with previous studies (e.g., Giorgi & Lionello, 2008; Seager et al., 2019; A. Tuel & Eltahir, 2020; Tuel et al., 2021). Surface temperature increases on average by 4.3°C. Alongside the warming, is a decrease in the zonal component of wind and an increase of the sea level pressure in the central Mediterranean, as detailed in the Text S1 and Figure S1 in Supporting Information S1.

2.4. Analyzed Rainfall Parameters and Statistical Methods

The parameters examined here are based on the 10-min rainfall fields from the innermost domain of both the historic and the “future” (PGW) simulations. Rainfall parameters from the historic and future simulations were compared both through their entire distribution across all events, and through an event-based paired comparison (historic-future). To enable comparison between events of different magnitudes, in many instances we normalize the quantity examined to its historic value: $100 \times \frac{\overline{\text{future}_i} - \overline{\text{historic}_i}}{\overline{\text{historic}_i}}$, where i indicates a specific event for which “future” and “historic” quantities are spatially averaged. We distinguished between conditional and unconditional rain rates. For *conditional* rain rate, we used only pixels and time steps where the rate was higher than 0.1 mm hr⁻¹; while for *unconditional* rain rate all cases (including 0) were considered.

The following rainfall parameters were considered for each event:

1. Rainfall accumulation for each pixel.
2. Areal mean rainfall accumulation, which is the average of (1) over the region of interest.

3. Factors affecting areal mean rainfall accumulation: the value in (2) above can be obtained by integrating rain rates over all rainy pixels and time steps and divide by the area of the region. Therefore, it is possible to consider three factors affecting the areal mean accumulation:
 - (a) The mean conditional rain rate (i.e., averaging rain rates $>0.1 \text{ mm hr}^{-1}$).
 - (b) The duration of the events is defined here as the time it took the central 90% of rainfall mass to precipitate.
 - (c) The rain area of the event is the time-average of the area covered by 10-min rain rates $>0.1 \text{ mm hr}^{-1}$ along the event. We also examine the rain area for higher rain rate thresholds in the range of $0.5\text{--}100 \text{ mm hr}^{-1}$.
4. Maximal rain rates for durations of 10, 20 and 30 min, 1, 3, 6, 12 and 24 hr for each pixel. To diminish the effect of single outlier pixels, the rain field is first smoothed spatially using a 3×3 pixel moving average window.
5. Regionally maximal rain rates for the same durations as in (4), which are taken as the maximal value from (4) over all pixels in the region.

The analyses above were evaluated both over the entire study region and over four sub-regions. These are the Mediterranean Sea area, the land area, and a division of the latter to the area north to the 200 mm isohyet roughly corresponding to the Mediterranean climate zone, and the area south of the 200 mm isohyet roughly corresponding to the desert climate zone (Figure 1b).

To analyze changes in the spatial structure of precipitation we compared the spatial autocorrelation structure of the 10-min rain fields whenever these were considered as having convective elements, following Marra and Morin (2018). Convective elements are defined here as spatially connected regions of area $\geq 3 \text{ km}^2$ with rain rates $>10 \text{ mm hr}^{-1}$ that include at least one pixel with rain rate $>25 \text{ mm hr}^{-1}$. Following Peleg et al. (2013), the spatial autocorrelation was calculated through fitting a three-parameter exponential function to the 2D spatial autocorrelation field (e.g., Nerini et al., 2017) of each of the convective rain fields as in Equation 2:

$$r(h) = ae^{-\left(\frac{h}{b}\right)^c} \quad (2)$$

where h is the lag distance, b , termed the correlation distance, is the distance at which the correlation decreases to $r = ae^{-1}$, a is the nugget (intercept) parameter and c is the shape parameter. For each event, the representative parameters of Equation 2 are the intra-event medians over all convective rain fields. The comparison of the autocorrelation structure between historic and future events is based on the inter-event medians of these representative parameters. This analysis can help in understanding the change in spatial structure of the rain fields, where larger values point to either larger rain cells or precipitation which is more spatially coherent, or both.

Statistical significance of the changes in *pixel*-based parameters is determined through the paired t -test with 5% significance level. For event-based parameters, statistical significance is declared if both paired t -test and paired Wilcoxon signed-rank tests are statistically significant at the 5% level.

3. Results

To have a better understanding of the changes between “future” (PGW) and historic simulations, we first present an examination of the first HPE in our collection, which exhibits many of the features observed throughout the events. It is followed by the results obtained throughout the HPEs collection.

3.1. Case Study #1

The first HPE in our collection (2–5 Nov 1991) is characterized by the passage of a MC, triggering numerous rain cells crossing the region with a general SW-NE track. These rain cells contributed $>100 \text{ mm}$ of accumulated rainfall mainly to the north coast and mountainous areas of the study region (Figure 2a, Movie S1 in Supporting Information S1). The areal average rainfall accumulation simulated over the entire domain for the historic event is 21.9 mm . Compared to the historic event, the future event exhibits a pronounced (-20%) decrease in precipitation with areal average rainfall accumulation summing to 17.5 mm (Figures 2b and 2c). The decrease is more pronounced over the land area (-28%) compared to the sea area (-16%), and is similar between the desert and Mediterranean regions of the land area (-29% and -27% , respectively).

In contrast to the decrease in total rain amounts, short duration (10-min) rain rates reveal a more complicated pattern (Figure 2d). When considering the distribution of all 10-min time steps and pixels, including those with




Figure 2. Rainfall in HPE#1 (2–5 Nov 1991). (a) Total rainfall for the historic simulation (see Armon et al., 2020: Figure 7 for a comparison with precipitation measured by the weather radar). (b) Total rainfall for the future (PGW) simulation. (c) Difference between historic and future simulations (future–historic). (d) Upper 1% quantile of the cumulative frequency curve of unconditional 10-min rain rates throughout the event (i.e., including no-rain intervals and pixels) for the historic (blue line) and future (orange) events. The dashed yellow line marks their intersection. The small inset presents the full cumulative frequency curve of conditional 10-min rain rates (where rain rate > 0.1 mm hr⁻¹), with mean rain rates in dashed lines. (e) Time series of the areal coverage of rainfall (% of the area covered by rain rate > 0.1 mm hr⁻¹). Please note the larger historic relative to the future unconditional rain rates (up to the 99.75% quantile [marked with yellow dashed line]; panel d) and the rain area (e) while future conditional rain rates are larger throughout the distribution (inset in d and Figure S2a in Supporting Information S1).

no-rain (i.e., unconditional rain rates), most of the distribution presents decreased rain rates and only the uppermost quantiles (>99.75%) of future rain rates increase compared to the historic ones. For example, the 99.99% quantile (corresponding to $\sim 1.4 \cdot 10^4$ pixel-timestep values of 10-min rain rates), is increased by 21% (from 77 mm hr⁻¹ to 93 mm hr⁻¹). However, the decrease in most of the unconditional rain rate quantiles is very much affected by the change in the spatiotemporal coverage of the event, namely the wet-frequency. Conversely, considering the distribution of the rainy pixels and time steps, that is, the conditional 10-min rain rate, quantiles of the future HPE are increasing throughout the distribution (Figure 2d inset). The mean value of the conditional rain rate increases from 2.64 mm hr⁻¹ for the historic event to 3.43 mm hr⁻¹ for the future one (+30%).

In addition to the mean conditional rain rate, two other factors affect the areal mean rainfall (Section 2.4), the duration and the rain area (Figure 2e). The duration of the event (Figure S2a Supporting Information S1) decreased from 2,440 min to 1,850 min (–24%) between the historic and future simulations. This reduction reflects a delayed start of the “core” of the rainfall during the passage of the MC, and an earlier termination (Movie S1 in Supporting Information S1). The rain area (Figure 2e) exhibits a major contraction (–38%) between the historic and future simulations, from $31.9 \cdot 10^3$ km² (10.5% of the study region) to $19.7 \cdot 10^3$ km² (6.5%) in historic and future simulations, respectively. This major decrease in rain area reflects the decrease in the area of precipitating rain cells, seen clearly in Movie S1 in Supporting Information S1, as well as in their number. However, it is important to note that we leave for future work a quantitative assessment or tracking of individual rain cells (e.g., Belachsen et al., 2017; Peleg & Morin, 2012). Nevertheless, we did compute the spatial autocorrelation of convective rainfall (Section 2.4). The spatial autocorrelation distance is 7 and 5 km, respectively for the historic and future events (Figure S2b Supporting Information S1) implying either that rainfall in the future event consists of smaller rain cells, or that the rain field is less spatially coherent. In addition, the number of 10-min convective time steps decreases by 5.1% (from 429 to 407).




Figure 3. Total rainfall summed over the 41 HPEs historic (a) and future (PGW; (b) simulations. (c) Difference between historic and future simulations (future–historic). Statistically significant differences are demarcated by gray lines. (d) Same as (c) but in relative terms: $\left(100 \times \frac{\text{future} - \text{historic}}{\text{historic}}\right)$.

In summary, this case study of the first HPE in our collection indicates that, moving from historic to future climates, areal mean rainfall accumulation decreases whereas conditional 10-min rain rates increase. This opposing behavior is caused by the decrease in the duration of the rainfall and even a greater decrease in the rain area, where the latter is probably due to the reduction in the area of precipitating rain cells and possibly in their number. The decrease in duration and in rain area, which means a decrease in wet-frequency, leads also to a decrease in almost all quantiles (except the uppermost ones) of the unconditional rain rate distribution, while the conditional rain rate distribution presents an increase in all quantiles.

3.2. Decreased Rainfall Accumulation Throughout Events

In general, future simulations show a significant decrease in rainfall accumulation compared with historic simulations (Figure 3), with a sum of the areal average over all 41 events of 485 mm, compared to 601 mm, respectively (–19%). This decrease is seen throughout the region, with >90% of the area exhibiting decreased rainfall. Given the large variability in rainfall in the region (inter-annual, inter- and intra-event), the fact that 35% of the area shows a significant change in rainfall accumulation highlights the robustness of the results. Out of the portion of pixels showing a statistically significant change, 99.97% exhibit a decrease in total rainfall. In absolute terms, this decrease is most severe in the wetter part of the region (the northern area and the mountains); 1% of pixels show a




Figure 4. Comparison of future (PGW) and historic areal mean rainfall accumulation for all 41 HPEs analyzed (a and b, Section 3.2), and the regional maximal 10-min rain rates (i.e., along all the pixels in the region and throughout time steps) (c)–(d, Section 3.4). Scatter plots (a), (c) compare the historic (horizontal axis) and future (vertical axis) values. Violin plots (b), (d) show the distribution of historic (pale colors) and future (bold colors) quantities for the different subregions (Figure 1b). Each colored dot along the violin represents a value of one HPE. Similar to boxplots, white dots are median values, and gray boxes are the inter quartile range. In addition, the colored region, mirrored in both sides of the boxplot, represents the kernel density estimation of the data (i.e., a non-parametric estimate of the probability density function). The change between medians of historic and future events is marked with dashed lines. Statistically significant differences between the paired-event populations for each region (presented also as scatter plots in Figures S3 and S7 in Supporting Information S1) are marked here with ** (see also Table 1).

decrease of ≥ 618 mm (Figure 3c). In relative terms, the decrease is most drastic over the Sinai-Negev desert (Figure 1b) and throughout the eastern part of the domain; 1% of pixels show a decrease of at least 55% (Figure 3d).

The decrease in rainfall accumulation recognized above for the first HPE is preserved among most of the analyzed HPEs. More than 90% of events feature smaller rainfall accumulations in future compared to historic simulations (Figure 4a, Table 1) with a significant inter-event average decrease of -2.8 mm (equivalent to -30%). Like the first case study, the decrease is more pronounced over land compared to the sea with a significant change of -36% and -26% in average precipitation, respectively. Similarly, 95% and 83% of HPEs exhibited less rainfall in future compared to historic simulations over land and sea, respectively. The decreased precipitation is significant in each of the sub-regions considered here (Figures 4b and S3, Table 1).

3.3. Opposing Changes in Rainfall Properties: Increased Conditional Rain Rates, Decreased Duration and Areal Coverage

Changes in the event-based areal mean rainfall accumulation are examined through changes in three rainfall components: mean conditional rain rate, event duration, and rain area (Figure 5, Figure S4 and S5 in Supporting Information S1, Table 1). The mean conditional rain rate significantly increases with an average change of 15%, and $>85\%$ of the events show higher average conditional rain rate in the future compared with the historic simulations. Although there is some correlation between the changes in mean conditional rain rate and total rainfall (Spearman's correlation: $\rho = 0.37$; Figure S5 in Supporting Information S1), increases as large as 47% (event

Table 1
Changes in Rainfall Properties Between Future and Historic Simulations

Region	Fraction of HPEs exhibiting larger values in future [%]				Average normalized change: $\frac{1}{n} \sum_{i=1}^n \left(\frac{future_i - historic_i}{historic_i} \right) [%]$			
	Areal mean				Areal mean			
	Land	Sea	Med	Des	Land	Sea	Med	Des
Areal average rainfall accumulation			10					-30
	5	17	10	12	-36	-26	-34	-37
Regionally max 10-min rain rate			85					22
	80	83	83	63	18	23	21	11
Mean conditional rain rate			85					15
	71	85	66	59	6	13	8	5
Duration			24					-9
	24	29	24	37	-8	-7	-8	2
Rain area			2					-40
	2	2	2	2	-42	-38	-41	-42

Note. Bolded numbers are statistically significant (i.e., on both paired t- and paired Wilcoxon signed-rank tests, Section 2.4). Italics are significant on the paired t-test only. The upper value represents the entire region and below are values for each sub-region (land, sea, med = Mediterranean, and des = desert, Figure 1b).

#16) are observed in events with a reduction of the rainfall accumulation. This suggests that the change in the mean conditional rain is rather weakly related to rainfall accumulation.

While mean conditional rain rate is increasing, the duration of events shows a significant negative change (-9%), from a mean of 3290 min–2980 min, and >75% of the events are longer in historic compared to future simulations (Table 1). Most of these cases exhibit a delayed rainfall initiation in the future (example: Figure S2a in Supporting Information S1, Movie S2 in Supporting Information S1) compared to the historic simulations (not shown). The




Figure 5. Normalized change between future and historic simulations $\left(100 \times \frac{future - historic}{historic} \right)$ for the 41 HPEs analyzed. Events are sorted by the change in rainfall accumulation (blue bars). Dashed lines represent average inter-event values (written also inside the legend). The non-normalized changes are shown in Figure S4 in Supporting Information S1. Correlations between the change in rainfall accumulation and the three other parameters are in Figure S5 in Supporting Information S1.




Figure 6. Changes in areal coverage for different rain rate thresholds. (a) Areal mean coverage along each event (small dots) for different rain intensity thresholds (different colors) in historic events (horizontal axis) and future (PGW) events (vertical axis). Values are normalized by the maximal value observed for each rain rate threshold. Average inter-event values are marked with large circles. Filled circles demonstrate statistically significant changes between future and historic values, and one hollow circle shows an insignificant difference. (b) Fraction of events with larger areal coverage in historic compared to future simulations for each of the inspected rain rate thresholds.

reduction in durations is not, however, a good predictor for the reduction in total rainfall accumulation, with a low, non-significant negative correlation between the two (Figure S5 in Supporting Information S1).

The third component in this analysis, the rain area, reveals the largest relative change out of all three components. It also presents the highest correlation with the change in rainfall accumulation ($\rho = 0.93$; Figure S5 in Supporting Information S1). The rain area is decreased in all but one of the events (>97% of events) with a significant and substantial average decrease (−40%). Events with the lowest reduction (or even slight increase) in rainfall accumulation are only those in which the reduction in the rain area is relatively small, while events with a large decrease in rainfall accumulation are accompanied by a large decrease in rain area. These results indicate that the factor most heavily related to changes in rainfall accumulation is the rain area.

3.3.1. Spatial Concentration of Future Rainstorms

To better understand the changes in the rain area, we examine its changes using a range of rain rate thresholds. Divergent changes in the extent of areas exceeding various rain rates are essential in understanding possible hydrological responses to climate change (Bacchi & Ranzi, 1996; Fowler, Lenderink, et al., 2021; Peleg et al., 2018). Figure 6a displays the event-average areal rainfall coverage for each HPE with different rain rate thresholds, normalized by the largest coverage for each intensity for both historic and future simulations. The fraction of events with larger areal coverage for each rain rate threshold in historic events, that is, the relative number of points below the 1:1 line, is displayed in Figure 6b. The areal coverage of rainfall with relatively low rain rates (0.1–5 mm h⁻¹) is reduced significantly for future compared to historic simulations. The opposite case is true for high rain rates (20–100 mm hr⁻¹). This change occurs at rain rates of ~10 mm hr⁻¹, where no significant change is documented between rain area above this threshold in historic and in future simulations.

These different trends imply that when compared with historic rainstorms, the total “wet” area in future events is lower, and the storms are more concentrated around the higher rain rates. This conclusion is further strengthened by the change in the autocorrelation pattern of convective rainfall, which demonstrates a much sharper decrease with distance in future compared with historic simulations and, accordingly, the median of the autocorrelation distance decreases from 8 to 5 km (Figure S6 in Supporting Information S1).




Figure 7. Changes in maximal rain rates averaged over all events for durations of 10-min (a), 1 hr (b), 6 hr (c) and 24-hr (d) between future and historic simulations (future-historic). Statistically significant differences are circumscribed in gray.

3.4. Changes in Extreme Rain Rates for Different Durations

The change in maximal rain rate for each pixel displays different behavior along the study area and between durations. The maps in Figure 7 show the difference between the inter-event average of the maximum rain rate per event for each pixel, over durations of 10-min to 24-hr. For short durations (10–60 min, Figures 7a, 7b, and Figure S7 in Supporting Information S1), significant decreases are identified mainly over the southern and eastern desert areas and far into the sea. In contrast, a positive change sub-parallel to the coastline and over the northern part of the region is present. It is observed mainly a few kilometers offshore and over the mountains at the north of the study region. Over longer durations (a few hours to one day), a larger portion of the region exhibits a significant change in maximal rain intensities (Figures 7c, 7d, and Figure S7 in Supporting Information S1). This change is almost exclusively negative, focusing over both the desert area and most of the northern land region (excluding the shoreline and the upslopes of the Lebanon Mountains). In relative terms, for the longer durations, almost 25% of the area exhibits a decrease of more than 40% compared with areal-average maximal rain rate in historic events (Figure S7 in Supporting Information S1). It must be noted, however, that the spatial perspective presented in Figure 7 involves both the increase in rain rates, and the decrease in wet-frequency caused by smaller areal coverage of these intensities and shorter event durations, resulting in a mixture of increased and decreased maximal rain rates.

In contrast to the decrease in rainfall accumulation, and as exemplified by the first case study (Section 3.1), regionally maximal 10-min rain rates (maximum along all pixels and time steps) in future simulations are significantly higher than in historic simulations (Figures 4c and 4d, Table 1) with an average increase of 22%. This conclusion holds for all sub-regions inspected here, except for the desert sub-region, in which the increase (11%) is non-significant (Figures 4d and Figure S8 in Supporting Information S1, Table 1). Increases of the regionally maximal 10-min rain rates over both the Mediterranean climate and Sea sub-regions are on average >21%, and the increase over land, as a result of the small increase over the desert, is 18%. Furthermore, most of the events (85%) have higher values in future compared to historic simulations and this is rather consistent among the different sub-regions (Table 1). The increase in regionally maximal rain rates between historic and future simulations holds for longer durations as well (Figure S9 in Supporting Information S1).

4. Summary and Discussion

This study shows the changes in rainfall patterns between paired simulations of historic and future HPEs, with the objectives of identifying whether common changes in rainfall patterns exist, and characterizing these changes. The collection of objectively identified 41 HPEs was simulated twice, and the results of the two simulations were compared. The first simulation is based on historic conditions, and the second applies expected changes in various meteorological parameters from the RCP 8.5 scenario for the end of the 21st century on top of historic initial and boundary conditions. Selected events represent some of the heaviest precipitation events in the region around the end of the 20th century. Our results, shown first for a case study, and then for the full collection of HPEs, demonstrate the added value of using event-based simulations, and provide high resolution projections of future changes in rainfall patterns, highlighting the importance of changes in specific rainfall constituents, as discussed below.

4.1. Opportunities Gained by the Event-Based Approach and Their Implications

Large-scale and long-term CPM simulations are becoming increasingly attainable, allowing to better characterize precipitation extremes in future climate scenarios (e.g., Coppola et al., 2020; Kendon et al., 2018). However, there are still difficulties in providing reliable projections of rainfall during HPEs (Kendon et al., 2021); the computational and power consumption costs of these simulations are huge (Fuhrer et al., 2018; Loft, 2020), and the rarest of extremes are difficult to characterize even in runs extending for many years. Therefore, if the purpose of a study is to identify potential changes in only a subset of the climate, for example, HPEs, a full-climate run should be used prudently.

Here, using an event-based approach, we were able to show plausible impacts of climate change on some of the heaviest rainstorms in the eastern Mediterranean. Furthermore, we show that many “plausible” instances (i.e., individual HPE events) point in the same direction; therefore, the plausible scenario may be considered as the probable scenario. Even if the entire variance of possibilities is not perfectly represented using this method, the emerging similar response enables us to garner insight on “climate questions,” such as projections of future precipitation patterns, using a *weather* model. We showed that rainfall accumulation under global warming conditions decreases over >90% of the simulated HPEs and analyzed the properties of rainfall accounting for this decrease. The rain area exhibits the largest and most consistent decrease and is heavily associated with the decrease in rainfall accumulation, while increased conditional rain rate is only weakly related to rainfall accumulation and cannot counteract the decreased rain area.

The simulated change in rainfall patterns can have considerable implications both on water resources and on natural hazards, which can be illuminated if we focus on specific rainstorms. For example, event #8 (1–7 Nov 1994) is an infamous ARST storm in which more than 500 people lost their lives, and extensive floods and damages occurred in Egypt and Israel (De Vries et al., 2013; Krichak et al., 2000). This event shows a substantial reduction in total rainfall under future-simulated conditions (–51%; Figures 5 and Figure S10 in Supporting Information S1). Such a reduction would probably lead to a reduced risk of flash flooding, especially at the northern part of the region. However, while in many places total rainfall decreased in the simulation, few high-intensity rain cells still impacted the Sinai desert (Movie S2 in Supporting Information S1), with total rainfall of >100 mm, which would undoubtedly cause substantive floods in this region.

HPE #12 (Figure S11 in Supporting Information S1) triggered a major streamflow increase and raised the level of the Sea of Galilee, the largest surface freshwater reservoir in the region, by 45 cm within a week (compared with <10 cm rise the week before the storm occurred). This rise is equivalent to the yearly industrial consumption of freshwater in Israel at that time ($\sim 90,10^6 \text{ m}^3$) and constitutes more than a fifth of the annual water rise of the lake. The simulation of the future event indicates a substantial decrease in total rainfall (-27%). As the hydrological response to decreases in rainfall is non-linear (e.g., Peleg et al., 2014), this would probably lead to an even larger decrease in freshwater recharge with major implications on water resources. While a hydrological simulation of the different events is out of the scope of this paper, we stress that to have better insights about the hydrological response, a comparison of historic and future simulations of specific events through a hydrological model is highly desired.

It is important to note that the frequency of events (e.g., Myhre et al., 2019) is not implicitly considered in our simulations. Rain events in the region are projected to have a reduced frequency ($\sim -20\%$; Hochman et al., 2018; Zappa et al., 2015), and thus, the decreased rainfall we show here for the specific simulated events, may be considered as an underestimation of the projected changes in total precipitation from HPEs accounting for event occurrences.

A possible limitation of this study is the use of a single climate scenario forcing for the PGW and as such it will give only plausible results, rather probable. However, (a) this single scenario is the ensemble mean of CMIP5 models, which can be considered as a best estimate, to date, of large scale future changes, though work currently in progress shows that CMIP6 models generally simulate similar, and if anything more severe, changes to CMIP5 in this region (not shown), (b) we use a collection of many objectively identified events that constitutes some of the highest magnitude HPEs in the region. Results for this large set of paired-simulations show a similar behavior of different events representing different synoptic-scale conditions. Therefore, we claim that the sign and magnitude of the changes that emerge from these simulations should be considered as a probable projection of HPEs in the region.

Indeed, the PGW event-based methodology provides us with projections for HPEs in a warmer climate. However, it must be noted we do not attempt to provide a climatology of HPEs in the future, nor give updated extreme event levels and frequencies. While these can be obtained using a framework which accounts for the frequency of events (Marra et al., 2019), the results we obtain have significance in drawing possible future scenarios for some of the heaviest precipitation events in the region. High resolution rainfall projections can also help improve future predictions in approaches requiring a changed rainfall distribution (Marra et al., 2021).

4.2. Changes in Rainfall Patterns During Rainstorms

Future rainstorms simulated in this work show quite a difference in rainfall patterns compared to historic rainstorms, mainly being more concentrated in both space and time. Given that the conditional rain rate increases, one might expect an increase in total precipitation during HPEs, as projected, for example, over Europe (e.g., Y. Chen, Paschalis, et al., 2020; Hawcroft et al., 2018; Kendon et al., 2014). However, two other factors, less often addressed, negatively affect total rainfall: the size of the rain area, and the duration of the events. Among these two, we find that the rain area is the main contributor to decreased rainfall accumulation, which decreases, on average, by 40%. Furthermore, the rain area has a high correlation with the changes in rainfall accumulation, while the event duration decreases on average by 9% and has a low correlation with rainfall accumulation changes.

It must be noted, however, that the changes in the rain area are not constant over different rain rates thresholds. The baseline 0.1 mm hr^{-1} intensity is a good proxy for the total storm area. Going to larger thresholds, the area represented is a better indicator for the intense “core” of the storm, namely the inner part of convective cells during the storm. In fact, we found an increase in the rain area for thresholds of $>10 \text{ mm hr}^{-1}$. This means that, although the total rain area of HPEs shrinks, their cores are getting larger in future simulations. Similar findings were reported by Peleg et al. (2018) using historic radar observations, and temperature as a proxy, over the eastern Mediterranean and by Wasko et al. (2016) using rain gauges in Australia. Both studies showed that total rain area and the convective core area scale with temperature in opposite directions: total area exhibits a negative scaling, while the area of the convective cores is positively scaled with temperature; this is probably related to an enhanced moisture convergence into the convective cores from the total storm extent. In contrast, results from studies of future extreme precipitation in the Netherlands and in the UK show the area of the storms is expected to

increase with global warming (Y. Chen, Paschalis, et al., 2020; Lochbihler et al., 2017, 2019), which may indicate a regional dependence in the scaling of the rain area, but this topic should be addressed in future studies (Fowler, Lenderink, et al., 2021).

Since the hydrological response to HPEs is heavily related to space-time precipitation characteristics, the results shown above would have an immense impact on the hydrology of future rainstorms. Larger storm cores, having increased short duration rain rates may increase the risk of urban flooding and short-lived, fast responding flash floods (e.g., Tarasova et al., 2019), as well as soil erosion (e.g., Shmilovitz et al., 2021). However, this effect is expected to be mitigated by the decreased rainfall frequency caused by the shorter storm duration and smaller overall area. Combined, a possible conclusion could be that over the affected (rainy) area, the risk of short-duration natural hazards is higher, while over the entire domain this is uncertain. Yet, a clearer conclusion can be drawn for the detrimental effects of the changes in rainfall patterns over the entire storm through longer-duration processes: mean rain rates and amounts are expected to dramatically decrease. Therefore, the expected hydrological impact would include a further reduction of streamflow and a decline in freshwater resources, which requires immediate attention by policy makers.

Two key aspects are missing from the results presented here: a detailed analysis of the meteorological factors affecting the modeled change in rainfall patterns and their scaling with temperature, and a modeling of the hydrological impact of these changes. These two prospective aspects are currently being further studied. We call for a continued use of the PGW methodology as a relatively easy-to-implement experiment, with results relevant to events of specific interest such as HPEs.

5. Conclusions

Through high-resolution event-based simulations of eastern Mediterranean HPEs in present and future climate, we show that in the future: (a) event rainfall accumulations decrease substantially (inter-event average = -30%), throughout the study region, (b) mean conditional rain rate is increased ($+15\%$), (c) event duration is getting shorter (-9%), and (d) rain area becomes dramatically smaller (-40%). The areal coverage for various rain rates shows opposing changes for lower and higher rain rates: it is reduced for low rain rate thresholds, and expanded for high rate thresholds. Thus, rainstorms become more concentrated in future simulations, with convective cores that exhibit shorter autocorrelation distance and higher regionally maximal rain rates ($+22\%$). Furthermore, some increases in local short duration rain rates are seen mostly over the coastal region, but long duration rain rates are decreased throughout the region. The changes found are rather consistent across events, suggesting that these event-based conclusions may actually be probable. Changes in rainfall properties identified here reveal the dominance of the rain area in determining the decrease in total rainfall, with potentially significant implications for future hydrological processes.

Data Availability Statement

Shacham radar data for the 41 HPEs are available online (<https://doi.org/10.5281/zenodo.5353714>). ERA-Interim data were downloaded from the Research Data Archive at the National Center for Atmospheric Research, Computational and Information Systems Laboratory (<https://doi.org/10.5065/D6CR5RD9>). CMIP5 data were downloaded from the ESGF Node at DKRZ (<https://esgf-data.dkrz.de/projects/esgf-dkrz/tou>). The WRF namelist input file can be found in the Supporting Information S1.

Acknowledgments

The authors thank Y. Shmilovitz and R. Dann for both fruitful discussions and help with coding issues. This research has been supported by the Israel Ministry of Science and Technology (grant no. 61792) and the Israel Water Authority, it was also partially funded by DFG (PALEX project, grant no. BR2208/13-1/-2). FM was supported by the Institute of Atmospheric Sciences and Climate (ISAC) of the National Research Council of Italy (CNR).

References

- Alpert, P., Ben-Gai, T., Baharad, A., Benjamini, Y., Yekutieli, D., Colacino, M., et al. (2002). The paradoxical increase of Mediterranean extreme daily rainfall in spite of decrease in total values. *Geophysical Research Letters*, 29(11), 1536. <https://doi.org/10.1029/2001GL013554>
- Alpert, P., Osetinsky, I., Ziv, B., & Shafir, H. (2004). Semi-objective classification for daily synoptic systems: Application to the eastern Mediterranean climate change. *International Journal of Climatology*, 24(8), 1001–1011. <https://doi.org/10.1002/joc.1036>
- Alpert, P., & Shay-EL, Y. (1994). The moisture source for the winter cyclones in the Eastern Mediterranean. *Israel Meteorological Research Papers*, 5, 20–27.
- Armon, M., Dente, E., Smith, J. A., Enzel, Y., & Morin, E. (2018). Synoptic-scale control over modern rainfall and flood patterns in the Levant drylands with implications for past climates. *Journal of Hydrometeorology*, 19(6), 1077–1096. <https://doi.org/10.1175/JHM-D-18-0013.1>
- Armon, M., Marra, F., Enzel, Y., Rostkier-Edelstein, D., & Morin, E. (2020). Radar-based characterisation of heavy precipitation in the eastern Mediterranean and its representation in a convection-permitting model. *Hydrology and Earth System Sciences*, 24(3), 1227–1249. <https://doi.org/10.5194/hess-24-1227-2020>

- Armon, M., Morin, E., & Enzel, Y. (2019). Overview of modern atmospheric patterns controlling rainfall and floods into the Dead Sea: Implications for the lake's sedimentology and paleohydrology. *Quaternary Science Reviews*, 216, 58–73. <https://doi.org/10.1016/j.quascirev.2019.06.005>
- Ashbel, D. (1938). Great floods in Sinai Peninsula, Palestine, Syria and the Syrian Desert, and the influence of the Red Sea on their formation. *Quarterly Journal of the Royal Meteorological Society*, 64(277), 635–639. <https://doi.org/10.1002/qj.49706427716>
- Atlas of Israel. (2011). *The new atlas of Israel: The national atlas*. Survey of Israel; The Hebrew University of Jerusalem.
- Bacchi, B., & Ranzi, R. (1996). On the derivation of the areal reduction factor of storms. *Atmospheric Research*, 42(1–4), 123–135. [https://doi.org/10.1016/0169-8095\(95\)00058-5](https://doi.org/10.1016/0169-8095(95)00058-5)
- Ban, N., Schmidli, J., & Schär, C. (2014). Evaluation of the convection-resolving regional climate modeling approach in decade-long simulations. *Journal of Geophysical Research*, 119(13), 7889–7907. <https://doi.org/10.1002/2014JD021478>
- Belachsen, I., Marra, F., Peleg, N., & Morin, E. (2017). Convective rainfall in a dry climate: Relations with synoptic systems and flash-flood generation in the Dead Sea region. *Hydrology and Earth System Sciences*, 21(10), 5165–5180. <https://doi.org/10.5194/hess-21-5165-2017>
- Borga, M., Stoffel, M., Marchi, L., Marra, F., & Jakob, M. (2014). Hydrogeomorphic response to extreme rainfall in headwater systems: Flash floods and debris flows. *Journal of Hydrology*, 518, 194–205. <https://doi.org/10.1016/j.jhydrol.2014.05.022>
- Brogli, R., Kröner, N., Sørland, S. L., Lüthi, D., & Schär, C. (2019). The role of Hadley circulation and lapse-rate changes for the future European summer climate. *Journal of Climate*, 32(2), 385–404. <https://doi.org/10.1175/JCLI-D-18-0431.1>
- Cannon, A. J., & Innocenti, S. (2019). Projected intensification of sub-daily and daily rainfall extremes in convection-permitting climate model simulations over North America: Implications for future intensity-duration-frequency curves. *Natural Hazards and Earth System Sciences*, 19(2), 421–440. <https://doi.org/10.5194/nhess-19-421-2019>
- Chan, S. C., Kendon, E. J., Giorgia, B., Elizabeth, F., & Fowler, H. J. (2020). Europe-wide precipitation projections at convection permitting scale with the Unified Model. *Climate Dynamics*, 1–23. <https://doi.org/10.1007/s00382-020-05192-8>
- Chen, G., Wang, W.-C., Cheng, C.-T., & Hsu, H.-H. (2021). Extreme snow events along the coast of the northeast United States: Potential changes due to global warming. *Journal of Climate*, 34(6), 2337–2353. <https://doi.org/10.1175/jcli-d-20-0197.1>
- Chen, J., Wang, Z., Tam, C. Y., Lau, N. C., Lau, D. S. D., & Mok, H. Y. (2020). Impacts of climate change on tropical cyclones and induced storm surges in the Pearl River Delta region using pseudo-global-warming method. *Scientific Reports*, 10(1), 1–10. <https://doi.org/10.1038/s41598-020-58824-8>
- Chen, Y., Paschalis, A., Kendon, E., Kim, D., & Onof, C. (2020). Changing spatial structure of summer heavy rainfall, using convection-permitting ensemble. *Geophysical Research Letters*, 1–12. <https://doi.org/10.1029/2020gl090903>
- Coppola, E., Sobolowski, S., Pichelli, E., Raffaele, F., Ahrens, B., Anders, I., et al. (2020). A first-of-its-kind multi-model convection permitting ensemble for investigating convective phenomena over Europe and the Mediterranean. *Climate Dynamics*, 55. <https://doi.org/10.1007/s00382-018-4521-8>
- Crook, J., Klein, C., Folwell, S., Taylor, C. M., Parker, D. J., Stratton, R., & Stein, T. (2019). Assessment of the Representation of West African Storm Lifecycles in Convection-Permitting Simulations. *Earth and Space Science*, 6(5), 818–835. <https://doi.org/10.1029/2018EA000491>
- Dayan, U., & Abramski, R. (1983). Heavy rain in the Middle East related to unusual jet stream properties. *Bulletin of the American Meteorological Society*, 64(10), 1138–1140. [https://doi.org/10.1175/1520-0477\(1983\)064<1138:hritme>2.0.co;2](https://doi.org/10.1175/1520-0477(1983)064<1138:hritme>2.0.co;2)
- Dayan, U., Lensky, I. M., Ziv, B., & Khain, P. (2021). Atmospheric conditions leading to an exceptional fatal flash flood in the Negev Desert, Israel. *Natural Hazards and Earth System Sciences*, 21(5), 1583–1597. <https://doi.org/10.5194/nhess-21-1583-2021>
- Dayan, U., & Morin, E. (2006). Flash flood—Producing rainstorms over the Dead Sea: A review. *New Frontiers in Dead Sea Paleoenvironmental Research: Geological Society of America Special Paper*, 401(04), 53–62. [https://doi.org/10.1130/2006.2401\(04\)](https://doi.org/10.1130/2006.2401(04))
- De Vries, A. J., Tyrils, E., Edry, D., Krichak, S. O., Steil, B., & Lelieveld, J. (2013). Extreme precipitation events in the Middle East: Dynamics of the active Red Sea trough. *Journal of Geophysical Research: Atmospheres*, 118(13), 7087–7108. <https://doi.org/10.1002/jgrd.50569>
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., et al. (2011). The ERA-Interim reanalysis: Configuration and performance of the data assimilation system. *Quarterly Journal of the Royal Meteorological Society*, 137(656), 553–597. <https://doi.org/10.1002/qj.828>
- Dirmeyer, P. A., Brubaker, K. L., & DelSole, T. (2009). Import and export of atmospheric water vapor between nations. *Journal of Hydrology*, 365(1–2), 11–22. <https://doi.org/10.1016/j.jhydrol.2008.11.016>
- Donat, M. G., Lowry, A. L., Alexander, L. V., O’Gorman, P. A., & Maher, N. (2016). More extreme precipitation in the world's dry and wet regions. *Nature Climate Change*, 6(5), 508–513. <https://doi.org/10.1038/nclimate2941>
- El-Fandy, M. G. (1946). Barometric lows of cyprus. *Quarterly Journal of the Royal Meteorological Society*, 72(314), 291–306. <https://doi.org/10.1002/qj.49707231406>
- Enzel, Y., Bookman (Ken Tor), R., Sharon, D., Gvirtzman, H., Dayan, U., Ziv, B., & Stein, M. (2003). Late Holocene climates of the Near East deduced from Dead Sea level variations and modern regional winter rainfall. *Quaternary Research*, 60(3), 263–273. <https://doi.org/10.1016/j.yqres.2003.07.011>
- Fatichi, S., Ivanov, V. Y., Paschalis, A., Peleg, N., Molnar, P., Rimkus, S., et al. (2016). Uncertainty partition challenges the predictability of vital details of climate change. *Earth's Future*, 4(5), 240–251. <https://doi.org/10.1002/2015EF000336>
- Ferreira, R. N. (2021). Cut-off lows and extreme precipitation in eastern Spain: Current and future climate. *Atmosphere*, 12(7), 835. <https://doi.org/10.3390/atmos12070835>
- Flaounas, E., Röthlisberger, M., Boettcher, M., Sprenger, M., & Wernli, H. (2021). Extreme wet seasons – Their definition and relationship with synoptic-scale weather systems. *Weather and Climate Dynamics*, 2(1), 71–88. <https://doi.org/10.5194/wcd-2-71-2021>
- Fosser, G., Khodayar, S., & Berg, P. (2015). Benefit of convection permitting climate model simulations in the representation of convective precipitation. *Climate Dynamics*, 44(1–2), 45–60. <https://doi.org/10.1007/s00382-014-2242-1>
- Fowler, H. J., Ali, H., Allan, R. P., Ban, N., Barbero, R., Berg, P., et al. (2021). Towards advancing scientific knowledge of climate change impacts on short-duration rainfall extremes. *Philosophical Transactions of the Royal Society A: Mathematical, Physical & Engineering Sciences*, 379(2195). <https://doi.org/10.1098/rsta.2019.0542>
- Fowler, H. J., Lenderink, G., Prein, A. F., Westra, S., Allan, R. P., Ban, N., et al. (2021). Anthropogenic intensification of short-duration rainfall extremes. *Nature Reviews Earth & Environment*, 2(2), 107–122. <https://doi.org/10.1038/s43017-020-00128-6>
- Fowler, H. J., Wasko, C., & Prein, A. F. (2021). Intensification of short-duration rainfall extremes and implications for flood risk: Current state of the art and future directions. *Philosophical Transactions of the Royal Society A: Mathematical, Physical & Engineering Sciences*, 379(2195). <https://doi.org/10.1098/rsta.2019.0541>
- Fuhrer, O., Chadha, T., Hoefler, T., Kwasniewski, G., Lapillonne, X., Leutwyler, D., et al. (2018). Near-global climate simulation at 1 km resolution: Establishing a performance baseline on 4888 GPUs with COSMO 5.0. *Geoscientific Model Development*, 11(4), 1665–1681. <https://doi.org/10.5194/gmd-11-1665-2018>

- Garfinkel, C. I., Adam, O., Morin, E., Enzel, Y., Elbaum, E., Bartov, M., et al. (2020). The role of zonally averaged climate change in contributing to intermodel spread in CMIP5 predicted local precipitation changes. *Journal of Climate*, 33(3), 1141–1154. <https://doi.org/10.1175/JCLI-D-19-0232.1>
- Giorgi, F., & Lionello, P. (2008). Climate change projections for the Mediterranean region. *Global and Planetary Change*, 63(2–3), 90–104. <https://doi.org/10.1016/j.gloplacha.2007.09.005>
- Goldreich, Y. (1994). The spatial distribution of annual rainfall in Israel—A review. *Theoretical and Applied Climatology*, 50(1–2), 45–59. <https://doi.org/10.1007/BF00864902>
- Goldreich, Y. (1995). Temporal variations of rainfall in Israel. *Climate Research*, 5, 167–179. <https://doi.org/10.3354/cr005167>
- Goldreich, Y. (2012). *The climate of Israel: Observation, research and application*. Springer Science & Business Media. <https://doi.org/10.1007/978-1-4615-0697-3>
- Gómez-Navarro, J. J., Raible, C. C., García-Valero, J. A., Messmer, M., Montávez, J. P., & Martius, O. (2019). Event selection for dynamical downscaling: A neural network approach for physically-constrained precipitation events. *Climate Dynamics*. <https://doi.org/10.1007/s00382-019-04818-w>
- Gutmann, E. D., Rasmussen, R. M., Liu, C., Ikeda, K., Bruyere, C. L., Done, J. M., et al. (2018). Changes in hurricanes from a 13-yr convection-permitting pseudo-global warming simulation. *Journal of Climate*, 31(9), 3643–3657. <https://doi.org/10.1175/JCLI-D-17-0391.1>
- Hawcroft, M., Walsh, E., Hodges, K., & Zappa, G. (2018). Significantly increased extreme precipitation expected in Europe and North America from extratropical cyclones. *Environmental Research Letters*, 13(12). <https://doi.org/10.1088/1748-9326/aaed59>
- Hochman, A., Harpaz, T., Saaroni, H., & Alpert, P. (2018). The seasons' length in 21st century CMIP5 projections over the eastern Mediterranean. *International Journal of Climatology*, 1–11. <https://doi.org/10.1002/joc.5448>
- Iacono, M. J., Delamere, J. S., Mlawer, E. J., Shephard, M. W., Clough, S. A., & Collins, W. D. (2008). Radiative forcing by long-lived greenhouse gases: Calculations with the AER radiative transfer models. *Journal of Geophysical Research*, 113(D13). <https://doi.org/10.1029/2008JD009944>
- Janjić, Z. I. (1994). The step-mountain ETA coordinate model: Further developments of the convection, viscous sublayer, and turbulence closure schemes. *Monthly Weather Review*, 122(5), 927–945. [https://doi.org/10.1175/1520-0493\(1994\)122<0927:TSMECM>2.0.CO;2](https://doi.org/10.1175/1520-0493(1994)122<0927:TSMECM>2.0.CO;2)
- Kahana, R., Ziv, B., Enzel, Y., & Dayan, U. (2002). Synoptic climatology of major floods in the Negev Desert, Israel. *International Journal of Climatology*, 22(7), 867–882. <https://doi.org/10.1002/joc.766>
- Kawase, H., Yoshikane, T., Hara, M., Kimura, F., Yasunari, T., Ailikun, B., et al. (2009). Intermodel variability of future changes in the Baiu rainband estimated by the pseudo global warming downscaling method. *Journal of Geophysical Research*, 114(24), 1–14. <https://doi.org/10.1029/2009JD011803>
- Keller, M., Kröner, N., Fuhrer, O., Lüthi, D., Schmidli, J., Stengel, M., et al. (2018). The sensitivity of Alpine summer convection to surrogate climate change: An intercomparison between convection-parameterizing and convection-resolving models. *Atmospheric Chemistry and Physics*, 18(8), 5253–5264. <https://doi.org/10.5194/acp-18-5253-2018>
- Kendon, E. J., Blenkinsop, S., & Fowler, H. J. (2018). When will we detect changes in short-duration precipitation extremes? *Journal of Climate*, 31(7), 2945–2964. <https://doi.org/10.1175/JCLI-D-17-0435.1>
- Kendon, E. J., Prein, A. F., Senior, C. A., & Stirling, A. (2021). Challenges and outlook for convection-permitting climate modelling. *Philosophical Transactions of the Royal Society A: Mathematical, Physical & Engineering Sciences*. <https://doi.org/10.1098/rsta.2019.0547>
- Kendon, E. J., Roberts, N. M., Fowler, H. J., Roberts, M. J., Chan, S. C., & Senior, C. A. (2014). Heavier summer downpours with climate change revealed by weather forecast resolution model. *Nature Climate Change*, 4(7), 570–576. <https://doi.org/10.1038/nclimate2258>
- Krichak, S. O., Tsidulko, M., & Alpert, P. (2000). November 2, 1994, severe storms in the southeastern Mediterranean. *Atmospheric Research*, 53(1–3), 45–62. [https://doi.org/10.1016/S0169-8095\(99\)00045-9](https://doi.org/10.1016/S0169-8095(99)00045-9)
- Kushnir, Y., Dayan, U., Ziv, B., Morin, E., & Enzel, Y. (2017). Climate of the Levant: Phenomena and Mechanisms. In Y. Enzel & B.-Y. Ofer (Eds.), *Quaternary of the Levant: Environments, climate change, and humans* (pp. 31–44). Cambridge University Press.
- Li, J., Wasko, C., Johnson, F., Evans, J. P., & Sharma, A. (2018). Can regional climate modeling capture the observed changes in spatial organization of extreme storms at higher temperatures? *Geophysical Research Letters*, 45(9), 4475–4484. <https://doi.org/10.1029/2018GL077716>
- Lochbihler, K., Lenderink, G., & Siebesma, A. P. (2017). The spatial extent of rainfall events and its relation to precipitation scaling. *Geophysical Research Letters*, 44(16), 8629–8636. <https://doi.org/10.1002/2017GL074857>
- Lochbihler, K., Lenderink, G., & Siebesma, A. P. (2019). Response of extreme precipitating cell structures to atmospheric warming. *Journal of Geophysical Research: Atmospheres*, 124(13), 6904–6918. <https://doi.org/10.1029/2018JD029954>
- Loft, R. (2020). Earth system modeling must become more energy efficient. *Eos*. <https://doi.org/10.1029/2020EO147051>
- Marra, F., Armon, M., Adam, O., Zoccatelli, D., Gazal, O., Garfinkel, C. I., et al. (2021). Toward narrowing uncertainty in future projections of local extreme precipitation. *Geophysical Research Letters*, 48(5), e2020GL091823. <https://doi.org/10.1029/2020GL091823>
- Marra, F., & Morin, E. (2015). Use of radar QPE for the derivation of intensity–duration–frequency curves in a range of climatic regimes. *Journal of Hydrology*, 531, 427–440. <https://doi.org/10.1016/j.jhydrol.2015.08.064>
- Marra, F., & Morin, E. (2018). Autocorrelation structure of convective rainfall in semiarid-arid climate derived from high-resolution X-Band radar estimates. *Atmospheric Research*, 200, 126–138. <https://doi.org/10.1016/j.atmosres.2017.09.020>
- Marra, F., Zoccatelli, D., Armon, M., & Morin, E. (2019). A simplified MEV formulation to model extremes emerging from multiple nonstationary underlying processes. *Advances in Water Resources*, 127, 280–290. <https://doi.org/10.1016/j.advwatres.2019.04.002>
- Meredith, E. P., Ulbrich, U., & Rust, H. W. (2020). Subhourly rainfall in a convection-permitting model. *Environmental Research Letters*, 15(3). <https://doi.org/10.1088/1748-9326/ab6787>
- Morin, E. (2011). To know what we cannot know: Global mapping of minimal detectable absolute trends in annual precipitation. *Water Resources Research*, 47(7), 1–9. <https://doi.org/10.1029/2010WR009798>
- Morrison, A., Villarini, G., Zhang, W., & Scoccimarro, E. (2019). Projected changes in extreme precipitation at sub-daily and daily time scales. *Global and Planetary Change*, 182, 103004. <https://doi.org/10.1016/j.gloplacha.2019.103004>
- Moustakis, Y., Papalexiou, S. M., Onof, C. J., & Paschalis, A. (2021). Seasonality, intensity, and duration of rainfall extremes change in a warmer climate. *Earth's Future*, 9(3), 1–15. <https://doi.org/10.1029/2020EF001824>
- Myhre, G., Alterskjær, K., Stjern, C. W., Hodnebrog Marelle, L., Samset, B. H., Sillmann, J., et al. (2019). Frequency of extreme precipitation increases extensively with event rareness under global warming. *Scientific Reports*, 9(1), 1–10. <https://doi.org/10.1038/s41598-019-52277-4>
- Nasta, P., Adane, Z., Lock, N., Houston, A., & Gates, J. B. (2018). Links between episodic groundwater recharge rates and rainfall events classified according to stratiform-convective storm scoring: A plot-scale study in eastern Nebraska. *Agricultural and Forest Meteorology*, 259, 154–161. <https://doi.org/10.1016/j.agrformet.2018.05.003>
- Nerini, D., Besic, N., Sideris, I., Germann, U., & Foresti, L. (2017). A non-stationary stochastic ensemble generator for radar rainfall fields based on the short-space Fourier transform. *Hydrology and Earth System Sciences*, 21(6), 2777–2797. <https://doi.org/10.5194/hess-21-2777-2017>

- O'Gorman, P. A. (2015). Precipitation extremes under climate change. *Current Climate Change Reports*, 1(2), 49–59. <https://doi.org/10.1007/s40641-015-0009-3>
- Peleg, N., Bartov, M., & Morin, E. (2014). CMIP5-predicted climate shifts over the East Mediterranean: Implications for the transition region between Mediterranean and semi-arid climates. *International Journal of Climatology*, 2153, 2144–2153. <https://doi.org/10.1002/joc.4114>
- Peleg, N., Ben-Asher, M., & Morin, E. (2013). Radar subpixel-scale rainfall variability and uncertainty: Lessons learned from observations of a dense rain-gauge network. *Hydrology and Earth System Sciences*, 17(6), 2195–2208. <https://doi.org/10.5194/hess-17-2195-2013>
- Peleg, N., Marra, F., Faticchi, S., Molnar, P., Morin, E., Sharma, A., & Burlando, P. (2018). Intensification of convective rain cells at warmer temperatures observed from high-resolution weather radar data. *Journal of Hydrometeorology*. <https://doi.org/10.1175/JHM-D-17-0158.1>
- Peleg, N., & Morin, E. (2012). Convective rain cells: Radar-derived spatiotemporal characteristics and synoptic patterns over the eastern Mediterranean. *Journal of Geophysical Research*, 117(15), 1–17. <https://doi.org/10.1029/2011JD017353>
- Pfahl, S., O'Gorman, P. A., & Fischer, E. M. (2017). Understanding the regional pattern of projected future changes in extreme precipitation. *Nature Climate Change*, 7(6), 423–427. <https://doi.org/10.1038/nclimate3287>
- Picard, L., & Mass, C. (2017). The sensitivity of orographic precipitation to flow direction: An idealized modeling approach. *Journal of Hydrometeorology*, 18(6), 1673–1688. <https://doi.org/10.1175/JHM-D-16-0209.1>
- Pichelli, E., Coppola, E., Sobolowski, S., Ban, N., Giorgi, F., Stocchi, P., et al. (2021). The first multi-model ensemble of regional climate simulations at kilometer-scale resolution part 2: Historical and future simulations of precipitation. *Climate Dynamics*, 56(11–12), 3581–3602. <https://doi.org/10.1007/s00382-021-05657-4>
- Poujol, B., Prein, A. F., & Newman, A. J. (2020). Kilometer-scale modeling projects a tripling of Alaskan convective storms in future climate. *Climate Dynamics*, 0123456789. <https://doi.org/10.1007/s00382-020-05466-1>
- Prein, A. F., Langhans, W., Fosser, G., Ferrone, A., Ban, N., Goergen, K., et al. (2015). A review on regional convection-permitting climate modeling: Demonstrations, prospects, and challenges. *Reviews of Geophysics*, 53(2), 323–361. <https://doi.org/10.1002/2014RG000475>
- Prein, A. F., Rasmussen, R. M., Ikeda, K., Liu, C., Clark, M. P., & Holland, G. J. (2017). The future intensification of hourly precipitation extremes. *Nature Climate Change*, 7(1), 48–52. <https://doi.org/10.1038/nclimate3168>
- Rasmussen, R., Liu, C., Ikeda, K., Gochis, D., Yates, D., Chen, F., et al. (2011). High-resolution coupled climate runoff simulations of seasonal snowfall over Colorado: A process study of current and warmer climate. *Journal of Climate*, 24(12), 3015–3048. <https://doi.org/10.1175/2010JCLI3985.1>
- Raveh-Rubin, S., & Wernli, H. (2016). Large-scale wind and precipitation extremes in the Mediterranean: Dynamical aspects of five selected cyclone events. *Quarterly Journal of the Royal Meteorological Society*, 142(701), 3097–3114. <https://doi.org/10.1002/qj.2891>
- Rinat, Y., Marra, F., Armon, M., Metzger, A., Levi, Y., Khain, P., et al. (2021). Hydrometeorological analysis and forecasting of a 3 d flash-flood-triggering desert rainstorm. *Natural Hazards and Earth System Sciences*, 21(3), 917–939. <https://doi.org/10.5194/nhess-21-917-2021>
- Romine, G. S., Schwartz, C. S., Snyder, C., Anderson, J. L., & Weisman, M. L. (2013). Model bias in a continuously cycled assimilation system and its influence on convection-permitting forecasts. *Monthly Weather Review*, 141(4), 1263–1284. <https://doi.org/10.1175/MWR-D-12-00112.1>
- Rostkier-Edelstein, D., Liu, Y., Wu, W., Kunin, P., Givati, A., & Ge, M. (2014). Towards a high-resolution climatology of seasonal precipitation over Israel. *International Journal of Climatology*, 34(6), 1964–1979. <https://doi.org/10.1002/joc.3814>
- Rubin, S., Ziv, B., & Paldor, N. (2007). Tropical plumes over Eastern North Africa as a source of rain in the Middle East. *Monthly Weather Review*, 135(12), 4135–4148. <https://doi.org/10.1175/2007MWR1919.1>
- Samuels, R., Hochman, A., Baharad, A., Givati, A., Levi, Y., Yosef, Y., et al. (2017). Evaluation and projection of extreme precipitation indices in the Eastern Mediterranean based on CMIP5 multi-model ensemble. *International Journal of Climatology*. <https://doi.org/10.1002/joc.5334>
- Samuels, R., Rimmer, A., & Alpert, P. (2009). Effect of extreme rainfall events on the water resources of the Jordan River. *Journal of Hydrology*, 375(3–4), 513–523. <https://doi.org/10.1016/j.jhydrol.2009.07.001>
- Sato, T., Kimura, F., & Kitoh, A. (2007). Projection of global warming onto regional precipitation over Mongolia using a regional climate model. *Journal of Hydrology*, 333(1), 144–154. <https://doi.org/10.1016/j.jhydrol.2006.07.023>
- Schär, C., Frei, C., Lüthi, D., & Davies, H. C. (1996). Surrogate climate-change scenarios for regional climate models. *Geophysical Research Letters*, 23(6), 669–672. <https://doi.org/10.1029/96GL00265>
- Schwartz, C. S., Romine, G. S., Sobash, R. A., Fossell, K. R., & Weisman, M. L. (2015). NCAR's experimental real-time convection-allowing ensemble prediction system. *Weather and Forecasting*, 30(6), 1645–1654. <https://doi.org/10.1175/WAF-D-15-0103.1>
- Seager, R., Osborn, T. J., Kushnir, Y., Simpson, I. R., Nakamura, J., & Liu, H. (2019). Climate variability and change of Mediterranean-type climates. *Journal of Climate*, 32(10), 2887–2915. <https://doi.org/10.1175/JCLI-D-18-0472.1>
- Shepherd, T. G. (2014). Atmospheric circulation as a source of uncertainty in climate change projections. *Nature Geoscience*, 7(10), 703–708. <https://doi.org/10.1038/NGEO2253>
- Shepherd, T. G. (2019). Storyline approach to the construction of regional climate change information. *Proceedings of the Royal Society A: Mathematical, Physical & Engineering Sciences*, 475(2225). <https://doi.org/10.1098/rspa.2019.0013>
- Shmilovitz, Y., Marra, F., Wei, H., Argaman, E., Nearing, M., Goodrich, D., et al. (2021). Frequency analysis of storm-scale soil erosion and characterization of extreme erosive events by linking the DWEPP model and a stochastic rainfall generator. *The Science of the Total Environment*, 787, 147609. <https://doi.org/10.1016/j.scitotenv.2021.147609>
- Sillmann, J., Shepherd, T. G., van den Hurk, B., Hazeleger, W., Martius, O., Slingo, J., & Zscheischler, J. (2021). Event-based storylines to address climate risk. *Earth's Future*, 9(2), 1–6. <https://doi.org/10.1029/2020EF001783>
- Tarasova, L., Merz, R., Kiss, A., Basso, S., Blöschl, G., Merz, B., et al. (2019). Causative classification of river flood events. *WIREs Water*, 6(4), 1–23. <https://doi.org/10.1002/wat2.1353>
- Taylor, K. E., Stouffer, R. J., & Meehl, G. A. (2012). An overview of CMIP5 and the experiment design. *Bulletin of the American Meteorological Society*. <https://doi.org/10.1175/BAMS-D-11-00094.1>
- Taylor, R. G., Todd, M. C., Kongola, L., Maurice, L., Nahozya, E., Sanga, H., & Macdonald, A. M. (2013). Evidence of the dependence of groundwater resources on extreme rainfall in East Africa. *Nature Climate Change*, 3(4), 374–378. <https://doi.org/10.1038/nclimate1731>
- Tebaldi, C., & Knutti, R. (2007). The use of the multi-model ensemble in probabilistic climate projections. *Philosophical Transactions of the Royal Society A: Mathematical, Physical & Engineering Sciences*, 365(1857), 2053–2075. <https://doi.org/10.1098/rsta.2007.2076>
- Tewari, M., Chen, F., Wang, W., Dudhia, J., LeMone, M. A., Mitchell, K., et al. (2004). Implementation and verification of the unified NOAA land surface model in the WRF model (Formerly Paper Number 17.5). In *20th Conference on weather analysis and forecasting/16th Conference on numerical weather prediction* (pp. 11–15).
- Thompson, G., Field, P. R., Rasmussen, R. M., & Hall, W. D. (2008). Explicit forecasts of winter precipitation using an improved bulk microphysics scheme. Part II: Implementation of a new snow parameterization. *Monthly Weather Review*, 136(12), 5095–5115. <https://doi.org/10.1175/2008MWR2387.1>

- Tiedtke, M. (1989). A comprehensive mass flux scheme for cumulus parameterization in large-scale models. *Monthly Weather Review*, *117*(8), 1779–1800. [https://doi.org/10.1175/1520-0493\(1989\)117<1779:ACMFSF>2.0.CO;2](https://doi.org/10.1175/1520-0493(1989)117<1779:ACMFSF>2.0.CO;2)
- Trenberth, K. E., Fasullo, J. T., & Shepherd, T. G. (2015). Attribution of climate extreme events. *Nature Climate Change*, *5*(8), 725–730. <https://doi.org/10.1038/nclimate2657>
- Tubi, A., Dayan, U., & Lensky, I. M. (2017). Moisture transport by tropical plumes over the Middle East: A 30-year climatology. *Quarterly Journal of the Royal Meteorological Society*, *143*(709), 3165–3176. <https://doi.org/10.1002/qj.3170>
- Tuel, A., & Eltahir, E. A. B. (2020). Why is the Mediterranean a climate change hot spot? *Journal of Climate*, *33*(14), 5829–5843. <https://doi.org/10.1175/JCLI-D-19-0910.1>
- Tuel, A., O’Gorman, P. A., & Eltahir, E. A. B. (2021). Elements of the dynamical response to climate change over the Mediterranean. *Journal of Climate*, *34*(3), 1135–1146. <https://doi.org/10.1175/JCLI-D-20-0429.1>
- Warner, T. T. (2011). *Numerical weather and climate prediction*. Cambridge University Press. <https://doi.org/10.1017/CBO9780511763243>
- Wasko, C., Sharma, A., & Westra, S. (2016). Reduced spatial extent of extreme storms at higher temperatures. *Geophysical Research Letters*, *43*(8), 4026–4032. <https://doi.org/10.1002/2016GL068509>
- Westra, S., Fowler, H. J., Evans, J. P., Alexander, L. V., Berg, P., Johnson, F., et al. (2014). Future changes to the intensity and frequency of short-duration extreme rainfall. *Reviews of Geophysics*, *52*, 522–555. <https://doi.org/10.1002/2014RG000464>
- Zappa, G., Hawcroft, M. K., Shaffrey, L., Black, E., & Brayshaw, D. J. (2015). Extratropical cyclones and the projected decline of winter Mediterranean precipitation in the CMIP5 models. *Climate Dynamics*, *45*(7–8), 1727–1738. <https://doi.org/10.1007/s00382-014-2426-8>
- Ziv, B., Dayan, U., Kushnir, Y., Roth, C., & Enzel, Y. (2006). Regional and global atmospheric patterns governing rainfall in the southern Levant. *International Journal of Climatology*, *26*(1), 55–73. <https://doi.org/10.1002/joc.1238>