

LYMPHOID NEOPLASIA

T $\gamma\delta$ LGLL identifies a subset with more symptomatic disease: analysis of an international cohort of 137 patients

Gregorio Barilà,^{1,2} Angela Grassi,³ HeeJin Cheon,⁴ Antonella Teramo,^{1,2} Giulia Calabretto,^{1,2} Jasmanet Chahal,⁴ Cristina Vicenzetto,^{1,2} Julia Almeida,⁵ Bryna C. Shemo,⁴ Min Shi,⁶ Vanessa Rebecca Gasparini,^{1,2} Noemi Munoz-Garcia,⁵ Cédric Pastoret,⁷ Hideyuki Nakazawa,⁸ Kazuo Oshimi,⁹ Lubomir Sokol,¹⁰ Fumihiro Ishida,⁸ Thierry Lamy,⁷ Alberto Orfao,⁵ William G. Morice,⁶ Thomas P. Loughran Jr,⁴ Gianpietro Semenzato,^{1,2} and Renato Zambello^{1,2}

¹Department of Medicine, Hematology and Clinical Immunology Section, Padua University School of Medicine, Padua, Italy; ²Veneto Institute of Molecular Medicine, Padua, Italy; ³Immunology and Molecular Oncology Unit, Veneto Institute of Oncology, IOV-IRCCS, Padua, Italy; ⁴Department of Medicine and University of Virginia Cancer Center, Division of Hematology & Oncology, University of Virginia School of Medicine, Charlottesville, VA; ⁵Department of Medicine and Cytometry Service, Cancer Research Center (Institute of Molecular and Cell Biology of Cancer–Consejo Superior de Investigaciones Científicas/Universidad de Salamanca), University of Salamanca, Instituto de Investigación Biomédica de Salamanca and Centro de Investigación Biomédica en Red en Oncología, Salamanca, Spain; ⁶Department of Laboratory Medicine and Pathology, Mayo Clinic, Rochester, NY; ⁷Department of Hematology, Rennes University Hospital, Rennes, France; ⁸Department of Biomedical Laboratory Sciences, Shinshu University School of Medicine, Matsumoto, Japan; ⁹Department of Medicine, Division of Hematology, Kushirosai Hospital, Kushirosai, Japan; and ¹⁰Department of Malignant Hematology, Moffitt Cancer Center, Tampa, FL

KEY POINTS

- **STAT3 mutations and V δ 2 status are needed to properly stratify T $\gamma\delta$ LGLL patients.**
- **Independently from STAT3 mutations, T $\gamma\delta$ LGLL represents a subset of T-LGLL characterized by dismal outcome as compared with T $\alpha\beta$ LGLL.**

T $\gamma\delta$ large granular lymphocyte leukemia (LGLL) is a rare variant of T-cell LGLL (T-LGLL) that has been less investigated as compared with the more frequent T $\alpha\beta$ LGLL, particularly in terms of frequency of STAT3 and STAT5b mutations. In this study, we characterized the clinical and biological features of 137 patients affected by T $\gamma\delta$ LGLL; data were retrospectively collected from 1997 to 2020 at 8 referral centers. Neutropenia and anemia were the most relevant clinical features, being present in 54.2% and 49.6% of cases, respectively, including severe neutropenia and anemia in ~20% of cases each. Among the various treatments, cyclosporine A was shown to provide the best response rates. DNA samples of 97 and 94 cases were available for STAT3 and STAT5b mutation analysis, with 38.1% and 4.2% of cases being mutated, respectively. Clinical and biological features of our series of T $\gamma\delta$ cases were also compared with a recently published T $\alpha\beta$ cohort including 129 cases. Though no differences in STAT3 and STAT5b mutational frequency were found,

T $\gamma\delta$ cases more frequently presented with neutropenia ($P = .0161$), anemia ($P < .0001$), severe anemia ($P = .0065$), and thrombocytopenia ($P = .0187$). Moreover, V δ 2⁻ cases displayed higher frequency of symptomatic disease. Overall, T $\gamma\delta$ cases displayed reduced survival with respect to T $\alpha\beta$ cases ($P = .0017$). Although there was no difference in STAT3 mutation frequency, our results showed that T $\gamma\delta$ LGLL represents a subset of T-LGLL characterized by more frequent symptoms and reduced survival as compared with T $\alpha\beta$ LGLL.

Introduction

Large granular lymphocyte leukemia (LGLL) is a rare and heterogeneous chronic lymphoproliferative disorder characterized by the clonal expansion of large granular lymphocytes (LGLs).^{1,2} The etiology of LGLL is unknown, but a constitutive activation of JAK/STAT pathway is involved in the pathogenesis of LGL proliferation,³ further supported by the discovery of somatic STAT3 and STAT5b mutations in ~40% of patients.⁴⁻⁹ Among LGLs, the latest World Health Organization classification recognizes a CD3⁺ T-cell LGLL (T-LGLL) and CD3⁻ natural killer (NK)-LGLL, accounting for 85% and 15% of cases, respectively. Moreover, based on surface

T-cell receptor expression, T $\alpha\beta$ and T $\gamma\delta$ subsets of LGLL can be identified.¹⁰

Although LGLL incidence ranges between 0.2 and 0.72 cases per 1 million individuals per year,¹ the frequency of T $\gamma\delta$ proliferation is still not well defined, and most information has been collected through small retrospective studies. As compared with the more frequent T $\alpha\beta$ LGLL, T $\gamma\delta$ LGLL has been less investigated. First reported by Oshimi et al in 1988 in a 60-year-old woman exposed to the radiation in Nagasaki in 1945,¹¹ T $\gamma\delta$ LGLL has been described in a sizable number of patients in 2006 by Sandberg et al, who reported an immunophenotypical analysis

of 44 cases.¹² Up to now, only 4 retrospective studies including more than 200 LGLL patients are available^{7,13-15}; however, few cases of T $\gamma\delta$ LGLL were included and only in the Italian cohort.⁷ Consequently, the clinical features of T $\gamma\delta$ LGLL and information on the efficacy of treatments in this LGLL variant are still missing. Furthermore, data on the frequencies of *STAT3* and *STAT5b* mutations are nowadays available for T $\alpha\beta$ LGLL, but still limited and controversial for T $\gamma\delta$ LGLL. The Italian group recently reported 25% and 19% of T $\gamma\delta$ cases mutated in *STAT3* and *STAT5b* genes, respectively,⁷ and *STAT3* mutations were found in all patients included in a small Japanese T $\gamma\delta$ LGLL cohort.¹⁶

With this as a background and lacking large cohorts of T $\gamma\delta$ patients, major referral groups dealing with LGLL were invited to join this collaborative study aimed at better characterizing T $\gamma\delta$ LGLL patients, pointing to the evaluation of putative correlations among mutations, phenotype, and clinical presentation, and the comparison of the clinical behavior of T $\gamma\delta$ LGLL with respect to the more common T $\alpha\beta$ variant. This large series of cases for the first time shows the dismal outcome of T $\gamma\delta$ LGLL with respect to T $\alpha\beta$ LGLL.

Methods

Study patients

The study cohort included 137 patients affected by T $\gamma\delta$ LGLL who were followed from 1997 to 2020 at 8 referral centers across the world (France, Italy, Japan, Spain, United States). All patients met the currently approved World Health Organization diagnostic criteria for T-LGLL.^{2,17} T-LGL clonality was assessed by TCR γ gene rearrangement.

Demographic and clinical features, including presence of cytopenias, concomitant autoimmune/inflammatory diseases, secondary primary malignancies (SPMs), treatment requirement, and response, were collected. Response to treatment was evaluated based on periodical clinical and laboratory examinations after at least 4 to 6 months of therapy, using the currently accepted response criteria for LGLL.¹⁸ The frequency of LGLs positive for the characteristic antigens was assessed by flow cytometry using direct immunofluorescence assays combining up to 6 markers per tube, according to standard operating procedures of individual centers. The investigation for LGL surface markers was performed on whole peripheral blood anticoagulated with EDTA or anticoagulant citrate dextrose and on purified peripheral blood mononuclear cells. The commercially available fluorescein isothiocyanate-conjugated; phycoerythrin (PE)-, PE-Cy5-, and PE-Cy7-conjugated; and allophycocyanin- and allophycocyanin-Cy7-conjugated mouse monoclonal antibodies used included anti-CD3, anti-CD4, anti-CD8, anti-CD16, anti-CD56 and anti-CD57, anti-TCR $\gamma\delta$, anti-KIRs (killer immunoglobulin-like receptors: CD158a, CD158b, CD158e), anti-NKG2A, anti-NKG2C, anti-V γ 9, anti-V δ 1, and anti-V δ 2 from Becton Dickinson (Sunnyvale, CA).

This international T $\gamma\delta$ LGL leukemia cohort was compared with a recently reported equal-size Italian T $\alpha\beta$ LGL leukemia cohort.⁷

This study was performed according to the Helsinki Declaration, and patients gave their written informed consent prior to inclusion in the study. The protocol and informed consent form were approved by the Padua ethics committee (approval number 4213/AO/17).

Screening for *STAT3* and *STAT5b* mutations

STAT3 and *STAT5b* sequencing was performed by Sanger Sequencing or Next Generation Sequencing according to local practice. For the screening of *STAT3* and *STAT5b* mutations by Sanger Sequencing, we used the set of primers reported by Koskela et al⁴ and by Rajala et al,⁸ respectively, to amplify the hot spot regions for mutations (exons 19-21 for *STAT3* and exons 16-18 for *STAT5b*).

Statistical analysis

Patients' demographic, clinical, and biological features expressed as categorical variables were compared using the Fisher exact test. Patient overall survival (OS) was calculated from the date of diagnosis to death by any cause or the last-known follow-up visit for censored patients. Survival curves were estimated using the Kaplan-Meier method and compared with respect to the patients' demographic and clinical characteristics using the log-rank test. Schoenfeld residual testing was applied to assess the proportional hazards assumption. A univariate Cox proportional hazards regression analysis was employed to evaluate the prognostic relevance of each variable. Results for significant variables were presented as hazard ratios (HRs) and 95% confidence intervals (CIs).

To determine the effect of response to first-line treatment on progression-free survival (PFS) and OS, we performed a 6-month landmark analysis in treated patients categorized by their response status (at least partial response vs stable disease or progressive disease) at 6 months after the start of therapy. The 6-month landmark time was selected a priori, before the beginning of data analysis, since at least 4 to 6 months of treatment are recommended before correctly assessing the response. For landmark analyses, PFS and OS were recalculated by shifting the time origin to 6 months after the start of therapy, and patients who experienced the event of progression or death before this time were excluded from the PFS or OS landmark analyses, respectively.

A restricted mean survival time (RMST) analysis was also performed to compare the T $\gamma\delta$ and T $\alpha\beta$ LGLL cohorts. RMST is a robust and clinically interpretable summary measure of the survival time distribution, estimable even under heavy censoring and when the proportional hazards assumption is not satisfied, as an alternative to the HR approach.^{19,20} This analysis depends on the truncation time point fixed for the RMST calculation. Four different truncation time points (100, 120, 140, and 160 months) were evaluated for the comparison of T $\gamma\delta$ and T $\alpha\beta$ LGLL cohorts. *P* values < .05 were considered significant. Statistical analysis was conducted using R version 3.6.2.

Results

Clinical and immunophenotypic features of T $\gamma\delta$ LGLL patients

Clinical and biological features of cases under study are summarized in supplemental Table 1, available on the *Blood* website. Median age at diagnosis was 58.5 years (range, 18-92), with 29.4% of subjects being >65 years old. No relevant gender prevalence was clearly demonstrated (male 55.9%, female 44.1%). By immunophenotype, all cases showed an expansion of CD3⁺ TCR $\gamma\delta$ ⁺ T cells, demonstrated to be clonal on

molecular grounds. T $\gamma\delta$ LGLs usually displayed CD8 positivity (64/105, 61.0%), with 23 of 105 (21.9%) cases showing partial CD8 expression; otherwise, CD4 was mostly absent, with only 3 cases showing partial expression. CD16 and CD57 were typical LGL markers, and they were expressed on the expanded T $\gamma\delta$ cells at the highest frequency (72.3% and 78.4%, respectively); CD56 was present in 31.1% of cases. A dominant KIR expression was demonstrated in 23 of 56 cases (41.1%), with CD158b being the most frequently expressed marker (13/56, 23.2%), followed by CD158a (8/56, 14.3%) and CD158e (5/56, 8.9%). CD94/NKG2 receptor expression was found in 32 of 75 cases (42.7%), with 12 cases displaying NKG2A (12/54, 22.2%) and 3 cases showing NKG2C positivity (3/30, 10%).

Not being part of the workup for the diagnosis of LGLL, bone marrow evaluation, either by flow cytometry or immunohistochemistry, was available for only 40 of 137 (29.2%) cases, showing variable degree of infiltration with a range from less than 1% to 60% of bone marrow cellularity.

Neutropenia (absolute neutrophil count [ANC] < 1500/mm³) and mild anemia (hemoglobin [Hb] < 120g/L) were the main relevant clinical features of the entire cohort, being present in 54.2% (65/120) and 49.6% (59/119) of cases, respectively. Severe neutropenia (ANC < 500/mm³) and severe anemia (Hb < 90 g/L) were observed in 25 of 120 cases (20.8%) and in 25 of 119 cases (21%), respectively. Thrombocytopenia (platelets [PLTs] < 100 000/mm³) and splenomegaly were detected in 18 of 119 (15.1%) and in 31 of 122 (21.4%) cases, respectively. Forty-nine cases (41.5%) were affected by concurrent autoimmune/inflammatory diseases, mostly rheumatoid arthritis (16/49), autoimmune hemolytic anemia (5/49), and pure red cell aplasia (PRCA) (5/49). Finally, SPMs were detected in 17 of 84 cases (20.2%), either at the time of diagnosis or during the follow-up. Seven SPMs were hematological (3 marginal zone lymphoma, 1 chronic lymphocytic leukemia, 1 myelodysplastic syndrome, 1 plasma cell dyscrasia, and 1 systemic mastocytosis) and 10 were nonhematological neoplasms, including 3 cases of thymoma, 3 cases of thyroid neoplasms, 1 lung cancer, 1 prostatic cancer, 1 cervical cancer, and 1 skin cancer.

Treatment of T $\gamma\delta$ patients

Overall, more than half (53.7%) of patients required therapy during the natural history of the disease. All these patients were treated according to currently accepted indications.^{2,21} In detail,

8 of 58 (13.8%) patients started therapy due to severe neutropenia, 4 of 58 (6.9%) due to symptomatic neutropenia, 14 of 58 (24.1%) for transfusion-dependent anemia, 13 of 58 (22.4%) for symptomatic anemia, 6 of 58 (10.4%) due to combined severe neutropenia and symptomatic anemia, and the remaining 5 of 58 (8.6%) for symptomatic concomitant autoimmune diseases. In 8 patients (13.8%) the primary diagnosis was settled by hematology centers without experience in LGLL, and subsequently the patients were moved to the referral centers. Consequently, a clear treatment indication was not available.

Considering first-line treatment, most patients (34/57, 59.6%) received methotrexate (MTX), 26.3% (15/57) were treated with cyclosporine A (CyA), and only 10.5% (6/57) received cyclophosphamide (CTX). The remaining 2 patients received cladribine and splenectomy as first-line treatment. Response rates and the absolute numbers of cases are reported in supplemental Figure 1 and Table 1. Overall response (ORR) and complete response (CR) rates were lower in MTX-treated patients (26.9% and 7.7%, respectively) compared with patients who received CyA and CTX (ORR: 53.9% and 40%, respectively; CR: 23.1% and 40%, respectively), although the latter therapies were used in lower numbers of cases, particularly CTX. Four patients treated with MTX discontinued the treatment due to toxicity.

Among patients requiring treatment (n = 57), landmark analyses for PFS and OS were performed according to response status at 6 months since therapy initiation, only in the subsets of patients for whom precise timing of response was available (n = 20 for PFS and n = 29 for OS). Irrespective from the type of first-line treatment, responders (patients reaching at least partial response) after 6 months from the start of therapy were characterized by an increase in PFS with respect to nonresponders (HR = 6.16, 95% CI: 0.77-50.00; log-rank test P = .05) (Figure 1A). Notably, although with a P value not statistically significant, responders at 6 months showed also longer OS as compared with nonresponders (log-rank test P = .13) (Figure 1B). These results suggest a possible prognostic role of early response to first-line therapy that should be further addressed in future prospective studies by systematically collecting response times.

STAT3 and STAT5b mutation analysis


DNA samples of 97 and 94 cases were available for STAT3 and STAT5b mutational analyses, respectively. STAT3 mutations

Table 1. Response to first-line treatment

Treatment	n	Response rate to first-line therapy, n/N (%)			Treatment interruption
		ORR	CR	N/A	Toxicity
MTX	34	7/26 (26.9)	2/26 (7.7)	4	4
CyA	15	7/13 (53.9)	3/13 (23.1)	2	—
CTX	6	2/5 (40)	2/5 (40)	1	—
Cladribine	1	1/1 (100)	0/1 (0)	—	—
Splenectomy	1	0/1 (0)	0/1 (0)	—	—

N/A, not available.

Figure 1. PFS and OS landmark analysis of patients treated for T γ δ LGLL. Kaplan-Meier curves showing 6-month landmark analysis for PFS (A) and OS (B) of T γ δ LGLL patients achieving at least a partial response to first-line therapy (Responders) compared with nonresponding patients (Non-responders) at 6 months from the start of therapy. Curves were compared by log-rank test.



were detected in 37 cases (38.1%), with a prevalence of variants as follows: Y640F was detected in 16 cases (43.2%), D661Y in 9 cases (24.4%), D661V and S614R in 2 cases each (5.4%), and the H410R, Q448E, G618R, E638Q, K658F, and N647I variants were found in 1 case each (2.7%). In the Italian cohort, 2 cases showed the A662_N663delinsH deletion and insertion and an in-frame insertion, G656_Y657ins, as previously reported.⁷ In contrast, *STAT5b* mutations were found in only 4 cases (4.2%), of whom 3 carried the N642H variant and 1 had the Y665F mutation. Of note, *STAT3* and *STAT5b* mutations were mutually exclusive in T γ δ LGLL cases, never being detected concurrently in the same patient.

From the phenotypic point of view, cases with *STAT3* mutations were characterized by lower frequency of expression of CD56 (3.8% vs 56.1%, $P < .0001$), V δ 2 (0% vs 50%, $P = .0003$), and V γ 9

(25% vs 57.1%, $P = .04$). In addition, they showed a higher frequency of neutropenia (65.7% vs 40.8%, $P = .0288$), severe neutropenia (31.4% vs 12.2%, $P = .0519$), anemia (55.9% vs 34.7%, $P = .0726$), and autoimmune/autoinflammatory disorders (59.4% vs 31.5%, $P = .0139$). They more frequently required therapy (67.9% vs 37.5%, $P = .0169$) (Table 2). In contrast, no significant differences were found between *STAT3*-mutated and wild-type T γ δ LGLL patients regarding the frequency of cases with LGL counts $> 2000/\text{mm}^3$ (25% vs 15.2%, $P = .3824$), expression of KIRs (20% vs 50%, $P = .1413$) and CD94 (38.9% vs 57.1%, $P = .2542$), thrombocytopenia (17.6% vs 14.3%, $P = .7628$), splenomegaly (22.9% vs 20.8%, $P > .9999$), and SPM (21.4% vs 22.7%, $P > .9999$) (Table 2). Unlike cases with *STAT3* mutations, cases with *STAT5b* mutations were mostly asymptomatic, with only 1 case experiencing mild neutropenia and splenomegaly.

Table 2. Biological and clinical features of STAT3-mutated and STAT3 wild-type T γ δ LGLL patients

	STAT3 mutated, n/N (%)	STAT3 wild-type, n/N (%)	P value
Age > 65 y	15/37 (40.5)	11/59 (18.6)	.0324
LGL > 2000/mm ³	8/32 (25.0)	7/46 (15.2)	.3824
CD16 expression	18/27 (66.7)	29/40 (72.5)	.7860
CD56 expression	1/26 (3.8)	23/41 (56.1)	<.0001
CD57 expression	21/27 (77.8)	34/41 (82.9)	.7541
KIR expression	2/10 (20.0)	12/24 (50.0)	.1413
CD158a expression	0/10 (0)	3/24 (12.5)	.5388
CD158b expression	1/10 (10.0)	8/24 (33.3)	.2250
CD158e expression	1/10 (10.0)	3/24 (12.5)	>.9999
CD94 expression	7/18 (38.9)	20/35 (57.1)	.2542
NKG2A expression	1/10 (10.0)	10/24 (41.7)	.1133
NKG2C expression	1/8 (12.5)	2/22 (9.1)	>.9999
V δ 1 ⁺	7/12 (58.3)	9/24 (37.5)	.2983
V δ 2 ⁺	0/17 (0)	17/34 (50.0)	.0003
V γ 9 ⁺	4/16 (25.0)	20/35 (57.1)	.0400
ANC < 1500/mm ³	23/35 (65.7)	20/49 (40.8)	.0288
ANC < 500/mm ³	11/35 (31.4)	6/49 (12.2)	.0519
Hb < 120 g/L	19/34 (55.9)	17/49 (34.7)	.0726
Hb < 90 g/L	10/34 (29.4)	6/49 (12.2)	.0874
PLTs < 100 000/mm ³	6/34 (17.6)	7/49 (14.3)	.7628
Splenomegaly	8/35 (22.9)	11/53 (20.8)	>.9999
Autoimmune/inflammatory diseases	19/32 (59.4)	17/54 (31.5)	.0139
SPMs	6/28 (21.4)	10/44 (22.7)	>.9999
Need for treatment	19/28 (67.9)	18/48 (37.5)	.0169

P values are calculated using Fisher exact test. Significant P values are reported in bold.

V δ pattern of expression analysis

T γ δ cells usually express 5 different V δ receptor families (from V δ 1 to V δ 5), V δ 2 being generally expressed in blood circulating T γ δ cells, and the other subsets are typically enriched in epithelia, liver, and spleen.²² In our cohort, flow cytometric V δ analysis was available in 51 cases; 17 cases (33.3%) were V δ 2⁺ and the remaining 34 (66.7%) were V δ 2⁻. Within this latter subset of cases, 16 of 34 (47.1%) were V δ 1⁺ and 18 cases were neither V δ 1⁺ nor V δ 2⁺ (Table 3).

V δ 2⁺ cases displayed a higher frequency of expression of CD56 (100% vs 9.1%, $P < .0001$), KIR (64.3% vs 18.8%, $P = .0236$), CD94 (76.5% vs 42.9%, $P = .0351$), and NKG2A (71.4% vs 6.2%, $P = .0004$), and no significant differences were found (vs V δ 2⁻ cases) regarding CD16 and CD57 expression (100% vs 81.8%, $P = .1412$, and 100% vs 81.8%, $P = .1412$, respectively). Interestingly, all V δ 2⁺ cases showed concomitant V γ 9 expression (100%), and only a small fraction of V δ 2⁻ cases was also V γ 9⁺ (18.2%, $P < .0001$).

From the clinical point of view, V δ 2⁺ cases displayed a more indolent LGLL. They rarely presented with symptomatic disease including neutropenia (5.9% vs 65.6%, $P < .0001$), severe neutropenia (0% vs 31.2%, $P = .0094$), anemia (0% vs 56.2%, $P < .0001$), severe anemia (0% vs 34.4%, $P = .0090$), splenomegaly (0% vs 26.7%, $P = .0371$), and concurrent autoimmune/inflammatory disease (6.2% vs 48.4%, $P = .0039$), in the absence of treatment requirement (0% vs 54.5%, $P = .0007$). Interestingly, STAT mutations were mutually exclusive in V δ 2⁻ and V δ 2⁺ cases, all cases with STAT5b mutation being V δ 2⁺ ($P = .0327$), whereas all cases with STAT3 mutations were V δ 2⁻ ($P = .0003$) (Table 3).

Clinical and biological features of T γ δ vs T α β LGLL

To get further insight into the unique clinical and biological features of T γ δ LGLL, we compared our cohort of patients with a recently published T α β LGLL cohort of comparable size⁷ (Table 4). No significant differences in gender and age were

Table 3. Biological and clinical features of T γ δ LGLL patients according to V δ 2 status

	V δ 2 ⁺ , n/N (%)	V δ 2 ⁻ , n/N (%)	P value
LGL > 2000/mm ³	2/17 (11.8)	9/29 (31.0)	.1723
CD16 expression	14/14 (100)	18/22 (81.8)	.1412
CD56 expression	14/14 (100)	2/22 (9.1)	<.0001
CD57 expression	14/14 (100)	18/22 (81.8)	.1412
KIR expression	9/14 (64.3)	3/16 (18.8)	.0236
CD158a expression	1/14 (7.1)	1/16 (6.2)	>.9999
CD158b expression	6/14 (42.9)	2/16 (12.5)	.1010
CD158e expression	3/14 (21.4)	1/16 (6.2)	.3155
CD94 expression	13/17 (76.5)	12/28 (42.9)	.0351
NKG2A expression	10/14 (71.4)	1/16 (6.2)	.0004
NKG2C expression	0/14 (0)	3/16 (18.8)	.2276
V γ 9 ⁺	17/17 (100)	6/33 (18.2)	<.0001
STAT3 mutated	0/17 (0)	17/34 (50)	.0003
STAT5b mutated	3/17 (17.6)	0/34 (0)	.0327
ANC < 1500/mm ³	1/17 (5.9)	21/32 (65.6)	<.0001
ANC < 500/mm ³	0/17 (0)	10/32 (31.2)	.0094
Hb < 120 g/L	0/17 (0)	18/32 (56.2)	<.0001
Hb < 90 g/L	0/17 (0)	11/32 (34.4)	.0090
PLTs < 100 000/mm ³	1/17 (5.9)	4/32 (12.5)	.6463
Splenomegaly	0/16 (0)	8/30 (26.7)	.0371
Autoimmune/inflammatory diseases	1/16 (6.2)	15/31 (48.4)	.0039
SPMs	4/14 (28.6)	6/24 (25)	>.9999
Need for treatment	0/14 (0)	12/22 (54.5)	.0007

P values are calculated using Fisher exact test. Significant P values are reported in bold.

found between the 2 disease subtypes ($P = .3906$ and $P = .2408$, respectively), while T $\alpha\beta$ LGLL cases generally showed higher LGL counts than T $\gamma\delta$ LGLL cases (LGL count > 2000/mm³ in 54.3% vs 22% cases, respectively; $P < .0001$). By immunophenotype, T $\gamma\delta$ LGLL displayed a significantly higher frequency of expression of CD16 (72.3% vs 45.7%, $P < .0001$), CD94 (42.7% vs 14%, $P < .0001$), NKG2A (22.2% vs 10.1%, $P = .0355$), and CD158a (14.3% vs 4.7%, $P = .0330$) together with an increased KIR expression (41.1% vs 27.9%, $P = .0876$), and they showed a lower frequency of CD56 (31.1% vs 48.1%, $P = .0106$) and CD57 expression (78.4% vs 94.6%, $P = .0003$). Regarding STAT mutations, no significant differences were found between T $\gamma\delta$ and T $\alpha\beta$ LGLL cases in the frequency of STAT3 (38.1% vs 37.9%, respectively; $P > .9999$) and STAT5b mutations (4.8% vs 12.5%, respectively; $P = .1130$).

From the clinical point of view, T $\gamma\delta$ LGLL cases more frequently showed symptomatic disease in terms of neutropenia (54.2% vs 38.8%, $P = .0161$), anemia (49.6% vs 11.6%, $P < .0001$), severe

anemia (21% vs 8.5%, $P = .0065$), thrombocytopenia (15.1% vs 5.4%, $P = .0187$), and concurrent autoimmune/inflammatory diseases (41.5% vs 21.7%, $P = .0009$) (Table 4).

The markedly different observation times of T $\gamma\delta$ -LGLL and T $\alpha\beta$ -LGLL cases prevented use of a Fisher exact test for the comparison of time-dependent factors since this could lead to major bias due to lack of consideration of the time variable. Consequently, for SPMs and need for treatment, the data and the related P value were not available.

Survival analysis

All demographic, clinical, and biological features were evaluated for association with OS in T $\gamma\delta$ LGLL cases. After a median follow-up of 48 months, the median OS of our cohort was not reached. Splenomegaly was the only variable significantly associated with a shortened OS (log-rank test $P = .0012$), with an HR = 0.18 (95% CI: 0.06-0.59) (Figure 2A), and other clinical and biological features of the disease had no significant impact on patient OS,

Table 4. Biological and clinical features of the Tαβ and the Tγδ LGLL cohorts

	Tαβ LGLL, * n/N (%)	Tγδ LGLL, n/N (%)	P value
Gender male	65/129 (50.4)	76/136 (55.9)	.3906
Age > 65 y	47/129 (36.4)	40/136 (29.4)	.2408
LGL > 2000/mm ³	70/129 (54.3)	24/109 (22.0)	<.0001
CD16 expression	59/129 (45.7)	73/101 (72.3)	<.0001
CD56 expression	62/129 (48.1)	32/103 (31.1)	.0106
CD57 expression	122/129 (94.6)	80/102 (78.4)	.0003
KIR expression	36/129 (27.9)	23/56 (41.1)	.0876
CD158a expression	6/129 (4.7)	8/56 (14.3)	.0330
CD158b expression	29/129 (22.5)	13/56 (23.2)	>.9999
CD158e expression	6/129 (4.7)	5/56 (8.9)	.3124
CD94 expression	18/129 (14.0)	32/75 (42.7)	<.0001
NKG2A expression	13/129 (10.1)	12/54 (22.2)	.0355
NKG2C expression	5/129 (3.9)	3/30 (10.0)	.1740
STAT3 mutated	39/103 (37.9)	37/97 (38.1)	>.9999
STAT5b mutated	12/96 (12.5)	4/94 (4.8)	.1130
ANC < 1500/mm ³	50/129 (38.8)	65/120 (54.2)	.0161
ANC < 500/mm ³	29/129 (22.5)	25/120 (20.8)	.7611
Hb < 120 g/L	15/129 (11.6)	59/119 (49.6)	<.0001
Hb < 90 g/L	11/129 (8.5)	25/119 (21.0)	.0065
PLTs < 100 000/mm ³	7/129 (5.4)	18/119 (15.1)	.0187
Splenomegaly	22/129 (17.5)	31/122 (21.4)	.1225
Autoimmune/inflammatory diseases	28/129 (21.7)	49/118 (41.5)	.0009

P values are calculated using Fisher exact test. Significant P values are reported in bold.

The markedly different observation times of Tγδ LGLL and Tαβ LGLL patients prevented use of Fisher exact test for the comparison of time-dependent factors since this could lead to major bias due to lack of consideration of the time variable. Consequently, for SPMs and need for treatment, the data and the related P value were not available.

*Tαβ LGLL cohort of comparable size.⁷

including those previously found to be relevant for Tαβ LGLL patients⁷ (ie, *STAT3* and *STAT5b* mutation status or the presence of severe neutropenia or anemia) (supplemental Figure 2).


Direct comparison of patients' OS between Tγδ LGLL and the more common Tαβ LGLL is likely to prove a poorer overall outcome for Tγδ LGLL cases vs Tαβ LGLL cases (log-rank test $P = .017$) (Figure 2B). This result must be interpreted with caution, since the 2 cohorts have different median follow-up times (Tγδ LGLL, 4 years, vs Tαβ LGLL, 9 years), and the proportional hazards assumption seems not to be fully satisfied due to the lack of events in the Tγδ cohort from 143 months onward. Given the rarity of Tγδ LGLL, it was not possible to increase the cohort size; consequently, we provided a supplementary analysis using a different measure of the effect that does not require the proportional hazards assumption (ie, the RMST). This analysis confirms a significant disadvantage in terms of survival of Tγδ LGLL patients with respect to Tαβ LGLL (supplemental Table 2).

Discussion

Here we report on the largest cohort of Tγδ LGL leukemia patients described so far in the literature with data collected between 1997 and 2020, as the result of a collaborative study involving 8 LGLL referral centers across the world. For the first time, we evaluated the clinical and biological features of this rare subset of T-LGLL on a large number of patients, screened for *STAT3* and *STAT5b* mutations. Overall, our results showed that Tγδ LGLL represents a variant with higher frequency of symptomatic disease and reduced survival compared with the most common Tαβ LGLL subtype, despite a similar frequency of *STAT3* and to a less extent of *STAT5b* mutations.

In the past, LGLL was considered a unique chronic and indolent disease, except for a few patients presenting with very aggressive disease.²³ In recent years, however, a better understanding of this disorder has been achieved, pointing out the need for therapy in a

Figure 2. OS analysis of T $\gamma\delta$ LGLL patients. (A) OS analysis of the T $\gamma\delta$ LGLL cohort with respect to presence/absence of splenomegaly. Survival curves were estimated using the Kaplan-Meier method and compared by log-rank test. (B) OS comparison between T $\alpha\beta$ and T $\gamma\delta$ cohorts. With a median follow-up of 108 months (T $\alpha\beta$) and of 48 months (T $\gamma\delta$), median OS was not reached in both the cohorts. Survival curves were estimated using the Kaplan-Meier method and compared by log-rank test.



significant fraction of LGLL patients.^{6,7,24} Data provided in this study further encourage distinguishing T $\gamma\delta$ LGLL from T $\alpha\beta$ LGLL, since T $\gamma\delta$ LGLL patients showed unique clinical and biological features. Even though characterized by lower LGL counts, T $\gamma\delta$ LGLs more frequently express the CD16 and CD94 receptors, and the CD56 adhesion molecule and the CD57 immunosenescence-associated protein are less commonly expressed. Most important, T $\gamma\delta$ LGLL patients more frequently displayed symptomatic disease due to anemia (often transfusion dependent), potentially partially explained by an increased frequency of autoimmune hemolytic anemia and PRCA,²⁵ and concomitant autoimmune diseases. Altogether, this translates into a poorer outcome as compared with that from the more common T $\alpha\beta$ subtype of LGLL.

Overall these results are not consistent with previously reported data that did not show clear clinical differences between T $\gamma\delta$ LGLL and T $\alpha\beta$ LGLL²⁶; however, the T-LGLL cohort reported by Bourgault-Rouxel et al included only a small number of T $\gamma\delta$ patients (20 cases) compared with the almost 200 T $\alpha\beta$ reported cases, which limits the robustness of the conclusions raised.²⁶ A possible limitation to be considered in the explanation of the worst outcome in T $\gamma\delta$ LGLL could be related to a high frequency of late-stage diseases due to the challenging diagnosis. As a matter of fact, in our series T $\gamma\delta$ patients showed lower LGL counts and CD57 expression as compared with the those in the more common T $\alpha\beta$ patients. The high frequency of symptomatic patients herein reported within the T $\gamma\delta$ LGLL cohort may

account for the reduced OS in this LGLL subtype. Aside from this potential bias in survival analysis, our data point to the recommendation to include the T $\gamma\delta$ immunophenotype in the diagnostic workup of unexplained cytopenia.

Despite the comparable size, the T $\gamma\delta$ and T $\alpha\beta$ LGLL cohorts we studied are characterized by different median follow-up (48 vs 108 months, respectively); moreover, the T $\gamma\delta$ LGLL cohort, due to its retrospective nature, suffers for the presence of several censored data. These findings led to certain limitations in the interpretation of results. For this reason, an additional RMST analysis has been provided to mitigate these limitations, confirming a significant survival disadvantage for T $\gamma\delta$ LGLL patients with respect to T $\alpha\beta$ LGLL. In future perspective studies aimed at comparing the 2 cohorts, it could be interesting to carefully plan the data collection to analyze variables that may depend on observation time (eg, SPM or need for treatment) with a more appropriate time-to-event approach, thus minimizing any bias due to different follow-up lengths.

According to retrospective studies including few and heterogeneous series of patients,²⁷⁻²⁹ treatment of LGLL still relies on immunosuppressive therapy, where MTX and CTX are used upfront, and CyA is generally reserved for relapsed or refractory patients.^{1,2,21} To date, only 1 published prospective trial evaluating the efficacy of immunosuppressive therapy in LGLL is available,³⁰ and 1 prospective and randomized trial comparing MTX and CTX as first-line therapy in LGLL is currently ongoing (NCT01976182). However, all these studies do not report on the frequency of T $\gamma\delta$ LGLL analyzed and their specific response to therapy. Unexpectedly, MTX treatment led to unsatisfactory response rates in our series of T $\gamma\delta$ LGLL patients, with ORR being observed in less than a third of patients, including CR in a very limited number of cases (7.7%). In contrast, first-line therapy with CyA turned out to provide higher efficacy, with almost half the patients responding, of whom 23.1% reached CR.

An association between T $\gamma\delta$ LGLL and PRCA has been widely described, and it is also known that PRCA patients benefit from CyA treatment. In our cohort, treatment indication for the CyA cohort was available for 14 patients, and 12 patients started therapy due to anemia, in 8 cases transfusion dependent; the remaining 2 patients had a concomitant diagnosis of PRCA. It can be argued that PRCA has been underestimated in T $\gamma\delta$ LGLL with anemia or severe anemia, thus explaining the high overall and CR rates obtained with CyA in this subgroup of patients. Of notice, the choice of the appropriate therapy is of utmost clinical relevance since we demonstrated here that responding patients were also characterized by a prolonged PFS and an improved OS. These data could offer a rationale for investigating CyA in the first-line treatment of T $\gamma\delta$ LGLL (eg, in new prospective trials).

Altogether, the results indicate that, besides the distinction between T-LGLL and natural killer-LGLL, further dissection of T-LGLL into the T $\alpha\beta$ and T $\gamma\delta$ LGLL disease variants is of clinical relevance due to the poorer outcome and distinct treatment response profile of the latter patients. It is also worth noting that T $\gamma\delta$ LGLL cases did not appear as a homogeneous disease entity. V δ 2 positivity was associated with an immunophenotype

characterized by V γ 9, CD56, KIR, and CD94/NKG2A expression and, on clinical grounds, by lower frequency of symptomatic disease in terms of neutropenia, anemia, splenomegaly concomitant autoimmune/inflammatory disease, and need of treatment compared with that of V δ 2⁻ patients. Furthermore, the V δ 2 expression profile also correlated with the STAT mutational status since all STAT3-mutated cases were V δ 2⁻, and the 3 patients with STAT5b mutations were V δ 2⁺. Interestingly, the 2 subsets of T $\gamma\delta$ LGLL defined by the V δ 2 expression profile are likely to identify distinct cells of origin of T $\gamma\delta$ LGLL.²² In line with this hypothesis, V δ 2⁺ T $\gamma\delta$ LGLL might represent the neoplastic counterpart of blood circulating T $\gamma\delta$ cells, and V δ 2⁻ T $\gamma\delta$ LGLL might mostly originate from tissue-derived T $\gamma\delta$ cells, with potential pathogenic implications.

Accumulating evidence indicates that the association between STAT3 mutation and symptomatic disease is already recognized in T $\alpha\beta$ LGLL.^{6,7,31} Recent data also support a reduced survival for STAT3-mutated vs STAT3 wild-type cases.⁷ In contrast, the clinical impact of STAT5b mutations is still matter of debate; this mutation is present in the rare aggressive variants of LGLL⁸ as well as in indolent CD4⁺ T-LGLL.^{7,32} In the T $\gamma\delta$ LGLL setting, the real incidence of STATs gene mutations is still unknown, being studied up to now only in small cohorts of patients.^{7,16,33} In our study, mutations in STAT3 and STAT5b were screened in nearly 100 T $\gamma\delta$ LGLL cases, and a frequency of STAT3 mutations was found to be comparable with previously reported data in LGLL.⁴⁻⁶ Moreover, we also detected 3 T $\gamma\delta$ LGLL cases harboring STAT5b mutations who displayed an indolent disease as observed in CD4⁺ T $\alpha\beta$ LGLL. In our cohort, we confirm the association between STAT3 mutation and symptomatic disease, particularly with neutropenia, and increased need for therapy, although we did not observe a reduced OS for STAT3-mutated cases. These results support a more aggressive disease behavior of T $\gamma\delta$ LGLL, particularly for cases who do not show V δ 2 expression, independently from the STAT3 mutational status.

In conclusion, data from this large multicentric cohort of T $\gamma\delta$ LGLL highlight the unique biological and clinical hallmarks of this rare variant of T-LGLL, likely associated with a discrete treatment response profile. Despite the similar frequency of STAT3 and STAT5b, T $\gamma\delta$ LGLL cases in general, and V δ 2⁻ T $\gamma\delta$ LGLL in particular, showed more symptomatic disease and a poorer outcome compared with those with T $\alpha\beta$ LGLL. At the same time, T $\gamma\delta$ LGLL patients appear to mostly benefit from CyA as first-line therapy. Altogether, these results underly the relevance of a precise characterization and subclassification of LGLL.

Acknowledgment

The authors thank the Associazione Italiana per la Ricerca sul Cancro (grant IG 2017-20216).

Authorship

Contribution: G.B. designed the research, analyzed data, and wrote the manuscript; A.G. analyzed data, performed statistical analysis, and wrote the manuscript; H.J.C., A.T., G.C., J.C., C.V., B.C.S., V.R.G., N.M.-G., H.N., and C.P. provided patient samples and patient data; J.A., M.S., K.O., L.S., F.I., T.P.L., A.O., W.G.M., and T.L. participated in the analysis of data and critically reviewed and edited the manuscript; G.S. provided

funding, participated in the analysis of data, and critically reviewed and edited the manuscript; R.Z. designed the study, analyzed data, wrote the manuscript, and supervised the study.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

ORCID profiles: G.B., 0000-0003-3430-6115; A.G., 0000-0002-8313-8384; H.C., 0000-0003-0149-9651; A.T., 0000-0002-2754-699X; J.C., 0000-0002-0397-6418; C.V., 0000-0002-2548-4596; J.A., 0000-0003-3124-8917; N.M.-G., 0000-0002-8381-301X; H.N., 0000-0002-6457-5052; W.G.M., 0000-0002-5801-2501; G.S., 0000-0002-6061-4595.

Correspondence: Renato Zambello, Padua University School of Medicine, Department of Medicine, Via Giustiniani 2, 35128 Padua, Italy; email: r.zambello@unipd.it; and Gianpietro Semenzato, University of Padua, Veneto Institute of Molecular Medicine, Via Orus 2, 35129 Padua, Italy; email: g.semenzato@unipd.it.

REFERENCES

1. Barila G, Calabretto G, Teramo A, et al. T cell large granular lymphocyte leukemia and chronic NK lymphocytosis. *Best Pract Res Clin Haematol*. 2019;32(3):207-216.
2. Lamy T, Moignet A, Loughran TP Jr. LGL leukemia: from pathogenesis to treatment. *Blood*. 2017;129(9):1082-1094.
3. Epling-Burnette PK, Liu JH, Catlett-Falcone R, et al. Inhibition of STAT3 signaling leads to apoptosis of leukemic large granular lymphocytes and decreased Mcl-1 expression. *J Clin Invest*. 2001;107(3):351-362.
4. Koskela HL, Eldfors S, Ellonen P, et al. Somatic STAT3 mutations in large granular lymphocytic leukemia. *N Engl J Med*. 2012;366(20):1905-1913.
5. Jerez A, Clemente MJ, Makishima H, et al. STAT3 mutations indicate the presence of subclinical T-cell clones in a subset of aplastic anemia and myelodysplastic syndrome patients. *Blood*. 2013;122(14):2453-2459.
6. Teramo A, Barila G, Calabretto G, et al. STAT3 mutation impacts biological and clinical features of T-LGL leukemia. *Oncotarget*. 2017;8(37):61876-61889.
7. Barilà G, Teramo A, Calabretto G, et al. Stat3 mutations impact on overall survival in large granular lymphocyte leukemia: a single-center experience of 205 patients. *Leukemia*. 2020;34(4):1116-1124.
8. Rajala HL, Eldfors S, Kuusanmaki H, et al. Discovery of somatic STAT5b mutations in large granular lymphocytic leukemia. *Blood*. 2013;121(22):4541-4550.
9. Teramo A, Barila G, Calabretto G, et al. Insights into genetic landscape of large granular lymphocyte leukemia. *Front Oncol*. 2020;10:152.
10. Alaggio R, Amador C, Anagnostopoulos I, et al. The 5th edition of the World Health Organization classification of

haematolymphoid tumours: lymphoid neoplasms. *Leukemia*. 2022;36(7):1720-1748.

11. Oshimi K, Hoshino S, Takahashi M, et al. T_H1 (WT31)-negative, CD3-positive, large granular lymphocyte leukemia with nonspecific cytotoxicity. *Blood*. 1988;71(4):923-931.
12. Sandberg Y, Almeida J, Gonzalez M, et al. TCR γ delta+ large granular lymphocyte leukemias reflect the spectrum of normal antigen-selected TCR γ delta+ T-cells. *Leukemia*. 2006;20(3):505-513.
13. Barea B, Rey J, Hamidou M, et al. Analysis of a French cohort of patients with large granular lymphocyte leukemia: a report on 229 cases. *Haematologica*. 2010;95(9):1534-1541.
14. Sanikommu SR, Clemente MJ, Chomczynski P, et al. Clinical features and treatment outcomes in large granular lymphocytic leukemia (LGL). *Leuk Lymphoma*. 2018;59(2):416-422.
15. Dong N, Castillo Tokumori F, Isenlumhe L, et al. Large granular lymphocytic leukemia—a retrospective study of 319 cases. *Am J Hematol*. 2021;96(7):772-780.
16. Yamane T, Kawakami T, Sekiguchi N, et al. High frequency of STAT3 gene mutations in T-cell receptor (TCR) γ delta-type T-cell large granular lymphocytic leukaemia: implications for molecular diagnostics. *Br J Haematol*. 2020;190(5):e301-e304.
17. Swerdlow SH, Campo E, Harris NL, et al. *WHO classification of tumours of haematopoietic and lymphoid tissues*. IARC Press; 2017.
18. Poullot E, Zambello R, Leblanc F, et al. Chronic natural killer lymphoproliferative disorders: characteristics of an international cohort of 70 patients. *Ann Oncol*. 2014;25(10):2030-2035.
19. Royston P, Parmar MK. Restricted mean survival time: an alternative to the hazard ratio for the design and analysis of

randomized trials with a time-to-event outcome. *BMC Med Res Methodol*. 2013;13:152.

20. Uno H, Claggett B, Tian L, et al. Moving beyond the hazard ratio in quantifying the between-group difference in survival analysis. *J Clin Oncol*. 2014;32(22):2380-2385.
21. Lamy T, Loughran TP Jr. How I treat LGL leukemia. *Blood*. 2011;117(10):2764-2774.
22. Lalor SJ, McLoughlin RM. Memory gamma delta T cells—newly appreciated antagonists in infection and immunity. *Trends Immunol*. 2016;37(10):690-702.
23. Semenzato G, Pandolfi F, Chisesi T, et al. The lymphoproliferative disease of granular lymphocytes. A heterogeneous disorder ranging from indolent to aggressive conditions. *Cancer*. 1987;60(12):2971-2978.
24. Barila G, Teramo A, Calabretto G, et al. Dominant cytotoxic NK cell subset within CLPD-NK patients identifies a more aggressive NK cell proliferation. *Blood Cancer J*. 2018;8(6):51.
25. Chen YH, Chadburn A, Evens AM, et al. Clinical, morphologic, immunophenotypic, and molecular cytogenetic assessment of CD4-/CD8-gamma delta T-cell large granular lymphocytic leukemia. *Am J Clin Pathol*. 2011;136(2):289-299.
26. Bourgault-Rouxel AS, Loughran TP Jr, Zambello R, et al. Clinical spectrum of gamma delta+ T cell LGL leukemia: analysis of 20 cases. *Leuk Res*. 2008;32(1):45-48.
27. Loughran TP Jr, Kidd PG, Starkebaum G. Treatment of large granular lymphocyte leukemia with oral low-dose methotrexate. *Blood*. 1994;84(7):2164-2170.
28. Moignet A, Hasanali Z, Zambello R, et al. Cyclophosphamide as a first-line therapy in LGL leukemia. *Leukemia*. 2014;28(5):1134-1136.
29. Battiwalla M, Melenhorst J, Sauntharajah Y, et al. HLA-DR4 predicts haematological

Footnotes

Submitted 4 August 2021; accepted 2 August 2022; prepublished online on *Blood* First Edition 12 September 2022. <https://doi.org/10.1182/blood.2021013489>.

For original data, please contact r.zambello@unipd.it or g.semenzato@unipd.it. In this research article, we compared the international T γ δ LGLL cohort with a recently published T α β LGLL cohort of comparable size (Barilà et al⁷).

The online version of this article contains a data supplement.

There is a [Blood Commentary](#) on this article in this issue.

The publication costs of this article were defrayed in part by page charge payment. Therefore, and solely to indicate this fact, this article is hereby marked "advertisement" in accordance with 18 USC section 1734.

- response to cyclosporine in T-large granular lymphocyte lymphoproliferative disorders. *Br J Haematol.* 2003;123(3):449-453.
30. Loughran TP Jr, Zickl L, Olson TL, et al. Immunosuppressive therapy of LGL leukemia: prospective multicenter phase II study by the Eastern Cooperative Oncology Group (E5998). *Leukemia.* 2015;29(4):886-894.
31. Munoz-Garcia N, Jara-Acevedo M, Caldas C, et al. STAT3 and STAT5B mutations in T/NK-cell chronic lymphoproliferative disorders of large granular lymphocytes (LGL): association with disease features. *Cancers (Basel).* 2020;12(12).
32. Andersson EI, Tanahashi T, Sekiguchi N, et al. High incidence of activating STAT5B mutations in CD4-positive T-cell large granular lymphocyte leukemia. *Blood.* 2016; 128(20):2465-2468.
33. Shi M, He R, Feldman AL, et al. STAT3 mutation and its clinical and histopathologic correlation in T-cell large granular lymphocytic leukemia. *Hum Pathol.* 2018;73:74-81.

© 2023 by The American Society of Hematology.
 Licensed under Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0), permitting only noncommercial, nonderivative use with attribution. All other rights reserved.