

Large $\nu_\mu \rightarrow \nu_\tau$ and $\nu_e \rightarrow \nu_\tau$ transitions in short-baseline experiments?

Carlo Giunti

*INFN, Sezione di Torino, and
Dipartimento di Fisica Teorica, Università di Torino
I-10125 Torino, Italy, and
School of Physics, Korea Institute for Advanced Study
Seoul 130-012, Korea
E-mail: giunti@to.infn.it*

Marco Laveder

*Dipartimento di Fisica “G. Galilei”, Università di Padova, and
INFN, Sezione di Padova
I-35131 Padova, Italy
E-mail: marco.laveder@pd.infn.it*

ABSTRACT: Considering four-neutrino schemes of type $3 + 1$, we identify four small regions of the neutrino mixing parameter space compatible with all data. Assuming a small mixing between the sterile neutrino and the isolated mass eigenstate we show that large $\nu_\mu \rightarrow \nu_\tau$ and $\nu_e \rightarrow \nu_\tau$ transitions are predicted in short-baseline experiments and could be observed in the near future in dedicated experiments. We discuss also implications for solar, atmospheric and long-baseline neutrino experiments and we present a formalism that allows to describe in $3 + 1$ schemes atmospheric neutrino oscillations, long-baseline ν_μ disappearance and $\nu_\mu \rightarrow \nu_\tau$ transitions in matter.

KEYWORDS: Solar and Atmospheric Neutrinos, Neutrino Physics.

Contents

1. Introduction	1
2. $3 + 1$ schemes with $U_{s4} ^2 \ll 1$	5
3. General bounds in $3 + 1$ schemes and $\nu_\mu \rightarrow \nu_e$ short-baseline transitions	5
4. Large $\nu_e \rightarrow \nu_\tau$ and $\nu_\mu \rightarrow \nu_\tau$ short-baseline transitions	8
5. Conclusions	13
A. Formalism of atmospheric neutrino oscillations	14

1. Introduction

Neutrino oscillation experiments are powerful probes of neutrino masses (see [1]–[6]). At present three types of neutrino oscillation experiments have obtained positive results: solar and atmospheric experiments and one short-baseline experiment.

All solar neutrino experiments (Homestake [7], Kamiokande [8], GALLEX [9], SAGE [10], Super-Kamiokande [11]) have found a deficit in the flux of electron neutrinos on Earth with respect to the Standard Solar Model prediction [12], which constitutes an indication in favor of oscillations of electron neutrinos into other states. Although no direct proof of these oscillations exist at present, there is a convincing evidence that the deficit of solar ν_e 's is due to neutrino physics, of which neutrino oscillations is the simplest and most natural phenomenon. Hopefully, the issue will be definitively settled in a few years by the new generation of neutrino oscillation experiments (GNO [13], SNO [14], Borexino [15], ICARUS [16] and others [17]).

Several atmospheric neutrino experiments (Kamiokande [18], IMB [19], Super-Kamiokande [20], Soudan 2 [21], MACRO [22]) have found an anomalous ratio of the events generated by muon and electron neutrinos and an anomalous angular dependence of the events generated by muon neutrinos. Although so far the oscillation pattern has not been observed, this is considered as an evidence in favor of oscillations of muon neutrinos into tau or sterile neutrinos. Transitions of muon neutrinos into electron neutrinos are strongly disfavored by the bounds established by the long-baseline $\bar{\nu}_e$ disappearance experiments CHOOZ [23] and Palo Verde [24] and by the fact that Super-Kamiokande data do not show any anomalous angular dependence of the events generated by atmospheric electron neutrinos. The present




Figure 1: Qualitative illustration of the possible four-neutrino schemes.

data of Super-Kamiokande [20] and MACRO [22] disfavor dominant transitions of atmospheric muon neutrinos into sterile states. There are good chances that the issue will be clarified in a definite way in the near future by long-baseline experiments with muon neutrino beams (K2K [25], MINOS [26], OPERA [27], ICARUS [16]) and by new atmospheric neutrino experiments (MONOLITH [28] and others [29]). The K2K experiment have already obtained some preliminary indication of a possible transition of muon neutrinos into other states [30].

The third evidence in favor of neutrino oscillations has been found in the short-baseline LSND experiment [31], where an excess of e^+ events have been observed. If interpreted in terms of $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ neutrino oscillations, this excess corresponds to an oscillation probability of $(2.5 \pm 0.6 \pm 0.4) \times 10^{-3}$. This result has not been confirmed by other experiments (many other less sensitive short-baseline experiments have not found any signal of neutrino oscillations; the strongest bounds have been obtained in the Bugey [32], CDHS [33], CCFR [34], BNL-E776 [35], KARMEN [36], CHORUS [37], NOMAD [38] experiments), but it is very interesting and important, because it is the only existing evidence of neutrino flavor transition from one state to another and because such transitions could be explored with high accuracy in future short-baseline experiments. In a few years the MiniBooNE experiment [39] will check the LSND signal in terms of neutrino oscillations.

It is well known (see [40]–[42]) that the three indications in favor of neutrino oscillations need at least three different neutrino mass-squared differences (Δm^2 's), arranged in the hierarchical order $\Delta m_{\text{SUN}}^2 \ll \Delta m_{\text{ATM}}^2 \ll \Delta m_{\text{SBL}}^2$. This means that at least four massive neutrinos must exist. Here we consider the minimal case of four massive neutrinos, that has been considered recently by many authors (see [43]–[54], [6, 40, 42] and references therein), whose flavor basis is constituted by the standard active neutrinos ν_e, ν_μ, ν_τ , and a sterile neutrino ν_s . Figure 1 shows the six possible four-neutrino schemes that can accommodate the observed hierarchy of Δm^2 's. These six schemes are divided in two classes: $3 + 1$ and $2+2$. In the $3 + 1$ schemes there is a group of three neutrino masses separated from an isolated mass by the LSND gap of the order of 1 eV, such that the largest mass-squared

difference, Δm_{41}^2 (where $\Delta m_{kj}^2 \equiv m_k^2 - m_j^2$ and m_k, m_j are neutrino masses, with $k, j = 1, \dots, 4$), generates the oscillations observed in the LSND experiment. In the 2+2 schemes there are two couples of close mass eigenstates separated by the LSND gap. The numbering of the mass eigenvalues in figure 1 is conveniently chosen in order to have always solar neutrino oscillations generated by $\Delta m_{21}^2 = \Delta m_{\text{SUN}}^2$ and short-baseline (SBL) oscillations generated by $|\Delta m_{41}^2| \simeq |\Delta m_{42}^2| = \Delta m_{\text{SBL}}^2$ (we have also $|\Delta m_{41}^2| \simeq |\Delta m_{43}^2|$ in 3 + 1 schemes and $|\Delta m_{41}^2| \simeq |\Delta m_{31}^2| \simeq |\Delta m_{32}^2|$ in 2+2 schemes). In 3 + 1 schemes atmospheric neutrino oscillations are generated by $|\Delta m_{31}^2| \simeq |\Delta m_{32}^2| = \Delta m_{\text{ATM}}^2$, whereas in 2+2 schemes they are generated by $|\Delta m_{43}^2| = \Delta m_{\text{ATM}}^2$.

It has been shown that in the framework of four-neutrino mixing the 3+1 schemes are disfavored by the experimental data, with respect to the 2 + 2 schemes [43]–[45], [40]. However, in a recent paper Barger, Kayser, Learned, Weiler and Whisnant [53] noticed that the new 99% CL allowed region obtained recently in the LSND experiment and presented at the Neutrino 2000 conference allows the existence 3 + 1 schemes (see also [50]) in four small regions of the amplitude $A_{\mu e}$ of $\nu_\mu \rightarrow \nu_e$ and $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ oscillations ($A_{\mu e}$ is equivalent to the usual $\sin^2 2\vartheta$ in the two-generation case). These regions, to be derived later (eqs. (3.1)–(3.6)) are shown in figure 2, enclosed by very thick solid lines. They lay at

$$\begin{aligned}
 \text{R1: } & |\Delta m_{41}^2| \simeq 0.25 \text{ eV}^2, \\
 \text{R2: } & |\Delta m_{41}^2| \simeq 0.9 \text{ eV}^2, \\
 \text{R3: } & |\Delta m_{41}^2| \simeq 1.7 \text{ eV}^2, \\
 \text{R4: } & |\Delta m_{41}^2| \simeq 6 \text{ eV}^2.
 \end{aligned}
 \tag{1.1}$$

Barger, Kayser, Learned, Weiler and Whisnant explored the phenomenological consequences of the assumption

$$1 - |U_{s4}|^2 \ll 1
 \tag{1.2}$$

(here $U_{\alpha k}$ are the elements of the 4×4 neutrino mixing matrix with $\alpha = s, e, \mu, \tau$ and $k = 1, \dots, 4$). Such a scheme is attractive because it represents a perturbation of the standard three-neutrino mixing in which a mass eigenstate is added, that mixes mainly with the new sterile neutrino ν_s and very weakly with the standard active neutrinos ν_e, ν_μ and ν_τ . In this case, the usual phenomenology of three-neutrino mixing in solar and atmospheric neutrino oscillation experiments is practically unchanged. The atmospheric neutrino anomaly would be explained by dominant $\nu_\mu \rightarrow \nu_\tau$ transitions, with possible sub-dominant $\nu_\mu \leftrightarrow \nu_e$ transitions constrained by the CHOOZ bound [55, 56, 51]. The solar neutrino problem would be explained by an approximately equal mixture of $\nu_e \rightarrow \nu_\mu$ and $\nu_e \rightarrow \nu_\tau$ transitions [56, 57].

Here, we consider another possibility that, as we will see, predicts relatively large $\nu_\mu \rightarrow \nu_\tau$ and $\nu_e \rightarrow \nu_\tau$ transitions in short-baseline neutrino oscillation experiments,




Figure 2: *Very Thick Solid Line:* Allowed regions. *Thick Solid Line:* Disappearance bound (3.5). *Dotted Line:* LSND 2000 allowed regions at 90% CL [31]. *Solid Line:* LSND 2000 allowed regions at 99% CL [31]. *Broken Dash-Dotted Line:* Bugey exclusion curve at 90% CL [32]. *Vertical Dash-Dotted Line:* CHOOZ exclusion curve at 90% CL [23]. *Long-Dashed Line:* KARMEN 2000 exclusion curve at 90% CL [36]. *Short-Dashed Line:* BNL-E776 exclusion curve at 90% CL [35].

that could be observed in the near future. We consider the 3 + 1 schemes with

$$|U_{s4}|^2 \ll 1. \tag{1.3}$$

This could be obtained, for example, in the hierarchical scheme I (see figure 1) with an appropriate symmetry keeping the sterile neutrino very light, i.e. mostly mixed with the lightest mass eigenstates. Notice that nothing forbids $|U_{s4}|^2$ to be even zero exactly. The only unwelcomed consequence of equation (1.3) is a large contribution of $\nu_\mu \rightarrow \nu_s$ transitions to the atmospheric neutrino anomaly, as discussed in section 4. These transitions are dominant in the regions R2, R3, R4, whereas they mix with $\nu_\mu \rightarrow \nu_\tau$ transitions in the region R1. As mentioned above, dominant $\nu_\mu \rightarrow \nu_s$ transitions of atmospheric neutrinos are disfavored by the present Super-Kamiokande and MACRO data [20, 22]. On the other hand, a mixture of $\nu_\mu \rightarrow \nu_s$ and $\nu_\mu \rightarrow \nu_\tau$ transitions is allowed by the data [49, 51, 52].

2. 3 + 1 schemes with $|U_{s4}|^2 \ll 1$

From the assumption (1.3), since the amplitude of $\nu_\alpha \rightarrow \nu_\beta$ and $\nu_\beta \rightarrow \nu_\alpha$ oscillations in short-baseline neutrino oscillation experiments (equivalent to the usual $\sin^2 2\vartheta$ in the two-generation case) is given by [43, 6]

$$A_{\alpha\beta} = A_{\beta\alpha} = 4|U_{\alpha 4}|^2|U_{\beta 4}|^2 \quad (\alpha, \beta = s, e, \mu, \tau), \quad (2.1)$$

we have

$$A_{\alpha s} \ll 1 \quad (\alpha = e, \mu, \tau), \quad (2.2)$$

i.e. the transitions from active to sterile neutrinos in short-baseline experiments are strongly suppressed. In the following we will neglect them.

Let us consider now the oscillation amplitude (equivalent to the usual two-generation $\sin^2 2\vartheta$) in short-baseline ν_α disappearance experiments, which is given by [43, 6]

$$B_\alpha = 4|U_{\alpha 4}|^2(1 - |U_{\alpha 4}|^2). \quad (2.3)$$

In general the oscillation amplitude in ν_α disappearance experiments is related to the amplitude of $\nu_\alpha \rightarrow \nu_\beta$ oscillations by the relation

$$B_\alpha = \sum_{\beta \neq \alpha} A_{\alpha\beta}, \quad (2.4)$$

that quantify the conservation of probability. Using equation (2.2), equation (2.4) gives

$$A_{e\tau} \simeq B_e - A_{\mu e}. \quad (2.5)$$

$$A_{\mu\tau} \simeq B_\mu - A_{\mu e}. \quad (2.6)$$

Therefore, we have

$$B_e^{\min} - A_{\mu e}^{\max} \lesssim A_{e\tau} \lesssim B_e^{\max} - A_{\mu e}^{\min}. \quad (2.7)$$

$$B_\mu^{\min} - A_{\mu e}^{\max} \lesssim A_{\mu\tau} \lesssim B_\mu^{\max} - A_{\mu e}^{\min}. \quad (2.8)$$

Let us determine B_e^{\min} , B_e^{\max} , B_μ^{\min} , B_μ^{\max} , $A_{\mu e}^{\min}$, $A_{\mu e}^{\max}$ from the results of short-baseline experiments.

3. General bounds in 3 + 1 schemes and $\nu_\mu \rightarrow \nu_e$ short-baseline transitions

The values of B_e^{\max} and B_μ^{\max} are given by the exclusion plots of $\bar{\nu}_e$ and ν_μ disappearance experiments (notice that B_e^{\max} and B_μ^{\max} depend on $|\Delta m_{41}^2|$). The most stringent bounds for $|\Delta m_{41}^2|$ in the LSND-allowed region are given by the exclusion

curves of the Bugey [32] and CHOOZ [23] reactor $\bar{\nu}_e$ disappearance experiments and the exclusion curve of the CDHS accelerator ν_μ disappearance experiment [33].

The bounds $B_e \leq B_e^{\max}$ and $B_\mu \leq B_\mu^{\max}$, together with the results of solar and atmospheric neutrino experiments imply that $|U_{e4}|^2$ and $|U_{\mu4}|^2$ are small [43, 45, 40]:

$$|U_{e4}|^2 \leq |U_{e4}|_{\max}^2 \quad \text{and} \quad |U_{\mu4}|^2 \leq |U_{\mu4}|_{\max}^2, \quad (3.1)$$

where $|U_{e4}|_{\max}^2$ and $|U_{\mu4}|_{\max}^2$ are given by

$$|U_{e4}|_{\max}^2 = \frac{1}{2} \left(1 - \sqrt{1 - B_e^{\max}} \right), \quad (3.2)$$

$$|U_{\mu4}|_{\max}^2 = \min \left[\frac{1}{2} \left(1 - \sqrt{1 - B_\mu^{\max}} \right), 0.55 \right]. \quad (3.3)$$

The number 0.55 comes [45] from the up-down asymmetry of multi-GeV muon-like events measured in the Super-Kamiokande experiment [20]. The values of $|U_{e4}|_{\max}^2$ and $|U_{\mu4}|_{\max}^2$ as functions of $|\Delta m_{41}^2|$ are shown, respectively, in figures 3 and 4.

In the 3 + 1 schemes we have

$$A_{\mu e} = 4|U_{\mu4}|^2|U_{e4}|^2, \quad (3.4)$$

and the bounds (3.1) imply that [43]

$$A_{\mu e} \leq 4|U_{\mu4}|_{\max}^2|U_{e4}|_{\max}^2. \quad (3.5)$$

As one can see from figure 2, from the LSND region constrained by the exclusion curves of KARMEN and BNL-E776 and by the bound (3.5), there are four allowed regions for $A_{\mu e}$, R1, R2, R3, R4 in equation (1.1), where we have

$$A_{\mu e}^{\min} \leq 4|U_{\mu4}|^2|U_{e4}|^2 \leq A_{\mu e}^{\max}. \quad (3.6)$$

These regions could be explored in the near future by the MiniBooNE experiment [39]. Region R4 is at the limit of the final NOMAD sensitivity in the $\nu_\mu \rightarrow \nu_e$ channel [58].

From the lower bound in equation (3.6) and the bounds (3.1), we obtain

$$|U_{e4}|^2 \geq \frac{A_{\mu e}^{\min}}{4|U_{\mu4}|^2} \geq \frac{A_{\mu e}^{\min}}{4|U_{\mu4}|_{\max}^2}, \quad (3.7)$$

$$|U_{\mu4}|^2 \geq \frac{A_{\mu e}^{\min}}{4|U_{e4}|^2} \geq \frac{A_{\mu e}^{\min}}{4|U_{e4}|_{\max}^2}. \quad (3.8)$$

Therefore, we have

$$\frac{A_{\mu e}^{\min}}{4|U_{\mu4}|_{\max}^2} \leq |U_{e4}|^2 \leq |U_{e4}|_{\max}^2, \quad (3.9)$$

$$\frac{A_{\mu e}^{\min}}{4|U_{e4}|_{\max}^2} \leq |U_{\mu4}|^2 \leq |U_{\mu4}|_{\max}^2. \quad (3.10)$$




Figure 3: *Dotted Line:* $|U_{e4}|^2_{\max}$ obtained from the 90% CL exclusion curve of the Bugey short-baseline reactor $\bar{\nu}_e$ disappearance experiment [32] (equation (3.2)). *Dashed Line:* $|U_{e4}|^2_{\max}$ obtained from the 90% CL exclusion curve of the CHOOZ long-baseline reactor $\bar{\nu}_e$ disappearance experiment [23] (equation (3.2)). *Solid Line:* Allowed regions (equation (3.9))

Since these bounds have been obtained without any assumption on the value of $|U_{s4}|^2$, they are generally valid in any $3 + 1$ scheme. The corresponding four allowed regions for $|U_{e4}|^2$ and $|U_{\mu4}|^2$ are shown, respectively, in figures 3 and 4.

The lower bounds for $|U_{e4}|^2$ and $|U_{\mu4}|^2$ in equations (3.7) and (3.8) imply lower bounds for the oscillation amplitudes $B_e = 4|U_{e4}|^2(1 - |U_{e4}|^2)$ and $B_\mu = 4|U_{\mu4}|^2 \times (1 - |U_{\mu4}|^2)$. Let us derive these bounds.

Since $|U_{e4}|^2_{\max}$ is always smaller than $1/2$, the lower bound for B_e is

$$B_e^{\min} = \frac{A_{\mu e}^{\min}}{|U_{\mu4}|^2_{\max}} \left(1 - \frac{A_{\mu e}^{\min}}{4|U_{\mu4}|^2_{\max}} \right). \quad (3.11)$$

On the other hand, since $|U_{\mu4}|^2_{\max}$ can be bigger than $1/2$, the lower bound for B_μ is

$$B_\mu^{\min} = \min \left[\frac{A_{\mu e}^{\min}}{|U_{e4}|^2_{\max}} \left(1 - \frac{A_{\mu e}^{\min}}{4|U_{e4}|^2_{\max}} \right), |U_{\mu4}|^2_{\max} (1 - |U_{\mu4}|^2_{\max}) \right]. \quad (3.12)$$




Figure 4: *Dotted Line:* $|U_{\mu 4}|_{\max}^2$ obtained from the exclusion curve of the CDHS accelerator ν_μ disappearance experiment [33] (equation (3.3)). *Dashed Line:* $|U_{\mu 4}|_{\max}^2$ obtained from the up-down asymmetry of multi-GeV muon-like events measured in the Super-Kamiokande experiment [20] (equation (3.3)). *Solid Line:* Allowed regions (equation (3.10)).

Figure 5 and 6 show the allowed regions in the B_e - $|\Delta m_{41}^2|$ and B_μ - $|\Delta m_{41}^2|$ planes given by the bounds

$$B_\alpha^{\min} \leq B_\alpha \leq B_\alpha^{\max} \quad (\alpha = e, \mu). \quad (3.13)$$

One can see that these regions lie just on the left of the Bugey+CHOOZ (figure 5) and CDHS (figure 6) exclusion curves and could be observed in the near future [59]. Let us emphasize that these regions are generally predicted in any $3 + 1$ scheme, since they have been obtained independently on any assumption on the mixing (as equation (1.2) or equation (1.3)).

4. Large $\nu_e \rightarrow \nu_\tau$ and $\nu_\mu \rightarrow \nu_\tau$ short-baseline transitions

We have now all the elements to calculate the bounds (2.7) and (2.8) on the amplitudes of short-baseline $\nu_e \rightarrow \nu_\tau$ and $\nu_\mu \rightarrow \nu_\tau$ oscillations that follow from the assumption (1.3), $|U_{s4}|^2 \ll 1$.




Figure 5: *Solid Line:* Allowed regions. *Dotted Line:* Bugey exclusion curve at 90% CL [32]. *Dashed Line:* CHOOZ exclusion curve at 90% CL [23].

Figure 7 shows the allowed regions in the $A_{\mu\tau}-|\Delta m_{41}^2|$ plane given by the bounds (2.8). One can see that the region R4 is excluded by the negative results of the CHORUS [37] and NOMAD [38] experiments. The other three regions are possible and predict relatively large oscillation amplitudes that could be observed in the near future, especially the two regions R2 and R3 in which $A_{\mu\tau} \sim 4 \times 10^{-2} - 10^{-1}$.

Figure 8 shows the allowed regions in the $A_{e\tau}-|\Delta m_{41}^2|$ plane given by the bounds (2.7). One can see that these regions predict relatively large oscillation amplitudes, but unfortunately lie rather far from the CHORUS and NOMAD exclusion curves (except the region R4 that is excluded by figure 7). Therefore, even under the favorable assumption (1.3), it will be very difficult to observe $\nu_e \rightarrow \nu_\tau$ transitions in short baseline experiments with conventional neutrino beams, but large transitions could be observed with a ν_e beam from a neutrino factory [60].

As one can see from figure 3, the bound in equation (3.1) imply that $|U_{e4}|^2$ is very small in all the three allowed regions in figure 7. On the other hand, one can see from figure 4 that $|U_{\mu4}|^2$ is small in the two allowed regions R2 and R3, but it is large in the region R1. As a consequence of the unitarity of the mixing matrix and the assumption (1.3), we have $1 - |U_{\tau4}|^2 \ll 1$ in R2 and R3, whereas $|U_{\tau4}|^2$ can be as small as about 1/2 in R1. The prediction for solar, atmospheric and long-baseline experiments depend on the value of $|U_{\mu4}|^2$. There are two possibilities:




Figure 6: *Solid Line:* Allowed regions. *Dotted Line:* CDHS exclusion curve at 90% CL [33].

1. $|U_{\mu 4}|^2 \ll 1$ in regions R2 and R3 in figure 4. In this case $1 - |U_{\tau 4}|^2 \ll 1$ and the atmospheric neutrino anomaly is due to dominant $\nu_\mu \rightarrow \nu_s$ transitions. This possibility is disfavored by present Super-Kamiokande and MACRO data [20, 22], but still it is not completely excluded (see [61, 62, 52]). Obviously, in this case long-baseline $\nu_\mu \rightarrow \nu_\tau$ transitions are suppressed with respect to the dominant $\nu_\mu \rightarrow \nu_s$ transitions, that are almost entirely responsible of the disappearance of ν_μ 's.

The solar neutrino problem is due to an approximately equal mixture of $\nu_e \rightarrow \nu_\mu$ and $\nu_e \rightarrow \nu_s$ transitions, which is allowed by the data. This has been shown in refs. [47, 48] in the framework of the 2 + 2 schemes in figure 1. However, the results obtained in refs. [47, 48] are valid also in the 3 + 1 schemes, because the formalism of solar neutrino oscillations in 3 + 1 schemes is identical to that in 2 + 2 schemes [46]–[48]. Indeed, from eqs. (3.1) and (3.2) we already know that the results of the Bugey and CHOOZ experiment imply that $|U_{e4}|^2$ is very small. Furthermore, taking into account the hierarchy

$$\Delta m_{21}^2 = \Delta m_{\text{SUN}}^2 \ll |\Delta m_{31}^2| = \Delta m_{\text{ATM}}^2 \ll |\Delta m_{41}^2| = \Delta m_{\text{SBL}}^2, \quad (4.1)$$




Figure 7: *Solid Line:* Allowed regions. *Long Dashed Line:* CHORUS exclusion curve at 90% CL [37]. *Short Dashed Line:* NOMAD exclusion curve at 90% CL [38]. *Dotted Line:* CDHS exclusion curve at 90% CL [33].

the effective survival probability of $\bar{\nu}_e$ and ν_e in long-baseline experiments is given by

$$P_{\nu_e \rightarrow \nu_e}^{\text{LBL}} = 1 - 4|U_{e3}|^2 (1 - |U_{e3}|^2) \sin^2 \left(\frac{\Delta m_{31}^2 L}{4E} \right) + |U_{e4}|^4, \quad (4.2)$$

where L is the propagation distance and E is the neutrino energy. Neglecting $|U_{e4}|^4$, equation (4.2) has the same structure as the usual two-generation survival probability (see [6]), with $\sin^2 2\vartheta$ replaced by $4|U_{e3}|^2 (1 - |U_{e3}|^2)$ and Δm^2 replaced by Δm_{31}^2 . The bound on $\sin^2 2\vartheta$ obtained in the CHOOZ experiment [23] for $|\Delta m_{31}^2| \gtrsim 10^{-3} \text{ eV}^2$ implies that $|U_{e3}|^2 \lesssim 2.6 \times 10^{-2}$. Therefore, both $|U_{e3}|^2$ and $|U_{e4}|^2$ are very small,

$$|U_{e3}|^2 \lesssim 3 \times 10^{-2} \quad |U_{e4}|^2 \lesssim 3 \times 10^{-2}, \quad (4.3)$$

and can be neglected in the study of solar neutrino oscillations, as done in refs. [46, 47, 48]. In other words, the production and detection of the mass eigenstates ν_3 and ν_4 is negligibly small in solar neutrino experiments. Moreover, because of the hierarchy (4.1), there are no matter-induced transitions from ν_1, ν_2 to ν_3, ν_4 . Hence, ν_3 and ν_4 effectively decouple in solar neutrino oscillations, leading to the formalism described in refs. [46]–[48].




Figure 8: *Solid Line:* Allowed regions. *Long Dashed Line:* CHORUS exclusion curve at 90% CL [37]. *Short Dashed Line:* NOMAD exclusion curve at 90% CL [38]. *Dotted Line:* Bugey exclusion curve at 90% CL [32]. *Dashed Line:* CHOOZ exclusion curve at 90% CL [23].

2. $|U_{\mu 4}|^2 \simeq 0.33 - 0.55$ in region R1 in figure 4. In this case the solar neutrino problem is due to a mixture of $\nu_e \rightarrow \nu_\mu$, $\nu_e \rightarrow \nu_\tau$ and $\nu_e \rightarrow \nu_s$ transitions, that is allowed by data as in the previous case.

The atmospheric neutrino anomaly is due to a mixture of $\nu_\mu \rightarrow \nu_\tau$ and $\nu_\mu \rightarrow \nu_s$ transitions. In refs. [49, 51, 52] it has been shown that a mixture of $\nu_\mu \rightarrow \nu_\tau$ and $\nu_\mu \rightarrow \nu_s$ transitions in the framework of 2 + 2 schemes is allowed by the atmospheric neutrino data. This indicates that such a mixture should be allowed also in the framework of 3 + 1 schemes. The groups specialized in the analysis of atmospheric neutrino data could check this possibility using the formalism presented in appendix A. The existence of mixed $\nu_\mu \rightarrow \nu_\tau$ and $\nu_\mu \rightarrow \nu_s$ transitions can also be checked by comparing the rates of ν_μ disappearance and $\nu_\mu \rightarrow \nu_\tau$ appearance in future long-baseline experiments with ν_μ beams. The formalism that allows to describe these oscillation channels in the framework of 3 + 1 four-neutrino schemes is presented in appendix A.

These predictions are testable in future experiments, especially measuring the percentage of transitions into active and sterile neutrinos in solar, atmospheric and long-baseline experiments (that should measure the same transitions observed in

atmospheric neutrino experiments). Taking into account also the prediction of large $\nu_\mu \rightarrow \nu_\tau$ and $\nu_e \rightarrow \nu_\tau$ in short-baseline experiments, the schemes under consideration can be checked and possibly falsified in the near future.

5. Conclusions

In conclusion, we have considered the four-neutrino $3 + 1$ schemes in figure 1, that are marginally allowed by present data (see figure 2). We have identified four small regions (R1, R2, R3, R4) in the parameter space of neutrino mixing compatible with all data and we have derived general upper and lower bounds for the elements U_{e4} and $U_{\mu 4}$ of the mixing matrix and on the oscillation amplitudes in short-baseline $\bar{\nu}_e$ and ν_μ disappearance experiments. The corresponding $\nu_\mu \rightarrow \nu_e$ transitions and ν_e and ν_μ disappearance in short-baseline experiments are relatively large and could be observed in future dedicated experiments.

Assuming a small mixing of the sterile neutrino with the isolated mass eigenstate, $|U_{s4}|^2 \ll 1$, we have shown that large $\nu_\mu \rightarrow \nu_\tau$ and $\nu_e \rightarrow \nu_\tau$ transitions are predicted in short-baseline experiments. In this case, the region R4 is already excluded by the results of the CHORUS [37] and NOMAD [38] $\nu_\mu \rightarrow \nu_\tau$ experiments.

We have also discussed the implications of $|U_{s4}|^2 \ll 1$ for solar, atmospheric and long-baseline neutrino oscillation experiments and we have presented in appendix A the formalism describing in general $3 + 1$ schemes the oscillations in matter of atmospheric neutrinos and neutrinos in long-baseline ν_μ disappearance and $\nu_\mu \rightarrow \nu_\tau$ appearance experiments. In the allowed regions R2 and R3 the atmospheric neutrino anomaly is due to dominant $\nu_\mu \rightarrow \nu_s$ transitions, that are disfavored by present Super-Kamiokande and MACRO data [20, 22]. On the other hand, in the allowed region R1 the atmospheric neutrino anomaly is due a mixture of $\nu_\mu \rightarrow \nu_\tau$ and $\nu_\mu \rightarrow \nu_s$ transitions, that is likely to be allowed by the data, as shown by the calculations in the framework of $2 + 2$ schemes presented in refs. [49, 51, 52].

Finally, let us remark that the four $3 + 1$ schemes in figure 1 are equivalent for solar, atmospheric and short-baseline neutrino oscillation experiments, but they may be distinguishable in long-baseline $\nu_\mu \rightarrow \nu_e$ experiments if $|U_{e3}|^2$ is not too small [63], or through their different effects in tritium β decay and neutrinoless double- β decay (see [3, 5, 6]) and through neutrino oscillations in supernovae (see [64, 65]) and in the early universe (see [66]–[70]).

Acknowledgments

C.G. would like to thank S. Bilenky, E. Lisi, O. Peres, F. Ronga, and A. Smirnov for useful discussions during the NOW2000 workshop. C.G. would also like to express his gratitude to the Korea Institute for Advanced Study (KIAS) for warm hospitality during the completion of this work.

A. Formalism of atmospheric neutrino oscillations

In this appendix we derive in a concise way the formalism in $3 + 1$ schemes of atmospheric neutrino oscillations and oscillations in long-baseline $\nu_\mu \rightarrow \nu_\tau$ and ν_μ disappearance experiments. We take into account matter effects, following a method similar to that used in ref. [46] in the case of $2 + 2$ schemes. As explained in the main text, the formalism of solar neutrino oscillations in $3 + 1$ schemes is identical to that in $2 + 2$ schemes [46]–[48].

From equation (4.3) we know that U_{e3} and U_{e4} can be neglected. Choosing the ordering $\nu_s, \nu_e, \nu_\mu, \nu_\tau$ for the flavor neutrino fields, the mixing matrix can be written as

$$U = V_{34} V_{14} V_{13} V_{12}, \quad (\text{A.1})$$

where

$$(V_{ij})_{ab} = \delta_{ab} + (\cos \vartheta_{ij} - 1) (\delta_{ia} \delta_{ib} + \delta_{ja} \delta_{jb}) + \sin \vartheta_{ij} (\delta_{ia} \delta_{jb} - \delta_{ja} \delta_{ib}) \quad (\text{A.2})$$

represents a rotation by an angle ϑ_{ij} in the i - j plane. We have also neglected possible CP-violating phases. The evolution of the neutrino flavor amplitudes ψ_α ($\alpha = s, e, \mu, \tau$) in vacuum and in matter is given by the MSW equation [71]

$$i \frac{d}{dx} \Psi = \mathcal{H} \Psi, \quad (\text{A.3})$$

where $\Psi = (\psi_s, \psi_e, \psi_\mu, \psi_\tau)^T$ and \mathcal{H} is the effective Hamiltonian

$$\mathcal{H} = \frac{1}{2p} (U \mathcal{M}_0^2 U^\dagger + \mathcal{A}). \quad (\text{A.4})$$

Here p is the neutrino momentum,

$$\mathcal{M}_0^2 = \text{diag}(0, 0, \Delta m_{31}^2, \Delta m_{41}^2) \quad (\text{A.5})$$

is the mass-squared matrix, in which we neglected Δm_{21}^2 according to the hierarchical relation (4.1), and \mathcal{A} is the matrix

$$\mathcal{A} = \text{diag}(-A_{NC}, A_{CC}, 0, 0), \quad (\text{A.6})$$

with $A_{CC} = 2pV_{CC}$ and $A_{NC} = 2pV_{NC}$, where V_{CC} and V_{NC} are, respectively, the matter-induced charged-current and neutral-current potentials (see [2, 3, 4, 5, 6]). For atmospheric neutrinos propagating in the Earth and accelerator neutrinos in long-baseline experiments $|A_{CC}| \sim |A_{NC}| \sim |\Delta m_{31}^2| \ll |\Delta m_{41}^2|$. In equation (A.4) we have neglected a common phase for the flavor amplitudes, that is irrelevant for neutrino oscillations.

The evolution equation (A.3) is most easily solved in the rotated basis $\Psi' = V_{14}^T V_{34}^T \Psi = (\psi'_1, \psi'_2, \psi'_3, \psi'_4)^T$ that obeys to a similar evolution equation, with \mathcal{H} replaced by

$$\mathcal{H}' = \frac{1}{2p} (V_{13} \mathcal{M}_0^2 V_{13}^T + V_{14}^T \mathcal{A} V_{14}) \quad (\text{A.7})$$

where we have taken into account the fact that $V_{12} \mathcal{M}_0^2 V_{12}^T = \mathcal{M}_0^2$ and $V_{34}^T \mathcal{A} V_{34} = \mathcal{A}$, which lead to a significant simplification of the evolution equation. The explicit form of \mathcal{H}' is

$$\mathcal{H}' = \frac{1}{4p} \begin{pmatrix} (1-c_{2\vartheta_{13}})\Delta m_{31}^2 - 2c_{\vartheta_{14}}^2 A_{NC} & 0 & s_{2\vartheta_{13}}\Delta m_{31}^2 & -s_{2\vartheta_{14}}A_{NC} \\ 0 & 2A_{CC} & 0 & 0 \\ s_{2\vartheta_{13}}\Delta m_{31}^2 & 0 & (1+c_{2\vartheta_{13}})\Delta m_{31}^2 & 0 \\ -s_{2\vartheta_{14}}A_{NC} & 0 & 0 & 2\Delta m_{41}^2 - 2s_{\vartheta_{14}}^2 A_{NC} \end{pmatrix}, \quad (\text{A.8})$$

with $c_\vartheta \equiv \cos \vartheta$ and $s_\vartheta \equiv \sin \vartheta$. This equation shows that the amplitudes ψ'_2 and ψ'_4 evolve independently (remember that $|\Delta m_{41}^2| \gg |A_{NC}|$), with phases given by the energy eigenvalues

$$E'_2 = A_{CC}/2p = V_{CC}, \quad E'_4 \simeq \Delta m_{41}^2/2p, \quad (\text{A.9})$$

whereas the evolutions of the amplitudes ψ'_1 and ψ'_3 are coupled and there is a resonance in the 1-3 sector for

$$-\cos^2 \vartheta_{14} A_{NC} = \cos 2\vartheta_{13} \Delta m_{31}^2. \quad (\text{A.10})$$

Notice that for neutrinos the left-hand side of equation (A.10) is positive, because $A_{NC} \leq 0$, whereas for antineutrinos it is negative because A_{NC} must be replaced by $\bar{A}_{NC} = -A_{NC}$. Hence, if $\Delta m_{31}^2 > 0$ (schemes I and IV in figure 1) and $\vartheta_{13} < \pi/4$ or $\Delta m_{31}^2 < 0$ (schemes II and III) and $\vartheta_{13} > \pi/4$, there can be a resonance for neutrinos, otherwise the resonance can be for antineutrinos.

One can calculate the evolution of the amplitudes ψ'_1 and ψ'_3 solving numerically the two coupled equations generated by the 1-3 sector of the effective Hamiltonian \mathcal{H}' . Another common method for the solution of the evolution equation is to divide the Earth interior into shell with constant density, calculate the evolution of the amplitudes Ψ' in each shell and match the amplitudes in the flavor basis at the shell boundaries. In this case, in each shell the amplitudes ψ'_1 and ψ'_3 can be written as

$$\psi'_1 = \cos \vartheta_{13}^M \psi_1^M + \sin \vartheta_{13}^M \psi_3^M, \quad \psi'_3 = -\sin \vartheta_{13}^M \psi_1^M + \cos \vartheta_{13}^M \psi_3^M, \quad (\text{A.11})$$

where ψ_1^M and ψ_3^M are the amplitudes of the energy eigenstates that evolve with phases given by the energy eigenvalues

$$E_{1,3}^M = \frac{1}{4p} \left[\Delta m_{31}^2 - \cos^2 \vartheta_{14} A_{NC} \mp \sqrt{(\cos 2\vartheta_{13} \Delta m_{31}^2 + \cos^2 \vartheta_{14} A_{NC})^2 + (\sin 2\vartheta_{13} \Delta m_{31}^2)^2} \right]. \quad (\text{A.12})$$

The effective mixing angle in matter ϑ_{13}^M is given by

$$\tan 2\vartheta_{13}^M = \frac{\sin 2\vartheta_{13}\Delta m_{31}^2}{\cos 2\vartheta_{13}\Delta m_{31}^2 + \cos^2 \vartheta_{14} A_{NC}}, \quad (\text{A.13})$$

from which one can see that $\vartheta_{13}^M = \pi/4$ when the resonance condition (A.10) is satisfied, and $\vartheta_{13}^M \rightarrow \pi/2$ in a very dense medium, where $|\cos^2 \vartheta_{14} A_{NC}| \gg |\cos 2\vartheta_{13}\Delta m_{31}^2|$ if $\cos 2\vartheta_{13}\Delta m_{31}^2 > 0$, whereas $\vartheta_{13}^M \rightarrow 0$ in a very dense medium if $\cos 2\vartheta_{13}\Delta m_{31}^2 < 0$.

The connection between the amplitudes Ψ' and the flavor amplitudes Ψ , needed for the calculation of the probability of flavor transitions, is given by

$$\begin{aligned} \psi_s &= \cos \vartheta_{14} \psi'_1 + \sin \vartheta_{14} \psi'_4, \\ \psi_e &= \psi'_2, \\ \psi_\mu &= -\sin \vartheta_{14} \sin \vartheta_{34} \psi'_1 + \cos \vartheta_{34} \psi'_3 + \cos \vartheta_{14} \sin \vartheta_{34} \psi'_4, \\ \psi_\tau &= -\sin \vartheta_{14} \cos \vartheta_{34} \psi'_1 - \sin \vartheta_{34} \psi'_3 + \cos \vartheta_{14} \cos \vartheta_{34} \psi'_4. \end{aligned} \quad (\text{A.14})$$

The second line of equation (A.14) imply that electron neutrinos do not oscillate in atmospheric neutrino experiments (remember that ψ'_2 evolves independently). This is due to the approximation $U_{e3} = 0$, motivated by equation (4.3), which also imply that the charged-current matter potential V_{CC} , felt only by ν_e , is irrelevant for atmospheric neutrino oscillations (only the neutral-current potential V_{NC} , felt by ν_μ and ν_τ , enter in eqs. (A.10)–(A.13)). On the other hand, simultaneous $\nu_\mu \rightarrow \nu_\tau$ and $\nu_\mu \rightarrow \nu_s$ transitions are allowed, with pure $\nu_\mu \rightarrow \nu_\tau$ transitions in the limit $\cos \vartheta_{14} = 0$, that corresponds to $U_{s4} = 1$, and pure $\nu_\mu \rightarrow \nu_s$ transitions in the limit $\cos \vartheta_{14} \cos \vartheta_{34} = 1$, that corresponds to $U_{\tau 4} = 1$.

Let us consider finally long baseline experiments in which the neutrino beam travels in the crust of the Earth, where the matter density is practically constant. The probability of $\nu_\alpha \rightarrow \nu_\beta$ transitions is given by

$$\begin{aligned} P_{\nu_\alpha \rightarrow \nu_\beta}^{\text{LBL}} &= \left| U_{\alpha 1}^M U_{\beta 1}^M + U_{\alpha 3}^M U_{\beta 3}^M \exp\left(-i \frac{\Delta_{31}^M L}{2p}\right) \right|^2 + |U_{\alpha 4}^M|^2 |U_{\beta 4}^M|^2 \\ &= -4U_{\alpha 1}^M U_{\beta 1}^M U_{\alpha 3}^M U_{\beta 3}^M \sin^2\left(\frac{\Delta_{31}^M L}{4p}\right) + 2|U_{\alpha 4}^M|^2 |U_{\beta 4}^M|^2, \end{aligned} \quad (\text{A.15})$$

where L is the propagation distance,

$$\Delta_{31}^M = 2p (E_3^M - E_1^M) = \sqrt{(\cos 2\vartheta_{13}\Delta m_{31}^2 + \cos^2 \vartheta_{14} A_{NC})^2 + (\sin 2\vartheta_{13}\Delta m_{31}^2)^2}, \quad (\text{A.16})$$

and U^M is the effective mixing matrix in matter,

$$U^M = V_{34} V_{14} V_{13}^M = \begin{pmatrix} c_{\vartheta_{14}} c_{\vartheta_{13}}^M & 0 & c_{\vartheta_{14}} s_{\vartheta_{13}}^M & s_{\vartheta_{14}} \\ 0 & 1 & 0 & 0 \\ -s_{\vartheta_{34}} s_{\vartheta_{14}} c_{\vartheta_{13}}^M - c_{\vartheta_{34}} s_{\vartheta_{13}}^M & 0 & -s_{\vartheta_{34}} s_{\vartheta_{14}} s_{\vartheta_{13}}^M + c_{\vartheta_{34}} c_{\vartheta_{13}}^M & s_{\vartheta_{34}} c_{\vartheta_{14}} \\ -c_{\vartheta_{34}} s_{\vartheta_{14}} c_{\vartheta_{13}}^M + s_{\vartheta_{34}} s_{\vartheta_{13}}^M & 0 & -c_{\vartheta_{34}} s_{\vartheta_{14}} s_{\vartheta_{13}}^M - s_{\vartheta_{34}} c_{\vartheta_{13}}^M & c_{\vartheta_{34}} c_{\vartheta_{14}} \end{pmatrix}, \quad (\text{A.17})$$

where V_{13}^M is equal to V_{13} with ϑ_{13} replaced by ϑ_{13}^M . The expression (A.15) can be used to analyze the data of long-baseline $\nu_\mu \rightarrow \nu_\tau$ and ν_μ disappearance experiments (the analysis of long-baseline $\nu_\mu \rightarrow \nu_e$ data requires the relaxation of the approximation $U_{e3} = 0$, leading to a significant complication of the formalism [63]).

References

- [1] S.M. Bilenky and B. Pontecorvo, *Lepton mixing and neutrino oscillations*, *Phys. Rep.* **41** (1978) 225.
- [2] S.P. Mikheev and A.Y. Smirnov, *Resonance oscillations of neutrinos in matter*, *Sov. Phys. Usp.*, **30** (1987) 759.
- [3] S.M. Bilenky and S.T. Petcov, *Massive neutrinos and neutrino oscillations*, *Rev. Mod. Phys.* **59** (1987) 671.
- [4] T.K. Kuo and J. Pantaleone, *Neutrino oscillations in matter*, *Rev. Mod. Phys.* **61** (1989) 937.
- [5] C.W. Kim and A. Pevsner, *Neutrinos in physics and astrophysics*, *Contemporary Concepts in Physics*, vol. 8, Harwood Academic Press, Chur, Switzerland 1993.
- [6] S.M. Bilenky, C. Giunti and W. Grimus, *Phenomenology of neutrino oscillations*, *Prog. Part. Nucl. Phys.* **43** (1999) 1 [hep-ph/9812360].
- [7] B.T. Cleveland et al., *Measurement of the solar electron neutrino flux with the Homestake Chlorine detector*, *Astrophys. J.* **496** (1998) 505.
- [8] KAMIOKANDE collaboration, *Solar neutrino data covering solar cycle 22*, *Phys. Rev. Lett.* **77** (1996) 1683.
- [9] GALLEX collaboration, *GALLEX solar neutrino observations: results for GALLEX IV*, *Phys. Lett.* **B 447** (1999) 127.
- [10] SAGE collaboration, *Measurement of the solar neutrino capture rate with gallium metal*, *Phys. Rev.* **D 60** (1999) 055801 [astro-ph/9907113].
- [11] SUPER-KAMIOKANDE collaboration, Y. Suzuki, *Solar neutrino results from Super-Kamiokande*, Talk presented at Neutrino 2000, Sudbury, Canada 16–21 June 2000, <http://nu2000.sno.laurentian.ca>; *Water-Cherenkov and radiochemical results*, Talk presented at NOW2000, Otranto, Italy September 2000, <http://www.ba.infn.it/~now2000>.
- [12] J.N. Bahcall, S. Basu and M.H. Pinsonneault, *How uncertain are solar neutrino predictions?*, *Phys. Lett.* **B 433** (1998) 1 [astro-ph/9805135].
- [13] GNO collaboration, M. Altmann et al., *GNO solar neutrino observations: results for GNO I*, *Phys. Lett.* **B 490** (2000) 16 [hep-ex/0006034].

- [14] SNO collaboration, A. Hallin, *The sudbury neutrino observatory*, *Nucl. Phys.* **A 663** (2000) 787.
- [15] BOREXINO collaboration, A. Ianni, *Status of the BOREXINO experiment*, *Nucl. Phys.* **A 663** (2000) 791.
- [16] ICARUS collaboration, *ICARUS 600 TON: a status report*, *Nucl. Phys.* **85** (*Proc. Suppl.*) (2000) 119; LNGS-94/99-I 1994, web page: <http://pcnometh4.cern.ch>.
- [17] F. von Feilitzsch, Talk presented at Neutrino 2000, Sudbury Canada, 16–21 June 2000 <http://nu2000.sno.laurentian.ca>
- [18] KAMIOKANDE collaboration, *Atmospheric $\nu_\mu\nu_e$ ratio in the multiGeV energy range*, *Phys. Lett.* **B 335** (1994) 237.
- [19] IMB collaboration, *Neutrino measurements with the IMB detector*, *Nucl. Phys. Proc. Suppl.*, **38** (1995) 331.
- [20] SUPER-KAMIOKANDE collaboration, *Tau neutrinos favored over sterile neutrinos in atmospheric muon neutrino oscillations*, *Phys. Rev. Lett.* **85** (2000) 3999 [[hep-ex/0009001](http://arxiv.org/abs/hep-ex/0009001)];
SUPER-KAMIOKANDE collaboration, T. Kajita, *Super-Kamiokande result*, talk presented at NOW2000, Otranto, Italy, September 2000 <http://www.ba.infn.it/~now2000>
- [21] SOUDAN 2 collaboration, *The atmospheric neutrino flavor ratio from a 3.9 fiducial kiloton year exposure of Soudan-2*, *Phys. Lett.* **B 449** (1999) 137.
- [22] MACRO collaboration, F. Ronga, *MACRO and Soudan 2*, talk presented at NOW2000, Otranto, Italy, September 2000 <http://www.ba.infn.it/~now2000>.
- [23] CHOOZ collaboration, *Initial results from the CHOOZ long baseline reactor neutrino oscillation experiment*, *Phys. Lett.* **B 420** (1998) 397 [[hep-ex/9711002](http://arxiv.org/abs/hep-ex/9711002)]; *Limits on neutrino oscillations from the CHOOZ experiment*, *Phys. Lett.* **B 466** (1999) 415 [[hep-ex/9907037](http://arxiv.org/abs/hep-ex/9907037)].
- [24] F. Boehm et al., *Results from the palo verde neutrino oscillation experiment*, *Phys. Rev.* **D 62** (2000) 072002 [[hep-ex/0003022](http://arxiv.org/abs/hep-ex/0003022)].
- [25] K2K collaboration, Y. Oyama, *K2K (KEK to Kamioka) neutrino oscillation experiment at KEK-PS*, [hep-ex/9803014](http://arxiv.org/abs/hep-ex/9803014); K2K web page: <http://pnahep.kek.jp/>.
- [26] MINOS collaboration, NuMI-L-476 March 1999, web page: <http://www.hep.anl.gov/ndk/hypertext/numi.html>.
- [27] OPERA collaboration, *The OPERA experiment at Gran Sasso*, *Nucl. Phys.* **85** (*Proc. Suppl.*) (2000) 125; *OPERA: an appearance experiment to search for $\nu_\mu \leftrightarrow \nu_\tau$ oscillations in the CNGS BEAM. Experimental proposal*, CERN-SPSC-2000-028.

- [28] MONOLITH collaboration, A. Geiser, [[hep-ex/0008067](#)]; web page: <http://www.desy.de/~hoepfner/Neutrino/Monolith>.
- [29] A. Geiser, *Future atmospheric neutrino detectors*, *Nucl. Phys.* **91** (*Proc. Suppl.*) (2000) 147 [[hep-ex/0009060](#)].
- [30] K2K collaboration, K. Nakamura, *K2K: status*, talk presented at Neutrino 2000, Sudbury, Canada, 16–21 June 2000, <http://nu2000.sno.laurentian.ca>;
K2K collaboration, M. Sakuda, talk presented at ICHEP2000, Osaka, Japan, July 2000 <http://ichep2000.hep.sci.osaka-u.ac.jp>.
- [31] LSND collaboration, G. Mills, *Final 1993-1998 LSND results*, talk presented at Neutrino 2000, Sudbury, Canada, 16–21 June 2000 <http://nu2000.sno.laurentian.ca>.
- [32] Y. Declais et al., *Search for neutrino oscillations at 15-meters, 40-meters and 95-meters from a nuclear power reactor at bugey*, *Nucl. Phys.* **B 434** (1995) 503.
- [33] F. Dydak et al., *A search for ν_μ oscillations in the Δm^2 range $0.3-90eV^2$* , *Phys. Lett.* **B 134** (1984) 281.
- [34] I.E. Stockdale et al., *Limits on muon neutrino oscillations in the mass range $55-eV^2 < \Delta m^2 < 800 - eV^2$* , *Phys. Rev. Lett.* **52** (1984) 1384.
- [35] L. Borodovsky et al., *Search for muon-neutrino oscillations $\nu_\mu \rightarrow \nu_e$ ($\bar{\nu}_\mu \rightarrow \bar{\nu}_e$) in a wide band neutrino beam. BNL-776 experiment*, *Phys. Rev. Lett.* **68** (1992) 274.
- [36] KARMEN collaboration, K. Eitel, *Latest results of the karmen2 experiment*, *Nucl. Phys.* **91** (*Proc. Suppl.*) (2000) 191 [[hep-ex/0008002](#)]; web page: <http://www-ik1.fzk.de/www/karmen>
- [37] CHORUS collaboration, L. Ludovici, Talk presented at Neutrino 2000, Sudbury, Canada, 16–21 June 2000 (<http://nu2000.sno.laurentian.ca>); web page: <http://choruswww.cern.ch>.
- [38] NOMAD collaboration, M. Mezzetto, Talk presented at Neutrino 2000, Sudbury, Canada, 16–21 June 2000 (<http://nu2000.sno.laurentian.ca>); web page: <http://nomadinfo.cern.ch>.
- [39] MINIBOONE collaboration, A. Bazarko, *Miniboone: status of the booster neutrino experiment*, *Nucl. Phys.* **91** (*Proc. Suppl.*) (2000) 210 [[hep-ex/0009056](#)];
Booster Neutrino Experiment (BooNE): <http://nu1.lampf.lanl.gov/BooNE>.
- [40] C. Giunti, *Four-neutrino mixing*, Proc. of the 23rd Johns Hopkins Workshop on Current Problems in Particle Theory, Neutrinos in the Next Millennium, Johns Hopkins University, Baltimore MD, June 10–12, 1999 [[hep-ph/9909395](#)].
- [41] G.L. Fogli, E. Lisi, A. Marrone and G. Scioscia, *Comments on atmospheric neutrino oscillation scenarios with large $\nu_\mu \leftrightarrow \nu_e$ transitions*, [[hep-ph/9906450](#)].

- [42] C. Giunti, *Four-neutrino scenarios*, talk presented at NOW 2000, Conca Specchiulla (Otranto, Italy), 9–16 Sep. 2000, [[hep-ph/0012236](#)].
- [43] S.M. Bilenky, C. Giunti and W. Grimus, *Neutrino mass spectrum from the results of neutrino oscillation experiments*, *Eur. Phys. J. C* **1** (1998) 247 [[hep-ph/9607372](#)]; *Neutrino oscillation experiments and the neutrino mass spectrum*, proc. of *Neutrino '96*, Helsinki, June 1996, K. Enqvist et al. eds., p. 174, World Scientific, 1997 [[hep-ph/9609343](#)].
- [44] V. Barger, S. Pakvasa, T.J. Weiler and K. Whisnant, *Variations on four-neutrino oscillations*, *Phys. Rev. D* **58** (1998) 093016 [[hep-ph/9806328](#)].
- [45] S.M. Bilenky, C. Giunti, W. Grimus and T. Schwetz, *Four-neutrino mass spectra and the Super-Kamiokande atmospheric up-down asymmetry*, *Phys. Rev. D* **60** (1999) 073007 [[hep-ph/9903454](#)].
- [46] D. Dooling, C. Giunti, K. Kang and C.W. Kim, *Matter effects in four-neutrino mixing*, *Phys. Rev. D* **61** (2000) 073011 [[hep-ph/9908513](#)].
- [47] C. Giunti, M.C. Gonzalez-Garcia and C. Peña-Garay, *Four-neutrino oscillation solutions of the solar neutrino problem*, *Phys. Rev. D* **62** (2000) 013005 [[hep-ph/0001101](#)].
- [48] C. Giunti, M.C. Gonzalez-Garcia and C. Peña-Garay, *Four-neutrino oscillations and the solar neutrino problem*, [hep-ph/0007154](#);
M.C. Gonzalez-Garcia and C. Peña-Garay, *Global and unified analysis of solar neutrino data*, *Nucl. Phys.* **91** (*Proc. Suppl.*) (2000) 80 [[hep-ph/0009041](#)].
- [49] O. Yasuda, *Four neutrino oscillation analysis of the Super-Kamiokande atmospheric neutrino data*, [[hep-ph/0006319](#)].
- [50] A.Y. Smirnov, *Neutrino mass spectrum and lepton mixing*, *Nucl. Phys.* **91** (*Proc. Suppl.*) (2000) 306, <http://nu2000.sno.laurentian.ca>, [[hep-ph/0010097](#)];
O. Peres, *4ν models: conciliating LSND results with solar and atmospheric neutrino data*, talk presented at NOW2000, Otranto, Italy, September 2000, <http://www.ba.infn.it/~now2000>.
- [51] G.L. Fogli, E. Lisi, A. Marrone and D. Montanino, *Analysis of oscillations of atmospheric neutrinos*, *Nucl. Phys.* **91** (*Proc. Suppl.*) (2000) 167, <http://nu2000.sno.laurentian.ca> [[hep-ph/0009269](#)].
- [52] G.L. Fogli, E. Lisi and A. Marrone, *Four-neutrino oscillation solutions of the atmospheric neutrino anomaly*, *Phys. Rev. D* **63** (2001) 053008 [[hep-ph/0009299](#)].
- [53] V. Barger, B. Kayser, J. Learned, T. Weiler and K. Whisnant, *Fate of the sterile neutrino*, *Phys. Lett. B* **489** (2000) 345 [[hep-ph/0008019](#)].
- [54] Unabridged reference list on four-neutrino mixing at <http://www.to.infn.it/~giunti/4nu>

- [55] G.L. Fogli, E. Lisi, A. Marrone and D. Montanino, *Zenith distribution of atmospheric neutrino events and electron neutrino mixing*, *Phys. Lett. B* **425** (1998) 341 [[hep-ph/9711421](#)].
- [56] S.M. Bilenky and C. Giunti, *Implications of CHOOZ results for the decoupling of solar and atmospheric neutrino oscillations*, *Phys. Lett. B* **444** (1998) 379 [[hep-ph/9802201](#)].
- [57] G.L. Fogli, E. Lisi, D. Montanino and A. Palazzo, *Three-flavor msw solutions of the solar neutrino problem*, *Phys. Rev. D* **62** (2000) 013002 [[hep-ph/9912231](#)].
- [58] NOMAD collaboration, Leslie Camilleri, *NOMAD, a status report*, Status Report to the SPSC, 14 september 1999
http://nomadinfo.cern.ch/Public/PUBLICATIONS/public_pub.html
- [59] L. Mikaelian, *Present and future oscillation experiments at reactors*, [[hep-ex/0008063](#)]; *Future reactor neutrino oscillation experiments at krasnoyarsk*, [[hep-ex/0008046](#)]; *Chooz, palo verde, krasnoyarsk*, *Nucl. Phys. B* (Proc. Suppl.) (2000) 284 [[hep-ex/9910042](#)];
L.A. Mikaelian and V.V. Sinev, *Neutrino oscillations at reactors: what next?*, *Phys. Atom. Nucl.* **63** (2000) 1002 [[hep-ex/9908047](#)].
- [60] S. Geer, *Neutrino beams from muon storage rings: characteristics and physics potential*, *Phys. Rev. D* **57** (1998) 6989 [[hep-ph/9712290](#)];
A. Blondel et al., *The neutrino factory : beam and experiment*, CERN-EP-2000-053;
C. Albright et al., *Physics at a neutrino factory*, [[hep-ex/0008064](#)].
- [61] N. Fornengo, M.C. Gonzalez-Garcia and J.W.F. Valle, *Updated global analysis of the atmospheric neutrino data in terms of neutrino oscillations*, *Nucl. Phys. B* **580** (2000) 58 [[hep-ph/0002147](#)].
- [62] R. Foot, *Is the $\nu_\mu \rightarrow \nu_s$ oscillation solution to the atmospheric neutrino anomaly excluded by the superkamiokande data?*, *Phys. Lett. B* **496** (2000) 169 [[hep-ph/0007065](#)].
- [63] D. Dooling, C. Giunti, K. Kang and C.W. Kim, in preparation
- [64] H. Nunokawa, J.T. Peltoniemi, A. Rossi and J.W.F. Valle, *Supernova bounds on resonant active-sterile neutrino conversions*, *Phys. Rev. D* **56** (1997) 1704 [[hep-ph/9702372](#)];
H. Athar and J.T. Peltoniemi, *Signatures of neutrino conversions to sterile states in collapsing stars*, *Phys. Rev. D* **51** (1995) 5785.
- [65] D.O. Caldwell, G.M. Fuller and Y.-Z. Qian, *Sterile neutrinos and supernova nucleosynthesis*, *Phys. Rev. D* **61** (2000) 123005 [[astro-ph/9910175](#)];
G.C. McLaughlin, J.M. Fetter, A.B. Balantekin and G.M. Fuller, *An active-sterile neutrino transformation solution for r-process nucleosynthesis*, *Phys. Rev. D* **59** (1999) 2873 [[astro-ph/9902106](#)].

- [66] R.R. Volkas, *Lepton asymmetries from neutrino oscillations*, *Nucl. Phys.* **91** (Proc. Suppl.) (2000) 487 [[hep-ph/0009162](#)];
P.D. Bari, R. Foot, R.R. Volkas and Y.Y.Y. Wong, *Comment on 'neutrino oscillations in the early universe: how can large lepton asymmetry be generated?'*, [[hep-ph/0008245](#)];
P.D. Bari and R. Foot, *Active-sterile neutrino oscillations and BBN+CMBR constraints*, *Phys. Rev.* **D 63** (2001) 043008 [[hep-ph/0008258](#)].
- [67] D.P. Kirilova and M.V. Chizhov, *Cosmological nucleosynthesis and active-sterile neutrino oscillations with small mass differences: the resonant case*, *Nucl. Phys.* **B 591** (2000) 457 [[hep-ph/9909408](#)]; *On neutrino mixing generated lepton asymmetry and the primordial helium-4 abundance*, [[hep-ph/9908525](#)]; *Neutrino degeneracy effect on neutrino oscillations and primordial helium yield*, *Nucl. Phys.* **B 534** (1998) 447 [[hep-ph/9806441](#)].
- [68] A.D. Dolgov, S.H. Hansen, S. Pastor and D.V. Semikoz, *Neutrino oscillations in the early universe: how large lepton asymmetry can be generated?*, *Astropart. Phys.* **14** (2000) 79 [[hep-ph/9910444](#)].
- [69] S. Esposito, G. Mangano, A. Melchiorri, G. Miele and O. Pisanti, *Testing standard and degenerate big bang nucleosynthesis with boomerang and maxima-1*, *Phys. Rev.* **D 63** (2001) 043004 [[astro-ph/0007419](#)].
- [70] X. dong Shi and G.M. Fuller, *Big bang nucleosynthesis and active sterile neutrino mixing: evidence for maximal $\nu_\mu \leftrightarrow \nu_\tau$ mixing in superkamiokande?*, *Phys. Rev.* **D 59** (1999) 063006 [[astro-ph/9810075](#)];
A reply to comment on 'big bang nucleosynthesis and active-sterile neutrino mixing: evidence for maximal $\nu_\mu \leftrightarrow \nu_\tau$ mixing in Super-Kamiokande?', [[astro-ph/9812232](#)];
K. Abazajian, X. dong Shi and G.M. Fuller, *Active-sterile neutrino mixing in the early universe and primordial nucleosynthesis*, [[astro-ph/9904052](#)];
Neutrino-mixing-generated lepton asymmetry and the primordial he-4 abundance, *Phys. Rev.* **D 60** (1999) 063002 [[astro-ph/9905259](#)].
- [71] S.P. Mikheyev and A.Yu. Smirnov, *Yad. Fiz.*, **42** (1985) 1441; *Resonance amplification of oscillations in matter and spectroscopy of solar neutrinos*, *Sov. J. Nucl. Phys.* **42** (1985) 913; *Il Nuovo Cim. C*, **9** (1986) 17;
L. Wolfenstein, *Neutrino oscillations in matter*, *Phys. Rev.* **D 17** (1978) 2369; *Neutrino oscillations and stellar collapse*, *Phys. Rev.* **D 20** (1979) 2634.