

## Administration of dexamethasone *per os* in finishing bulls. II. Effects on blood parameters used as indicators of animal welfare

A. Marin<sup>1</sup>, G. Pozza<sup>1</sup>, F. Gottardo<sup>2</sup>, L. Moro<sup>1</sup>, A. L. Stefani<sup>1†</sup>, G. Cozzi<sup>2</sup>, M. Brscic<sup>2</sup>, I. Andrighetto<sup>1</sup> and L. Ravarotto<sup>1</sup>

<sup>1</sup>Istituto Zooprofilattico delle Venezie, 35020 Legnaro (PD), Italy; <sup>2</sup>Department of Animal Science, University of Padua, Padua, Italy

(Received 2 August 2007; Accepted 3 March 2008)

---

A set of hormonal, haematological and biochemical parameters was used to evaluate the physiological response and welfare status of 14 finishing Marchigiana bulls treated for 49 days with a low daily dosage (0.75 mg/head per day) of dexamethasone *per os*. Compared to the Control group, dexamethasone decreased cortisol concentrations (42.3 v. 5.7 nmol/l; s.e.d. = 4.17;  $P < 0.001$ ), and led to the reversal of the leukocyte formula in the animals treated ( $P < 0.05$ ). Total serum proteins (70.2 v. 73.9 g/l; s.e.d. = 1.55;  $P < 0.05$ ), in particular  $\beta_1$  globulins (7.5 v. 9.1 g/l; s.e.d. = 0.24;  $P < 0.01$ ) and fibrinogen (199 v. 258 mg/dl; s.e.d. = 32.70;  $P < 0.05$ ), increased as a consequence of treatment. Prolonged dexamethasone administration led the bulls to an apparently chronic stress condition. Moreover, the study indicated various blood parameters that might be used by health officials as effective tools in identifying beef cattle suspected of being illegally treated with dexamethasone.

---

**Keywords:** beef cattle, biochemical parameters, dexamethasone, haematological parameters, hormonal parameters

### Introduction

Public opinion and health institutions are becoming increasingly alarmed by the illegal use of hormones for increased growth and slaughter performance (Courtheyn *et al.*, 2002). Consumers are primarily concerned for their own safety; public health authorities are also apprehensive about animal welfare. Given that growth promoters are currently banned within the European Union, the Report of the Scientific Committee on Animal Health and Animal Welfare (SCAHAW, 2001) did not identify illegal treatment as a potential risk factor for beef cattle welfare. Data from control plans conducted at the national level, however, show that illegal growth promoters are still being administered to fattening bulls. Residue controls conducted at Italian farms and slaughterhouses found percentages of samples positively targeted for corticosteroids of 0.66 and 1.08, respectively (Italian Ministry of Health, 2005).

As previously observed by other authors (Van der Wal *et al.*, 1975; Istasse *et al.*, 1989; Courtheyn *et al.*, 2002), the non-therapeutic use of dexamethasone affects beef cattle growth performance and carcass traits, while its effect on

hormonal, haematological and biochemical parameters is still unknown. Broom (1988), however, reported that the administration of exogenous corticosteroids to animals may trigger an adaptation process that might lead to significant alterations in metabolism and blood parameters.

The potential modification of haematological and endocrine parameters induced by treatment could be used to identify illegally treated animals at the farm level and subsequently confirmed by treatment residue investigations in organic matrices. Moreover, changes in beef cattle blood parameters caused by the administration of low-dosage dexamethasone also indicate animal welfare levels.

### Material and methods

In accordance with Decreto Legislativo n. 116/1992 (1992), the Italian Ministry of Health authorized this study following the submission of a detailed description of the experimental plan by the project's scientific coordinator.

#### *Animals and housing*

In this study, we used 15 finishing Marchigiana bulls with average body weight of  $487 \pm 14.9$  kg housed in five pens of three animals each.

---

† E-mail: astefani@izsvenezie.it

Nine animals in three pens were considered the Control group. One of these bulls was excluded from the experiment at day 30 because of an ocular trauma. The total number of animals considered for data collection was therefore 14. The remaining six bulls (Dexa group) were treated daily with 0.75 mg/head of dexamethasone *per os* (Desashock<sup>®</sup>; Fort Dodge Animal Health SpA, Bologna, Italy). The treatment period started on day 5 and was suspended on day 53, i.e. 3 days prior to slaughter. Further description of the experimental design and treatments, housing system and feeding management are reported in Gottardo *et al.* (2008).

#### *Blood sample collection and analysis*

Individual blood samples were taken from the jugular vein on day 6, 27, 48 and 56. Blood was collected in the morning prior to diet administration, and placed in four separate vacutainer tubes (Becton Dickinson, Meylan Cedex, France), each one dedicated to a specific set of analyses described below.

Test tubes of 10 ml containing K<sub>3</sub>EDTA were used for the collection of the samples submitted to haematological profile determination. The analyses were performed using a Cell Dyn 3500<sup>®</sup> automated analyzer (Abbott Laboratories; Abbott Park, IL, USA). A specific software for bovine blood was utilized to achieve total white blood cell counts (WBC) and differential leukocyte counts, and to evaluate the proportion of different leukocyte types divided in neutrophils (NEU), lymphocytes (LYM), monocytes (MONO), eosinophils (EOS) and basophils (BASO). The neutrophil/lymphocyte ratio (Neu/Lym) was also considered. Parameters correlated to red blood cells (RBC), such as haemoglobin concentration (HGB), haematocrit (HCT), mean corpuscular volume (MCV), mean corpuscular haemoglobin (MCH), mean corpuscular haemoglobin concentration (MCHC) and red cell distribution width (RDW), were investigated, as well as total platelet counts (PLT) and mean platelet volume (MPV).

Blood specimens were collected in heparinized test tubes to detect oxidative stress index glutathione peroxidase (GPx) using a commercial kit (Ransel, Randox Laboratories Ltd, Crumlin, Co. Antrim, UK).

Serum samples, obtained after clot formation in tubes without anticoagulant and centrifugation at 2500 × g at 20°C for 10 min in a refrigerated centrifuge, were used to determine total antioxidant status (TAS) using a commercial kit (Total Antioxidant Status; Randox Laboratories Ltd). Both parameters (GPx and TAS) were automatically analyzed using a BM Hitachi 911 analyzer (Roche, Basel, Switzerland).

The haematic concentration of adrenocorticotrophic hormone (ACTH) was analysed by means of a commercial kit (LKAC1, Medical System, Genoa, Italy) in plasma obtained from K<sub>3</sub>EDTA tubes that were centrifuged at 1500 × g at 4°C for 15 min.

Cortisol concentration was determined in serum blood specimens using a commercial kit (LKCO1; Medical System). Both hormones (ACTH and cortisol) were quantified by chemiluminescent immunometric assays performed with the automated analyzer, Immulite One (Medical System).

Total serum proteins were measured by a colorimetric assay with biuret reagent (TP; Roche) using the automated BM Hitachi 911 analyzer (Roche), while the concentration of albumin and α, β and γ globulin fractions was measured using an automated electrophoresis apparatus Hydrasis LC (Sebia, Issy-les-Moulineaux, France) on 0.8% agarose gel (Hydragel 30 Protein; Sebia).

Test tubes of 5 ml containing Na citrate were used to collect blood samples for fibrinogen determination. Tubes were centrifuged at 1000 × g at 4°C for 15 min and plasma was then separated to measure the fibrinogen concentration and coagulation time with an appropriate commercial kit (HemosIL<sup>™</sup> Fibrinogen-C; Instrumentation Laboratory, Barcelona, Spain). The analysis was performed with the automated coagulometer ACL 7000 (Instrumentation Laboratory).

#### *Statistical analysis*

Experimental data were submitted to statistical analysis using a linear model within PROC-GLM of SAS (1990). The model considered the effects of treatment, animal within-treatment, sample collection day, and the interaction between treatment and sample collection day. The effect of treatment was tested using animal within treatment variance as the error term. Means differences were considered statistically significant at  $P < 0.05$ .


## Results

#### *Endocrine parameters*

Bulls treated with dexamethasone showed significantly lower haematic cortisol concentrations than Control animals (5.7 v. 42.3 nmol/l; s.e.d. = 4.17). A significant treatment per sample collection day interaction (Figure 1) showed the peaking of cortisol concentration in the Control group, occurring in the first 2 days of sampling, before falling to the constantly low level of the bulls treated. Regardless of the day of sampling, however, the average cortisol concentration in the Dexa group was always very low in the vicinity of the instrumental detection threshold (5.5 nmol/l). Another significant treatment per sample collection day interaction was regarding ACTH concentration (Figure 1). In the Control bulls, ACTH levels were higher at first sampling and then decreased, while the bulls treated showed a significant peak after 48 days.

#### *Haematological profile*

Dexamethasone administration provided no significant effect on the variables regarding erythrocytes such as RBC, HGB, HCT, MCV, MCH, MCHC and RDW, or platelets such as PLT and MPV (Table 1). As shown in Figure 2, dexamethasone treatment did not affect the concentration of several leukocyte parameters, such as WBC, MONO, EOS or BASO. Instead, the lymphocytic count was always lower in the Dexa group than for the Control bulls. Neutrophil concentration showed an opposite trend until the third sampling day, whereas at the last collection, performed



**Figure 1** Plasma concentration of cortisol (s.e.d. = 4.17) and adrenocorticotrophic hormone (s.e.d. = 21.00) in Control bulls (—●—) and in animals treated with dexamethasone *per os* (- -□- -) from day 5 to day 53 of the trial. \*\*Least square means within sampling day are significantly different for  $P < 0.01$ . \*\*\*Least square means within sampling day are significantly different for  $P < 0.001$ .

**Table 1** Haematological profile in Control bulls and in animals treated with dexamethasone *per os* (Dexa)

Items	Unit	Treatment		Significance		s.e.d.
		Control	Dexa	Treatment	Day	
Red blood cell	M/ $\mu$ l	7.98	8.13	ns	***	0.33
Haemoglobin concentration	g/dl	12.4	12.3	ns	***	0.40
Haematocrit	%	35.4	35.9	ns	***	1.28
Mean corpuscular volume	fl	44.7	44.5	ns	***	2.08
Mean corpuscular haemoglobin	pg	15.6	15.2	ns	***	0.68
Mean corpuscular haemoglobin concentration	g/dl	34.8	34.2	ns	*	0.29
Red cells distribution width	%	22.5	23.9	ns	ns	0.76
Number of platelets	K/ $\mu$ l	414	414	ns	***	70.31
Mean platelet volume	fl	4.45	4.77	ns	***	0.29

No significant differences were observed for treatment  $\times$  day of sampling interaction.  
 ns = Least square means within row are not significantly different.  
 \*Least square means within row are significantly different for  $P < 0.05$ .  
 \*\*\*Least square means within row are significantly different for  $P < 0.001$ .

3 days after the suspension of treatment, no further difference was detected between the two groups. Treated animals showed an increase of the Neu/Lym ratio, which was significantly higher at 27 and 48 days of trial (Figure 2).

**Oxidative parameters**

Although corticosteroid treatment had no effect on GPx concentration (Figure 3), a significant treatment per sample collection day interaction was observed for TAS as a result of the increased concentration measured in the Dexa group at sampling day 27 (Figure 3).

**Serum protein profile**

Significant differences between the Control and Dexa groups were observed for a number of serum profile parameters (Table 2). Total serum protein concentration increased in bulls receiving treatment at the second sampling day (Figure 4). Regarding  $\beta_1$  globulins, apart from the first sampling performed the day after treatment began, there was difference between the two groups showing another significant treatment per sample collection day interaction (Figure 4). The remaining variables were not modified by treatment, and in most cases, only a sample collection day effect was observed (Table 2).

**Coagulation parameters**


Fibrinogen concentration was significantly higher in the Dexa group from the second sample collection day until the end of the experimental period, and this explains the significant treatment per sample collection day interaction ( $P < 0.05$ ) observed (Figure 5).

Fibrinogen coagulation time was lower in the animals treated and statistical differences occurred from the second sample until the end of the experimental period (Figure 5).


**Discussion**

A daily oral low dosage (0.75 mg/head) administration of dexamethasone to beef cattle has been shown to modify certain blood parameters used as animal welfare indicators (Broom and Johnson, 1993). Such treatment has been observed to lower the production of endogenous cortisol below physiological values in bovine species (Kaneko *et al.*, 1997) and this result is consistent with those obtained by Cagnasso *et al.* (1987) in calves, by Lopes *et al.* (2004) in growing pigs and by Allersmeier *et al.* (2005) in horses using pour-on posology. Dexamethasone, acting as endogenous cortisol, inhibits hypothalamus and hypophysis activities (Sacher and McPherson, 2000), exerting a negative effect on ACTH secretion (Rijnberk and Mol, 1997). Low

## Dexamethasone treated bulls: blood parameters



**Figure 2** Plasma concentration of white blood cells (s.e.d. = 0.74), neutrophils (s.e.d. = 0.44), lymphocytes (s.e.d. = 0.38), monocytes (s.e.d. = 0.14), eosinophils (s.e.d. = 0.06), basophils (s.e.d. = 0.01) and neutrophils/lymphocytes ratio (s.e.d. = 0.13) in Control bulls (—●—) and in animals treated with dexamethasone *per os* (- -□- -) from day 5 to day 53 of the trial. \*Least square means within sampling day are significantly different for  $P < 0.05$ . \*\*Least square means within sampling day are significantly different for  $P < 0.01$ . \*\*\*Least square means within sampling day are significantly different for  $P < 0.001$ .



**Figure 3** Plasma concentration of total antioxidant status (TAS; s.e.d. = 0.04) and glutathione peroxidase (GPx; s.e.d. = 82.03) in Control bulls (—●—) and in animals treated with dexamethasone *per os* (- -□- -) from day 5 to day 53 of the trial. \*\*Least square means within sampling day are significantly different for  $P < 0.01$ .

haematic cortisol levels usually indicate high-quality animal welfare and suggest satisfactory animal ability to cope with the surrounding environment (Moberg, 1985). This theory could not be applied in this study, however, because dexa-


methasone as a long-term stressor would have impaired adrenal function and animal welfare in consequence.

Moreover, both Cagnasso *et al.* (1987) and Allersmeier *et al.* (2005), working, respectively, with calves and horses,


**Table 2** Serum protein profile in Control bulls and in animals treated with dexamethasone per os (Dexa)

Items	Unit	Treatment			Significance		s.e.d.
		Control	Dexa	Treatment	Day	Treatment × Day	
TP	g/l	70.2	73.9	*	***	ns	1.55
Albumin	g/l	31.5	33.2	ns	***	ns	1.10
α <sub>1</sub>	g/l	4.1	3.9	ns	***	ns	0.13
α <sub>2</sub>	g/l	6.9	7.3	ns	ns	ns	0.33
β <sub>1</sub>	g/l	7.5	9.1	***	***	**	0.24
β <sub>2</sub>	g/l	4.3	4.0	ns	***	ns	0.32
γ	g/l	16.0	16.4	ns	*	ns	1.01
A/G		0.81	0.82	ns	*	ns	0.04

TP = total protein; A/G = albumin/globulin ratio.  
 ns = Least square means within row are not significantly different.  
 \*Least square means within row are significantly different for  $P < 0.05$ .  
 \*\*Least square means within row are significantly different for  $P < 0.01$ .  
 \*\*\*Least square means within row are significantly different for  $P < 0.001$ .



**Figure 4** Serum concentration of total protein (s.e.d. = 1.55) and β<sub>1</sub> globulin fraction (s.e.d. = 0.24) in Control bulls (—●—) and in animals treated with dexamethasone per os (- -□ - -) from day 5 to day 53 of the trial. \*Least square means within sampling day are significantly different for  $P < 0.05$ . \*\*Least square means within sampling day are significantly different for  $P < 0.01$ . \*\*\*Least square means within sampling day are significantly different for  $P < 0.001$ .



**Figure 5** Plasma concentration of fibrinogen (s.e.d. = 32.70) and fibrinogen related coagulation time (s.e.d. = 1.93) in Control bulls (—●—) and in animals treated with dexamethasone per os (- -□ - -) from day 5 to day 53 of the trial. \*Least square means within sampling day are significantly different for  $P < 0.05$ . \*\*Least square means within sampling day are significantly different for  $P < 0.01$ . \*\*\*Least square means within sampling day are significantly different for  $P < 0.001$ .

described a negative effect on ACTH levels ascribed to dexamethasone administration. This effect was not unequivocally observed in our study, in which a significantly decreased ACTH concentration in the Dexa group was observed only on the first sampling day.

Considering erythrocyte and platelet variables, in this study their mean values fell all within the reference ranges (Kramer, 2000) in both treated and untreated animals. Several authors reported that short-term stressors such as transport, lairage at the slaughterhouse and slaughtering procedures have been shown to affect erythrocyte-related

parameters in cattle (Dunn, 1990; Chacon *et al.*, 2005; Tadich *et al.*, 2005). If the prolonged low dosage dexamethasone administration used in this study is considered a long-term stressor, changes in these parameters would be logically expected in the bulls treated. No alteration in RBC parameters was observed, but this might be ascribed to either the low-dosage adopted or animal adaptation to the frequent handling required for daily oral corticosteroids administration. For this reason, haematological parameters do not appear to be a suitable welfare status indicator for cattle exposed to long-term stress. This hypothesis is

also supported by Gupta *et al.* (2005), whose results did not show significant alterations in steer haematological parameters when animals were subjected to frequent re-penning and re-grouping.

Activation of the hypothalamic–pituitary–adrenal (HPA) axis has been shown to weaken the immune system and this could have a significant impact on animal welfare through the hyperactivity of the adrenal cortex (Raussi *et al.*, 2006). As a corticosteroid, dexamethasone affects the HPA axis (Rijnberk and Mol, 1997), and several studies conducted on cattle and veal calves have shown that such molecules might lead to leukocytosis, neutrophilia, eosinopenia, lymphopenia and monocytosis, as well as to an increase in the Neu/Lym ratio (Lan *et al.*, 1995; Anderson *et al.*, 1999; Thanasak *et al.*, 2004). Oldham and Howard (1992) have shown that daily intramuscular injections of dexamethasone at 0.5 mg/kg per day in calves for 20 days resulted in neutrophilia and lymphopenia. In our study, the administration of Dexamethasone *per os* at a low dosage led to a higher Neu/Lym ratio as a consequence of an increased number of neutrophils and a significant decrease of lymphocytes, even if their absolute values are in the physiological range reported by Kramer (2000). The reversal of the leukocytic formula demonstrates that exogenous corticosteroid treatments can produce effects similar to those caused by environmental stress (Allen *et al.*, 2000).

Neutrophil concentration seems to respond to dexamethasone treatment because it decreased in the Dexamethasone group to the same level as that measured in Control bulls 3 days after the suspension of treatment. This result supports previous findings observed in calves by Lan *et al.* (1995), who demonstrated that the neutrophil count returned to physiological levels 3 days after the last 0.04 mg/kg injection of dexamethasone.

The limited effect of dexamethasone on both oxidative stress parameters considered in this study (GPx and TAS) might be explained by the animal's attempt to maintain the balance between plasma pro-oxidant and antioxidant constituents (Miller *et al.*, 1993; Bourdon and Blanche, 2001), which implies a dynamic response. For this reason, GPx and TAS do not appear to be useful biochemical tools for the detection of cattle treated illegally by corticosteroids.

Confirming previous findings in dairy cows by Furll and Leidel (2002), dexamethasone administration to the experimental bulls increased their total serum protein concentration. Our study showed that the highest levels of total protein in treated bulls might be related to the increase of  $\beta_1$  globulin fraction. This parameter might be affected either by the higher concentration of fibrinogen, an inflammatory acute phase protein stimulated by dexamethasone acting as a stressor or by an increase in other acute phase proteins (Sevaljevic *et al.*, 1998; Sorrells *et al.*, 2007).

## Conclusion

This study conducted on a large number of physiological parameters proposed as indicators of cattle welfare found

that only a few are significantly affected by the low-dosage administration of dexamethasone. The main alterations induced by treatment were decreased cortisol concentration to the sub-physiological level, the reversal of the leukocytic formula, and increased total serum protein promoted by  $\beta_1$  globulins and fibrinogen levels. All these parameters appear useful in identifying suspected illegally treated bulls at the farm level. Other blood variables such as erythrocyte-related parameters or neutrophil counts were either unaltered by treatment or returned rapidly to physiological levels after suspension of treatment.

Low-dosage dexamethasone administration for a prolonged period of around 50 days produced chronic stress conditions and left evident metabolic traces in the bulls tested. Beef farmers should therefore take notice that illegal non-therapeutic dexamethasone treatment during the finishing period is easily detectable and can be confirmed by treatment residue investigations in organic matrices.

## Acknowledgements

This study was financed by the Italian Ministry of Health. The authors are grateful to Dr Cristiano Ossensi, Mr Roberto Rampazzo and Mr Roberto Chimetto for their precious technical support.

## References

- Allen DG, Pringle JK and Smith DA 2000. Handbook of veterinary drugs. Lippincott Williams & Wilkins, Hagerstown, MD, USA.
- Allersmeier M, Abraham G, Schusser GF, Hoppen HO and Ungemach FR 2005. Effects of topically applied dexamethasone on ACTH and cortisol release in horses. *Tierärztliche Umschau* 60, 664–670.
- Anderson BH, Watson DL and Colditz IG 1999. The effect of dexamethasone on some immunological parameters in cattle. *Veterinary Research Communications* 23, 399–413.
- Bourdon E and Blanche D 2001. The importance of proteins in defence against oxidation. *Antioxidants and Redox Signaling* 3, 293–311.
- Broom DM 1988. The scientific assessment of animal welfare. *Applied Animal Behaviour Science* 20, 5–19.
- Broom DM and Johnson KG 1993. *Stress and Animal Welfare*, 1st edition. Chapman and Hall, London, UK.
- Cagnasso A, Dotta U, Aria G, Monti F and Prato S 1987. Evaluation of adrenal cortex function in pre-ruminant fattening calves treated with dexamethasone. *Schweizer Archiv für Tierheilkunde* 129, 429–435.
- Chacon G, Garcia-Belenguer S, Villarroel M and Maria GA 2005. Effect of transport stress on physiological responses of male bovines. *Deutsche Tierärztliche Wochenschrift* 112, 465–469.
- Courtheyn D, Le Bizec B, Brambilla G, De Brabander HF, Cobbeart E, Van de Wiele M, Vercammen J and De Wasch K 2002. Recent developments in the use and abuse of growth promoters. *Analytica Chimica Acta* 473, 71–82.
- Decreto Legislativo n. 116/1992 1992. Protezione degli animali utilizzati a fini sperimentali o ad altri fini scientifici. Supplemento Ordinario – *Gazzetta Ufficiale della Repubblica Italiana* n. 40.
- Dunn CS 1990. Stress reactions of cattle undergoing ritual slaughter using two methods of restraint. *The Veterinary Record* 126, 522–525.
- Furll M and Leidel I 2002. Studies on the stabilization of health status in periparturient dairy cows. *Tierärztliche Umschau* 57, 423–438.
- Gottardo F, Brscic M, Pozza G, Ossensi C, Contiero B, Marin A and Cozzi G 2008. Administration of dexamethasone *per os* in finishing bulls: I. Effects on productive traits, meat quality and cattle behaviour as indicator of welfare. *Animal* 2, 1073–1079.
- Gupta S, Earley B, Ting STL and Crowe MA 2005. Effect of repeated regrouping and relocation on the physiological, immunological, and

- haematological variables and performance of steers. *Journal of Animal Science* 83, 1948–1958.
- Istasse L, De Haan V, Van Eenaeme C, Buts B, Baldwin P, Gielen M, Demeyer D and Bienfait JM 1989. Effect of dexamethasone injections on performance in a pair of monozygotic cattle twin. *Journal of Animal Physiology and Animal Nutrition* 62, 150–158.
- Italian Ministry of Health 2005. Report of National Residues Plan of the year 2003. Retrieved January 20, 2005, from <http://www.regione.veneto.it/Servizi+alla+Persona/Sanita/Prevenzione/Sanit%C3%A0+veterinaria/Piano+Nazionale+Residui.htm>
- Kaneko JJ, Harvey JW and Bruss ML 1997. *Clinical Biochemistry of Domestic Animals*, 5th edition. Academic Press, San Diego, CA, USA.
- Kramer JW 2000. Normal Hematology of Cattle, Sheep, and Goats. In *Schalm's Veterinary Hematology*, 5th edition (ed. BF Felman, JG Zinkl and NC Jain), pp. 1075–1084. Lippincott Williams and Wilkins, Baltimore, MD, USA.
- Lan HC, Reddy PG, Chambers MA, Walker G, Srivastava KK and Ferguson JA 1995. Effect of stress on interleukin-2 receptor expression by bovine mononuclear leukocytes. *Veterinary Immunology and Immunopathology* 49, 241–249.
- Lopes SO, Claus R, Lacorn M, Wagner A and Mosenthin R 2004. Effects of dexamethasone application in growing pigs on hormones, N-retention and other metabolic parameters. *Journal of Veterinary Medicine Series A* 51, 97–105.
- Miller JK, Brzezinska-Slebodzinska E and Madsen FC 1993. Oxidative stress, antioxidants, and animal function. *Journal of Dairy Science* 76, 2812–2823.
- Moberg GP 1985. Biological response to stress: key to assessment of animal well-being. In *Animal stress* (ed. GP Moberg), pp. 27–49. American Physiological Society, Bethesda, MD, USA.
- Oldham G and Howard CJ 1992. Suppression of bovine lymphocyte responses to mitogens following in vivo and in vitro treatment with dexamethasone. *Veterinary Immunology and Immunopathology* 30, 161–177.
- Raussi S, Boissy A, Andanson S, Kaihilahti J, Pradel P and Veissier I 2006. Repeated regrouping of pair-housed heifers around puberty affects their behavioural and HPA axis reactivities. *Animal Research* 55, 131–144.
- Rijnberk A and Mol J 1997. Adrenocortical function. In *Clinical Biochemistry of Domestic Animals*, 5th edition (ed. JJ Kaneko, JW Harvey and MG Bruss), pp. 553–570. Academic Press, San Diego, CA, USA.
- Sacher RA and McPherson RA 2000. *Widmann's Clinical Interpretation of Laboratory Tests*, 11th edition. Davis FA Company, Philadelphia, PA, USA.
- SAS 1990. *User's Guide: Statistic*. SAS Institute, Inc., Cary, NC, USA.
- SCAHAW – Scientific Committee on Animal Health and Animal Welfare 2001. The welfare of cattle kept for beef production. Retrieved April 25, 2001, from [http://europa.eu.int/com/food/fs/sc/scsah/outcom\\_en.html](http://europa.eu.int/com/food/fs/sc/scsah/outcom_en.html)
- Sevaljevic L, Macvanin M, Zakula Z, Kanazir DT and Ribarac-Septic N 1998. Adrenalectomy and dexamethasone treatment alter the patterns of basal and acute phase response-induced expression of acute phase protein genes in rat liver. *Journal of Steroid Biochemistry and Molecular Biology* 66, 347–353.
- Sorrells AD, Eicher SD, Harris MJ, Pajor EA and Richert BT 2007. Periparturient cortisol, acute phase cytokine, and acute phase protein profiles of gilts housed in groups or stalls during gestation. *Journal of Animal Science* 85, 1750–1757.
- Tadich N, Gallo C, Bustamante H, Schwerter M and van Schaik G 2005. Effects of transport and lairage time on some blood constituents of Fresian-cross steers in Chile. *Livestock Production Sciences* 93, 223–233.
- Thanasak J, Jorritsma R, Hoek A, Noordhuizen JPTM, Rutten VPMG and Müller KE 2004. The effects of a single injection of dexamethasone-21-isonicotinate on the lymphocyte functions of dairy cows at two weeks post partum. *Veterinary Research* 35, 103–112.
- Van der Wal P, Berende PLM and Sprietsma JE 1975. Effect of anabolic agents on performance of calves. *Journal of Animal Science* 41, 978–985.