

On the scalar potential of minimal flavour violation

R. Alonso,^a M.B. Gavela,^a L. Merlo^{b,c} and S. Rigolin^d

^a*Departamento de Física Teórica, Universidad Autónoma de Madrid
and Instituto de Física Teórica IFT-UAM/CSIC,
Cantoblanco, 28049 Madrid, Spain*

^b*Physik-Department, Technische Universität München,
James-Franck-Strasse, D-85748 Garching, Germany*

^c*TUM Institute for Advanced Study, Technische Universität München,
Lichtenbergstrasse 2a, D-85748 Garching, Germany*

^d*Dipartimento di Fisica Galileo Galilei, Università di Padova and INFN, Sezione di Padova,
Via Marzolo 8, I-35131 Padua, Italy*

E-mail: rodrigo.alonso@uam.es, belen.gavela@uam.es,
luca.merlo@ph.tum.de, stefano.rigolin@pd.infn.it

ABSTRACT: Assuming the Minimal Flavour Violation hypothesis, we derive the general scalar potential for fields whose background values are the Yukawa couplings. We analyze the minimum of the potential and discuss the fine-tuning required to dynamically generate the mass hierarchies and the mixings between different quark generations. Two main cases are considered, corresponding to Yukawa interactions being effective operators of dimension five or six (or, equivalently, resulting from bi-fundamental and fundamental scalar fields, respectively). At the renormalizable and classical level, no mixing is naturally induced from dimension five Yukawa operators. On the contrary, from dimension six Yukawa operators one mixing angle and a strong mass hierarchy among the generations result.

KEYWORDS: Beyond Standard Model, Quark Masses and SM Parameters, Global Symmetries

ARXIV EPRINT: [1103.2915](https://arxiv.org/abs/1103.2915)

Contents

1	Introduction	1
2	Two family case	5
2.1	$d = 5$ Yukawa operators: the bi-fundamental approach	6
2.1.1	The scalar potential at the renormalizable level	7
2.1.2	The scalar potential at the non-renormalizable level	10
2.2	$d = 6$ Yukawa operator: the fundamental approach	11
2.2.1	The scalar potential	13
2.2.2	The first generation	14
3	The three-family case	15
3.1	$d = 5$ Yukawa operator: the bi-fundamental approach	15
3.2	$d = 6$ Yukawa operator: the fundamental approach	17
3.3	Combining fundamentals and bi-fundamentals	18
4	Conclusions	19
A	$d = 6$ operators in the bifundamental approach	21
B	A fine-tuned scalar potential in the bifundamental approach	22
B.1	Minimization of the scalar potential	23
B.2	Three family case	25
C	The scalar potential for the fundamental approach	28
D	Note added in proof	29

1 Introduction

After years of intense searches, all flavour processes observed in the hadronic sector, from rare decays measurements in the kaon and pion sectors to superB-factories results, are well in agreement with the expectations of the Standard Model of particle physics (SM). To say that all flavour processes are consistent with the SM predictions is tantamount to state that all flavour effects observed until now are consistent with being generated through the Yukawa couplings, which are the sole vehicles of flavour and CP violation in the SM.

Nevertheless, the origin of fermion masses and mixings remains the most unsatisfactory question in the visible sector of nature: it involves important fine-tunings and lack of predictivity, as essentially for each mass or mixing angle a new parameter is added by

hand to the SM. It is commonly expected that an underlying dynamics will provide a rationale for the observed patterns.

The hypothesis of *Minimal Flavour Violation* (MFV) [1] is a humble, matter-of-fact, and highly predictive working frame built only on: i) the assumption that, at low energies, the Yukawa couplings are the only sources of flavour and CP violation both in the SM and in whatever may be the flavour theory beyond it, abiding in this way to the experimental indications mentioned above; ii) the use of the flavour symmetries which the SM exhibits in the limit of vanishing Yukawa couplings.

Indeed, the hadronic part of the SM Lagrangian, in the absence of quark Yukawa terms, exhibits a flavour symmetry given by

$$G_f = \text{SU}(3)_{Q_L} \times \text{SU}(3)_{U_R} \times \text{SU}(3)_{D_R}, \quad (1.1)$$

plus three extra U(1) factors corresponding to the baryon number, the hypercharge and the Peccei-Quinn symmetry [2]. The non-abelian subgroup G_f controls the flavour structure of the Yukawa matrices, and we focus on it for the remainder of this paper. Under G_f , the $\text{SU}(2)_L$ quark doublet, Q_L , and the $\text{SU}(2)_L$ quark singlets, U_R and D_R , transform as:

$$Q_L \sim (3, 1, 1), \quad U_R \sim (1, 3, 1), \quad D_R \sim (1, 1, 3). \quad (1.2)$$

The SM Yukawa interactions break explicitly the flavour symmetry:

$$\mathcal{L}_Y = \bar{Q}_L Y_D D_R H + \bar{Q}_L Y_U U_R \tilde{H} + \text{h.c.} \quad (1.3)$$

The technical realization of the MFV ansatz promotes the Yukawa couplings $Y_{U,D}$ to be spurion fields which transform under G_f as

$$Y_U \sim (3, \bar{3}, 1), \quad Y_D \sim (3, 1, \bar{3}), \quad (1.4)$$

recovering the invariance under G_f of the full SM Lagrangian. Following the usual MFV convention for the Yukawas, one defines

$$Y_D = \begin{pmatrix} y_d & 0 & 0 \\ 0 & y_s & 0 \\ 0 & 0 & y_b \end{pmatrix}, \quad Y_U = \mathcal{V}_{\text{CKM}}^\dagger \begin{pmatrix} y_u & 0 & 0 \\ 0 & y_c & 0 \\ 0 & 0 & y_t \end{pmatrix}, \quad (1.5)$$

with \mathcal{V}_{CKM} being the usual quark mixing matrix, encoding three angles and one CP-odd phase.

MFV is not a model of flavour and the value of the new dynamical flavour scale Λ_f is not fixed: it does not determine the energy scale at which new flavour effects will show up. Nevertheless it is quite successful in predicting precise and constrained relations between different flavour transitions, to be observed whenever the new physics scale becomes experimentally accessible [3]. The reason is that in the MFV framework the coefficients of all SM-gauge invariant operators have a fixed flavour structure in terms of Yukawa couplings, so as to make the operator invariant under G_f , plus the fact that the top Yukawa coupling may dominate any coefficient in which it participates.¹

¹This is modified, though, in some MFV versions such as two-Higgs doublet models [1] with extra discrete symmetries [4], or in models with strong dynamics [5].

MFV sheds also an interesting light on the relative size of the electroweak and the flavour scale. The origin of all visible masses and the family structure are the two major unresolved puzzles of the SM and it is unknown whether a relation exists between the nature and size of those two scales. While the electroweak data, and the theoretical fine-tunings they require, suggest that new physics should appear around the TeV scale, traditional model-independent limits on the flavour scale Λ_f point to order(s) of magnitude larger values [6]. Within MFV both sizes could be reconciled instead around the TeV scale, due to the Yukawa suppression of the flavour-changing operator coefficients. This holds either assuming only the SM as the renormalizable theory [1] or in beyond the SM scenarios (BSM), such as supersymmetric [8] or extradimensional [9] versions of the MFV ansatz.²

It is unlikely that MFV holds at all scales [7]. MFV assumes a new dynamical scale Λ_f , which points to MFV being just an accidental low-energy property of the theory. In this sense, MFV implicitly points to a dynamical origin for the values of the Yukawa couplings. The latter may correspond to the vacuum expectation values (vevs) of elementary or composite fields or combinations of them. In other words, the spurions may be promoted to fields, usually called flavons. For instance, in the first formulation of MFV by Chivukula and Georgi [10], the Yukawa couplings corresponded to a fermion condensate. In this work, we further explore the dynamical character of the flavons, in a rather model-independent way.

The Yukawa interactions may be then seen as effective operators of dimension larger than four — denominated *Yukawa operators* in what follows — weighted down by powers of the large flavour scale³ Λ_f . The precise dimension d of the Yukawa operators is not determined, as illustrated in figure 1. As long as the vev to be taken by the flavon fields is smaller than Λ_f , an analysis ordered by inverse powers of this scale is a sensible approach. The simplest case is that of a $d = 5$ operator:

$$\mathcal{L}_Y = \bar{Q}_L \frac{\Sigma_d}{\Lambda_f} D_R H + \bar{Q}_L \frac{\Sigma_u}{\Lambda_f} U_R \tilde{H} + \text{h.c.} , \tag{1.6}$$

with the scalar flavons Σ_d and Σ_u being dynamical fields in the bi-fundamental representation of G_f (i.e. $\Sigma_u \sim (3, \bar{3}, 1)$ and $\Sigma_d \sim (3, 1, \bar{3})$, see eq. (1.4)) such that⁴

$$Y_D \equiv \frac{\langle \Sigma_d \rangle}{\Lambda_f} , \quad Y_U \equiv \frac{\langle \Sigma_u \rangle}{\Lambda_f} . \tag{1.7}$$

²The BSM theory may introduce more than one distinct flavour scale: this work sticks to a conservative and minimalist approach, focusing on the physics related to Λ_f as described above.

³For instance, a possible realization among many takes Λ_f to be the mass of heavy flavour mediators in some BSM theory [11, 12]: at energies $E < \Lambda_f$, they can be integrated out resulting in $d > 4$ operators involving the SM fields and the flavons.

⁴The Goldstone bosons that would result from the spontaneous breaking of a continuous global flavour symmetry, may be avoided for instance by gauging the symmetry. In practical realizations, this in turn tends to induce dangerous flavour-changing neutral currents mediated by the new gauge bosons. A new promising avenues to cope with this problem has been recently proposed in ref. [13, 14].




Figure 1. Effective Yukawa coupling.

An alternative realization, that we also explore below, is that of a $d = 6$ Yukawa operator, involving generically two scalar flavons for each spurion,

$$\mathcal{L}_Y = \bar{Q}_L \frac{\chi_d^L \chi_d^{R\dagger}}{\Lambda_f^2} D_R H + \bar{Q}_L \frac{\chi_u^L \chi_u^{R\dagger}}{\Lambda_f^2} U_R \tilde{H} + \text{h.c.}, \quad (1.8)$$

which provide the following relations between Yukawa couplings and vevs:

$$Y_D \equiv \frac{\langle \chi_d^L \rangle \langle \chi_d^{R\dagger} \rangle}{\Lambda_f^2}, \quad Y_U \equiv \frac{\langle \chi_u^L \rangle \langle \chi_u^{R\dagger} \rangle}{\Lambda_f^2}. \quad (1.9)$$

In this interesting case, the flavons are simply vectors under the flavour group, alike to quarks, with the simplest quantum number assignment being $\chi_{u,d}^L \sim (3, 1, 1)$, $\chi_u^R \sim (1, 3, 1)$ and $\chi_d^R \sim (1, 1, 3)$. Following this pattern, would the Yukawa couplings result from a condensate of fermionic flavons [10], a $d = 7$ Yukawa operator could be adequate

$$Y_D \equiv \frac{\langle \bar{\Psi}_d^L \Psi_d^R \rangle}{\Lambda_f^3}, \quad Y_U \equiv \frac{\langle \bar{\Psi}_u^L \Psi_u^R \rangle}{\Lambda_f^3}, \quad (1.10)$$

with fermions quantum numbers under G_f as in the previous case. Notice that these realizations in which the Yukawa couplings correspond to the vev of an aggregate of fields, rather than to a single field, are not the simplest realization of MFV as defined in ref. [28], while still corresponding to the essential idea that the Yukawa spurions may have a dynamical origin.

The goal of this work is to address the problem of the determination of the general scalar potential, compatible with the flavour symmetry G_f , for the flavon fields denoted above by Σ or χ . An interesting question is whether it is possible to obtain the SM Yukawa pattern — i.e. the observed values of quark masses and mixings — with a renormalizable potential. We derive the potential, analyze the possible vacua, and discuss the degree of “naturalness” of the possible solutions. It will be shown that the possibility of obtaining a large mass hierarchy and mixing at the renormalizable level varies much depending on the dimension of the Yukawa operator. The role played by non-renormalizable terms and the fine-tunings required to accommodate the full spectrum will be explored.

A relevant issue is what will be meant by natural: following 't Hooft's naturalness criteria, all dimensionless free parameters of the potential not constrained by the symmetry should be of order one, and all dimensionful ones are expected to be of the order of the scale(s) of the theory. We will thus explore in which cases — if any — those criteria allow that the minimum of the MFV potential corresponds automatically to mixings and large mass hierarchies. Stronger than $O(10\%)$ adjustments (typical Clebsh-Gordan values in any theory) will be considered fine-tuned.

It is worth to note that the structure of the scalar potentials constructed here is more general than the particular effective realization in eqs. (1.6) and (1.8). Indeed, it relies exclusively on invariance under the symmetry G_f and on the flavon representation, bi-fundamental or fundamental.⁵

We limit our detailed discussions below to the quark sector. The implementation of MFV in the leptonic sector [15, 16] requires some supplementary assumptions, as Majorana neutrino masses require to extend the SM and involve a new scale: that of lepton number violation. Due to the smallness of neutrino masses, the *effective* scale of lepton number violation must be distinct from the flavour and electroweak ones, if new observable flavour effects are to be expected [17]. Nevertheless, the analysis of the flavon scalar potential performed below may also apply when considering leptons, although the precise analysis and implications for the leptonic spectrum will be carried out elsewhere.

The structure of the manuscript is as follows. In section 2, for the two-family case we analyze the renormalizable potential for $d = 5$ and $d = 6$ Yukawa operators, or in other words of flavons in the bi-fundamental and in the fundamental of G_f , respectively, showing that in the latter case mixing and a strong hierarchy are intrinsically present. The corrections induced by non-renormalizable terms are also discussed. In section 3 the analogous analyses are carried out for the realistic three-family case and it is also discussed the qualitative new features appearing when considering simultaneously $d = 5$ and $d = 6$ Yukawa operators. The conclusions are presented in section 4. Details of the analytical and numerical discussions of the potential minimization can be found in the appendices.

2 Two family case

We start the discussion of the general scalar potential for the MFV framework by illustrating the two-family case, postponing the discussion of three families to the next section. Even if we restrict to a simplified case, with a smaller number of Yukawa couplings and mixing angles, it is a very reasonable starting-up scenario, that corresponds to the limit in which the third family is decoupled, as suggested by the hierarchy between quark masses and the smallness of the CKM mixing angles⁶ θ_{23} and θ_{13} . In this section, moreover, we will introduce most of the conventions and ideas to be used later on for the three-family analysis.

⁵For instance, the potential and the consequences for mixing obtained in this work will apply as well to the construction in ref. [13], notwithstanding the fact that there the flavon vevs show an inverse hierarchy than that for the minimal version of MFV, as they are proportional to the inverse of the SM Yukawa couplings.

⁶We follow in this paper the PDG [18] conventions for the CKM matrix parametrization.

With only two generations the non-Abelian flavour symmetry group, G_f , is reduced to

$$G_f = \text{SU}(2)_{Q_L} \times \text{SU}(2)_{U_R} \times \text{SU}(2)_{D_R}, \quad (2.1)$$

under which the quark fields transform as

$$Q_L \sim (2, 1, 1), \quad U_R \sim (1, 2, 1), \quad D_R \sim (1, 1, 2). \quad (2.2)$$

Following the MFV prescription, in order to preserve the flavour symmetry in the Lagrangian, the Yukawa spurions introduced in eq. (1.4) now transform under G_f as

$$Y_U \sim (2, \bar{2}, 1), \quad Y_D \sim (2, 1, \bar{2}). \quad (2.3)$$

The masses of the first two generations and the mixing angle among them arise once the spurions take the following values:

$$Y_D = \begin{pmatrix} y_d & 0 \\ 0 & y_s \end{pmatrix}, \quad Y_U = \mathcal{V}_C^\dagger \begin{pmatrix} y_u & 0 \\ 0 & y_c \end{pmatrix}, \quad (2.4)$$

where

$$\mathcal{V}_C = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} \quad (2.5)$$

is the usual Cabibbo rotation among the first two families.

2.1 $d = 5$ Yukawa operators: the bi-fundamental approach

The most intuitive approach, in looking for a dynamical origin of MFV, is probably to promote each Yukawa coupling from a simple spurion to a flavon field. In other words, to consider the effective $d = 5$ Lagrangian described above in eq. (1.6). The new fields — flavons — are singlets under the SM gauge group but have, for the two-family case, the non-trivial transformation properties under G_f given by

$$\Sigma_u \sim (2, \bar{2}, 1) \longrightarrow \Sigma'_u = \Omega_L \Sigma_u \Omega_{U_R}^\dagger, \quad \Sigma_d \sim (2, 1, \bar{2}) \longrightarrow \Sigma'_d = \Omega_L \Sigma_d \Omega_{D_R}^\dagger, \quad (2.6)$$

where Ω_X denotes the doublet transformation under the $\text{SU}(2)_X$ -component of the flavour group. Once these flavon fields develop vevs as in eq. (1.7) and eq. (2.4), the flavour symmetry is explicitly broken and quark masses and mixings are originated. The effective field theory obtained at the electroweak scale is exactly MFV [1] (restricted to the two-family case). Then, within this approach, the problem of the origin of flavour is replaced by the need to explain if and how this particular vev configuration can naturally arise from the minimization of the associated scalar potential.

This minimal framework can be easily extended in different ways, such as, for instance:

- Considering different scales for the Σ_u and Σ_d flavon vevs.

- Adding new representations. The most straightforward way to complete the basis in eq. (2.6), is to add a third flavon transforming as a bi-fundamental of the RH components:

$$\Sigma_R \sim (1, 2, \bar{2}) \longrightarrow \Sigma'_R = \Omega_{U_R} \Sigma_R \Omega_{D_R}^\dagger. \quad (2.7)$$

This new field does not contribute to the Yukawa terms, at least at the renormalizable level, but introduces new operators with respect to MFV, which induce flavour changing neutral currents (FCNC) mediating fully right-handed (RH) processes.⁷

- Adding new replicas of the bi-fundamental representations. This could be very helpful as a natural source of new scales and possible mixings.

The first two possibilities do not affect essentially the flavour structure of the quark Yukawa couplings, which is the focus of this work, and we will not consider them below. No further consideration is given either in this section to the third possibility, both for the sake of simplicity and because of the aesthetically unappealing aspect of being a trivial replacement of the puzzle of quark replication with that of flavon replication.

We will thus restrict the remaining of this section to the analysis of the potential for just one Σ_u and one Σ_d fields, eqs. (2.6). The general scalar potential, can then be written as a sum of two parts, the first dealing only with the SM Higgs fields and the second accounting also for the flavons interactions:

$$V \equiv V_H + V_\Sigma = -\mu^2 H^\dagger H + \lambda_H (H^\dagger H)^2 + \sum_{i=4}^{\infty} V^{(i)}[H, \Sigma_u, \Sigma_d]. \quad (2.8)$$

Inside $V^{(i)}$ all possible scalar potential terms of the effective field theory are included. In particular, $V^{(4)}$ contains all the renormalizable couplings written in terms of H and $\Sigma_{u,d}$ while $V^{(i>4)}$ incorporate all the non-renormalizable higher dimensional operators. There is no particular reason to impose that the EW and the flavour symmetry breaking should occur at the same scale. Indeed it is plausible that the flavour symmetry is broken by some new physics mechanism at a larger energy scale. Although it is true that the mixed Higgs-flavons terms could affect the value and location of the electroweak and flavour minima, the flavour composition of each term will not be modified by them. Once the flavour symmetry breaking occurs, all the terms in $V^{(i)}$ either contribute to the scalar potential as constants or can be redefined into μ^2 or λ_H . In what follows the analysis is restricted to consider only the flavon part of the scalar potential, $V^{(i)}[\Sigma_u, \Sigma_d]$.

2.1.1 The scalar potential at the renormalizable level

From the transformation properties in eq. (2.6), it is straightforward to write the most general independent invariants that enter in the scalar potential. At the renormalizable level

⁷The phenomenological impact of these operators has already been introduced and studied in the three-family case in ref. [19, 20], in a different context.

and for the case of two generations, five independent invariants can be constructed⁸ [21]:

$$\begin{aligned}
 A_u &= \text{Tr} \left(\Sigma_u \Sigma_u^\dagger \right), & B_u &= \det \left(\Sigma_u \right), \\
 A_d &= \text{Tr} \left(\Sigma_d \Sigma_d^\dagger \right), & B_d &= \det \left(\Sigma_d \right), \\
 A_{ud} &= \text{Tr} \left(\Sigma_u \Sigma_u^\dagger \Sigma_d \Sigma_d^\dagger \right).
 \end{aligned}
 \tag{2.9}$$

eqs. (1.7) and (2.4) allow to express these invariants in terms of physical observables, i.e. the four Yukawa eigenvalues and the Cabibbo angle:

$$\langle \Sigma_d \rangle = \Lambda_f \begin{pmatrix} y_d & 0 \\ 0 & y_s \end{pmatrix}, \quad \langle \Sigma_u \rangle = \Lambda_f \mathcal{V}_C^\dagger \begin{pmatrix} y_u & 0 \\ 0 & y_c \end{pmatrix},
 \tag{2.10}$$

leading to:

$$\begin{aligned}
 \langle A_u \rangle &= \Lambda_f^2 (y_u^2 + y_c^2), & \langle B_u \rangle &= \Lambda_f^2 y_u y_c, \\
 \langle A_d \rangle &= \Lambda_f^2 (y_d^2 + y_s^2), & \langle B_d \rangle &= \Lambda_f^2 y_d y_s, \\
 \langle A_{ud} \rangle &= \Lambda_f^4 [(y_c^2 - y_u^2)(y_s^2 - y_d^2) \cos 2\theta + (y_c^2 + y_u^2)(y_s^2 + y_d^2)] / 2.
 \end{aligned}
 \tag{2.11}$$

Notice that the mixing angle appears only in the vev of A_{ud} , which is the only operator that mixes the up and down flavon sectors. This is as intuitively expected: *the mixing angle describes the relative misalignment between two different directions in flavour space*. It is also interesting to notice that the expression for $\langle A_{ud} \rangle$ is related to the Jarlskog invariant for two families,

$$4J = 4 \det \left[Y_U Y_U^\dagger, Y_D Y_D^\dagger \right] = (\sin 2\theta)^2 (y_c^2 - y_u^2)^2 (y_s^2 - y_d^2)^2,$$

by the following relation:

$$\frac{1}{\Lambda_f^4} \frac{\partial \langle A_{ud} \rangle}{\partial \theta} = -2\sqrt{J}.
 \tag{2.12}$$

Using the invariants in eqs. (2.9), the most general renormalizable scalar potential allowed by the flavour symmetry reads:

$$V^{(4)} = \sum_{i=u,d} \left(-\mu_i^2 A_i - \tilde{\mu}_i^2 B_i + \lambda_i A_i^2 + \tilde{\lambda}_i B_i^2 \right) + g_{ud} A_u A_d + f_{ud} B_u B_d + \sum_{i,j=u,d} h_{ij} A_i B_j + \lambda_{ud} A_{ud},
 \tag{2.13}$$

where strict naturalness criteria would require all dimensionless couplings λ, f, g, h to be of order 1, and the dimensionful μ -terms to be smaller or equal than Λ_f although of the same order of magnitude. It is clear from the start that, with the only use of symmetry implemented here, a strict implementation of such criteria could lead at best to a strong hierarchy with some fields massless and the rest with masses of about the same scale. The ‘‘fan’’ structure of quark mass splittings observed clearly calls, instead, for a readjustment of the relative size of some μ parameters, at least when restraining to the

⁸Any other invariant operator can be expressed in terms of these five independent invariants. For example: $\text{Tr} \left(\Sigma_u \Sigma_u^\dagger \Sigma_u \Sigma_u^\dagger \right) = \text{Tr} \left(\Sigma_u \Sigma_u^\dagger \right)^2 - 2 \det \left(\Sigma_u \right)^2$.

analysis of the renormalizable and classic terms of the potential. One question is whether, in this situation, even further fine-tunings are required among the mass parameters in the potential to accommodate nature.

The relations in eq. (2.11) allow to determine the positions of the potential minima in terms of physical observables. A careful analytical and numerical study of the potential can be found in the appendices. Here we briefly comment on the most relevant physical results. Consider first the angular part of the potential. Deriving $V^{(4)}$ with respect to the angle θ , it follows that

$$\left. \frac{\partial V^{(4)}}{\partial \theta} \right|_{\min} \equiv \lambda_{ud} \frac{\partial \langle A_{ud} \rangle}{\partial \theta} \propto \lambda_{ud} \sin 2\theta (y_c^2 - y_u^2) (y_s^2 - y_d^2) \propto \lambda_{ud} \sqrt{J}. \quad (2.14)$$

The minimum of the scalar potential thus occurs when at least one of the following conditions is satisfied i) $\lambda_{ud} = 0$, ii) $\sin \theta = 0$, iii) $\cos \theta = 0$ or iv) two Yukawas in the same sector are degenerate. When condition i) is imposed, the angle remains undetermined; this assumption corresponds however to a severe fine-tuning on the model, as no symmetry protects this term from reappearing at the quantum level. Instead, due to the smallness of the Cabibbo angle, condition ii) can be interpreted as a first order solution which needs to be subsequently corrected, for example by the introduction of higher order operators. This possibility will be discussed in more detail in the next subsection. Finally, the last conditions, iii) and iv), are phenomenologically non representative of nature and large (higher order) corrections should be advocated in order to diminish the angle or to split the Yukawa degeneracy, respectively, making these solutions unattractive. All in all, the straightforward lesson that follows from eq. (2.14) is that, given the mass splittings observed in nature, *the scalar potential for bi-fundamental flavons does not allow mixing at leading order.*

From the requirement that the derivatives of the scalar potential with respect to $y_{u,d,c,s}$ also vanish at the minima, four additional independent relations on the physical parameters are obtained. As discussed above, to obtain simultaneously a sizeable mixing and a mass spectrum largely splitted in masses, instead of generically degenerate, it is necessary to (re-)introduce a large, and unnatural, hierarchy among the different operators appearing in the scalar potential (see appendix B for numerical details).

These observations can be summarized stating that, with a natural choice of the coefficients appearing in the renormalizable scalar potential $V^{(4)}$, after minimization one naturally ends up with a vanishing or undetermined mixing angle and with a naturally degenerate spectrum. In this respect we agree with a remark that can be found in refs. [13, 21]. It is, however, interesting to notice that if the invariants $B_{u,d}$ (i.e. the determinants of the flavons) are neglected, which could be justified for example introducing some *ad hoc* discrete symmetry, the minima equations would then allow, instead, solutions non-degenerate in mass for same-charge quarks, with (non-)vanishing Yukawa couplings for the first (second) quark generations. This may open the possibility to study a modified version of the scalar potential in eq. (2.13), that predicts a natural hierarchy among the Yukawas of different generations.

2.1.2 The scalar potential at the non-renormalizable level

Consider the addition of non-renormalizable operators to the scalar potential, $V^{(i>4)}$. It is very interesting to notice that this does *not* require the introduction of new invariants beyond those in eq. (2.9): all higher order traces and determinants can in fact be expressed in terms of that basis of five “renormalizable” invariants.

The lowest higher dimensional contributions to the scalar potential have dimension six (the complete list can be found in appendix A). At this order, the only terms affecting the mixing angle are

$$V^{(6)} \supset \frac{1}{\Lambda_f^2} \sum_{i=u,d} (\alpha_i A_{ud} B_i + \beta_i A_{ud} A_i) . \quad (2.15)$$

These terms, however, show the same dependence on the Cabibbo angle previously found in eq. (2.14) and, consequently, they can simply be absorbed in the redefinition of the lowest order parameter, λ_{ud} . In other words, even at the non-renormalizable level, the most favorable trend leads to no mixing. To find a non-trivial angular structure it turns out that terms in the potential of dimension eight (or higher) have to be considered, that is

$$V^{(8)} \supset \lambda_{udud} A_{ud}^2 , \quad (2.16)$$

and eq. (2.14) would be replaced by

$$\left. \frac{\partial V}{\partial \theta} \right|_{\min} \propto \sin 2\theta (y_c^2 - y_u^2) (y_s^2 - y_d^2) [\lambda_{ud} - 2 y_c^2 y_s^2 \lambda_{udud} \sin^2 \theta + \dots] , \quad (2.17)$$

implying

$$\sin^2 \theta \simeq \frac{\lambda_{ud}}{2 y_c^2 y_s^2 \lambda_{udud}} . \quad (2.18)$$

Using the experimental values of the Yukawa couplings y_s and y_c , a meaningful value for $\sin \theta$ can be obtained although at the price of assuming a highly fine-tuned hierarchy between the dimensionless coefficients of $d = 4$ and $d = 8$ terms, $\lambda_{ud}/\lambda_{udud} \sim 10^{-10}$, that cannot be naturally justified in an effective Lagrangian approach.

The remaining four equations defining the minima, obtained deriving the scalar potential with respect to $y_{u,d,c,s}$, lead to no improvement as compared to the renormalizable case: the Yukawa couplings are always given by general combinations of the coefficients of the scalar potential, underlining the complete absence of hierarchies among them. Realistic masses can be obtained at the classical level only when suitable fine-tunings are enforced.⁹

To summarize, it is possible to account for a non-vanishing mixing angle adding non-renormalizable terms to the scalar potential, although at the prize of introducing a large fine-tuning. This requirement comes in addition to the fact that the hierarchies among the Yukawa couplings can only be imposed by hand. Therefore the use of bi-fundamental scalar fields leads to an unsatisfactory answer to the problem of explaining the origin of flavour within the MFV hypothesis.

⁹See note added in proof.

For the sake of illustrating the argument with a practical exercise, we conclude this section showing, as an explicit example, a fine-tuned scalar potential which can allow hierarchical Yukawas and a non-vanishing mixing angle:

$$V = \sum_i \left(-\mu_i^2 A_i + \tilde{\lambda}_i B_i^2 + \lambda_i A_i^2 \right) + \frac{\lambda_{udud}}{\Lambda_f^4} (A_{udud} - 2A_{uudd}) - \epsilon_b \tilde{\mu}_d^2 B_d - \epsilon_u \tilde{\mu}_u^2 B_u + \epsilon_\theta \lambda_{ud} A_{ud}, \quad (2.19)$$

where $\epsilon_{u,d,\theta}$ are suppressing factors, possibly associated to some discrete symmetry, and A_{uudd} , A_{udud} , dimension eight invariants defined by the following relations:

$$A_{udud} = \text{Tr} \left(\Sigma_u \Sigma_u^\dagger \Sigma_d \Sigma_d^\dagger \Sigma_u \Sigma_u^\dagger \Sigma_d \Sigma_d^\dagger \right), \quad A_{uudd} = \text{Tr} \left(\Sigma_u \Sigma_u^\dagger \Sigma_u \Sigma_u^\dagger \Sigma_d \Sigma_d^\dagger \Sigma_d \Sigma_d^\dagger \right). \quad (2.20)$$

By minimizing the potential in eq. (2.19) one obtains the following values for the Yukawa eigenvalues and the Cabibbo angle:

$$\begin{aligned} y_u &\simeq \epsilon_u \frac{\sqrt{\lambda_u} \tilde{\mu}_u}{\sqrt{2} \tilde{\lambda}_u \mu_u} \frac{\tilde{\mu}_u}{\Lambda_f}, & y_d &\simeq \epsilon_d \frac{\sqrt{\lambda_d} \tilde{\mu}_d}{\sqrt{2} \tilde{\lambda}_d \mu_d} \frac{\tilde{\mu}_d}{\Lambda_f}, \\ y_c &\simeq \frac{\mu_u}{\sqrt{2} \Lambda_f \sqrt{\lambda_u}}, & y_s &\simeq \frac{\mu_d}{\sqrt{2} \Lambda_f \sqrt{\lambda_d}}, \\ \sin^2 \theta &\simeq \epsilon_\theta \frac{\lambda_{ud}}{\lambda_{udud} y_c^2 y_s^2}. \end{aligned} \quad (2.21)$$

Imposing for no good reason the values $\epsilon_u \sim 10^{-3}$, $\epsilon_d \sim 5 \times 10^{-2}$, $\epsilon_\theta \sim 10^{-10}$ and $\mu/(\sqrt{\lambda}\Lambda_f) \approx \tilde{\mu}/(\sqrt{\tilde{\lambda}}\Lambda_f) \sim 10^{-3}$, the correct hierarchies between the quark masses and the correct Cabibbo angle could be obtained (see details for this special case in appendix B).

The discussion about $d = 8$ terms presented above has pure illustrative purposes, as it may be a priori misleading to discuss the effects of $d = 8$ terms in the potential without simultaneously considering quantum or other higher-order sources of corrections, such as the possible impact of a Σ_R flavon¹⁰ — see eq. (2.7) — or other G_f representations.

2.2 $d = 6$ Yukawa operator: the fundamental approach

The identification of the Yukawa spurions as single flavon fields, transforming in the bi-fundamental representation of the flavour group (e.g. for a $d = 5$ Yukawa operator), is only one of the possible ways the MFV ansatz can be implemented. An attractive alternative is to consider the Yukawas as composite objects or aggregates of several fields, e.g. suggesting Yukawa operators with $d > 5$. In the simplest case, each Yukawa corresponds to two scalar fields χ transforming in the fundamental representation of G_f (e.g. $Y \sim \langle \chi \rangle \langle \chi^\dagger \rangle / \Lambda_f^2$, see eqs. (1.8) and (1.9)). This approach would *a priori* allow to introduce one new field for each component of the flavour symmetry: i.e. to reconstruct the spurions in eq. (1.4) just out of three vectors transforming as $(2, 1, 1)$, $(1, 2, 1)$ and $(1, 1, 2)$. However, such a minimal setup leads to an unsatisfactory realization of the flavour sector as no physical mixing angle is

¹⁰The impact of the fully RH bi-fundamental Σ_R is negligible: indeed it can enter in the scalar potential only as powers of $\Sigma_R \Sigma_R^\dagger$ or its hermitian conjugate, and in particular, being a singlet of $SU(2)_{Q_L}$, it cannot mix with the other flavons. As a result, its contributions can always be absorbed through a redefinition of the parameters and then the conclusions above still hold.

allowed at the renormalizable level.¹¹ The situation improves qualitatively, though, if two $(2, 1, 1)$ representations are introduced, one for the up and one for the down quark sectors. Consider then the following four fields:

$$\chi_u^L \in (2, 1, 1), \quad \chi_u^R \in (1, 2, 1), \quad \chi_d^L \in (2, 1, 1), \quad \chi_d^R \in (1, 1, 2). \quad (2.22)$$

The corresponding $d = 6$ effective Lagrangian and Yukawa couplings have been shown in eqs. (1.8) and (1.9). These flavons are vectors under the flavour symmetry. The only physical invariants that can be associated to vectors are the norm of the vectors and, eventually, their relative angles. Any matrix resulting from multiplying two vectors has only one non-vanishing eigenvalue, independently of the number of dimensions of the space. This fact alone already implies that, at the leading renormalizable order under discussion, just one “up”-type quark and one “down”-type quark are massive: a strong mass hierarchy between quarks of the same electric charge is thus automatic in this setup, which is a very promising first step in the path to explain the observed quark mass hierarchies.

More in detail, the resulting Yukawa matrices are general 2×2 matrices, containing many unphysical parameters. Without loss of generality, it is possible to express the Yukawa couplings in terms of physical quantities by choosing the flavon vevs as follows:

$$\langle \chi_i \rangle \equiv |\chi_i| \mathcal{V}_i \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad (2.23)$$

where by $|\chi_i|$ we denote the norm of the vev of χ , $|\chi_i| \equiv |\langle \chi_i \rangle|$, and \mathcal{V}_i are 2×2 unitary matrices. Redefining the quark fields as follows,

$$Q'_L = \mathcal{V}_L^{(d)\dagger} Q_L, \quad U'_R = \mathcal{V}_R^{(u)\dagger} U_R, \quad D'_R = \mathcal{V}_R^{(d)\dagger} D_R, \quad (2.24)$$

it results

$$\mathcal{L}_Y = \bar{Q}'_L Y_D D'_R H + \bar{Q}'_L Y_U U'_R \tilde{H} + \text{h.c.}, \quad (2.25)$$

with the corresponding Yukawa matrices given by¹²

$$Y_D = \frac{|\chi_d^L| |\chi_d^R|}{\Lambda_f^2} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \quad Y_U = \frac{|\chi_u^L| |\chi_u^R|}{\Lambda_f^2} \mathcal{V}_L^{(d)\dagger} \mathcal{V}_L^{(u)} \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}. \quad (2.26)$$

This illustrates explicitly that: i) *there is a natural hierarchy among the mass of the first and second generations, without imposing any constraint on the parameters of the scalar potential*; ii) *the product $\mathcal{V}_L^{(d)\dagger} \mathcal{V}_L^{(u)}$ is a non-trivial unitary matrix that contains all the information about the mixing angle* (the phase can be easily removed in the two-family case under discussion). There is now a clear geometrical interpretation of the Cabibbo angle: *the mixing angle between two generations of quarks is the misalignment of the χ^L*

¹¹ Because then the flavons associated to the up and down left-handed character are not misaligned in flavour space, but correspond instead to just one $(2, 1, 1)$ flavon.

¹²The cutoff scale Λ_f refers to the scale of the flavour dynamics. In principle we could have different scales for the left and right flavons as well as for the up and down ones, but for simplicity we assume that all the scales are close and Λ_f refers to the average value.

flavons in the flavour space, with the mixing matrix appearing in weak currents, eq. (2.5), given by

$$\mathcal{V}_C = \mathcal{V}_L^{(u)\dagger} \mathcal{V}_L^{(d)}. \quad (2.27)$$

Let us compare the phenomenology expected from bi-fundamental flavons (i.e. $d = 5$ Yukawa operator) with that from fundamental flavons (i.e. $d = 6$ Yukawa operators). For bi-fundamentals, the list of effective FCNC operators is exactly the same that in the original MFV proposal [1]. The case of fundamentals presents some differences: higher-dimension invariants can be constructed in this case, exhibiting lower dimension than in the bi-fundamental case. For instance, one can compare these two operators:

$$\bar{D}_R \Sigma_d^\dagger \Sigma_u \Sigma_u^\dagger Q_L \sim [\text{mass}]^6 \quad \longleftrightarrow \quad \bar{D}_R \chi_d^R \chi_u^{L\dagger} Q_L \sim [\text{mass}]^5, \quad (2.28)$$

where the mass dimension of the invariant is shown in brackets; with these two types of basic bilinear FCNC structures it is possible to build effective operators describing FCNC processes, but differing on the degree of suppression that they exhibit. This underlines the fact that the identification of Yukawa couplings with aggregates of two or more flavons is a setup which goes technically beyond the realization of MFV, resulting possibly in a distinct phenomenology which could provide a way to distinguish between fundamental and bi-fundamental origin

2.2.1 The scalar potential

The general scalar potential that can be written including flavons in the fundamental is analogous to that in eq. (2.8), replacing Σ_i with χ_i ,

$$V \equiv V_H + V_\chi. \quad (2.29)$$

Previous considerations regarding the scale separation between EW and flavour breaking scale hold also in this case, and in consequence the Higgs sector contributions will not be explicitly described.

Any flavour invariant operator can be constructed out of the following five independent building blocks:

$$\chi_u^{L\dagger} \chi_u^L, \quad \chi_u^{R\dagger} \chi_u^R, \quad \chi_d^{L\dagger} \chi_d^L, \quad \chi_d^{R\dagger} \chi_d^R, \quad \chi_u^{L\dagger} \chi_d^L. \quad (2.30)$$

From the expressions for the Yukawa matrices in eqs. (2.26), it follows that in this scenario the scalar potential depends only on three of the five physical parameters: one angle and the two (larger) Yukawa couplings

$$|\chi_u^L| |\chi_u^R| = \Lambda_f^2 y_c, \quad |\chi_d^L| |\chi_d^R| = \Lambda_f^2 y_s, \quad \chi_u^{L\dagger} \chi_d^L = \cos \theta_c |\chi_u^L| |\chi_d^L|, \quad (2.31)$$

given by the product of the left and right up (down) flavon moduli. As expected, the mixing angle is simply the angle defined in flavour space by the up and down left vectors. From the point of view of the measurable quantities, there is a certain parametrization freedom, and a possible convenient choice is given by¹³

$$\frac{|\chi_u^R|}{\Lambda_f} = 1 = \frac{|\chi_d^R|}{\Lambda_f}. \quad (2.32)$$

¹³See appendix C for a detailed discussion.

As a result, the invariants physically relevant for the flavour structure are:

$$|\chi_u^L| = \Lambda_f y_c, \quad |\chi_d^L| = \Lambda_f y_s, \quad \chi_u^{L\dagger} \chi_d^L = \Lambda_f^2 y_c y_s \cos \theta. \quad (2.33)$$

At the renormalizable level, the scalar potential is given by

$$V^{(4)} = - \sum_{i=u,d} \mu_i^2 \chi_i^{L\dagger} \chi_i^L - \sum_{i=u,d} \tilde{\mu}_i^2 \chi_i^{R\dagger} \chi_i^R - \mu_{ud}^2 \chi_u^{L\dagger} \chi_d^L + \dots, \quad (2.34)$$

where dots stand for all possible quartic couplings. The total number of operators that can be introduced at the renormalizable level is 20. However, as shown in appendix C, many of them (i.e. quartic couplings that mix different flavours) do not have any real impact on the existence and determination of the minima. Studying the latter, the following relations between the (large) up and down Yukawa eigenvalues and the Cabibbo angle follow:

$$\frac{y_s^2}{y_c^2} = \frac{\mu_d^2 \lambda_u}{\mu_u^2 \lambda_d}, \quad \cos \theta = \frac{\sqrt{\lambda_u \lambda_d} \mu_{ud}^2}{\lambda_{ud} \mu_u \mu_d} \quad (2.35)$$

which shows that without strong fine-tunings this scenario can explain the hierarchy between the first and second family, and account for a sizable Cabibbo angle.

2.2.2 The first generation

In this two-generation analysis, the first family has remained massless at the renormalizable level. A first possibility is that non-renormalizable corrections may induce this small masses. Non-renormalizable interactions manifest themselves in form of higher order contributions to the Yukawa operators and the flavon vevs and/or as non-renormalizable terms in the potential, which can modify its minima.

From eq. (1.8) and the flavon transformation properties, it follows that higher order contributions to the Yukawa operators can only be constructed by further insertions of $\chi^\dagger \chi$ inside the renormalizable operators. However, such kind of insertions do not modify the flavour structure of the Yukawa matrices, but simply redefine the two heavier couplings, y_c and y_s . On the other hand, the introduction of higher order operators in the scalar potential has the effect of modifying the vevs of the flavons, replacing the relation in eq. (2.23) with

$$\frac{\langle \chi_{u,d}^{L,R} \rangle}{\Lambda_f} \equiv \left| (1 + \mathcal{O}(\epsilon)) \chi_{u,d}^{L,R} \left(\mathcal{V}_{L,R}^{(u,d)} (1 + \mathcal{O}(\epsilon)) \right) \begin{pmatrix} \mathcal{O}(\epsilon) \\ 1 \end{pmatrix} \right|, \quad (2.36)$$

where $\epsilon \ll 1$ parametrizes the ratio among higher and leading order contributions. The only effect of these modifications is to redefine the mixing angle θ and the second family Yukawas, y_c and y_s , without changing the rank of the Yukawa matrices and leaving thus the first generation massless. In summary, non-renormalizable interactions cannot switch on additional (first family) Yukawas if they were absent at the renormalizable level.

An alternative can be built on the fact that each up-down set of fundamental flavons provides a supplementary scale, in addition to new sources of mixing from their misalignment. A possibility along this direction is to enlarge the number of flavons to six, made out of a set of three ($\chi_{u,d}^R$ plus just one χ^L) replicated: in total two $\chi^L \sim (2, 1, 1)$,

two $\chi_u^R \sim (1, 2, 1)$ and two $\chi_d^R \sim (1, 1, 2)$. In this case the Yukawa terms change in a non-trivial way:

$$Y_D \equiv \frac{\sum_{ij} \alpha_{ij}^d \langle \chi_i^L \rangle \langle \chi_j^{R\dagger} \rangle}{\Lambda_f^2}, \quad Y_U \equiv \frac{\sum_{ij} \alpha_{ij}^u \langle \chi_i^L \rangle \langle \chi_j^{R\dagger} \rangle}{\Lambda_f^2}, \quad (2.37)$$

with $\alpha_{i,j}$ numerical coefficients and i, j running over all flavons. An explicit computation reveals that, for generic values of $\alpha_{ij} (\neq 0)$, the rank of the Yukawa matrices is indeed two. However, in this case, the natural hierarchy between the first and second family is lost, being all the Yukawas of the same order unless the vevs of the new flavons are unnaturally smaller than those of the first replica. In conclusion, adding new RH flavon copies does not lead either to an appealing and natural source of masses for the first generation.

3 The three-family case

Let us extend the previous analysis to the three-family case. While most of the procedure, with both bi-fundamental and fundamental representations, follows straightforwardly, two main differences should be underlined. First of all, the top Yukawa coupling, y_t , is now a parameter which is of $\mathcal{O}(1)$. The fact that in the two-family case the largest Yukawa, y_c was much smaller than one, allowed us to safely retain only the lowest order terms in the (Yukawa) perturbative expansion. In the three-family scenario, in principle, one should include all orders in the expansion. However, in this case, the Cayley-Hamilton identity [22, 23] provides a way out, as it proves that a general 3×3 matrix X must satisfy the relation:

$$X^3 - \text{Tr}[X] X^2 + \frac{1}{2} X (\text{Tr}[X]^2 - \text{Tr}[X^2]) - \det[X] = 0, \quad (3.1)$$

which allows to express all powers X^n (with $n > 2$) in terms solely of $\mathbf{1}$, X and X^2 , with coefficients involving the traces of X and X^2 and the determinant of X . In the case under study, X corresponds to the invariant products $\Sigma^\dagger \Sigma$ or $\chi^\dagger \chi$, depending on whether bi-fundamental or fundamental representations are considered.

The second main difference with respect to the two-family case, is the appearance of a physical phase in the quark mixing matrix. For the sake of simplicity, in this paper we disregard CP-violation, deferring its discussion to a future work [24].

3.1 $d = 5$ Yukawa operator: the bi-fundamental approach

In this section we extend the approach discussed in section 2.1 to the three-family case. Consider two bi-triplets under the flavour symmetry G_f , see eq. (1.4),

$$\Sigma_u \sim (3, \bar{3}, 1) \longrightarrow \Sigma'_u = \Omega_L \Sigma_u \Omega_{U_R}^\dagger, \quad \Sigma_d \sim (3, 1, \bar{3}) \longrightarrow \Sigma'_d = \Omega_L \Sigma_d \Omega_{D_R}^\dagger, \quad (3.2)$$

where now the Ω_X matrices refer to the triplet transformations under the $SU(3)_X$ component of the flavour group. The Yukawa Lagrangian is the same as that in eq. (1.6). Once the flavons develop a vev as in eq. (1.7), the flavour symmetry is broken and one recovers the observed fermion masses and CKM matrix given in eq. (1.5). Recall that the present realization is the simplest realization of the original MFV approach [1]. Again, it would be possible

to extend it introducing a third RH flavon field, $\Sigma_R \sim (1, 3, \bar{3}) \longrightarrow \Sigma'_R = \Omega_{U_R} \Sigma_R \Omega_{D_R}^\dagger$. We do not further consider it when constructing the scalar potential, as it cannot contribute to the Yukawa spurions neither at $\mathcal{O}(1/\Lambda_f)$ nor $\mathcal{O}(1/\Lambda_f^2)$, that is, neither via $d = 5$ nor $d = 6$ Yukawa operators.

Restricting the explicit analysis to the part of the renormalizable scalar potential not containing the SM Higgs field, a complete and independent basis is given by the following seven invariant operators:

$$\begin{aligned}
 A_u &= \text{Tr} \left(\Sigma_u \Sigma_u^\dagger \right), & \langle A_u \rangle &= \Lambda_f^2 (y_t^2 + y_c^2 + y_u^2), \\
 B_u &= \det (\Sigma_u), & \langle B_u \rangle &= \Lambda_f^3 y_u y_c y_t, \\
 A_d &= \text{Tr} \left(\Sigma_d \Sigma_d^\dagger \right), & \langle A_d \rangle &= \Lambda_f^2 (y_b^2 + y_s^2 + y_d^2), \\
 B_d &= \det (\Sigma_d), & \langle B_d \rangle &= \Lambda_f^3 y_d y_s y_b, \\
 A_{uu} &= \text{Tr} \left(\Sigma_u \Sigma_u^\dagger \Sigma_u \Sigma_u^\dagger \right), & \langle A_{uu} \rangle &= \Lambda_f^4 (y_t^4 + y_c^4 + y_u^4), \\
 A_{dd} &= \text{Tr} \left(\Sigma_d \Sigma_d^\dagger \Sigma_d \Sigma_d^\dagger \right), & \langle A_{dd} \rangle &= \Lambda_f^4 (y_b^4 + y_s^4 + y_d^4), \\
 A_{ud} &= \text{Tr} \left(\Sigma_u \Sigma_u^\dagger \Sigma_d \Sigma_d^\dagger \right), & \langle A_{ud} \rangle &= \Lambda_f^4 (P_0 + P_{\text{int}}),
 \end{aligned} \tag{3.3}$$

where P_0 and P_{int} encode the angular dependence,

$$P_0 \equiv - \sum_{i < j} \left(y_{u_i}^2 - y_{u_j}^2 \right) \left(y_{d_i}^2 - y_{d_j}^2 \right) \sin^2 \theta_{ij}, \tag{3.4}$$

$$\begin{aligned}
 P_{\text{int}} &\equiv \sum_{i < j, k} \left(y_{d_i}^2 - y_{d_k}^2 \right) \left(y_{u_j}^2 - y_{u_k}^2 \right) \sin^2 \theta_{ik} \sin^2 \theta_{jk} + \\
 &\quad - \left(y_d^2 - y_s^2 \right) \left(y_c^2 - y_t^2 \right) \sin^2 \theta_{12} \sin^2 \theta_{13} \sin^2 \theta_{23} + \\
 &\quad + \frac{1}{2} \left(y_d^2 - y_s^2 \right) \left(y_c^2 - y_t^2 \right) \cos \delta \sin 2\theta_{12} \sin 2\theta_{23} \sin \theta_{13},
 \end{aligned} \tag{3.5}$$

with $i, j, k = 1, 2, 3$. P_0 generalizes the expression found in the two-family case — see eq. (2.11) — containing all the terms with a single angular dependence. The second piece, instead, P_{int} , contains all contributions that involve more than one mixing angle. Notice that in this case the Jarlskog invariant appears only at the non-renormalizable level.

The most general scalar potential at the renormalizable level is now given by

$$V^{(4)} = \sum_{i=u,d} \left(-\mu_i^2 A_i + \tilde{\mu}_i B_i + \lambda_i A_i^2 + \lambda'_i A_{ii} \right) + g_{ud} A_u A_d + \lambda_{ud} A_{ud}. \tag{3.6}$$

Notice that the invariants $B_{u,d}$ have mass dimension three (instead of two for the two-generation case), so that no $B_{u,d}^2$ term can be introduced at this level.

The solutions that minimize this scalar potential have a pattern very similar to that in the two-family case: i) no mixing is favored,¹⁴ ii) in most of the parameter space. Now

¹⁴However, due to the peculiar structure of the last term in eq. (3.6), minima with non-vanishing angles are now allowed, although leading to solutions which are both fine-tuned and overall physically incorrect.

however, there is a region in parameter space for which a hierachical solution is allowed for non strictly zero, but constrained, $\tilde{\mu}$. This solution has one non-vanishng Yukawa eigenvalue per up and down sectors, but to recover the hierarchy among top and bottom masses it is necessary to further demand $g_{ud} < y_b^2/y_t^2$ which, in the absence of ad hoc symmetries, results in a similar degree of fine-tunnig to that for the two-family case. Furthermore, alike to the case of an initial vanishing $\sin \theta$ at the renormalizable level for two families, it cannot be corrected by non-renormalizable terms in the potential.

As in section 2.1 for two generation, we studied the contributions of non-renormalizable operators in the scalar potential, with similar conclusion: the introduction of higher order terms does not lead to a more natural description of the physical parameters. Nevertheless, some improvement can be obtained when discussing the scenario with a fine-tuned choice of parameters g_{ud} , $\tilde{\mu}_i$. In this case, in fact, lighter Yukawas can be introduced through higher order operators, even if no natural hierarchy between the first two families can be obtained.

In summary, for three generations, to consider bi-fundamental scalars (as in the case of $d = 5$ Yukawa operator) alone as the possible dynamical origin of Yukawa couplings does not lead naturally to a satisfactory pattern of masses and mixings.¹⁵

3.2 $d = 6$ Yukawa operator: the fundamental approach

We deal now with the case of flavons transforming in the fundamental of the flavour group G_f . For most of the conventions we refer to the two-family treatment done in section 2.2. To account for non-trivial mixing, it is necessary to introduce at least four flavons, corresponding to up and down, left and right flavons:

$$\chi_u^L \in (3, 1, 1) , \quad \chi_u^R \in (1, 3, 1) , \quad \chi_d^L \in (3, 1, 1) , \quad \chi_d^R \in (1, 1, 3) . \quad (3.7)$$

When they develop vevs, the flavour symmetry is spontaneously broken and the Yukawa matrices are given as in eq. (1.9). Without loss of generality, it is possible to write:

$$\langle \chi_{u,d}^{L,R} \rangle \equiv |\chi_{u,d}^{L,R}| \mathcal{V}_{L,R}^{(u,d)} \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} , \quad (3.8)$$

where $\mathcal{V}_{L,R}^{(u,d)}$ are 3×3 unitary matrices. Similarly to what was shown in section 2.2, removing the unphysical parameters, the following expressions for the Yukawa matrices are obtained:

$$Y_D = \frac{|\chi_d^L| |\chi_d^R|}{\Lambda_f^2} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} , \quad Y_U = \frac{|\chi_u^L| |\chi_u^R|}{\Lambda_f^2} \mathcal{V}_L^{(d)\dagger} \mathcal{V}_L^{(u)} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} . \quad (3.9)$$

This illustrates that, independently of the parametrization chosen, Y_D and Y_U can have only one non-vanishing eigenvalue, as they result from multiplying two vectors. For obvious

¹⁵See note added in proof.

reasons, in eq. (3.9) the massive state is chosen to be that of the third generation. The flavon vevs have not broken completely the flavour symmetry, leaving a residual $SU(2)_{Q_L} \times SU(2)_{D_R} \times SU(2)_{U_R}$ symmetry group. As a consequence any rotation in the 12 sector is unphysical and the only physical angle, given by the misalignment between $\langle \chi_u^L \rangle$ and $\langle \chi_d^L \rangle$ in the flavour space, can be identified with the 23 CKM mixing angle:

$$\mathcal{V}_L^{(d)\dagger} \mathcal{V}_L^{(u)} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \theta_{23} & \sin \theta_{23} \\ 0 & -\sin \theta_{23} & \cos \theta_{23} \end{pmatrix}. \quad (3.10)$$

The analysis of the scalar potential follows exactly that in section 2.2 for two families (see for example eq. (2.35)), with the obvious replacement of y_c, y_s for y_t, y_b and with the physical mixing angle corresponding now to θ_{23} . Both the largest hierarchy and a $\cos \theta_{23}$ naturally of $\mathcal{O}(1)$ are beautifully explained without any fine-tuning. However, as in the two-family case, it is not possible to generate lighter fermion masses either introducing non-renormalizable interactions or adding extra RH flavons.

Nevertheless, the partial breaking of flavour symmetry provided by eq. (3.9) can open quite interesting possibilities from a model-building point of view. Consider as an example the following multi-step approach. In a first step, only the minimal number of fundamental fields are introduced: i.e. χ^L, χ_u^R and χ_d^R . Their vevs break $G_f = SU(3)^3$ down to $SU(2)^3$, originating non-vanishing Yukawa couplings only for the top and the bottom quarks, without any mixing angle (as we have only one left-handed flavon). As a second step, four new G_f -triplet fields $\chi_{u,d}^{\prime L,R}$ are added, whose contributions to the Yukawa terms are suppressed relatively to the previous flavons (i.e. $\langle \chi' \rangle \ll \langle \chi \rangle$). If their vevs point in the direction of the unbroken flavour subgroup $SU(2)^3$, then the residual symmetry is further reduced. As a result, non-vanishing charm and strange Yukawa couplings are generated together with a mixing among the first two generations:

$$Y_u \equiv \frac{\langle \chi^L \rangle \langle \chi_u^{R\dagger} \rangle}{\Lambda_f^2} + \frac{\langle \chi_u^{\prime L} \rangle \langle \chi_u^{\prime R\dagger} \rangle}{\Lambda_f^2} = \begin{pmatrix} 0 & \sin \theta y_c & 0 \\ 0 & \cos \theta y_c & 0 \\ 0 & 0 & y_t \end{pmatrix}, \quad (3.11)$$

$$Y_d \equiv \frac{\langle \chi^L \rangle \langle \chi_d^{R\dagger} \rangle}{\Lambda_f^2} + \frac{\langle \chi_d^{\prime L} \rangle \langle \chi_d^{\prime R\dagger} \rangle}{\Lambda_f^2} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & y_s & 0 \\ 0 & 0 & y_b \end{pmatrix}.$$

The relative suppression of the two sets of flavon vevs correspond to the hierarchy between y_c and y_t (y_s and y_b).¹⁶ Hopefully, a refinement of this argument would allow to explain the rest of the Yukawas and the remaining angles. The construction of the scalar potential for such a setup would be quite model dependent though, and beyond the scope of this paper.

3.3 Combining fundamentals and bi-fundamentals

Until now we have considered separately Yukawa operators of dimension $d = 5$ and $d = 6$. It is, however, interesting to explore if some added value from the simultaneous presence

¹⁶Alternatively, all flavon vevs of similar magnitude with different flavour scale would lead to the same pattern.

of both kinds of operators can be obtained. This is a sensible choice from the point of view of effective Lagrangians in which, working at $\mathcal{O}(1/\Lambda_f^2)$, contributions of four types may be included: i) the leading $d = 5$ $\mathcal{O}(1/\Lambda_f)$ operators; ii) renormalizable terms stemming from fundamentals (i.e. from $d = 6$ $\mathcal{O}(1/\Lambda_f^2)$ operators); iii) $\mathcal{O}(1/\Lambda_f^2)$ of the form $\Sigma_{u,d}\Sigma_R$ if Σ_R turns out to be present in the spectrum; iv) other corrections numerically competitive at the orders considered here. We focus here as illustration on the impact of i) and ii):

$$\mathcal{L}_Y = \bar{Q}_L \left[\frac{\Sigma_d}{\Lambda_f} + \frac{\chi_d^L \chi_d^{R\dagger}}{\Lambda_f^2} \right] D_R H + \bar{Q}_L \left[\frac{\Sigma_u}{\Lambda_f} + \frac{\chi_u^L \chi_u^{R\dagger}}{\Lambda_f^2} \right] U_R \tilde{H} + \text{h.c.}, \quad (3.12)$$

As the bi-fundamental flavons arise at first order in the $1/\Lambda_f$ expansion, it is suggestive to think of the fundamental contributions as a “higher order” correction. Let us then consider the case in which the flavons develop vevs as follows:

$$\frac{\langle \Sigma_{u,d} \rangle}{\Lambda_f} \sim \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & y_{t,b} \end{pmatrix}, \quad \frac{\langle \chi_{u,d}^L \rangle}{\Lambda_f^2} \sim \begin{pmatrix} 0 \\ y_{c,s} \\ 0 \end{pmatrix}, \quad (3.13)$$

and $\chi_{u,d}^R$ acquire arbitrary vev values, although $\mathcal{O}(1)$, for all components. Nevertheless, it is important to recall that the bi-fundamentals Σ point in most cases to degenerate Yukawa eigenvalues instead of the pattern in the left-hand side of eq. (3.13), and either restrictive conditions on the parameters, or an extra symmetry, have to be imposed to obtain it, see sects. 2.1.1 and 3.2. Finally,

$$Y_u = \begin{pmatrix} 0 & \sin \theta_c y_c & 0 \\ 0 & \cos \theta_c y_c & 0 \\ 0 & 0 & y_t \end{pmatrix}, \quad Y_d = \begin{pmatrix} 0 & 0 & 0 \\ 0 & y_s & 0 \\ 0 & 0 & y_b \end{pmatrix}. \quad (3.14)$$

This seems an appealing pattern, with masses for the two heavier generations and one sizable mixing angle, that we chose to identify here with the Cabibbo angle.¹⁷ As for the lighter family, non-vanishing masses for the up and down quarks could now result from non-renormalizable operators.

The drawback of these combined analysis is that the direct connection between the minima of the potential and the spectrum is lost and the analysis of the potential would be very involved.

4 Conclusions

The ansatz of MFV implicitly assumes a dynamical origin for the SM Yukawa couplings. In this paper we explored such a possibility. The simplest dynamical realization of MFV is to identify the Yukawa couplings with the vevs of some dynamical fields, the flavons. For instance, the Yukawa interactions themselves could result, after spontaneous symmetry breaking, from effective operators of dimension $d > 4$ invariant under the flavour symmetry, which involve one or more flavons together with the usual SM fields.

¹⁷Similar constructions have been suggested also in other contexts as in [25–27].

Only a scalar field (or an aggregate of fields in a scalar configuration) can get a vev, which should correspond to the minimum of a potential. What may be the scalar potential of the MFV flavons? May some of its minima naturally correspond to the SM spectra of masses and mixing angles? These are the questions addressed in this work.

First of all, we showed here that the underlying flavour symmetry — under which the terms in the potential have to be invariant — is a very restrictive constraint: at the renormalizable level only a few terms are allowed in the potential, and even at the non-renormalizable level quite constrained patterns have to be respected.

The simplest realization is obtained by a one-to-one correspondence of each Yukawa coupling with a single scalar field transforming in the bi-fundamental of the flavour group. In the language of effective Lagrangians, this may correspond to the lowest order terms in the flavour expansion: $d = 5$ effective Yukawa operators made out of one flavon field plus the usual SM fields. We have constructed the general scalar potential for bi-fundamental flavons, both for the case of two and three families. At the renormalizable level, at the minimum of the potential only vanishing or undetermined mixing angles are allowed. The introduction of either additional ad hoc symmetries or the restriction to a contrived region of the parameter domain could allow to obtain solutions with vanishing Yukawa couplings for all quarks but those in the heaviest family. Still, mixing would be absent. The addition of non-renormalizable terms to the potential would allow masses for the lighter families, although without providing naturally a correct pattern of masses and mixings. In resume, the sole consideration of flavons in the bi-fundamental representation of the flavour group does not naturally lead to a satisfactory dynamical description of the SM quark flavour sector, at least at the classical level.

Another avenue explored in this work associates two vector flavons to each Yukawa spurion, i.e. a Yukawa $Y \sim \langle \chi^L \rangle \langle \chi^{R\dagger} \rangle / \Lambda_f^2$. This is a very attractive scenario in that while Yukawas are composite objects, the new fields are in the fundamental representation of the flavour group, in analogy with the case of quarks. Those flavons could be scalars or fermions: we focused exclusively on scalars. From the point of view of effective Lagrangians, this case could correspond to the next-to leading order term in the expansion: $d = 6$ Yukawa operators. We have constructed the general scalar potential for scalar flavons in the fundamental representation, both for the case of two and three families of quarks. By construction, this scenario results unavoidably in a strong hierarchy of masses: at the renormalizable level only one quark gets mass in each quark sector: they could be associated with the top and bottom quark for instance. Non-trivial mixing requires as expected a misalignment between the flavons associated to the up and down left-handed quarks. In consequence, the minimal field content corresponds to four fields χ_u^L , χ_d^L , χ_u^R and χ_d^R , and the physics of mixing lies in the interplay of the first two. In resume, for fundamental flavons it follows in a completely natural way: i) a strong mass hierarchy between quarks of the same charge, pointing to a distinctly heavier quark in each sector; ii) one non-vanishing mixing angle, which can be identified with the Cabibbo angle in the case of two generations, and for instance with the rotation in the 23 sector of the CKM matrix in the case of three generations.

Nevertheless, to achieve non-vanishing Yukawa couplings for the lighter quarks and the full mixing pattern requires, at least at the classical level explored here, more complicated scenarios and variable degrees of fine-tuning. Interesting possibilities which we started to explore here include replicas of fundamental flavons, in several varieties. An intriguing one consists in considering the minimal set of only three fields, χ^L , χ_u^R and χ_d^R , plus their replicas: it allows a double step symmetry breaking mechanism, which may produce the hierarchical quark spectrum and the shell-like pattern of the CKM matrix.

Finally, we briefly explored the possibility of introducing simultaneously bi-fundamentals and fundamentals flavons. It is a very sensible possibility from the point of view of effective Lagrangians to consider both $d = 5$ and $d = 6$ Yukawa operators when working to $\mathcal{O}(1/\Lambda_f^2)$. It suggests that $d = 5$ operators, which bring in the bi-fundamentals, could give the dominant contributions, while the $d = 6$ operator — which brings in the fundamentals — should provide a correction inducing the masses of the two lighter families and the Cabibbo angle. It requires, though, to appeal to a discrete symmetry or to restrict the parameters of the potential to a contrived region to avoid quark mass degeneracies induced by the bi-fundamental flavons.

Overall, it is remarkable that the requirement of invariance under the flavour symmetry strongly constraints the scalar potential of MFV, up to the point that the obtention of quark mass hierarchies and mixing angles is far from trivial. Furthermore, besides exploring the — disappointing — impact in mixing of bi-fundamental flavons, this work has shown that flavons in the fundamental are instead a tantalizing avenue to induce hierarchies and non-trivial fermion mixing. A long path remains ahead, though, to naturally account for the complete observed fermion spectrum and mixings.

Acknowledgments

We are specially indebted to Enrico Nardi for fruitful discussions and suggestions. We also thank Alvaro de Rujula and Pilar Hernández for illuminating discussions. L. Merlo and S. Rigolin thank the Departamento de Física Teórica of the Universidad Autónoma de Madrid for hospitality during the development of this project. R. Alonso and M.B. Gavela acknowledge CICYT through the project FPA2009-09017 and by CAM through the project HEPHACOS, P-ESP-00346. R. Alonso acknowledges financial support from the MICINN grant BES-2010-037869. L. Merlo acknowledges the German ‘Bundesministerium für Bildung und Forschung’ under contract 05H09W0E. S. Rigolin acknowledges the partial support of an Excellence Grant of Fondazione Cariparo and of the European Program, Unification in the LHC era, under the contract PITN-GA-2009-237920 (UNILHC).

A $d = 6$ operators in the bifundamental approach

We give here a summary of the invariant operators that appear in the potential at dimension 6 for the Bi-fundamental approach. There are two types of operators: products of the invariants in eq. (2.9) and new operators written in terms of traces of flavons. The list of

the former invariants is:

$$\begin{aligned}
 & \text{Tr} \left(\Sigma_i \Sigma_i^\dagger \right) \text{Tr} \left(\Sigma_j \Sigma_j^\dagger \right) \text{Tr} \left(\Sigma_k \Sigma_k^\dagger \right), & \det \left(\Sigma_i \right) \text{Tr} \left(\Sigma_j \Sigma_j^\dagger \right) \text{Tr} \left(\Sigma_k \Sigma_k^\dagger \right), \\
 & \det \left(\Sigma_i \right) \det \left(\Sigma_j \right) \text{Tr} \left(\Sigma_k \Sigma_k^\dagger \right), & \det \left(\Sigma_i \right) \det \left(\Sigma_j \right) \det \left(\Sigma_k \right), \\
 & \text{Tr} \left(\Sigma_i \Sigma_i^\dagger \right) \text{Tr} \left(\Sigma_u \Sigma_u^\dagger \Sigma_d \Sigma_d^\dagger \right), & \det \left(\Sigma_i \right) \text{Tr} \left(\Sigma_u \Sigma_u^\dagger \Sigma_d \Sigma_d^\dagger \right).
 \end{aligned} \tag{A.1}$$

where i, j and k run over u,d. The new invariant operators that appear are of the form

$$\text{Tr} \left(\Sigma_i \Sigma_i^\dagger \Sigma_j \Sigma_j^\dagger \Sigma_k \Sigma_k^\dagger \right). \tag{A.2}$$

In the two family case the vevs of these operators are not independent and can be expressed as linear combinations of the lowest order (LO) ones. To understand it, notice that five parameters (four Yukawas and an angle) suffice to parametrize the vevs. Then the relations of these parameters with the first five LO invariants can be formally inverted and substituted in any higher dimension new invariant, to express them as functions of the five former invariants. As an example,

$$\begin{aligned}
 \text{Tr} \left(\langle \Sigma_u \rangle \langle \Sigma_u^\dagger \rangle \langle \Sigma_u \rangle \langle \Sigma_u^\dagger \rangle \langle \Sigma_d \rangle \langle \Sigma_d^\dagger \rangle \right) = & \text{Tr} \left(\langle \Sigma_u \rangle \langle \Sigma_u^\dagger \rangle \right) \text{Tr} \left(\langle \Sigma_u \rangle \langle \Sigma_u^\dagger \rangle \langle \Sigma_d \rangle \langle \Sigma_d^\dagger \rangle \right) + \\
 & - \det \left(\langle \Sigma_u \rangle \right)^2 \text{Tr} \left(\langle \Sigma_d \rangle \langle \Sigma_d^\dagger \rangle \right).
 \end{aligned}$$

B A fine-tuned scalar potential in the bifundamental approach

This appendix gives details on a particular scalar potential whose minimum sets the observed values of masses and mixings for the first two generations. Its purpose is to illustrate the theoretical prize to be paid in order to obtain a realistic solution, and to discuss its degree of naturalness. This ansatz for the potential is given in eq. (2.19). Rewriting it as a sum of four double well potential terms involving the Yukawa eigenvalues and terms involving the mixing angle, it reads:

$$V_\Sigma = \sum_{i=u,d} \left[\lambda_i \left(A_i - \frac{\mu_i^2}{2\lambda_i} \right)^2 + \tilde{\lambda}_i \left(B_i - \epsilon_i \frac{\tilde{\mu}_i^2}{2\tilde{\lambda}_i} \right)^2 \right] + \frac{\lambda_{udud}}{\Lambda_f^4} (A_{udud} - 2A_{uudd}) + \epsilon_\theta \lambda_{ud} A_{ud}. \tag{B.1}$$

Here ϵ_u, ϵ_d and ϵ_θ parametrize the suppression of the respective operators and will be defined when discussing the results. The invariants in the previous expression have already been defined in eqs. (2.9), (2.20), and the renormalizable operators have been expressed in terms of masses and mixings in eq. (2.11). At the minimum of the potential, the non-renormalizable term corresponds to:

$$\begin{aligned}
 \langle A_{udud} - 2A_{uudd} \rangle = & \Lambda_f^8 \left[(y_c^2 - y_u^2)^2 (y_s^2 - y_d^2)^2 \sin^4 \theta + \right. \\
 & \left. + 2 (y_c^2 y_d^2 + y_s^2 y_u^2) (y_c^2 - y_u^2) (y_s^2 - y_d^2) \sin^2 \theta - y_c^4 y_s^4 - y_d^4 y_u^4 \right].
 \end{aligned} \tag{B.2}$$

The first part of the potential being positive definite, it is minimized when vanishing, which implies $\langle A_u \rangle = \Lambda_f^2 (y_c^2 + y_u^2) = \mu_u^2 / 2\lambda_u$, $\langle B_u \rangle = \Lambda_f y_c y_u = \epsilon_u \tilde{\mu}_u^2 / 2\tilde{\lambda}_u$ and similar expressions for the down sector. These equations define a circle and an hyperbola in the (y_c, y_u) plane. Their intersection defines the minimum as depicted in figure 2.




Figure 2. Graphic determination of the minimum.

B.1 Minimization of the scalar potential

The explicit equations for the minimum of the scalar potential considered above are shown in what follows. In a first approximation we neglect all the terms suppressed by $\epsilon_{u,d,\theta}$.

- The equation associated with y_u is then given by:

$$\frac{\partial V_\Sigma}{\partial y_u} = 2y_u \Lambda_f^4 \left[-\frac{\mu_u^2}{\Lambda_f^2} + 2\lambda_u (y_u^2 + y_c^2) + \tilde{\lambda}_u y_c^2 - 2\lambda_{udud} F_u \right] = 0. \quad (\text{B.3})$$

The term F_u is a function of parameters that will not enter in the determination of y_u and vanishes in the limit of massless first family and no mixing. The physical choice in eq. (B.3) is to cancel the first factor taking $y_u = 0$, as the cancellation of the other factor would lead to $y_c = 0$. This is stable provided that $\tilde{\lambda}_u > 0$. In a similar way, $y_d = 0$ is a solution to the equation $\partial V_\Sigma / \partial y_d = 0$.

- When deriving with respect to the angle θ we find

$$\begin{aligned} \frac{\partial V_\Sigma}{\partial \theta} = & 2 \sin 2\theta \lambda_{udud} (y_c^2 - y_u^2) (y_s^2 - y_d^2) \times \\ & \times [(y_c^2 - y_u^2) (y_s^2 - y_d^2) \sin^2 \theta + (y_c^2 y_d^2 + y_s^2 y_u^2)] = 0. \end{aligned} \quad (\text{B.4})$$

Substituting the solutions to the previous minima equations considered, $y_u = 0 = y_d$, eq. (B.4) forces $\sin 2\theta = 0$.

- For the heavy Yukawa couplings, once $y_u = y_d = 0$ is chosen the equations take the form:

$$\frac{\partial V_\Sigma}{\partial y_c} = 2y_c \Lambda_f^4 \left(2\lambda_u y_c^2 - 2\lambda_{udud} y_c^2 y_s^4 - \frac{\mu_u^2}{\Lambda_f^2} \right) = 0. \quad (\text{B.5})$$

Neglecting the trivial solution, which is unstable for positive definite coefficients, this equation yields the expression for y_c :

$$y_c^2 = \frac{\mu_u^2}{2\Lambda_f^2 (\lambda_u - \lambda_{udud} y_s^4)} \simeq \frac{\mu_u^2}{2\Lambda_f^2 \lambda_u}, \quad (\text{B.6})$$

where the last equality holds when taking into account the observed value of the strange Yukawa coupling. A similar result can be found for y_s .

Summarizing, neglecting all terms suppressed by $\epsilon_{u,d,\theta}$, the minimum of the scalar potential is given by:

$$y_u = y_d = 0, \quad y_c = \frac{\mu_u}{\sqrt{2}\Lambda_f\sqrt{\lambda_u}}, \quad y_s = \frac{\mu_d}{\sqrt{2}\Lambda_f\sqrt{\lambda_d}}, \quad \sin\theta = 0. \quad (\text{B.7})$$

The observed values of y_c and y_s are understood as the outcome of the hierarchy among the vevs of the flavons, $\langle \Sigma \rangle \sim \mu$, and the flavour scale Λ_f . Note that the parameters $\epsilon_{u,d,\theta}$ do not enter into the definition of y_c and y_s , but control the hierarchy between the light and the heavy generations and the appearance of a non-trivial mixing angle. This solution is stable with all the coefficients in eq. (B.1) positive and furthermore the inclusion of the corrections given by the ϵ -terms will shift the minimum but will not change its stability.

We now discuss the changes of the solutions found above by the introduction of $\epsilon_{u,d,\theta}$.

- The corrections for the first family Yukawa couplings shift their values from zero by an amount $\epsilon_{u,d}$. Explicitly, once the leading order solution found for y_c and y_s is inserted into eq. (B.3), the dominant contributions are given by

$$\frac{\partial V_\Sigma}{\partial y_u} = \Lambda_f^4 \left[2y_u \tilde{\lambda}_u y_c^2 - \epsilon_u \frac{\tilde{\mu}_u^2}{\Lambda_f^2} y_c \right] = 0, \quad (\text{B.8})$$

which leads to a non-vanishing Yukawa coupling for the up quark:

$$y_u = \epsilon_u \frac{\sqrt{\lambda_u} \tilde{\mu}_u}{\sqrt{2} \tilde{\lambda}_u \mu_u} \frac{\tilde{\mu}_u}{\Lambda_f}. \quad (\text{B.9})$$

A similar result holds also for y_d .

- When considering the equation that determines the mixing angle, several corrections are present, although the dominant one is given by

$$\frac{\partial V_\Sigma}{\partial \theta} = 2 \sin\theta \cos\theta y_c^2 y_s^2 [2\lambda_{udud} (y_c^2 y_s^2 \sin^2\theta) - \epsilon_\theta \lambda_{ud}] = 0, \quad (\text{B.10})$$

and the corresponding non-trivial solution reads

$$\sin^2\theta = \epsilon_\theta \frac{\lambda_{ud}}{2 \lambda_{udud} y_s^2 y_c^2}. \quad (\text{B.11})$$

The minimum of the scalar potential proposed in eq. (B.1) is then given by

$$\begin{aligned}
 y_u &\simeq \epsilon_u \frac{\sqrt{\lambda_u} \tilde{\mu}_u}{\sqrt{2} \tilde{\lambda}_u \mu_u \Lambda_f}, & y_d &\simeq \epsilon_d \frac{\sqrt{\lambda_d} \tilde{\mu}_d}{\sqrt{2} \tilde{\lambda}_d \mu_d \Lambda_f}, \\
 y_c &\simeq \frac{\mu_u}{\sqrt{2} \Lambda_f \sqrt{\lambda_u}}, & y_s &\simeq \frac{\mu_d}{\sqrt{2} \Lambda_f \sqrt{\lambda_d}}, \\
 \sin^2 \theta &\simeq \epsilon_\theta \frac{\lambda_{ud}}{2 \lambda_{udud} y_s^2 y_c^2}.
 \end{aligned}
 \tag{B.12}$$

We can now specify the value of $\epsilon_{u,d,\theta}$ in order to accommodate the observed hierarchies and mixing for the first two generations: considering the ratios $\mu/(\sqrt{\lambda}\Lambda_f) \approx \tilde{\mu}/(\sqrt{\tilde{\lambda}}\Lambda_f) \sim 10^{-3}$, it follows that

$$\epsilon_u \sim 10^{-3}, \quad \epsilon_d \sim 5 \times 10^{-2}, \quad \epsilon_\theta \sim 10^{-10}, \tag{B.13}$$

must hold. A comment is in order: when discussing this special illustrative scalar potential, we considered up to dimension 8 operators, while neglecting many terms otherwise allowed by the symmetry. However, even such an arbitrary choice was not sufficient to recover realistic mass hierarchies and the mixing angle, and further fine-tunings were required, including ϵ values as tiny as 10^{-10} to recover the Cabibbo angle. These remarks should suffice to show how unnatural is the set up when trying to fix all observables from pure $d = 5$ Yukawa operators.

B.2 Three family case

The three family case involves a wider variety of operators. This is because some of the accidental simplifications in two families no longer hold for three. The analytic treatment to find the minima becomes more complicated as well, as the number of observables increases to six quark masses and three angles (obviating the CP-odd phase). We present a graphic analysis of the scalar potential in this case. This approach assumes a positive definite potential whose minimum is just the point in which the geometrical surfaces defined by constant invariant quantities meet. When focusing on the masses in either the up or the down sector, we project the parameter space to one that has as many dimensions as families. This means that instead of the curves in the (y_c, y_u) plane of figure 2 we will consider surfaces in (y_t, y_c, y_u) space.

The lowest dimension invariants that involve Yukawa eigenvalues only for the up sector correspond to:

$$\begin{aligned}
 \langle A_u \rangle &= \Lambda_f^2 (y_t^2 + y_c^2 + y_u^2), \\
 \langle B_u \rangle &= \Lambda_f^3 y_t y_c y_u, \\
 \langle A'_{uu} \rangle &= \langle A_u^2 - A_{uu} \rangle = 2 \Lambda_f^4 (y_t^2 y_c^2 + y_u^2 y_t^2 + y_c^2 y_u^2),
 \end{aligned}
 \tag{B.14}$$

where the last invariant is introduced as a linear combination of some of those in eq. (3.3). Notice that three independent invariants are necessary to fix the three different masses. We can study the intersection of the surfaces defined by giving fixed values to these operators.




Figure 3.

In view of these surfaces and the expressions of the invariants, the vevs of the fields shall satisfy the hierarchy:

$$\frac{\langle B_u \rangle}{\Lambda_f^3} \ll \frac{\langle A'_{uu} \rangle}{\Lambda_f^4} \ll \frac{\langle A_u \rangle}{\Lambda_f^2}. \quad (\text{B.15})$$

The same analysis for the down sector leads to the following relation:

$$\frac{\langle B_d \rangle}{\Lambda_f^3} \sim \frac{\langle A'_{dd} \rangle}{\Lambda_f^4} \ll \frac{\langle A_d \rangle}{\Lambda_f^2}. \quad (\text{B.16})$$

The geometrical analysis allows to interpret the vevs of the invariant operators as geometric quantities, assuming the hierarchy in eqs. (B.15), (B.16); as can be seen in figure 4:

1. $\langle A_i \rangle$ sets the radius of the sphere, therefore sets the value of the highest mass.
2. The value of $\langle A'_{ii} \rangle$ determines how close is the surface in figure 3(b) to the axis. The intersection of this curve and the sphere is a circle around the axis, and the radius of such circle is related to the second highest value of mass.
3. The quantity $\langle B_i \rangle$ sets the distance of the surface shown in figure 3(c) to the planes $y_t y_c$, $y_c y_u$ and $y_u y_t$. This surface, considered in the plane of the circle determined by the intersection of the previous surfaces, is an hyperbola so that the graphic image connects to that for the case of two families.

The requirements of eq. (B.15) are not naturally obtained from a general potential, the typical ansatz to fix the vev of the invariants being through a “double-well” potential of the type:

$$V_{f-t} = \lambda_u (A_u - v_{A_u})^2 + \frac{\gamma_u}{\Lambda_{fl}^3} (B_u - v_{B_u})^2 + \frac{\gamma'_u}{\Lambda_{fl}^4} (A'_{uu} - v_{A'_{uu}})^2.$$

However, for writing this kind of potential, one has to neglect many cross terms that would typically spoil the hierarchy. Again, the argument proposed here only illustrates a possible, clearly not natural, way to fix the quark masses.




Figure 4. Determination of the minimum for a positive definite potential constructed with the invariants in eq. (B.14).

The mixing angles appear in the potential through the operator A_{ud} :

$$\begin{aligned}
 V^{(4)} \supset \lambda_{ud} A_{ud} &= \lambda_{ud} (P_0 + P_{\text{int}}) , \\
 P_0 &= - \sum_{i < j} (y_{u_i}^2 - y_{u_j}^2) (y_{d_i}^2 - y_{d_j}^2) \sin^2 \theta_{ij} , \\
 P_{\text{int}} &= \sum_{i < j, k} (y_{d_i}^2 - y_{d_k}^2) (y_{u_j}^2 - y_{u_k}^2) \sin^2 \theta_{ik} \sin^2 \theta_{jk} + \\
 &\quad - (y_d^2 - y_s^2) (y_c^2 - y_t^2) \sin^2 \theta_{12} \sin^2 \theta_{13} \sin^2 \theta_{23} + \\
 &\quad + \frac{1}{2} (y_d^2 - y_s^2) (y_c^2 - y_t^2) \cos \delta \sin 2\theta_{12} \sin 2\theta_{23} \sin \theta_{13} .
 \end{aligned} \tag{B.17}$$

Neglecting the Yukawa couplings for the first family, the equations determining the angles at the minimum of the potential are given by

$$\begin{aligned}
 c_{12} c_{23} s_{12} s_{23} s_{13} \sin \delta &= 0 , \\
 s_{12} c_{12} [y_c^2 + y_t^2 (s_{23}^2 - s_{13}^2 c_{23}^2)] - y_t^2 s_{13} s_{23} c_{23} (c_{12}^2 - s_{12}^2) \cos \delta &= 0 , \\
 s_{23} c_{23} [y_b^2 - y_b^2 s_{13}^2 + y_s^2 s_{12}^2 (1 + s_{13}^2)] - y_s^2 s_{13} s_{12} c_{12} (c_{23}^2 - s_{23}^2) \cos \delta &= 0 , \\
 s_{13} c_{13} (1 - s_{23}^2) (y_b^2 - y_s^2 s_{12}^2) - y_s^2 s_{12} c_{12} s_{23} c_{23} c_{13} \cos \delta &= 0 ,
 \end{aligned} \tag{B.18}$$

where c_{ij} and s_{ij} stand for $\cos \theta_{ij}$ and $\sin \theta_{ij}$. The last three equations can be combined into:

$$s_{12}^2 c_{12}^2 [y_c^2 + y_t^2 (s_{23}^2 - s_{13}^2 c_{23}^2)]^2 = y_t^4 s_{13}^2 s_{23}^2 c_{23}^2 (1 - 2s_{12}^2)^2 \cos^2 \delta , \tag{B.19}$$

$$s_{23}^2 (y_b^2 + y_b^2 s_{13}^2 + y_s^2 s_{12}^2 c_{13}^2) = s_{13}^2 (y_b^2 - y_s^2 s_{12}^2) , \tag{B.20}$$

$$s_{13}^2 c_{13}^2 c_{23}^2 (y_b^2 - y_s^2 s_{12}^2) [y_b^2 - y_s^2 (\cos^2 \delta s_{12}^2 + \sin^2 \delta s_{12}^4)] = 0 . \tag{B.21}$$

From eq. (B.21) it follows that $\sin \theta_{13} = 0$ is a solution. Neglecting this angle, $\sin \theta_{23} = 0$ can be derived from eq. (B.20). Finally, from eq. (B.19), it would result $\sin \theta_{12} = 0$. The other alternatives: $\cos \theta_{13} = 0$ or $\cos \theta_{13} = 0$ lead to unphysical solutions but stand as nonvanishing angle configurations and therefore a novel -if unrealistic- possibility with respect to the two family case.

C The scalar potential for the fundamental approach

There are five independent invariant operators that can be constructed with four fields in fundamental representations of the flavour group, as shown in eqs. (2.22) and (3.7). These invariant operators can be arranged in a vector: denoting this vector by X^2 and by $\langle X^2 \rangle$ its vev,

$$\begin{aligned} X^2 &\equiv \left(\chi_u^{L\dagger} \chi_u^L, \chi_d^{L\dagger} \chi_d^L, \chi_u^{R\dagger} \chi_u^R, \chi_d^{R\dagger} \chi_d^R, \chi_d^{L\dagger} \chi_u^L \right)^T, \\ \langle X^2 \rangle &\equiv \left(|\chi_u^L|^2, |\chi_d^L|^2, |\chi_u^R|^2, |\chi_d^R|^2, \langle \chi_u^{L\dagger} \chi_d^L \rangle \right)^T. \end{aligned} \tag{C.1}$$

All these invariant operators have dimension two and the most general renormalizable scalar potential is given by:

$$V_\chi = -\frac{1}{2} \sum_i (\mu_i^2 X_i^2 + \text{h.c.}) + \sum_{i,j} \lambda_{ij} (X_i^2)^* X_j^2 = -\frac{1}{2} [\mu^2 X^2 + \text{h.c.}] + (X^2)^\dagger \lambda X^2, \tag{C.2}$$

where λ is a 5×5 hermitian matrix¹⁸ and the mass terms are arranged in the vector μ^2 . There are therefore a total of 20 invariant operators in the most general renormalizable potential. Assuming that λ is invertible and adding a constant term to the potential the above expression can be rewritten as:

$$V_\chi = \left(X^2 - \frac{1}{2} \lambda^{-1} \mu^2 \right)^\dagger \lambda \left(X^2 - \frac{1}{2} \lambda^{-1} \mu^2 \right). \tag{C.3}$$

For a bounded-from-below potential, λ has to be positive definite which implies that the minimum of the scalar potential is reached for:

$$\langle X^2 \rangle = \frac{1}{2} \lambda^{-1} \mu^2. \tag{C.4}$$

This is the formal expression for the minimum. Yukawa eigenvalues and the Cabibbo angle are related to the configuration of the potential minimum through eq. (2.31), which together with the previous equation yield:¹⁹

$$\begin{aligned} y_c^2 &= \frac{1}{4\Lambda_f^4} (\lambda^{-1} \mu^2)_{uL} (\lambda^{-1} \mu^2)_{uR}, & y_s^2 &= \frac{1}{4\Lambda_f^4} (\lambda^{-1} \mu^2)_{dL} (\lambda^{-1} \mu^2)_{dR}, \\ \cos \theta_c &= \frac{(\lambda^{-1} \mu^2)_{ud}}{\sqrt{(\lambda^{-1} \mu^2)_{dL} (\lambda^{-1} \mu^2)_{uL}}}. \end{aligned} \tag{C.5}$$

Remarkably, naturalness criteria imply $\cos \theta_c \sim \mathcal{O}(1)$ at this very general level. Yukawa eigenvalues are $\mathcal{O}(\mu^2/\lambda\Lambda_f^2)$, implying $\sqrt{(\lambda^{-1} \mu^2)_{uR,uL}} \sim 10^{-1} \Lambda_f$ in order to fix the charm Yukawa eigenvalue to the observed value and $\sqrt{(\lambda^{-1} \mu^2)_{dR,dL}} \sim 10^{-2} \Lambda_f$ to fix analogously the strange Yukawa eigenvalue.

For the sake of clarity and definiteness, we present next an example of a scalar potential whose mass parameters are directly connected to the Yukawa couplings. We assume that

¹⁸Indices run over the five values $\{uL,dL,uR,dR,ud\}$.

¹⁹ $(\lambda^{-1} \mu^2)_i = \sum_j (\lambda^{-1})_{ij} \mu_j^2$

RH flavons acquire vevs equal to Λ_f , then the parametrization in eq. (2.33) follows. Such assumption can be justified by naturalness arguments or simply fixing the parameters associated to $|\chi_{d,u}^R|$ through eq. (C.4). We can then concentrate only on the scalar potential for the LH flavons:

$$V'_\chi = \lambda_u \left(\chi_u^{L\dagger} \chi_u^L - \frac{\mu_u^2}{2\lambda_u} \right)^2 + \lambda_d \left(\chi_d^{L\dagger} \chi_d^L - \frac{\mu_d^2}{2\lambda_d} \right)^2 + \lambda_{ud} \left(\chi_u^{L\dagger} \chi_d^L - \frac{\mu_{ud}^2}{2\lambda_{ud}} \right)^2. \quad (\text{C.6})$$

As already stated, at the minimum the invariants in this potential can be written in terms of Yukawa eigenvalues and the Cabibbo angle:

$$V'_\chi = \lambda_u \left(\Lambda_f^2 y_c^2 - \frac{\mu_u^2}{2\lambda_u} \right)^2 + \lambda_d \left(\Lambda_f^2 y_s^2 - \frac{\mu_d^2}{2\lambda_d} \right)^2 + \lambda_{ud} \left(\Lambda_f^2 y_c y_s \cos \theta_c - \frac{\mu_{ud}^2}{2\lambda_{ud}} \right)^2. \quad (\text{C.7})$$

From this relation, the expression for Yukawa eigenvalues and the Cabibbo angle in terms of the parameters of the potential can be read:

$$y_c = \frac{\mu_u}{\sqrt{2\lambda_u}\Lambda_f}, \quad y_s = \frac{\mu_d}{\sqrt{2\lambda_d}\Lambda_f}, \quad \cos \theta_c = \frac{\sqrt{\lambda_u\lambda_d}\mu_{ud}^2}{\lambda_{ud}\mu_d\mu_u}. \quad (\text{C.8})$$

The resulting $\cos \theta_c$ is naturally of $\mathcal{O}(1)$, while correct charm and strange masses arise when $\mu_u \sim 10^{-2}\sqrt{\lambda_u}\Lambda_f$ and $\mu_d \sim 10^{-3}\sqrt{\lambda_d}\Lambda_f$. The differences with the bi-fundamental approach can be seen comparing the above equation with eq. (B.12). The example shown corresponds to a potential with some terms omitted,²⁰ whose mere purpose is to illustrate explicitly the mechanism of generation of Yukawa eigenvalues and mixing angle through a scalar potential for the $d = 6$ Yukawa operator.

Finally, notice that the extension to the three family case is trivial, substituting in the formulae above y_c and y_s by y_t and y_b , respectively, and the Cabibbo angle by θ_{23} . This stems from the fact that, considering the renormalizable scalar potential, only the heaviest Yukawas are non-vanishing, as discussed in the main text.

D Note added in proof

After this work was submitted, a paper appeared in the arXiv [28] where it has been suggested that the introduction of Coleman-Weinberg quantum corrections to our results for the bi-fundamental case could generate subdominant Yukawa splittings.

The author, using a slightly modified version of our notation, re-derived our renormalizable potential for the bi-fundamental case for three families. Looking at eq. (3.6) and eqs. (2.9)–(2.11), it is easy to identify the relations to move from one notation to the other:

$$\begin{aligned} A_u &\rightarrow T_u, & B_u &\rightarrow D_u, & A'_{uu} &= (A_u^2 - A_{uu}) \rightarrow 2A_u \\ \mu_u &\rightarrow m_u, & \tilde{\mu}_u &\rightarrow 2|\tilde{\mu}_u|, & \lambda_u &\rightarrow \lambda_u + \frac{1}{2}\tilde{\lambda}'_u, & \lambda'_u &\rightarrow -\frac{1}{2}\tilde{\lambda}'_u. \end{aligned}$$

The freedom on the relative sign between the determinant and $\tilde{\mu}$ terms allowed in our paper has been retaken in v2 of [28] as a cosine dependence, which now allows negative coefficients and redefines their norm.

²⁰Terms like $g_{ud}\chi_d^{L\dagger}\chi_d^L\chi_u^{L\dagger}\chi_u^L$ do not affect the position of the minimum provided $g_{ud} < y_s^2/y_c^2$.

The hierarchical solution in which that paper is based, was already identified in our work, together with the degenerate one. To quantify the validity range of the two solutions we found, we add here a detailed analysis of the stability of the potential, that was not included in our previous version.

A) Stability condition for two families. For the two-family case the extremality equations read:

$$\begin{aligned} \frac{\partial V_u}{\partial y_c} &= 4\lambda_u \Lambda_{fl}^2 y_c \left(A_u - \frac{\mu_u^2}{2\lambda_u} \right) + 2\tilde{\lambda}_u \Lambda_{fl}^2 y_u \left(B_u - \frac{\tilde{\mu}_u^2}{2\tilde{\lambda}_u} \right) - h_u \Lambda_{fl}^2 (y_u A_u + 2y_c B_u) = 0 \\ \frac{\partial V_u}{\partial y_u} &= 4\lambda_u \Lambda_{fl}^2 y_u \left(A_u - \frac{\mu_u^2}{2\lambda_u} \right) + 2\tilde{\lambda}_u \Lambda_{fl}^2 y_c \left(B_u - \frac{\tilde{\mu}_u^2}{2\tilde{\lambda}_u} \right) - h_u \Lambda_{fl}^2 (y_c A_u + 2y_u B_u) = 0. \end{aligned}$$

One can easily verify that the hierarchical pattern $(0, y)$ is not a solution of these equations, unless a severe fine-tuning on the parameters $\mu_u, \tilde{\mu}_u, \lambda_u$ and h_u is introduced. Only the symmetric solution (y, y) arises as a natural minimum.

B) Stability condition for three families. The conditions defining the minima now read as follows.

1. The parameter region in which only the symmetric solution (y, y, y) provides a stable minimum is defined by

$$\frac{\tilde{\mu}_u^2}{\mu_u^2} > \frac{8 \lambda_u'^2}{\lambda_u + \lambda_u'}$$

for $\lambda_u' < 0$. On the other hand, for $\lambda_u' > 0$, the configuration (y, y, y) is a stable minimum for any value of $\tilde{\mu}_u^2/\mu_u^2$.

2. The parameter region in which the symmetric solution is the absolute minimum, while the hierarchical configuration $(0, 0, y)$ is a local minimum corresponds to

$$8(\lambda_u + \lambda_u') \left(\left(4 - 2 \frac{\lambda_u'}{\lambda_u + \lambda_u'} \right)^{3/2} - \left(8 - 6 \frac{\lambda_u'}{\lambda_u + \lambda_u'} \right) \right) < \frac{\tilde{\mu}_u^2}{\mu_u^2} < \frac{8 \lambda_u'^2}{\lambda_u + \lambda_u'} \quad (\text{D.1})$$

3. The parameter region in which the symmetric solution is a local minimum, while the hierarchical solution is the absolute minimum is defined by

$$\frac{8 \lambda_u'^2}{3\lambda_u + 2\lambda_u'} < \frac{\tilde{\mu}_u^2}{\mu_u^2} < 8(\lambda_u + \lambda_u') \left(\left(4 - 2 \frac{\lambda_u'}{\lambda_u + \lambda_u'} \right)^{3/2} - \left(8 - 6 \frac{\lambda_u'}{\lambda_u + \lambda_u'} \right) \right). \quad (\text{D.2})$$

4. Finally, the parameter region in which only the hierarchical configuration is a minimum corresponds to

$$\frac{\tilde{\mu}_u^2}{\mu_u^2} < \frac{8 \lambda_u'^2}{3\lambda_u + 2\lambda_u'}. \quad (\text{D.3})$$

As illustrated in figure (5) (see also [29]), for a typical λ_u value in the perturbative regime the symmetric configuration is the absolute minimum for most of the parameter space (here shown in dark and light orange). However, even when the hierarchical solution is




Figure 5. Parameter space for the symmetric and hierarchical configurations. The dark-Orange corresponds to a region where the symmetric solution (y, y, y) is the stable absolute minimum, while the hierarchical solution $(0, 0, y)$ is a saddle point. In the light-Orange region, the symmetric configuration is the absolute minimum, while the hierarchical solution is a local one. On the contrary, in the light-Brown region the symmetric configuration is a local minimum, while the hierarchical solution is an absolute one. Finally, in the dark-Brown region only the hierarchical solution is a minimum. In the plot the value $\lambda_u = 1/2$ has been used for illustration.

preferred, yielding non-zero top and bottom Yukawa eigenvalues only, the hierarchy among up and down sectors must be considered. In particular the presence of the term $g_{ud} A_u A_d$ must be constrained by setting $g_{ud} < y_b^2 / y_t^2 \sim 10^{-3}$ to warranty the top-bottom mass hierarchy. As already stated before, such a fine-tuning can be justified through additional symmetries. In particular in [28], it is placed in the vev of flavons transforming under Abelian factors.

References

- [1] G. D’Ambrosio, G.F. Giudice, G. Isidori and A. Strumia, *Minimal flavour violation: An effective field theory approach*, *Nucl. Phys. B* **645** (2002) 155 [[hep-ph/0207036](#)] [[SPIRES](#)].
- [2] R.D. Peccei and H.R. Quinn, *Constraints imposed by CP conservation in the presence of instantons*, *Phys. Rev. D* **16** (1977) 1791 [[SPIRES](#)].
- [3] A.J. Buras, *Minimal flavour violation and beyond: Towards a flavour code for short distance dynamics*, *Acta Phys. Polon. B* **41** (2010) 2487 [[arXiv:1012.1447](#)] [[SPIRES](#)].
- [4] G.C. Branco, W. Grimus and L. Lavoura, *Relating the scalar flavour changing neutral couplings to the CKM matrix*, *Phys. Lett. B* **380** (1996) 119 [[hep-ph/9601383](#)] [[SPIRES](#)].
- [5] A.L. Kagan, G. Perez, T. Volansky and J. Zupan, *General minimal flavor violation*, *Phys. Rev. D* **80** (2009) 076002 [[arXiv:0903.1794](#)] [[SPIRES](#)].
- [6] G. Isidori, Y. Nir and G. Perez, *Flavor physics constraints for physics beyond the standard model*, [arXiv:1002.0900](#) [[SPIRES](#)].
- [7] G. Isidori, *The challenges of flavour physics*, [arXiv:1012.1981](#) [[SPIRES](#)].

- [8] Z. Lalak, S. Pokorski and G.G. Ross, *Beyond MFV in family symmetry theories of fermion masses*, *JHEP* **08** (2010) 129 [[arXiv:1006.2375](#)] [[SPIRES](#)].
- [9] A.L. Fitzpatrick, G. Perez and L. Randall, *Flavor from minimal flavor violation & a viable Randall-Sundrum model*, [arXiv:0710.1869](#) [[SPIRES](#)].
- [10] R.S. Chivukula and H. Georgi, *Composite technicolor standard model*, *Phys. Lett. B* **188** (1987) 99 [[SPIRES](#)].
- [11] C.D. Froggatt and H.B. Nielsen, *Hierarchy of quark masses, Cabibbo angles and CP-violation*, *Nucl. Phys. B* **147** (1979) 277 [[SPIRES](#)].
- [12] G. Altarelli and F. Feruglio, *Discrete flavor symmetries and models of neutrino mixing*, *Rev. Mod. Phys.* **82** (2010) 2701 [[arXiv:1002.0211](#)] [[SPIRES](#)].
- [13] B. Grinstein, M. Redi and G. Villadoro, *Low scale flavor gauge symmetries*, *JHEP* **11** (2010) 067 [[arXiv:1009.2049](#)] [[SPIRES](#)].
- [14] T. Feldmann, *See-saw masses for quarks and leptons in SU(5)*, *JHEP* **04** (2011) 043 [[arXiv:1010.2116](#)] [[SPIRES](#)].
- [15] V. Cirigliano, B. Grinstein, G. Isidori and M.B. Wise, *Minimal flavor violation in the lepton sector*, *Nucl. Phys. B* **728** (2005) 121 [[hep-ph/0507001](#)] [[SPIRES](#)].
- [16] S. Davidson and F. Palorini, *Various definitions of minimal flavour violation for leptons*, *Phys. Lett. B* **642** (2006) 72 [[hep-ph/0607329](#)] [[SPIRES](#)].
- [17] M.B. Gavela, T. Hambye, D. Hernandez and P. Hernández, *Minimal flavour seesaw models*, *JHEP* **09** (2009) 038 [[arXiv:0906.1461](#)] [[SPIRES](#)];
- [18] PARTICLE DATA GROUP collaboration, K. Nakamura et al., *Review of particle physics*, *J. Phys. G* **37** (2010) 075021 [[SPIRES](#)].
- [19] T. Feldmann and T. Mannel, *Minimal flavour violation and beyond*, *JHEP* **02** (2007) 067 [[hep-ph/0611095](#)] [[SPIRES](#)].
- [20] A.J. Buras, K. Gemmler and G. Isidori, *Quark flavour mixing with right-handed currents: an effective theory approach*, *Nucl. Phys. B* **843** (2011) 107 [[arXiv:1007.1993](#)] [[SPIRES](#)].
- [21] T. Feldmann, M. Jung and T. Mannel, *Sequential flavour symmetry breaking*, *Phys. Rev. D* **80** (2009) 033003 [[arXiv:0906.1523](#)] [[SPIRES](#)].
- [22] G. Colangelo, E. Nikolidakis and C. Smith, *Supersymmetric models with minimal flavour violation and their running*, *Eur. Phys. J. C* **59** (2009) 75 [[arXiv:0807.0801](#)] [[SPIRES](#)].
- [23] L. Mercolli and C. Smith, *EDM constraints on flavored CP-violating phases*, *Nucl. Phys. B* **817** (2009) 1 [[arXiv:0902.1949](#)] [[SPIRES](#)].
- [24] R. Alonso, M.B. Gavela, L. Merlo and S. Rigolin, in preparation.
- [25] Z. Berezhiani and F. Nesti, *Supersymmetric SO(10) for fermion masses and mixings: Rank-1 structures of flavour*, *JHEP* **03** (2006) 041 [[hep-ph/0510011](#)] [[SPIRES](#)].
- [26] L. Ferretti, S.F. King and A. Romanino, *Flavour from accidental symmetries*, *JHEP* **11** (2006) 078 [[hep-ph/0609047](#)] [[SPIRES](#)].
- [27] L. Calibbi, L. Ferretti, A. Romanino and R. Ziegler, *Consequences of a unified, anarchical model of fermion masses and mixings*, *JHEP* **03** (2009) 031 [[arXiv:0812.0087](#)] [[SPIRES](#)].
- [28] E. Nardi, *Naturally large Yukawa hierarchies*, [arXiv:1105.1770](#) [[SPIRES](#)].
- [29] B. Gavela, plenary talk at *Planck 2011*, Lisbon Portugal, 30 May – 3 June 2011.