

January 9, 2018

Recent Advances in Clinical Practice

Modern Diagnosis of GERD: The Lyon Consensus

C Prakash Gyawali<sup>1</sup>, Peter Kahrilas<sup>2</sup>, Edoardo Savarino<sup>3</sup>, Frank Zerbib<sup>4</sup>, Francois Mion<sup>5</sup>, André Smout<sup>6</sup>, Michael Vaezi<sup>7</sup>, Daniel Sifrim<sup>8</sup>, **Mark Fox<sup>9</sup>**, Marcelo F. Vela<sup>10</sup>, Radu Tutuian<sup>11</sup>, Jan Tack<sup>12</sup>, Albert J Bredenoord<sup>6</sup>, John Pandolfino<sup>2</sup>, Sabine Roman<sup>5</sup>

(4999 words)

<sup>1</sup>Division of Gastroenterology, Washington University School of Medicine, St. Louis, MO, USA

<sup>2</sup>Division of Gastroenterology, Department of Medicine, Northwestern University, Chicago, IL, USA

<sup>3</sup>Division of Gastroenterology, Department of Surgical, Oncological and Gastroenterological Sciences, Padua, Italy

<sup>4</sup>Department of Gastroenterology, Bordeaux University Hospital, and Université de Bordeaux, Bordeaux, France

<sup>5</sup> Université de Lyon, Hospices Civils de Lyon, Digestive Physiology, Hopital E Herriot, F-69437 LYON, France; Université de Lyon, Lyon I University, Digestive Physiology, F-69008 LYON, France; Université de Lyon, Inserm U1032, LabTAU, F-69008 LYON, France

<sup>6</sup>Department of Gastroenterology and Hepatology, Academic Medical Center, Amsterdam, The Netherlands

<sup>7</sup>Division of Gastroenterology, Vanderbilt University, Nashville, TN, USA

<sup>8</sup>Barts and The London School of Medicine and Dentistry Queen Mary, University of London, UK

<sup>9</sup>**Gastroenterology, St. Claraspital, Kleinriedenstrasse 30, Basel, Switzerland**

<sup>10</sup>Division of Gastroenterology and Hepatology, Mayo Clinic, Scottsdale, AZ, USA

<sup>11</sup>Division of Gastroenterology, University Clinics for Visceral Surgery and Medicine, Bern University Hospital, Bern, Switzerland

<sup>12</sup>Department of Gastroenterology, Catholic University of Leuven, Leuven, Belgium

Address all correspondence to

C. Prakash Gyawali, MD, MRCP  
Professor of Medicine  
Washington University School of Medicine  
660 S. Euclid Ave, Campus Box 8124  
St. Louis, MO 63110  
Email: [cprakash@wustl.edu](mailto:cprakash@wustl.edu)  
Phone: 314 454 8201  
Fax: 314 362 3643

Contributorship: All authors contributed to the content of the manuscript, and reviewed, edited and approved the final draft.

## DISCLOSURES

C. Prakash Gyawali: Consulting: Ironwood, Torax, Quintiles; Teaching and speaking: Medtronic, Diversatek, Reckitt-Benckiser

Edoardo Savarino: Consulting: Abbvie, Allergan, MSD, Takeda, Sofar, Janssen; Teaching and speaking: Medtronic, Reckitt-Benckiser, Malesci, Zambon

Frank Zerbib: Research support: Medtronic, Sandhill Scientific; Consulting: Allergan, Reckitt-Benckiser; Speaking and teaching: Ipsen Pharma, Biocodex, Coloplast, Takeda, Vifor Pharma, Mayoly Spindler

Peter Kahrilas: Consulting: Ironwood

Francois Mion: Teaching & speaking: Laborie, Medtronic; Consulting: Allergan, Endostim

André Smout: none

Michael Vaezi: Vanderbilt University and Diversatek co-own patent on mucosal impedance technology

Daniel Sifrim: Research support: Diversatek, Reckitt-Benckiser; OMOM®, Jinshan Science & Technology (Group) Co. Ltd., Chongqing, China

Mark Fox: Research support: Given Imaging / Covidien, Reckitt Benckiser, Mui Scientific. Educational events: Given Imaging / Covidien, MMS, Sandhill Scientific Instruments. Speaking and teaching: Given Imaging / Covidien, Reckitt Benckiser, Shire, Almirall.

Marcelo Vela: consulting: Torax

Radu Tutuian: Teaching: Laborie

Jan Tack: Consulting: Ironwood

Arjan Bredenoord: Research support: Danone, Bayer; speaking and/or consulting: MMS, Astellas, AstraZeneca, Bayer, Almirall and Allergan

John Pandolfino: Research support: Impleo; speaking and/or consulting: Medtronic, Diversatek, Torax, Ironwood, Takeda, Astra Zeneca; stock options: Crospon

Sabine Roman: Research support: Sandhill Scientific, Crospon; Teaching: Medtronic; Speaker: Mayoly Spindler

**ABSTRACT**

Clinical history, questionnaire data and response to antisecretory therapy are insufficient to make a conclusive diagnosis of GERD in isolation, but are of value in determining need for further investigation. Conclusive evidence for reflux on esophageal testing include advanced grade erosive esophagitis (LA grades C and D), long-segment Barrett's mucosa or peptic strictures on endoscopy, or distal esophageal acid exposure time (AET) >6% on ambulatory pH or pH-impedance monitoring. A normal endoscopy does not exclude GERD, but provides supportive evidence refuting GERD in conjunction with distal AET <4% and <40 reflux episodes on pH-impedance monitoring off PPI. Reflux-symptom association on ambulatory reflux monitoring provides supportive evidence for reflux triggered symptoms, and may predict a better treatment outcome when present. When endoscopy and pH- or pH-impedance monitoring are inconclusive, adjunctive evidence from biopsy findings (histopathology scores, dilated intercellular spaces), motor evaluation (hypotensive lower esophageal sphincter, hiatus hernia and esophageal body hypomotility on high-resolution manometry) and novel impedance metrics (baseline impedance, post-reflux swallow induced peristaltic wave index) can add confidence for a GERD diagnosis; however, diagnosis cannot be based on these findings alone. An assessment of anatomy, motor function, reflux burden and symptomatic phenotype will therefore help direct management. Future GERD management strategies should focus on defining individual patient phenotypes based on the level of refluxate exposure, mechanism of reflux, efficacy of clearance, underlying anatomy of the esophagogastric junction, and psychometrics defining symptomatic presentations.

Keywords: gastroesophageal reflux disease; pH-impedance monitoring; manometry; endoscopy

## INTRODUCTION

Gastroesophageal reflux disease (GERD) has an estimated worldwide prevalence of 8-33%, involves all age groups and both genders<sup>1</sup>, and carries a price tag estimated at >\$9-10 billion/year in the US alone, largely related to proton pump inhibitors (PPI) use and diagnostic testing<sup>2</sup>. The current paradigm of GERD diagnosis hinges on the identification of esophageal mucosal lesions or troublesome symptoms caused by gastroesophageal reflux<sup>3</sup>. A putative GERD diagnosis is bolstered by a favorable response to PPI therapy<sup>4</sup>. The primary determinant of mucosal injury is excessive esophageal acid exposure attributable to anatomical or physiological defects of the esophagogastric junction (EGJ) and esophageal peristalsis<sup>5</sup>. GERD symptoms, however, have multiple potential determinants including the number of reflux episodes, the proximal extent to which the refluxate migrates, the acidity of the refluxate, esophageal hypersensitivity, and cognitive hypervigilance. Consequently, depending on the clinical context, the defining features of GERD can be pathology, physiology, or symptomatology. In this paradigm, esophageal testing is often undertaken to define optimal management, be that PPI therapy, antireflux surgery (ARS), or cognitive behavioral therapy.

The aim of the GERD consensus process was to determine modern indications for esophageal testing in GERD, and as an extension to that aim, to define criteria for a clinical diagnosis of GERD. The consensus process started in 2014, when the primary aims were formulated, followed by a literature search and grading of evidence. There was extensive discourse within a multi-national group of GERD experts over 2 years, following which consensus statements were developed and published<sup>6-8</sup>. These consensus statements were adapted for the practicing gastroenterologist by a cohort of international experts at the Lyon GERD consensus meeting in November 2017, the conclusions from which are presented in this manuscript.

## DIAGNOSIS OF GERD

GERD is empirically diagnosed and treated in clinical practice based on the clinician's symptom assessment. Indications for testing include treatment failure, diagnostic uncertainty, and treating (or preventing) complications of GERD. However, diagnostic testing may or may not support the initial diagnosis, as the criteria defining GERD are specific to each testing modality. Consequently, understanding the performance characteristics of each diagnostic modality and recognizing evidence that supports or refutes the clinical impression of GERD is crucial. The Lyon Consensus evaluated GERD diagnostic tests from that perspective, and test results were categorized as being adequate to establish or refute a GERD diagnosis or inconclusive in the absence of additional supportive evidence.

The primary focus of esophageal testing has hitherto been restricted to detection of excessive acid reflux as indicative of pathologic GERD, supported by reflux-symptom association analysis. **The threshold value discriminating abnormal from normal esophageal acid exposure with 24-hour pH-metry was initially selected based on evidence that higher acid exposure is associated with the presence of reflux esophagitis<sup>9</sup>. However, the focus of current GERD management is not only on healing mucosal disease but also on managing symptoms. The association of esophageal acid exposure with patient symptoms is weak<sup>10, 11</sup>, making sole reliance of this metric problematic.** Expansion of testing to include esophageal mucosal impedance, manometry, histopathology, and psychometrics may help in this regard, potentially identifying distinct GERD phenotypes with unique management implications. Each test adds a piece to the overall puzzle of symptom generation, disease pathophysiology, and precision management.

### **Clinical history and questionnaires**

Typical GERD symptoms (heartburn and acid regurgitation) are more likely than atypical symptoms to respond to treatment, emphasizing the value of an accurate clinical history<sup>5</sup>. However, when compared to objective evidence of GERD defined by pH-metry or endoscopy, even an expert history by a

gastroenterologist has only 70% sensitivity and 67% specificity<sup>12</sup>, reiterating the distinction between a physiology-based and a symptom-based GERD diagnosis. Likewise, questionnaires such as the reflux disease questionnaire (RDQ) and GERDQ have similar limitations when compared to physiologic testing<sup>12-14</sup>. However, in clinical practice, diagnosing and treating GERD based on typical symptoms is pragmatic and endorsed by societal guidelines<sup>4</sup>, even though these symptoms are neither sensitive nor specific for objectively defined GERD<sup>12</sup>.

### **Proton pump inhibitor (PPI) trial**

Although pragmatic, symptomatic response to PPI therapy does not equate to a GERD diagnosis, exhibiting an imperfect correspondence with objectively defined disease. On average, 69% of esophagitis patients, 49% of non-erosive reflux disease (NERD) patients, and 35% of patients with normal endoscopy and pH-metry gain symptom relief from a PPI trial<sup>15</sup>. Hence, when evaluated as a diagnostic test for GERD among heartburn patients, an empiric PPI trial has a sensitivity of 71% and specificity of only 44% compared to the combination of endoscopy and pH-metry<sup>12,16</sup>. With atypical symptoms (chest pain, chronic cough, laryngitis, etc), PPI response rates are much lower than with heartburn, thereby diminishing the utility of that approach to diagnosis<sup>17</sup>. A major limitation of the “PPI test” is the strong modulation of symptoms by esophageal hypersensitivity<sup>18,19</sup>; there is also variation in PPI dosing and duration of the test<sup>20</sup>. Nonetheless, despite low specificity and high placebo response<sup>21</sup>, the empiric PPI treatment approach is less costly than diagnostic testing<sup>22</sup> and is endorsed by societal guidelines<sup>4, 23</sup> undoubtedly leading to the over-diagnosis of GERD and overuse of PPIs.

## **Endoscopy and biopsy**

When putative GERD symptoms do not respond to empiric PPI therapy, upper endoscopy (EGD) is advised both to evaluate for GERD complications and to detect potential alternative diagnoses that might redirect therapy. High-grade esophagitis (LA Grades C or D), Barrett's esophagus, or peptic stricturing are considered confirmatory evidence for GERD<sup>6</sup>. However, erosive esophagitis is found in only 30% of treatment-naive heartburn patients and in <10% when already taking a PPI.<sup>24, 25</sup> Furthermore, most of that is low-grade, and lower grades of esophagitis, particularly LA Grade A, are nonspecific, found in 5-7.5% of asymptomatic controls<sup>26-28</sup>. When accurately defined, LA Grade B esophagitis provides adequate evidence for initiation of medical management of GERD, but problems with inter-observer variability led an expert panel to conclude that additional pH-metry evidence is requisite prior to pursuing anti-reflux surgery (ARS)<sup>29</sup>. Barrett's esophagus is observed in 5-15% of chronic GERD patients<sup>30-32</sup>, but histological confirmation is documented in only 50% of these. In summary, EGD findings can be clinically important and specific for GERD, but EGD has a low sensitivity in GERD diagnosis.

The Rome IV consensus recommended esophageal biopsies during EGD to rule out eosinophilic esophagitis<sup>33</sup>. Biopsies may also have value in differentiating NERD (with positive pH-metry) from reflux hypersensitivity, functional heartburn and controls when scored using a structured histopathological protocol evaluating papillary elongation, basal cell hyperplasia, dilated intercellular spaces, intraepithelial inflammatory cells, necrosis and erosions<sup>34, 35</sup>; changes that resolve following adequate GERD therapy<sup>36</sup>. However, histopathological findings can overlap between the groups studied, and are not conclusive of GERD. Identification of dilated intercellular spaces on electron microscopy suggests mucosal injury from reflux<sup>37, 38</sup>, but clinical application is limited. The widespread adoption of histopathological examination for GERD injury is hindered by the cumbersome protocol and need for a dedicated esophageal pathologist<sup>34, 39, 40</sup>.

### **Ambulatory reflux monitoring**

Ambulatory reflux monitoring can provide confirmatory evidence of GERD, in patients with normal endoscopy, atypical symptoms and/or when contemplating ARS<sup>6</sup>. Reflux monitoring demonstrates the consequence of GERD pathophysiology, evident as either excessive esophageal acid exposure time (AET) or reflux episodes, rather than the mechanism by which that occurs. Reflux-symptom association utilizes simple ratios and statistical tests to determine whether reflux episodes co-occur with symptoms, and adds value to ambulatory reflux monitoring. Hence, reflux monitoring can confirm or exclude pathologic GERD, albeit not always conclusively.

The primary outcome of a 24-hour pH-metry study is the AET. Extending recording time to 48 or 96 hours with the wireless pH monitoring system increases the diagnostic yield<sup>41-43</sup> and test reproducibility<sup>44</sup>, and is particularly useful when a transnasal catheter was not tolerated or yielded a negative result despite high suspicion of GERD<sup>41, 45</sup>. However, wireless pH monitoring is expensive, limiting its availability. Another variation on reflux monitoring is pH-impedance monitoring, which characterizes reflux events with both a pH electrode and a series of impedance electrodes. Since pH-impedance detects all reflux (liquid, gas or mixed) regardless of acidity, and defines the direction of flow, it is considered the gold standard<sup>6, 46</sup>. However, the added yield is limited<sup>47, 48</sup>, the test is not widely available, and the interpretation is laborious.

Reflux monitoring can be done ‘on’ or ‘off’ PPI therapy in patients with persistent and/or atypical symptoms despite PPI therapy. The Lyon Consensus proposes that testing always be performed off therapy to demonstrate baseline AET in ‘unproven GERD’, meaning no (or low grade) esophagitis at endoscopy, and no prior positive pH testing<sup>8</sup>. Testing off therapy is also recommended when done to evaluate for ARS<sup>6, 33</sup>. In contrast, the Lyon Consensus proposes that patients with ‘proven GERD’ (prior LA Grade C or D esophagitis, long segment Barrett’s esophagus, or prior abnormal pH-metry) be evaluated on double-dose PPI therapy to establish correlation between refractory symptoms and reflux



episodes and/or to exclude inadequate acid suppression or poor compliance as the mechanism of persisting symptoms. This assessment requires pH-impedance (versus pH) monitoring since most reflux episodes on PPI therapy are weakly acidic (pH 4-7) <sup>49, 50</sup>.

Assessment of proximal esophageal or pharyngeal reflux has also been proposed, but methodology and interpretation have not been standardized, and outcome studies are currently lacking <sup>51-54</sup>. An additional problem with the pharyngeal pH probe designed to evaluate both aerosolized and liquid acid reflux <sup>55</sup> is that it detects pharyngeal pH drops in the absence of concomitant esophageal pH-impedance events <sup>56, 57</sup>, even in post-gastrectomy patients, raising questions about its accuracy <sup>58</sup>.

#### *Interpretation of pH and pH-Impedance Monitoring*

Among the pH monitoring metrics, AET is the most reproducible <sup>59</sup>, is reliably extracted from automated analysis, and is predictive of response from medical and surgical reflux therapy <sup>60, 61</sup>. However, the significance of an abnormal AET is proportionate to the degree of abnormality, and the Lyon Consensus proposes that AET less than 4% be considered definitively normal (physiologic) and greater than 6% be considered definitively abnormal <sup>6</sup> with intermediate values between these limits being inconclusive. Another outcome metric of pH-impedance monitoring is the number of reflux episodes (acidic, weakly acidic or weakly alkaline) with the caveat that this is overestimated by the automated analysis and requires manual review of the tracing <sup>62</sup>. The Lyon GERD Consensus proposes that > 80 reflux episodes per 24 hours are definitively abnormal, while a number <40 is physiologic <sup>6</sup> and intermediate values inconclusive. The clinical relevance of an abnormal number of reflux episodes remains incompletely defined, although recent preliminary data demonstrate improvement of regurgitation verified by increased reflux episodes following magnetic sphincter augmentation <sup>63</sup>. Consequently, this is considered an adjunctive measure to be utilized when AET is inconclusive (i.e. between 4-6%). pH-impedance monitoring assists diagnosis of belching disorders and rumination, which

can mimic reflux disease. Additional impedance parameters such as bolus exposure, baseline impedance and post reflux swallow-induced peristaltic wave (PSPW) also have potential as reflux metrics, but outcome data are currently limited.

### *Reflux-Symptom Association*

Both pH monitoring and combined pH-impedance monitoring provide analysis of the temporal association between symptoms with a crisp onset (e.g. heartburn, regurgitation, chest pain, cough or belching) and reflux episodes<sup>64</sup>. The time window applied for reflux-symptom association analysis is 2 minutes<sup>65,66</sup>. The Symptom Index (SI) is the percentage of symptom events preceded by reflux episodes<sup>67</sup>, and the optimal SI threshold for heartburn is 50%<sup>68</sup>. The disadvantage of the SI is that the number of reflux episodes is not considered leaving open the possibility of chance association. The Symptom Association Probability (SAP) and the Ghillebert Probability Estimate (GPE), also known as Binomial Symptom Index (BSI), use more complex statistical calculations to express the probability that symptom events and reflux episodes are associated<sup>69,70</sup> and are considered positive if the probability (p value) of the observed association occurring by chance is < 5%. **Both SAP and GPE/BSI take all relevant components, i.e. total numbers of symptom events, reflux episodes and reflux-related symptom events, into account. In summary, SI is a measure of “effect size”, whereas SAP is a measure of probability. As such the two metrics are complementary, measure different things and cannot be compared to each other. The combination of a positive SI and positive SAP provides the best evidence of a clinically relevant association between reflux episodes and symptoms<sup>6,71,72</sup>. Both the SI and SAP are predictive of the effect of medical and surgical anti-reflux therapy, independent of AET<sup>73-76</sup>.**

The reliability of reflux-symptom association analysis is critically dependent on proper execution of the reflux monitoring procedure and meticulous analysis protocols including careful selection of symptoms of interest. Patients must be instructed to use the symptom event button on the portable data logger accurately and to fill in the symptom diary accurately. The outcome of symptom association

analysis is more reliable when at least three symptom events occur during the test<sup>6</sup>. **When cough is the symptom of interest, an automated acoustic or manometric cough monitor is necessary to accurately capture cough events<sup>77, 78</sup>.** Prolonged wireless pH monitoring increases the yield of symptom association analysis<sup>41, 42</sup>, as does combined pH-impedance monitoring with the detection of weakly acidic reflux episodes<sup>79</sup>. This requires manual analysis, as automated analysis significantly over-detects weakly acidic reflux episodes and inaccurately reports association with non-acid reflux events in nearly 20% of cases<sup>62</sup>. However, “rapid” visual analysis limited to the 2-minute window preceding each symptom event yields SI and SAP values concordant with a full visual analysis, with excellent intra- and inter-observer agreement<sup>80</sup>. Although pH-impedance monitoring provides analysis of symptom-reflux association on PPI therapy, testing off PPI therapy increases the number of symptoms reported, which increases the chance of a positive symptom-reflux association<sup>81</sup>.

Reflux-symptom association analysis has a high degree of reproducibility, the SI being somewhat less reproducible than the SAP<sup>82</sup>. Monte Carlo simulations have identified limitations of reflux-symptom association analysis, especially when acid exposure is low and symptom events are few<sup>83, 84</sup>. If new methods for reflux-symptom association are developed in the future, rigorous outcome testing will be needed to define superiority to the existing metrics.

### *Novel Metrics*

Two novel impedance-detected parameters, the PSPW index and baseline impedance have been investigated within GERD phenotypes<sup>85, 86</sup>. These metrics may augment the diagnostic value of impedance-pH testing, especially in discriminating patients with GERD from those with functional heartburn<sup>87-89</sup>. In health, reflux episodes trigger primary peristalsis to neutralize acidified esophageal mucosa with saliva. This is evident as the antegrade progression of impedance decline within 30 seconds of a reflux episode (PSPW) on a pH-impedance study<sup>88</sup>. **The PSPW index, which currently requires cumbersome manual calculation as it is not programmed into the analysis software, consists of the**

proportion of reflux episodes on pH-impedance monitoring followed by a PSPW. The PSPW index reflects the integrity of primary peristalsis stimulated by reflux episodes, correlates with contraction reserve assessed using multiple rapid swallows (MRS)<sup>90</sup>, and has excellent performance characteristics in differentiating erosive esophagitis and pathological acid exposure from functional heartburn and controls (sensitivity 99-100%, specificity 92%)<sup>88,91</sup>.

Baseline impedance values reflect the permeability of the esophageal mucosa, both in animal models and healthy volunteers, with lower values found in erosive and non-erosive GERD<sup>92, 93</sup>. Low baseline esophageal mucosal impedance has been linked to alteration in intercellular space and tight junctions<sup>94, 95</sup> and to reflux symptoms<sup>96</sup>. Since frequent swallows and reflux events impact measurement, baseline impedance is best measured from pH-impedance tracings during sleep, termed mean nocturnal baseline impedance (MNBI) when averaged from three 10-minute periods spaced an hour apart<sup>85</sup>. MNBI is lower in persisting erosive esophagitis compared to healed esophagitis<sup>97</sup>, in PPI-responsive NERD and chronic cough compared to PPI-refractory states<sup>85, 98</sup>, and in erosive esophagitis, NERD and reflux hypersensitivity compared to functional heartburn and healthy controls<sup>87, 88, 94</sup>. Low MNBI (<2292 ohms) independently predicts response to antireflux therapy<sup>99</sup>, links PPI responsive heartburn to reflux better than AET<sup>91</sup>, and improves with healing esophagitis<sup>100, 101</sup>.

Baseline impedance can also be directly measured from the esophageal mucosa. Initially, this was done using probes with impedance sensors that were passed through the endoscope. Subsequently, the design has improved using two radial sensors mounted on a 10-cm balloon that is inflated to insure optimal contact with a long segment of esophageal mucosa. Baseline mucosal impedance values correlate with esophageal mucosal inflammation, differentiating erosive and non-erosive GERD from eosinophilic esophagitis and normal patients with better specificity (95% vs. 64%) and positive predictive value (96% vs. 40%) compared to pH monitoring.<sup>102-104</sup> Similar to baseline impedance from pH-impedance monitoring, values normalize following PPI therapy. Although normative values are not yet available, ongoing studies should clarify the role of mucosal impedance measurements in GERD management.

### **Esophageal high-resolution manometry**

A common indication for high-resolution manometry (HRM) is to accurately place pH or pH-impedance catheters. HRM is also used to assess peristalsis and to detect alternative major motor disorders prior to ARS or when symptoms do not improve with GERD therapy. Consequently, HRM studies are often performed in the setting of GERD. Nonetheless, although fundamental to GERD pathophysiology, a pathophysiologic classification of motor findings in GERD was only recently described<sup>7</sup>.

#### *EGJ Barrier Function*

The most fundamental abnormality in GERD is incompetence of the EGJ as an antireflux barrier, making quantifying EGJ competence an attractive biomarker. However, the EGJ is a complex sphincter composed of both the crural diaphragm (CD) and lower esophageal sphincter (LES), the relative dominance of which varies with circumstance. Not only does EGJ pressure vary with time, respiration and swallowing, but EGJ morphology can also vary over time, transitioning between superimposed and separated CD and LES elements<sup>105</sup>. Furthermore, some degree of EGJ incompetence is physiological, evident by the phenomenon of transient LES relaxation (TLESR), reflex relaxation of both the LES and CD that facilitates gas venting from the stomach<sup>106</sup>. Clearly, there are challenges to quantifying EGJ barrier function.

No single HRM metric adequately summarizes EGJ competence. Hence, the Lyon Consensus proposes adopting two metrics, one expressing the anatomical morphology of the EGJ and the second summarizing its contractile vigor. EGJ morphology, defined by the relationship between the LES and CD, has been characterized into three subtypes on HRM<sup>107, 108</sup>: type 1 with superimposed LES and CD, type 2 with axially separated LES and CD pressure signals separated by less than 3 cm, and type 3 with a  $\geq 3$  cm

separation between the LES and CD pressure signatures (Figure 1). Type 3 EGJ morphology is associated with reduced LES pressure and lower inspiratory augmentation, which correlates with reflux severity<sup>107, 109, 110</sup>. However, this relationship is not linear and there are clear exceptions.

The second HRM metric to quantify EGJ barrier function is the EGJ contractile integral (EGJ-CI) (Figure 2). The EGJ-CI is calculated using methodology analogous to that for calculating the distal contractile integral (DCI) with the DCI box set to encompass the LES and CD over a period of three respiratory cycles above a threshold of gastric pressure. The calculated “DCI” is then divided by the duration of the three respiratory cycles to make it independent of time and expressed in units of mmHg•cm<sup>111</sup>. Several groups of investigators have subsequently tested the performance of the EGJ-CI in segregating GERD populations (Table 1)<sup>111-115</sup>, with general agreement that this metric identifies a subset of patients with severe barrier dysfunction prone to either endoscopic esophagitis or unequivocally abnormal reflux testing. However, it is also evident from the spread of normal ranges reported among these studies that there are likely methodological discrepancies in exactly how the EGJ-CI is calculated. In view of this, the Lyon Consensus concluded that the EGJ-CI is a promising metric, but needs further research before widespread adoption. In an attempt to standardize methodology among groups, they recommended exclusion of CD component of the EGJ in instances of type 3 EGJ morphology and calculation of EGJ-CI above the gastric baseline pressure.

Further insight into the genesis of the EGJ-CI has been gleaned through studies using 3D-HRM<sup>116</sup>. Isolation of the CD component of the composite EGJ signal on 3D-HRM concluded that approximately 85% of overall EGJ contractility was attributable to the CD<sup>117</sup>. Further, analysis of 3D-HRM recordings differentiating the CD and LES constituents of the EGJ pressure complex demonstrated that the CD component correlated strongly with the EGJ-CI, suggesting that both are largely determined by CD contractility<sup>118</sup>. Together, these studies provide physiological support for adopting the EGJ-CI as a good summary metric of EGJ barrier function, albeit with the caveat that the metric is largely an indicator of CD contractility.

### *Esophageal Peristaltic Function*

Esophageal peristalsis can be characterized by the DCI which summarizes the vigor of post transition zone contraction (**Figure 2**)<sup>108</sup>. A DCI threshold of 450 mmHg•cm•s correlates with an averaged distal peristaltic amplitude of 30 mmHg<sup>119</sup>, the original manometric threshold defining ineffective swallows. When abnormal, esophageal peristalsis is often weak in GERD<sup>120-122</sup>, with poor and/or delayed formation of post transition zone contraction segment<sup>123</sup>. This can result in major breaks (>5 cm) in the peristaltic contour even when contraction vigor is preserved, a condition termed fragmented peristalsis when  $\geq 50\%$  of test swallows demonstrate this finding<sup>108</sup>.

Peristaltic dysfunction becomes progressively more common going from NERD to erosive esophagitis, to Barrett's esophagus<sup>124, 125</sup>. High proportions of ineffective contractions increase the likelihood of abnormal AET, particularly while supine<sup>126</sup>, and increase the likelihood of reflux symptoms<sup>127</sup>. The Chicago Classification defines ineffective esophageal motility (IEM) as  $\geq 50\%$  of test swallows with DCI<450 mmHg•cm•s, inclusive of any combination of weak (DCI 100-450 mmHg•cm•s) or failed (DCI<100 mmHg•cm•s) sequences<sup>108</sup>. Failed sequences are more predictive of an abnormal AET than a similar proportion of weak sequences<sup>128</sup>. The greatest reflux burden is seen with absent contractility (100% of test swallows with DCI<100 mmHg•cm•s)<sup>129</sup>.

### *Provocative Tests*

The physiological phenomenon of deglutitive inhibition is more pronounced with multiple swallows in rapid succession such that the esophagus remains in inhibition until after the final swallow, which is then followed by a peristaltic contraction<sup>130</sup>. Multiple rapid swallows (MRS) and rapid drink challenge (RDC) are two provocative tests of the integrity of deglutitive inhibition during HRM<sup>131</sup>. With

MRS, five 2-ml swallows are taken less than 4 seconds apart and with RDC 200 ml of water is swallowed within 30 seconds. The Lyon Consensus proposes that every HRM study should be accompanied by at least one of these provocative tests.

Post-MRS contractions are an indicator of ‘contraction reserve’ in the esophagus, the phenomenon wherein the post-MRS contraction has greater DCI than the preceding test swallows (Figure 2)<sup>132</sup>. Recent data suggests three MRS sequences for reliable assessment of contraction reserve<sup>133</sup>. The absence of contraction reserve in IEM is predictive of the poor efficacy of promotility drugs<sup>134</sup>, higher AET in NERD<sup>90</sup>, outlet obstruction and subsequent benefit from dilation following ARS<sup>135 136</sup>, and persistence or development of IEM after ARS<sup>137</sup>. Absent contraction reserve is also the most common manometric finding in systemic sclerosis.<sup>138</sup> The Lyon Consensus accepted the value of adopting MRS into HRM protocols for determining contraction reserve in IEM or absent contractility<sup>108</sup>, acknowledging that MRS is the most widely studied provocative test<sup>132, 136, 139</sup>, provides a computationally simple endpoint (peristaltic augmentation ratio: post-MRS versus pre-MRS)<sup>138</sup>, and is quick and easy to perform.

In contrast to MRS, the most important clinical application of RDC is in distinguishing EGJ obstruction from achalasia, by identifying LES relaxation in the former, and an exaggerated pressure gradient across a non-relaxed EGJ in the latter<sup>140 141, 142</sup>. Therefore, RDC is most helpful in detecting pan-esophageal pressurization in achalasia, identifying increased resistance to EGJ outflow, and uncovering latent hypercontractility<sup>140</sup>. Additionally, RDC may offer supportive evidence for erosive GERD; effective post-RDC peristalsis was seen in 83% of healthy controls compared to 70% of patients with NERD and only 30% of patients with erosive esophagitis<sup>143</sup>. Solid test meals have also been utilized as provocative tests during HRM, mainly in evaluating transit symptoms<sup>143-146</sup>. Normal values of esophageal pressure responses to RDC and solid meals have been recently reported in normal healthy volunteers<sup>146,</sup>



HRM studies performed during the post-prandial period could be of interest for identification of pathophysiological mechanisms in GERD<sup>148, 149</sup>, particularly reflux episodes that tend to be post-prandial. A reflux episode may occur during a TLESR, from low LES pressure, or in conjunction with rumination (increased gastric pressure with or without decreased thoracic pressure) or supragastric belching (air swallowing to initiate belching)<sup>148</sup>. Post-prandial HRM has also been used to evaluate the efficacy of drugs targeted to TLESRs, rumination, and supragastric belching<sup>150, 151</sup>. However, several limitations exist, including lack of normative postprandial HRM data, difficulties with standardization of the test meal, and unclear optimal duration of the recording period.

#### *Classification of Motility Findings in GERD*

The most common motility pattern in GERD is a normal study. However, either the EGJ or the esophageal body, or both can be abnormal. The EGJ can be hypotensive, with or without a hiatus hernia. Peristalsis can be fragmented, ineffective or absent, with or without contraction reserve. The Lyon Consensus endorses the hierarchical classification of motility findings in GERD first evaluating EGJ morphology and function with LES-CD separation and the EGJ-CI, secondly characterizing the integrity of peristalsis as normal, weak, fragmented or absent, and third, evaluating for contraction reserve<sup>7</sup> (Table 2). This classification is intended to be used in conjunction with the Chicago Classification.

#### ADVANCES SINCE THE PORTO CONSENSUS

The Lyon Consensus builds on the Porto Consensus of 2002<sup>46</sup>, providing recommendations for the use and interpretation of reflux testing techniques in 2017 including esophageal HRM and baseline impedance measurement that were not widely available in 2002 (Table 3). The primary indication for reflux testing is in distinguishing among patients with pathologic reflux burden, reflux mediated hypersensitivity, and functional syndromes (Table 4)<sup>152</sup>. The Lyon Consensus attempts to augment this

approach by stratifying the significance of findings into those that are conclusive of pathologic GERD, as opposed to suggestive of the diagnosis (Figure 3). The Lyon Consensus also proposes the concept of ‘borderline’ or inconclusive evidence when additional evidence can sway the final judgment toward or away from GERD. This is an area where novel metrics and diagnostic techniques may prove helpful. Conditions that can mimic GERD, such as achalasia, supragastric belching and rumination syndrome need to be excluded with appropriate testing.

### *Optimization of GERD Testing*

GERD symptoms are diverse, response to treatment is variable, pathogenesis is heterogeneous, and mechanistic phenotypes are heavily influenced by hypersensitivity and hypervigilance. Because simple algorithms starting with a PPI trial do not consider these complex phenotypes of GERD, they often lead to inappropriate PPI utilization, delayed diagnosis, and inaccurate diagnoses<sup>153</sup>. The Lyon Consensus opines that the optimal initial testing for PPI non-responders with no prior endoscopic or pH-metry demonstration of GERD is pH- or pH-impedance monitoring done withholding antisecretory therapy. A key potential outcome of that testing is to rule out GERD and to redirect management toward weaning off PPIs, using neuromodulators and/or cognitive behavioral therapy as appropriate. In contrast, optimal testing in poorly responsive patients with a prior demonstration of GERD is the combination of EGD, HRM, and pH-impedance monitoring done on twice-daily PPI therapy. This combination of tests serves both to redirect therapy toward alternative diagnoses and to mechanistically subtype patients in terms of poor clearance, excessive reflux episodes and hypersensitivity (Table 4), each of which can trigger specific management options. The precise roles of baseline impedance, PSPW index and provocative maneuvers on HRM remain to be clarified with future research.

### *Outcome Measures in GERD*

The optimal use of diagnostic testing may translate into better therapeutic outcomes, but appropriate outcome measures are necessary to properly evaluate that improvement. Esophagitis healing is a common measure for therapeutic trials, but visible esophagitis is rare in patients with refractory GERD symptoms<sup>25</sup>, and the objective of the evaluation is to determine if refractory symptoms are attributable to GERD or not. Hence, potentially relevant outcome measures are symptom description<sup>154</sup>, symptom questionnaires<sup>155</sup>, pictograms with visual depictions of symptoms<sup>156, 157</sup>, and ambulatory reflux monitoring with analysis of reflux-symptom association<sup>75, 158</sup>. Outcome measures typically utilized in GERD therapeutic trials have included individual symptom assessment with Likert or visual analog scales<sup>159</sup>, global outcome evaluations on Likert scales<sup>160</sup>, adequate vs. inadequate relief, disease-specific questionnaires<sup>161</sup>, and quality of life questionnaires. Looking to the future, regulatory agencies have stipulated that validated patient reported outcome questionnaires (PROs) and quality of life questionnaires will be requisite in future therapeutic trials. Such PROs require a fastidious approach that includes item generation, testing for reliability, responsiveness, validity and interpretability, and finally, cross-cultural adaptation when applicable, for each diagnostic category<sup>162</sup>.

### CONCLUSIONS AND FUTURE DIRECTION

GERD is a complex disease with a heterogeneous symptom profile and a multifaceted pathogenic basis that defies a simple diagnostic algorithm or categorical classification. The Lyon Consensus defines parameters on esophageal testing that conclusively establish the presence of GERD and characteristics that rule out GERD. Additional evidence from reflux-symptom association, motor findings on HRM, novel metrics from pH-impedance monitoring, baseline mucosal impedance and PPI response complement esophageal testing when pH-metry is borderline or inconclusive. While acknowledging the

limitations of currently available esophageal testing in GERD, the Lyon Consensus proposes this model as a guide to direct management.

The future approach to phenotyping GERD patients should focus on assessing important physiologic biomarkers and PROs to categorize patients based on the severity of refluxate exposure, mechanism of reflux, effectors of clearance, and underlying EGJ pathophysiology (Table 4), while recognizing that no single approach is perfect. Novel metrics assessing tissue resistance, esophageal clearance, peripheral and central neural integration, and psychometrics will allow for a tailored therapeutic approach including pharmacologic treatments, surgical/endoscopic interventions, and behavioral strategies targeting the underlying defect(s) in the antireflux barrier, esophageal clearance, visceral sensitivity, and cognitive response to reflux. As newer metrics emerge, the Lyon Consensus plans future meetings to update and adapt the Consensus conclusions. Collaboration between high volume medical centers involved in GERD testing has opened possibilities for more robust normative data and for validation of conclusions and recommendations from the Lyon Consensus. As the GERD diagnostic paradigm evolves, utilizing diagnostic testing to define a precision approach tailored to the individual patient becomes possible. The goals of evaluation should therefore transition toward defining GERD phenotypes to facilitate tailored treatment.

## ACKNOWLEDGEMENT

The authors wish to acknowledge the International Working Group for Gastrointestinal Motility and Function for initiating the consensus meetings and for providing material support for the consensus process.

## REFERENCES

1. El-Serag HB, Sweet S, Winchester CC, et al. Update on the epidemiology of gastro-oesophageal reflux disease: a systematic review. *Gut* 2014;63:871-80.
2. Shaheen NJ, Hansen RA, Morgan DR, et al. The burden of gastrointestinal and liver diseases, 2006. *Am J Gastroenterol* 2006;101:2128-38.
3. Vakil N, van Zanten SV, Kahrilas P, et al. The Montreal definition and classification of gastroesophageal reflux disease: a global evidence-based consensus. *Am J Gastroenterol* 2006;101:1900-20; quiz 1943.
4. Katz PO, Gerson LB, Vela MF. Guidelines for the diagnosis and management of gastroesophageal reflux disease. *Am J Gastroenterol* 2013;108:308-28; quiz 329.
5. Kahrilas PJ, Boeckxstaens G, Smout AJ. Management of the patient with incomplete response to PPI therapy. *Best Pract Res Clin Gastroenterol* 2013;27:401-14.
6. Roman S, Gyawali CP, Savarino E, et al. Ambulatory reflux monitoring for diagnosis of gastro-oesophageal reflux disease: Update of the Porto consensus and recommendations from an international consensus group. *Neurogastroenterol Motil* 2017;29:1-15.
7. Gyawali CP, Roman S, Bredenoord AJ, et al. Classification of esophageal motor findings in gastro-oesophageal reflux disease: Conclusions from an international consensus group. *Neurogastroenterol Motil* 2017;29:e13104.
8. Savarino E, Bredenoord AJ, Fox M, et al. Expert consensus document: Advances in the physiological assessment and diagnosis of GERD. *Nat Rev Gastroenterol Hepatol* 2017;14:665-676.
9. Demeester TR, Johnson LF, Joseph GJ, et al. Patterns of gastroesophageal reflux in health and disease. *Ann Surg* 1976;184:459-70.
10. Lundell LR, Dent J, Bennett JR, et al. Endoscopic assessment of oesophagitis: clinical and functional correlates and further validation of the Los Angeles classification. *Gut* 1999;45:172-80.
11. Mainie I, Tutuian R, Shay S, et al. Acid and non-acid reflux in patients with persistent symptoms despite acid suppressive therapy: a multicentre study using combined ambulatory impedance-pH monitoring. *Gut* 2006;55:1398-402.
12. Dent J, Vakil N, Jones R, et al. Accuracy of the diagnosis of GORD by questionnaire, physicians and a trial of proton pump inhibitor treatment: the Diamond Study. *Gut* 2010;59:714-21.
13. Jones R, Junghard O, Dent J, et al. Development of the GerdQ, a tool for the diagnosis and management of gastro-oesophageal reflux disease in primary care. *Aliment Pharmacol Ther* 2009;30:1030-8.
14. Bolier EA, Kessing BF, Smout AJ, et al. Systematic review: questionnaires for assessment of gastroesophageal reflux disease. *Dis Esophagus* 2015;28:105-20.
15. Bytzer P, Jones R, Vakil N, et al. Limited ability of the proton-pump inhibitor test to identify patients with gastroesophageal reflux disease. *Clin Gastroenterol Hepatol* 2012;10:1360-6.
16. Sifrim D, Zerbib F. Diagnosis and management of patients with reflux symptoms refractory to proton pump inhibitors. *Gut* 2012;61:1340-54.
17. Gyawali CP, Fass R. Management of Gastroesophageal Reflux Disease. *Gastroenterology* 2018;(in press).
18. Roman S, Keefer L, Imam H, et al. Majority of symptoms in esophageal reflux PPI non-responders are not related to reflux. *Neurogastroenterol Motil* 2015;27:1667-74.
19. Weijenborg PW, Smout AJ, Verseijden C, et al. Hypersensitivity to acid is associated with impaired esophageal mucosal integrity in patients with gastroesophageal reflux disease with and without esophagitis. *Am J Physiol Gastrointest Liver Physiol* 2014;307:G323-9.
20. de Leone A, Tonini M, Dominici P, et al. The proton pump inhibitor test for gastroesophageal reflux disease: optimal cut-off value and duration. *Dig Liver Dis* 2010;42:785-90.

21. Numans ME, Lau J, de Wit NJ, et al. Short-term treatment with proton-pump inhibitors as a test for gastroesophageal reflux disease: a meta-analysis of diagnostic test characteristics. *Ann Intern Med* 2004;140:518-27.
22. Jonasson C, Moum B, Bang C, et al. Randomised clinical trial: a comparison between a GerdQ-based algorithm and an endoscopy-based approach for the diagnosis and initial treatment of GERD. *Aliment Pharmacol Ther* 2012;35:1290-300.
23. Savarino E, Bredenoord AJ, Fox M, et al. Expert consensus document: Advances in the physiological assessment and diagnosis of GERD. *Nat Rev Gastroenterol Hepatol* 2017;14:665-676.
24. Savarino E, Zentilin P, Savarino V. NERD: an umbrella term including heterogeneous subpopulations. *Nat Rev Gastroenterol Hepatol* 2013;10:371-80.
25. Poh CH, Gasiorowska A, Navarro-Rodriguez T, et al. Upper GI tract findings in patients with heartburn in whom proton pump inhibitor treatment failed versus those not receiving antireflux treatment. *Gastrointest Endosc* 2010;71:28-34.
26. Akdamar K, Ertan A, Agrawal NM, et al. Upper gastrointestinal endoscopy in normal asymptomatic volunteers. *Gastrointest Endosc* 1986;32:78-80.
27. Takashima T, Iwakiri R, Sakata Y, et al. Endoscopic reflux esophagitis and *Helicobacter pylori* infection in young healthy Japanese volunteers. *Digestion* 2012;86:55-8.
28. Zagari RM, Fuccio L, Wallander MA, et al. Gastro-oesophageal reflux symptoms, oesophagitis and Barrett's oesophagus in the general population: the Loiano-Monghidoro study. *Gut* 2008;57:1354-9.
29. Jobe BA, Richter JE, Hoppo T, et al. Preoperative diagnostic workup before antireflux surgery: an evidence and experience-based consensus of the Esophageal Diagnostic Advisory Panel. *J Am Coll Surg* 2013;217:586-97.
30. Rex DK, Cummings OW, Shaw M, et al. Screening for Barrett's esophagus in colonoscopy patients with and without heartburn. *Gastroenterology* 2003;125:1670-7.
31. Sharma P. Review article: prevalence of Barrett's oesophagus and metaplasia at the gastro-oesophageal junction. *Aliment Pharmacol Ther* 2004;20 Suppl 5:48-54; discussion 61-2.
32. Johansson J, Hakansson HO, Mellblom L, et al. Prevalence of precancerous and other metaplasia in the distal oesophagus and gastro-oesophageal junction. *Scand J Gastroenterol* 2005;40:893-902.
33. Aziz Q, Fass R, Gyawali CP, et al. Functional Esophageal Disorders. *Gastroenterology* 2016;150:1368-79.
34. Savarino E, Zentilin P, Mastracci L, et al. Microscopic esophagitis distinguishes patients with non-erosive reflux disease from those with functional heartburn. *J Gastroenterol* 2013;48:473-82.
35. Kandulski A, Jechorek D, Caro C, et al. Histomorphological differentiation of non-erosive reflux disease and functional heartburn in patients with PPI-refractory heartburn. *Aliment Pharmacol Ther* 2013;38:643-51.
36. Calabrese C, Bortolotti M, Fabbri A, et al. Reversibility of GERD ultrastructural alterations and relief of symptoms after omeprazole treatment. *Am J Gastroenterol* 2005;100:537-42.
37. Vela MF, Craft BM, Sharma N, et al. Refractory heartburn: comparison of intercellular space diameter in documented GERD vs. functional heartburn. *Am J Gastroenterol* 2011;106:844-50.
38. van Malenstein H, Farre R, Sifrim D. Esophageal dilated intercellular spaces (DIS) and nonerosive reflux disease. *Am J Gastroenterol* 2008;103:1021-8.
39. Zentilin P, Savarino V, Mastracci L, et al. Reassessment of the diagnostic value of histology in patients with GERD, using multiple biopsy sites and an appropriate control group. *Am J Gastroenterol* 2005;100:2299-306.
40. Caviglia R, Ribolsi M, Gentile M, et al. Dilated intercellular spaces and acid reflux at the distal and proximal oesophagus in patients with non-erosive gastro-oesophageal reflux disease. *Aliment Pharmacol Ther* 2007;25:629-36.

41. Sweis R, Fox M, Anggiansah A, et al. Prolonged, wireless pH-studies have a high diagnostic yield in patients with reflux symptoms and negative 24-h catheter-based pH-studies. *Neurogastroenterol Motil* 2011;23:419-26.
42. Prakash C, Clouse RE. Value of extended recording time with wireless pH monitoring in evaluating gastroesophageal reflux disease. *Clin Gastroenterol Hepatol* 2005;3:329-34.
43. Ayazi S, Lipham JC, Portale G, et al. Bravo catheter-free pH monitoring: normal values, concordance, optimal diagnostic thresholds, and accuracy. *Clin Gastroenterol Hepatol* 2009;7:60-7.
44. Scarpulla G, Camilleri S, Galante P, et al. The impact of prolonged pH measurements on the diagnosis of gastroesophageal reflux disease: 4-day wireless pH studies. *Am J Gastroenterol* 2007;102:2642-7.
45. Penagini R, Sweis R, Mauro A, et al. Inconsistency in the Diagnosis of Functional Heartburn: Usefulness of Prolonged Wireless pH Monitoring in Patients With Proton Pump Inhibitor Refractory Gastroesophageal Reflux Disease. *J Neurogastroenterol Motil* 2015;21:265-72.
46. Sifrim D, Castell D, Dent J, et al. Gastro-oesophageal reflux monitoring: review and consensus report on detection and definitions of acid, non-acid, and gas reflux. *Gut* 2004;53:1024-31.
47. Savarino E, Marabotto E, Zentilin P, et al. The added value of impedance-pH monitoring to Rome III criteria in distinguishing functional heartburn from non-erosive reflux disease. *Dig Liver Dis* 2011;43:542-7.
48. Zerbib F, Roman S, Ropert A, et al. Esophageal pH-impedance monitoring and symptom analysis in GERD: a study in patients off and on therapy. *Am J Gastroenterol* 2006;101:1956-63.
49. Charbel S, Khandwala F, Vaezi MF. The role of esophageal pH monitoring in symptomatic patients on PPI therapy. *Am J Gastroenterol* 2005;100:283-9.
50. Vela MF, Camacho-Lobato L, Srinivasan R, et al. Simultaneous intraesophageal impedance and pH measurement of acid and nonacid gastroesophageal reflux: effect of omeprazole. *Gastroenterology* 2001;120:1599-606.
51. Vaezi MF, Schroeder PL, Richter JE. Reproducibility of proximal probe pH parameters in 24-hour ambulatory esophageal pH monitoring. *Am J Gastroenterol* 1997;92:825-9.
52. McCollough M, Jabbar A, Cacchione R, et al. Proximal sensor data from routine dual-sensor esophageal pH monitoring is often inaccurate. *Dig Dis Sci* 2004;49:1607-11.
53. Williams RB, Ali GN, Wallace KL, et al. Esophagopharyngeal acid regurgitation: dual pH monitoring criteria for its detection and insights into mechanisms. *Gastroenterology* 1999;117:1051-61.
54. Zerbib F, Roman S, Bruley Des Varannes S, et al. Normal values of pharyngeal and esophageal 24-hour pH impedance in individuals on and off therapy and interobserver reproducibility. *Clin Gastroenterol Hepatol* 2013;11:366-72.
55. Ayazi S, Lipham JC, Hagen JA, et al. A new technique for measurement of pharyngeal pH: normal values and discriminating pH threshold. *J Gastrointest Surg* 2009;13:1422-9.
56. Hayat JO, Yazaki E, Moore AT, et al. Objective detection of esophagopharyngeal reflux in patients with hoarseness and endoscopic signs of laryngeal inflammation. *J Clin Gastroenterol* 2014;48:318-27.
57. Ummarino D, Vandermeulen L, Roosens B, et al. Gastroesophageal reflux evaluation in patients affected by chronic cough: Restech versus multichannel intraluminal impedance/pH metry. *Laryngoscope* 2013;123:980-4.
58. Wilhelm D, Jell A, Feussner H, et al. Pharyngeal pH monitoring in gastrectomy patients - what do we really measure? *United European Gastroenterol J* 2016;4:541-5.
59. Wiener GJ, Morgan TM, Copper JB, et al. Ambulatory 24-hour esophageal pH monitoring. Reproducibility and variability of pH parameters. *Dig Dis Sci* 1988;33:1127-33.
60. Patel A, Sayuk GS, Gyawali CP. Parameters on Esophageal pH-Impedance Monitoring That Predict Outcomes of Patients With Gastroesophageal Reflux Disease. *Clin Gastroenterol Hepatol* 2015;13:884-91.



61. Patel A, Sayuk GS, Gyawali CP. Prevalence, characteristics, and treatment outcomes of reflux hypersensitivity detected on pH-impedance monitoring. *Neurogastroenterol Motil* 2016;28:1382-90.
62. Roman S, Bruley des Varannes S, Poudoux P, et al. Ambulatory 24-h oesophageal impedance-pH recordings: reliability of automatic analysis for gastro-oesophageal reflux assessment. *Neurogastroenterol Motil* 2006;18:978-86.
63. Bell R, Lipham JC, Louie BE, et al. Randomized Controlled Trial of Magnetic Sphincter Augmentation (MSA) VS. Omeprazole in Gerd Patients with Regurgitation: Initial Results from the Caliber Trial. *Gastroenterology* 2017;152:S-1309.
64. Bredenoord AJ, Weusten BL, Smout AJ. Symptom association analysis in ambulatory gastro-oesophageal reflux monitoring. *Gut* 2005;54:1810-7.
65. Lam HG, Breumelhof R, Roelofs JM, et al. What is the optimal time window in symptom analysis of 24-hour esophageal pressure and pH data? *Dig Dis Sci* 1994;39:402-9.
66. Herregods TVK, Pauwels A, Tack J, et al. Reflux-cough syndrome: Assessment of temporal association between reflux episodes and cough bursts. *Neurogastroenterol Motil* 2017; e13129.
67. Wiener GJ, Richter JE, Copper JB, et al. The symptom index: a clinically important parameter of ambulatory 24-hour esophageal pH monitoring. *Am J Gastroenterol* 1988;83:358-61.
68. Singh S, Richter JE, Bradley LA, et al. The symptom index. Differential usefulness in suspected acid-related complaints of heartburn and chest pain. *Dig Dis Sci* 1993;38:1402-8.
69. Weusten BL, Roelofs JM, Akkermans LM, et al. The symptom-association probability: an improved method for symptom analysis of 24-hour esophageal pH data. *Gastroenterology* 1994;107:1741-5.
70. Ghillebert G, Janssens J, Vantrappen G, et al. Ambulatory 24 hour intraoesophageal pH and pressure recordings v provocation tests in the diagnosis of chest pain of oesophageal origin. *Gut* 1990;31:738-44.
71. Kushnir VM, Sathyamurthy A, Drapekin J, et al. Assessment of concordance of symptom reflux association tests in ambulatory pH monitoring. *Aliment Pharmacol Ther* 2012;35:1080-7.
72. Kushnir VM, Sayuk GS, Gyawali CP. Abnormal GERD parameters on ambulatory pH monitoring predict therapeutic success in noncardiac chest pain. *Am J Gastroenterol* 2010;105:1032-8.
73. Patel A, Sayuk GS, Gyawali CP. Acid-based parameters on pH-impedance testing predict symptom improvement with medical management better than impedance parameters. *Am J Gastroenterol* 2014;109:836-44.
74. Watson RG, Tham TC, Johnston BT, et al. Double blind cross-over placebo controlled study of omeprazole in the treatment of patients with reflux symptoms and physiological levels of acid reflux--the "sensitive oesophagus". *Gut* 1997;40:587-90.
75. Taghavi SA, Ghasedi M, Saberi-Firoozi M, et al. Symptom association probability and symptom sensitivity index: preferable but still suboptimal predictors of response to high dose omeprazole. *Gut* 2005;54:1067-71.
76. Aanen MC, Weusten BL, Numans ME, et al. Effect of proton-pump inhibitor treatment on symptoms and quality of life in GERD patients depends on the symptom-reflux association. *J Clin Gastroenterol* 2008;42:441-7.
77. Rosen R, Amirault J, Heinz N, et al. The sensitivity of acoustic cough recording relative to intraesophageal pressure recording and patient report during reflux testing. *Neurogastroenterol Motil* 2014;26:1635-41.
78. Xiao Y, Carson D, Boris L, et al. The acoustic cough monitoring and manometric profile of cough and throat clearing. *Dis Esophagus* 2014;27:5-12.
79. Bredenoord AJ, Weusten BL, Timmer R, et al. Addition of esophageal impedance monitoring to pH monitoring increases the yield of symptom association analysis in patients off PPI therapy. *Am J Gastroenterol* 2006;101:453-9.

80. Tenca A, Campagnola P, Bravi I, et al. Impedance pH Monitoring: Intra-observer and Inter-observer Agreement and Usefulness of a Rapid Analysis of Symptom Reflux Association. *J Neurogastroenterol Motil* 2014;20:205-11.
81. Hemmink GJ, Bredenoord AJ, Weusten BL, et al. Esophageal pH-impedance monitoring in patients with therapy-resistant reflux symptoms: 'on' or 'off' proton pump inhibitor? *Am J Gastroenterol* 2008;103:2446-53.
82. Aanen MC, Bredenoord AJ, Numans ME, et al. Reproducibility of symptom association analysis in ambulatory reflux monitoring. *Am J Gastroenterol* 2008;103:2200-8.
83. Slaughter JC, Goutte M, Rymer JA, et al. Caution about overinterpretation of symptom indexes in reflux monitoring for refractory gastroesophageal reflux disease. *Clin Gastroenterol Hepatol* 2011;9:868-74.
84. Barriga-Rivera A, Elena M, Moya MJ, et al. The binomial symptom index: toward an optimal method for the evaluation of symptom association in gastroesophageal reflux. *Neurogastroenterol Motil* 2013;25:664-9.
85. Martinucci I, de Bortoli N, Savarino E, et al. Esophageal baseline impedance levels in patients with pathophysiological characteristics of functional heartburn. *Neurogastroenterol Motil* 2014;26:546-55.
86. Frazzoni M, Bertani H, Manta R, et al. Impairment of chemical clearance is relevant to the pathogenesis of refractory reflux oesophagitis. *Dig Liver Dis* 2014;46:596-602.
87. de Bortoli N, Martinucci I, Savarino E, et al. Association between baseline impedance values and response proton pump inhibitors in patients with heartburn. *Clin Gastroenterol Hepatol* 2015;13:1082-8 e1.
88. Frazzoni M, Savarino E, de Bortoli N, et al. Analyses of the Post-reflux Swallow-induced Peristaltic Wave Index and Nocturnal Baseline Impedance Parameters Increase the Diagnostic Yield of Impedance-pH Monitoring of Patients With Reflux Disease. *Clin Gastroenterol Hepatol* 2016;14:40-6.
89. Frazzoni M, de Bortoli N, Frazzoni L, et al. Impairment of chemical clearance and mucosal integrity distinguishes hypersensitive esophagus from functional heartburn. *J Gastroenterol* 2017;52:444-451.
90. Martinucci I, Savarino EV, Pandolfino JE, et al. Vigor of peristalsis during multiple rapid swallows is inversely correlated with acid exposure time in patients with NERD. *Neurogastroenterol Motil* 2016;28:243-50.
91. Frazzoni L, Frazzoni M, de Bortoli N, et al. Postreflux swallow-induced peristaltic wave index and nocturnal baseline impedance can link PPI-responsive heartburn to reflux better than acid exposure time. *Neurogastroenterol Motil* 2017;29:e13116.
92. Farre R, Blondeau K, Clement D, et al. Evaluation of oesophageal mucosa integrity by the intraluminal impedance technique. *Gut* 2011;60:885-92.
93. Kessing BF, Bredenoord AJ, Weijenberg PW, et al. Esophageal acid exposure decreases intraluminal baseline impedance levels. *Am J Gastroenterol* 2011;106:2093-7.
94. Kandulski A, Weigt J, Caro C, et al. Esophageal intraluminal baseline impedance differentiates gastroesophageal reflux disease from functional heartburn. *Clin Gastroenterol Hepatol* 2015;13:1075-81.
95. Zhong C, Duan L, Wang K, et al. Esophageal intraluminal baseline impedance is associated with severity of acid reflux and epithelial structural abnormalities in patients with gastroesophageal reflux disease. *J Gastroenterol* 2013;48:601-10.
96. Woodland P, Al-Zinaty M, Yazaki E, et al. In vivo evaluation of acid-induced changes in oesophageal mucosa integrity and sensitivity in non-erosive reflux disease. *Gut* 2013;62:1256-61.
97. Frazzoni M, de Bortoli N, Frazzoni L, et al. The added diagnostic value of postreflux swallow-induced peristaltic wave index and nocturnal baseline impedance in refractory reflux disease studied with on-therapy impedance-pH monitoring. *Neurogastroenterol Motil* 2017;29:e12947.

98. Ribolsi M, Savarino E, De Bortoli N, et al. Reflux pattern and role of impedance-pH variables in predicting PPI response in patients with suspected GERD-related chronic cough. *Aliment Pharmacol Ther* 2014;40:966-73.
99. Patel A, Wang D, Sainani N, et al. Distal mean nocturnal baseline impedance on pH-impedance monitoring predicts reflux burden and symptomatic outcome in gastro-oesophageal reflux disease. *Aliment Pharmacol Ther* 2016;44:890-8.
100. van Rhijn BD, Weijnenborg PW, Verheij J, et al. Proton pump inhibitors partially restore mucosal integrity in patients with proton pump inhibitor-responsive esophageal eosinophilia but not eosinophilic esophagitis. *Clin Gastroenterol Hepatol* 2014;12:1815-23 e2.
101. Rinsma NF, Farre R, Bouvy ND, et al. The effect of endoscopic fundoplication and proton pump inhibitors on baseline impedance and heartburn severity in GERD patients. *Neurogastroenterol Motil* 2015;27:220-8.
102. Saritas Yuksel E, Higginbotham T, Slaughter JC, et al. Use of direct, endoscopic-guided measurements of mucosal impedance in diagnosis of gastroesophageal reflux disease. *Clin Gastroenterol Hepatol* 2012;10:1110-6.
103. Katzka DA, Ravi K, Geno DM, et al. Endoscopic Mucosal Impedance Measurements Correlate With Eosinophilia and Dilation of Intercellular Spaces in Patients With Eosinophilic Esophagitis. *Clin Gastroenterol Hepatol* 2015;13:1242-1248 e1.
104. Ates F, Yuksel ES, Higginbotham T, et al. Mucosal impedance discriminates GERD from non-GERD conditions. *Gastroenterology* 2015;148:334-43.
105. Bredenoord AJ, Weusten BL, Timmer R, et al. Intermittent spatial separation of diaphragm and lower esophageal sphincter favors acidic and weakly acidic reflux. *Gastroenterology* 2006;130:334-40.
106. Roman S, Holloway R, Keller J, et al. Validation of criteria for the definition of transient lower esophageal sphincter relaxations using high-resolution manometry. *Neurogastroenterol Motil* 2017;29:e12920.
107. Pandolfino JE, Kim H, Ghosh SK, et al. High-resolution manometry of the EGJ: an analysis of crural diaphragm function in GERD. *Am J Gastroenterol* 2007;102:1056-63.
108. Kahrilas PJ, Bredenoord AJ, Fox M, et al. The Chicago Classification of esophageal motility disorders, v3.0. *Neurogastroenterol Motil* 2015;27:160-74.
109. Ham H, Cho YK, Lee HH, et al. Esophagogastric junction contractile integral and morphology: Two high-resolution manometry metrics of the anti-reflux barrier. *J Gastroenterol Hepatol* 2017;32:1443-1449.
110. Tolone S, de Cassan C, de Bortoli N, et al. Esophagogastric junction morphology is associated with a positive impedance-pH monitoring in patients with GERD. *Neurogastroenterol Motil* 2015;27:1175-82.
111. Nicodeme F, Pipa-Muniz M, Khanna K, et al. Quantifying esophagogastric junction contractility with a novel HRM topographic metric, the EGJ-Contractile Integral: normative values and preliminary evaluation in PPI non-responders. *Neurogastroenterol Motil* 2014;26:353-60.
112. Tolone S, De Bortoli N, Marabotto E, et al. Esophagogastric junction contractility for clinical assessment in patients with GERD: a real added value? *Neurogastroenterol Motil* 2015;27:1423-31.
113. Jasper D, Freitas-Queiroz N, Hollenstein M, et al. Prolonged measurement improves the assessment of the barrier function of the esophago-gastric junction by high-resolution manometry. *Neurogastroenterol Motil* 2016;29:e12925.
114. Wang D, Patel A, Mello M, et al. Esophagogastric junction contractile integral (EGJ-CI) quantifies changes in EGJ barrier function with surgical intervention. *Neurogastroenterol Motil* 2016;28:639-46.
115. Xie C, Wang J, Li Y, et al. Esophagogastric junction contractility integral reflected the anti-reflux barrier dysfunction in GERD patients. *J Neurogastroenterol Motil* 2017;23:27-33.

116. Kwiatek MA, Pandolfino JE, Kahrilas PJ. 3D-high resolution manometry of the esophagogastric junction. *Neurogastroenterol Motil* 2011;23:e461-9.
117. Nicodeme F, Soper NJ, Lin Z, et al. Calculation of esophagogastric junction vector volume using three-dimensional high-resolution manometry. *Dis Esophagus* 2015;28:684-90.
118. Lin Z, Xiao Y, Li Y, et al. Novel 3D high-resolution manometry metrics for quantifying esophagogastric junction contractility. *Neurogastroenterol Motil* 2017;29:e13054.
119. Xiao Y, Kahrilas PJ, Kwasny MJ, et al. High-resolution manometry correlates of ineffective esophageal motility. *Am J Gastroenterol* 2012;107:1647-54.
120. Ho SC, Chang CS, Wu CY, et al. Ineffective esophageal motility is a primary motility disorder in gastroesophageal reflux disease. *Dig Dis Sci* 2002;47:652-6.
121. Diener U, Patti MG, Molena D, et al. Esophageal dysmotility and gastroesophageal reflux disease. *J Gastrointest Surg* 2001;5:260-5.
122. Chan WW, Haroian LR, Gyawali CP. Value of preoperative esophageal function studies before laparoscopic antireflux surgery. *Surg Endosc* 2011;25:2943-9.
123. Kumar N, Porter RF, Chanin JM, et al. Analysis of intersegmental trough and proximal latency of smooth muscle contraction using high-resolution esophageal manometry. *J Clin Gastroenterol* 2012;46:375-81.
124. Meneghetti AT, Tedesco P, Damani T, et al. Esophageal mucosal damage may promote dysmotility and worsen esophageal acid exposure. *J Gastrointest Surg* 2005;9:1313-7.
125. Savarino E, Gemignani L, Pohl D, et al. Oesophageal motility and bolus transit abnormalities increase in parallel with the severity of gastro-oesophageal reflux disease. *Aliment Pharmacol Ther* 2011;34:476-86.
126. Fornari F, Blondeau K, Durand L, et al. Relevance of mild ineffective oesophageal motility (IOM) and potential pharmacological reversibility of severe IOM in patients with gastro-oesophageal reflux disease. *Aliment Pharmacol Ther* 2007;26:1345-54.
127. Blonski W, Vela M, Safder A, et al. Revised criterion for diagnosis of ineffective esophageal motility is associated with more frequent dysphagia and greater bolus transit abnormalities. *Am J Gastroenterol* 2008;103:699-704.
128. Rengarajan A, Bolkhir A, Gor P, et al. Esophagogastric junction and esophageal body contraction metrics on high resolution manometry predict esophageal acid burden. *Neurogastroenterol Motil* 2018;(in press).
129. Reddy CA, Patel A, Gyawali CP. Impact of Symptom Burden and Health-Related Quality of Life (HRQOL) on Esophageal Motor Disorders. *Neurogastroenterol Motil* 2017:e12970.
130. Sifrim D, Jafari J. Deglutitive inhibition, latency between swallow and esophageal contractions and primary esophageal motor disorders. *J Neurogastroenterol Motil* 2012;18:6-12.
131. Savarino E, de Bortoli N, Bellini M, et al. Practice guidelines on the use of esophageal manometry - A GISMAD-SIGE-AIGO medical position statement. *Dig Liver Dis* 2016;48:1124-35.
132. Shaker A, Stoikes N, Drapekin J, et al. Multiple rapid swallow responses during esophageal high-resolution manometry reflect esophageal body peristaltic reserve. *Am J Gastroenterol* 2013;108:1706-12.
133. Mauro A, Savarino E, De Bortoli N, et al. Optimal number of multiple rapid swallows needed during high-resolution esophageal manometry for accurate prediction of contraction reserve. *Neurogastroenterol Motil* 2018;(in press).
134. Jafari J, Yazaki E, Woodland P, et al. Effect of Azithromycin on Esophageal Hypomotility (EH) and Prediction of Response by Esophageal Stimulations Tests During High Resolution Manometry. *Gastroenterology* 2015;148:S-75.
135. Stoikes N, Drapekin J, Kushnir V, et al. The value of multiple rapid swallows during preoperative esophageal manometry before laparoscopic antireflux surgery. *Surg Endosc* 2012;26:3401-7.

136. Wang YT, Tai LF, Yazaki E, et al. Investigation of Dysphagia After Antireflux Surgery by High-resolution Manometry: Impact of Multiple Water Swallows and a Solid Test Meal on Diagnosis, Management, and Clinical Outcome. *Clin Gastroenterol Hepatol* 2015;13:1575-83.
137. Mello MD, Shriver AR, Li Y, et al. Ineffective esophageal motility phenotypes following fundoplication in gastroesophageal reflux disease. *Neurogastroenterology & Motility* 2016;28:292-298.
138. Carlson DA, Crowell MD, Kimmel JN, et al. Loss of Peristaltic Reserve, Determined by Multiple Rapid Swallows, Is the Most Frequent Esophageal Motility Abnormality in Patients With Systemic Sclerosis. *Clin Gastroenterol Hepatol* 2016; 14:1502-6.
139. Price LH, Li Y, Patel A, et al. Reproducibility patterns of multiple rapid swallows during high resolution esophageal manometry provide insights into esophageal pathophysiology. *Neurogastroenterol Motil* 2014;26:646-53.
140. Marin I, Serra J. Patterns of esophageal pressure responses to a rapid drink challenge test in patients with esophageal motility disorders. *Neurogastroenterol Motil* 2016;28:543-53.
141. Ang D, Hollenstein M, Misselwitz B, et al. Rapid Drink Challenge in high-resolution manometry: an adjunctive test for detection of esophageal motility disorders. *Neurogastroenterol Motil* 2017;29: e12902.
142. Elvevi A, Mauro A, Pugliese D, et al. Usefulness of low- and high-volume multiple rapid swallowing during high-resolution manometry. *Digestive and Liver Disease*;47:103-107.
143. Daum C, Sweis R, Kaufman E, et al. Failure to respond to physiologic challenge characterizes esophageal motility in erosive gastro-esophageal reflux disease. *Neurogastroenterol Motil* 2011;23:517-e200.
144. Sweis R, Anggiansah A, Wong T, et al. Assessment of esophageal dysfunction and symptoms during and after a standardized test meal: development and clinical validation of a new methodology utilizing high-resolution manometry. *Neurogastroenterol Motil* 2014;26:215-28.
145. Ang D, Misselwitz B, Hollenstein M, et al. Diagnostic yield of high-resolution manometry with a solid test meal for clinically relevant, symptomatic oesophageal motility disorders: serial diagnostic study. *Lancet Gastroenterol Hepatol* 2017;2:654-661.
146. Hollenstein M, Thwaites P, Butikofer S, et al. Pharyngeal swallowing and oesophageal motility during a solid meal test: a prospective study in healthy volunteers and patients with major motility disorders. *Lancet Gastroenterol Hepatol* 2017;2:644-653.
147. Marin I, Cisternas D, Abrao L, et al. Normal values of esophageal pressure responses to a rapid drink challenge test in healthy subjects: results of a multicenter study. *Neurogastroenterol Motil* 2017;29:e13021.
148. Yadlapati R, Tye M, Roman S, et al. Postprandial High-Resolution Impedance Manometry Identifies Mechanisms of Nonresponse to Proton Pump Inhibitors. *Clin Gastroenterol Hepatol* 2018;(in press).
149. Ribolsi M, Holloway RH, Emerenziani S, et al. Impedance-high resolution manometry analysis of patients with nonerosive reflux disease. *Clin Gastroenterol Hepatol* 2014;12:52-7.
150. Boeckxstaens GE, Hirsch DP, Verkleij CB, et al. Reproducibility of meal-induced transient lower oesophageal sphincter relaxations in patients with gastro-oesophageal reflux disease. *Neurogastroenterol Motil* 2005;17:23-8.
151. Kessing BF, Smout AJ, Bredenoord AJ. Clinical applications of esophageal impedance monitoring and high-resolution manometry. *Curr Gastroenterol Rep* 2012;14:197-205.
152. Galmiche JP, Zerbib F, des Varannes SB. Treatment of GORD: Three decades of progress and disappointments. *United European Gastroenterol J* 2013;1:140-50.
153. Lee WC, Yeh YC, Lacy BE, et al. Timely confirmation of gastro-esophageal reflux disease via pH monitoring: estimating budget impact on managed care organizations. *Curr Med Res Opin* 2008;24:1317-27.

154. Hatlebakk JG, Katz PO, Camacho-Lobato L, et al. Proton pump inhibitors: better acid suppression when taken before a meal than without a meal. *Aliment Pharmacol Ther* 2000;14:1267-72.
155. Carlsson R, Dent J, Watts R, et al. Gastro-oesophageal reflux disease in primary care: an international study of different treatment strategies with omeprazole. International GORD Study Group. *Eur J Gastroenterol Hepatol* 1998;10:119-24.
156. Tack J, Carbone F, Holvoet L, et al. The use of pictograms improves symptom evaluation by patients with functional dyspepsia. *Aliment Pharmacol Ther* 2014;40:523-30.
157. Tack J, Caenepeel P, Arts J, et al. Prevalence of acid reflux in functional dyspepsia and its association with symptom profile. *Gut* 2005;54:1370-6.
158. Desjardin M, Luc G, Collet D, et al. 24-hour pH-impedance monitoring on therapy to select patients with refractory reflux symptoms for antireflux surgery. A single center retrospective study. *Neurogastroenterol Motil* 2016;28:146-52.
159. Mee AS, Rowley JL. Rapid symptom relief in reflux oesophagitis: a comparison of lansoprazole and omeprazole. *Aliment Pharmacol Ther* 1996;10:757-63.
160. Richter JE, Kahrilas PJ, Johanson J, et al. Efficacy and safety of esomeprazole compared with omeprazole in GERD patients with erosive esophagitis: a randomized controlled trial. *Am J Gastroenterol* 2001;96:656-65.
161. Carlsson R, Dent J, Bolling-Sternevald E, et al. The usefulness of a structured questionnaire in the assessment of symptomatic gastroesophageal reflux disease. *Scand J Gastroenterol* 1998;33:1023-9.
162. Alrubaiy L, Hutchings HA, Williams JG. Assessing patient reported outcome measures: A practical guide for gastroenterologists. *United European Gastroenterol J* 2014;2:463-70.

Table 1 Studies that have compared the EGJ-CI among patient and control populations. Values reported at median (IQR).

| Study                        | Subject groups  | EGJ-CI<br>(mmHg•cm)                                     | Notes   |
|------------------------------|---|---|---|
| Nicodème 2014 <sup>111</sup> | Controls (n=75)<br>GERD (n=7)<br>Functional (n=45)                            | 39 [25-55]<br>18* [8-30]<br>27 [17-69]                  | GERD had +++ abnormal pH-impedance studies vs partial + or - for functional |
| Tolone 2015 <sup>112</sup>   | Functional (n=39)<br>GERD (n=91)  | 22 [10-41]<br>11* [3-21]                                | GERD or functional by endoscopy and pH-impedance testing                    |
| Jasper 2016 <sup>113</sup>   | Controls (n=65)<br>GERD (n=116)   | 63 [50-90]<br>50* [28-70]                               | GERD by pH-metry  |
| Wang 2016 <sup>114</sup>     | Controls (n=21)<br>GERD (n=68)  | 35 [26-58]<br>30* [15-53]                               | GERD patients underwent fundoplication                                      |
| Xie 2017 <sup>115</sup>      | Controls (n=21)<br>Esophagitis (n=39)<br>NERD (n=38)<br>Hypersensitive (n=21) | 63 [38-83]<br>22* [20-31]<br>26* [15-38]<br>30* [19-44] | Patients differentiated by pH-impedance and symptom correlation             |
| Ham 2017 <sup>109</sup>      | Controls (n=23)<br>Esophagitis (n=25)<br>NERD (n=16)<br>Non-GERD ((n=91)      | 67 [27-79]<br>28* [4-63]<br>26* [15-32]<br>51 [3-153]   | Non-GERD patients had negative pH-impedance studies                         |

\*p<0.05 vs controls or comparator

Methods of EGJ-CI computation were not uniform between these studies, and this might explain differences in calculated thresholds. NERD: non-erosive reflux disease

Table 2. Classification of motor function in GERD using esophageal high-resolution manometry

|  | Metrics  | Description   |
|--|--|---|
| EGJ barrier function   |  |   |
| Morphology   | Separation between LES and crural diaphragm (CD)   | Type 1: superimposed LES and CD<br>Type 2: axially separated LES and CD pressure signals separated by less than 3 cm<br>Type 3a: $\geq 3$ cm separation between the LES and CD pressure signatures with respiratory inversion point at the level of the CD<br>Type 3b: $\geq 3$ cm separation between the LES and CD pressure signatures with respiratory inversion point at the level of the LES |
| Vigor  | EGJ-CI (mmHg•cm)   | DCI box set to encompass the LES and CD over a period of three complete respiratory cycles above a threshold pressure of the gastric baseline   |
| Esophageal body motor function                                       |  |   |
|  | Distal contractile integral, DCI (mmHg•cm•s)<br><br>Defect (measure at 20-mmHg isobaric contour) | Intact: $\geq 50\%$ of contractions with DCI $> 450$ mmHg•cm•s and no defect<br><br>Fragmented: $\geq 50\%$ of contractions with DCI $> 450$ mmHg•cm•s and defect $> 5$ cm<br><br>Ineffective esophageal motility: $\geq 50\%$ of contractions with DCI $< 450$ mmHg•cm•s<br><br>Absent peristalsis: 100% of contractions with DCI $< 100$ mmHg•cm•s  |
| Provocative tests  |  |   |
| MRS (5 liquid swallows - 2ml each - taken less than 4 seconds apart) | Contractile response   | Post MRS DCI augmentation   |
|  | Failure of contractile response  | Absent post MRS contraction   |
| RDC (free water drinking of 200ml of water within 30 seconds)        | Panesophageal pressurization   |   |
|  | LES relaxation   |   |
|  | Effective post RDC contraction   |   |



Table 3: Comparison of the Porto and the Lyon consensus conclusions.

| Porto Consensus   | Lyon consensus  |
|---|---|
| <i>No discussion of endoscopy</i>   | Conclusive endoscopic criteria for GERD<br>- LA Grade C or D esophagitis<br>- Biopsy-proven Barrett's esophagus<br>- Peptic stricture   |
| Esophageal impedance monitoring is the only recording method that can achieve high sensitivity for detection of all types of reflux episodes while pH monitoring is required for characterization of reflux acidity. However, the role of impedance monitoring in the management of patients with GERD still needs to be defined. | pH-impedance monitoring is the gold standard for detection and characterization of reflux episodes but is expensive, not widely available and interpretation is time consuming.<br><br>When reflux monitoring is indicated on PPI, pH-impedance should be performed.<br>When reflux monitoring is indicated off PPI, the choice between catheter based pH-monitoring, wireless pH monitoring and pH-impedance monitoring is dependent of cost and availability. |
| <i>No discussion of the conditions (off or on PPI) to perform reflux testing</i>  | Reflux monitoring is recommended off PPI in instances of "unproven" GERD and on PPI in instances of "proven GERD" (previous LA grade C or D esophagitis, biopsy proven Barrett's esophagus, peptic stricture or AET off PPI > 6%)   |
| <i>No discussion of normal values</i>   | An AET < 4% is normal and an AET > 6% is abnormal (whatever the type of reflux monitoring and whether the study was performed off or on PPI)  |
| <i>No discussion of normal values</i>   | Reflux episodes > 80 / 24 hours is abnormal and < 40 is physiologic on pH-impedance performed off or on PPI. Number of reflux episodes is an adjunctive metric to be utilized when AET is borderline or inconclusive  |
| Basal intraluminal impedance is abnormally low in patients with esophageal mucosal abnormalities such as Barrett's esophagus or esophagitis.  | Measurement of baseline mucosal impedance (using either through the scope device or MNBI during ambulatory pH-impedance monitoring) is an adjunctive metric for the diagnosis of GERD   |
| <i>No discussion of reflux-symptom association</i>  | A combination of a positive SI and positive SAP provides the best evidence of clinically relevant association between reflux episodes and symptoms.   |

|   |   |
|---|---|
| <p>Using manometry, common cavities occur during a higher proportion of reflux episodes in neonates and infants than in adults</p> <p><i>No discussion of esophageal motor function in GERD</i></p> | <p>Esophageal high-resolution manometry is not useful for the direct diagnosis of GERD but can provide adjunctive information</p> <ul style="list-style-type: none"> <li>- to assess EGJ barrier function including its morphology (type I to III) and its vigor (using EGJ-CI)</li> <li>- to evaluate esophageal body motor function (intact, ineffective, fragmented or absent contractility) that correlates with esophageal reflux burden</li> </ul> <p>Adjunctive tests should be included in the HRM protocol</p> <ul style="list-style-type: none"> <li>- to evaluate the contractile response (multiple rapid swallow)</li> <li>- to evaluate EGJ obstruction (rapid drink challenge test)</li> </ul> |
| <p>Bilitec™ is a monitoring system that can detect duodeno-gastro-oesophageal reflux (DGOR) by utilizing the optical properties of bilirubin</p>  | <p><i>Bilitec™ is no longer considered a reliable diagnostic tool for GERD and was not discussed</i></p>  |

Table 4. GERD phenotypes predicting abnormal reflux burden from clinical evaluation and esophageal testing

|                                    | Pathologic GERD   |   |   | Modifiers  |
|------------------------------------|---|---|---|--|
|                                    | High likelihood   | Intermediate likelihood   | Low likelihood  |  |
| <b>CLINICAL PHENOTYPES</b>         |   |   |   |  |
| Symptoms                           | Heartburn, acid regurgitation                                   | Chest pain  | Cough, laryngeal symptoms   | Hypersensitivity and hypervigilance                                |
| Endoscopy                          | High-grade esophagitis, Barrett's mucosa, peptic stricture      | Low-grade esophagitis, Normal exam on PPI therapy                 |   | Hiatus hernia, ongoing PPI therapy                                 |
| ROME IV                            | NERD (abnormal pH metry)*                                       | Symptom response to PPI therapy                                   | Reflux hypersensitivity functional heartburn, functional chest pain | Hypersensitivity and hypervigilance                                |
| Lyon Consensus*                    | Conclusive evidence of GERD                                     | Borderline or inconclusive evidence                               | Physiologic reflux parameters                                       | Novel metrics<br>Motor classification                              |
| <b>MECHANISTIC PHENOTYPES</b>      |   |   |   |  |
| Pattern of reflux                  | Increased acid exposure ± increased numbers of reflux episodes* | Borderline acid exposure ± borderline numbers of reflux episodes* | Normal reflux metrics   | pH of refluxate, baseline impedance, hypochlorhydria, achlorhydria |
| Mechanism of reflux                | TLESR<br>Hypotensive EGJ<br>Abnormal EGJ morphology             | Supragastric belch<br>Rumination                                  | Normal EGJ morphology and function                                  | Obesity, increased abdominal girth                                 |
| Clearance of refluxate             | Absent contractility<br>Hiatus hernia                           | Minor motor disorder ± contraction reserve                        | Normal peristalsis  | Xerostomia, baseline impedance, PSPW index, motor classification   |
| Cognition, perception of sensation | Appropriate symptom perception, symptom reflux association      | Increased perception  | Visceral hypersensitivity, hypervigilance                           | Anxiety, depression<br>Panic disorder                              |

\*as described by the Lyon Consensus, Figure 3

NERD: nonerosive reflux disease; EGD: esophagogastroduodenoscopy; TLESR: transient lower esophageal sphincter relaxation; EGJ: esophagogastric junction; PSPW: post-reflux swallow induced peristaltic wave

## FIGURE LEGENDS

Figure 1. Esophagogastric junction morphology as depicted in HRM. With type 1 morphology the crural diaphragm (CD) component, evident during inspiration (I), is completely superimposed of the lower esophageal sphincter (LES) component such that the magnitude of the actual LES pressure is not discernible. With Type 2 morphology, there is partial separation of the LES and CD constituents, but the respiratory inversion point (RIP) remains at the level of the CD, evident by the decrease observed in the LES pressure band during inspiration. Other characteristics of Type 2 morphology are that the LES-CD separation is  $<3$  cm and that the pressure trough between the LES and CD is greater than intragastric pressure. With Type 3 morphology, there is  $\geq 3$  cm separation between the LES and CD and the pressure trough between the two is equal to intragastric pressure during expiration (E). However, the RIP remains at the level of the CD in type 3a and elevated to the level of the LES pressure band with 3b. This is evident by the decreases in LES pressure during inspiration in IIIa and increases in LES pressure during inspiration in 3b.

Figure 2. High-resolution manometry metrics used in the motor classification of GERD. The esophagogastric junction contractile integral (EGJ-CI) measures vigor of the EGJ barrier using a software tool that encompasses length and vigor of the EGJ above the gastric baseline. The measurement is made over three respiratory cycles during quiet rest, and corrected for duration of respiration. The distal contractile integral (DCI) measures vigor of smooth muscle contraction taking length, duration and amplitude of contraction into consideration. Following a series of repetitive swallows (multiple rapid swallows, MRS), DCI augments higher than mean DCI from single swallows when there is contraction reserve.

Figure 3. Interpretation of esophageal test results in the context of GERD. Any one conclusive finding provides strong evidence for the presence of GERD. While a normal EGD does not exclude GERD on its own, this provides strong evidence against GERD when combined with acid exposure time (AET)  $<4\%$

and <40 reflux episodes on pH-impedance monitoring off PPI therapy. When evidence is inconclusive or borderline, adjunctive or supportive findings can add confidence to the presence or absence of GERD. Histopathology as an adjunctive measure requires a dedicated scoring system (incorporating papillary elongation, basal cell hyperplasia, dilated intercellular spaces (DIS), intraepithelial inflammatory cells, necrosis and erosions) or evidence of DIS on electron microscopy. However, adjunctive findings, particularly histopathology and motor findings in isolation, are not enough to diagnose GERD. HRM: high-resolution manometry; AET: acid exposure time; MNBI: mean nocturnal baseline impedance; PSPWI index: post-reflux swallow-induced peristaltic wave index; EGJ: esophagogastric junction.