

Milk protein composition in purebred Holsteins and in first/second-generation crossbred cows from Swedish Red, Montbeliarde and Brown Swiss bulls

A. Maurmayr¹, S. Pegolo^{1†}, F. Malchiodi², G. Bittante¹ and A. Cecchinato¹

¹Department of Agronomy, Food, Natural Resources, Animals and Environment (DAFNAE), University of Padua, Viale dell'Università 16, 35020 Legnaro, Italy;

²Department of Animal Biosciences, Centre for Genetic Improvement of Livestock, University of Guelph, Guelph, Ontario, Canada, N1G 2W1

(Received 24 August 2017; Accepted 11 December 2017)

The aim of this study was to analyze milk protein composition in purebred and crossbred dairy cattle and estimate the effects of individual sources of variation on the investigated traits. Milk samples were collected from 505 cows from three commercial farms located in Northern Italy, some of which had originated from crossbreeding programs, although most were purebred Holsteins (HO). The basic crossbreeding scheme was a three-breed rotational system using Swedish Red (SR) semen on HO cows (SR × HO), Montbeliarde (MO) semen on SR × HO cows (MO × (SR × HO)) and HO semen again on MO × (SR × HO) cows. A smaller number of purebred HO from each of the herds were mated inverting the breed order (MO × HO and SR × (MO × HO)) or using Brown Swiss (BS) bulls (BS × HO) then MO bulls (MO × (BS × HO)). Milk samples were analyzed by reverse-phase HPLC to obtain protein fraction amounts (g/l) and proportions (% of total true protein). Traits were analyzed using a linear model, which included the fixed effects of herd-test-day (HTD), parity, days in milk and breed combination. Results showed that milk protein fractions were influenced by HTD, stage of lactation, parity and breed combination. The increase in protein concentration during lactation was due in particular to β-casein (β-CN), α_{S1}-CN and β-lactoglobulin (β-LG). The higher protein content of primiparous milk was mainly due to higher concentrations of all casein fractions. The milk from crossbred cows had higher contents and proportions of κ-CN and α-lactalbumin (α-LA), lower proportions of β-LG and greater proportion of caseins/smaller in whey proteins on milk true protein than purebred HO. The three-way crossbreds differed from two-way crossbreds only in having greater proportions of α-LA in their milk. Of the three-way crossbreds, the SR sired cows yielded milk with a smaller content and proportion of β-LG than the MO sired cows, and, consequently, a higher proportion of caseins than whey proteins. Results from this study support the feasibility of using crossbreeding programs to alter milk protein profiles with the aim of improving milk quality and cheese-making properties.

Keywords: crossbreeding, milk protein fractions, Swedish Red, Brown Swiss, Montbeliarde

Implications

Crossbreeding can be a practical way to improve not only functional traits of dairy cows, but also milk quality and technological properties through the modification of milk protein profile. In this study, stage of lactation, parity and herd-test-day (HTD; which includes feeding and management practices) affected milk protein composition. Compared with purebred Holsteins (HO), cows crossbred from Swedish Red (SR), Montbeliarde (MO) and Brown Swiss (BS) sires yielded milk with a protein composition more favorable to cheese production. Few differences were found between first- and second-generation and within second-generation

crossbred cows. Crossbreeding programs might therefore be effectively used to improve milk quality and technological characteristics.

Introduction

The proteins in bovine milk consist of almost 80% caseins (CN), corresponding to 2.5% to 2.8% w/v (Holland *et al.*, 2010). Caseins are the most important protein fractions as they are directly responsible for curd formation and cheese yield (Wedholm *et al.*, 2006), even though they have different characteristics and roles. For instance, κ-CN, which constitutes ~12% of total CN, is responsible for the stability of CN micelles and facilitates micelle coagulation (Creamer *et al.*, 1998). Whey proteins account for ~20% of total milk

† E-mail: sara.pegolo@unipd.it

proteins, and, despite having less economic importance, they have high nutritional value and are much more quickly digested and absorbed than CN (McSweeney and Fox, 2013). Therefore, modification of the content, proportion and daily yield of milk protein fractions may have very important nutritional, technological and economic implications. Indeed, the presence of genetic polymorphisms within CN and whey proteins encoding genes influences the amount (and the proportion) of milk protein fractions and consequently affects milk coagulation properties (Bonfatti *et al.*, 2010; Bittante *et al.*, 2012; Jensen *et al.*, 2012). This is highly significant for some countries, particularly in the European Union where the majority of the milk production goes to the cheese-making industry (Wedholm *et al.*, 2006).

The success of selection for milk production has contributed to the spread of the HO breed throughout the world, but has negatively affected important traits, like milk quality, productive life, fertility and longevity. Nowadays, different selection indices and mating schemes ensure that dairy cows and their genetic richness, including milk quality and functional traits, are better managed. Crossbreeding may be a way of overcoming most of these problems by providing new genetic combinations, and more robust crossbred animals compared with the parental breeds (Sørensen *et al.*, 2008). Crossbreeding has been described as a useful strategy to improve the fitness, fertility and longevity of dairy cows, and to genetically alter the yield of milk and its components (Lopez-Villalobos *et al.*, 2000). Indeed, first-generation offspring (F1) of HO crossbred with MO and Scandinavian Red exhibited better functional traits but lower fat and protein contents and milk yields than pure HO (Heins *et al.*, 2006; Heins and Hansen, 2012). More recently, however, MO × HO and Viking Red × HO crossbreds, despite a lower productivity in terms of milk yield, were collectively found to have higher fat and protein yields and percentages than purebred HO cows (Hazel *et al.*, 2017). Likewise, BS × HO crosses had higher fat and protein yields than HO purebreds, although the milk production of the latter was still reported to be higher (Dechow *et al.*, 2007). Several other studies have been carried out on the effects of crossbreeding on functional traits, such as fertility and udder health, and on production traits, such as milk yield, fat and protein contents, and cheese-making properties (e.g., Malchiodi *et al.*, 2011, 2014a and 2014b; de Haas *et al.*, 2013; Dezetter *et al.*, 2015) but there is little information on individual milk protein fractions behavior in crossbred cows, which is particularly important in countries, like many in Europe, where a large proportion of milk production is destined for cheese-making.

The objectives of this study, therefore, were (i) to determine the content and proportions of milk protein fractions in individual milk samples of purebred HO, two-way crossbred cows from BS (BS × HO), MO (MO × HO) or SR (SR × HO) bulls, and three-way crossbred cows from the same breeds (MO × (BS × HO); MO × (SR × HO); and SR × (MO × HO)); and (ii) to estimate the effects of HTD, parity and stage of lactation on these traits.

Material and methods

Animals and milk sampling

Milk samples were collected once during the evening milking from individual purebred HO cows and crossbred cows (n = 544) reared in three herds located in the north of Italy (2 to 3 sampling days per herd). The herds were following the same crossbreeding scheme (Genesis Project SRL, Castelnuovo Sotto, Reggio Emilia, Italy) starting from purebred H cows, as detailed in Malchiodi *et al.* (2014b). The basic scheme was a three-breed rotational system using SR semen on HO cows (SR × HO), MO semen on SR × HO cows (MO × (SR × HO)) and HO semen again on MO × (SR × HO). A smaller number of purebred HO from each herd were mated inverting the breed order (MO × HO and SR × (MO × HO)) or using BS bulls instead of SR (BS × HO) and then MO bulls (MO × (BS × HO)). Figure 1 shows the crossbreeding scheme together with the number of cows sampled from each breed combination. Herds were managed in accordance with EU regulations for the production of Protected Designation of Origin (PDO) Parmigiano-Reggiano cheese, meaning that silage, pasture and fresh herbage were not allowed, and the rations (fed as a total mixed diet) consisted of dry roughage, concentrates and added water.

A preservative (Bronopol, 0.6:100 v/v; Sigma-Aldrich, St. Louis, MO, USA) was added to the milk immediately after collection to prevent microbial growth. Milk samples were refrigerated at 4°C, then transferred to the Laboratory of the Department of Agronomy, Food, Natural Resources, Animals and Environment (DAFNAE) of the University of Padua (Legnaro, Padua, Italy), where they were stored at -80°C until protein fraction analysis.

Analysis of milk quality traits and protein fractions by RP-HPLC

Individual milk samples were analyzed for fat and protein with a MilkoScan FT6000 (Foss, Hillerød, Denmark). Milk pH was determined using a Crison Basic 25 electrode (Crison, Barcelona, Spain). Somatic cell counts were obtained with a Fossomatic FC counter (Foss, Hillerød, Denmark) and then log transformed to obtain somatic cell scores (SCS) (Ali *et al.*, 1980).

The α_{S1} -CN, α_{S2} -CN, β -CN, κ -CN, β -lactoglobulin (β -LG) and α -lactalbumin (α -LA) contents were measured in milk samples from 505 animals using the RP-HPLC method proposed by Maurmayr *et al.* (2013), which allows CN and whey proteins to be separated in one run, thereby ensuring rapid analysis (<20 min) and good peak resolution. A single




Figure 1 Crossbreeding scheme. The number of cows sampled for milk quality traits and milk protein fractions analyses is reported in brackets. HO = Holstein; MO = Montbeliarde; SR = Swedish Red; BS = Brown Swiss.

mother solution of purified proteins of commercial standards, including κ -CN (>80%), α -CN (>70%), β -CN (>90%), α -LA (~85%), β -lactoglobulin B variant (>90%), β -lactoglobulin A variant (>90%) (Sigma-Aldrich), was used to draw specific calibration curves. The γ -CN fraction, consisting of proteolytic products of β -CN, was not detectable with the current method.

To carry out protein quantification using the current RP-HPLC method, total casein content (caseins, g/l) was calculated as the sum of the κ -CN, α_{S1} -CN, α_{S2} -CN and β -CN contents in the milk, and total whey protein content (g/l) was calculated as the sum of β -LG and α -LA. The proportions of protein fractions were expressed as the percentage (%) of total true protein content (g/l), calculated as the sum of the casein and whey protein contents.

Statistical analysis

Data were analyzed using the SAS GLM procedure (SAS Institute Inc., Cary, NC, USA) according to the following linear model:

$$Y_{ijklm} = \mu + HTD_i + Parity_j + DIM_k + Breed_l + e_{ijklm}$$

where Y_{ijkl} is the measure of a trait; μ the general mean of the model; HTD_i is the fixed effect of HTD i ($i = 1, \dots, 7$); $Parity_j$ the fixed effect of parity j of the cow ($j = 1$: first parity, $j = 2$: second parity, $j = 3$: third and subsequent parities); DIM_k the fixed effect of days in milk (DIM) class k (five classes of 60-day intervals, except the last class, which included samples collected at DIM 240 or more days); $Breed_l$ the fixed effect of the l th breed combination ($l = 1$ to 7); and e_{ijklm} the random residual assumed to have a normal distribution with $e_{ijklm} \sim N(0, \sigma_e^2)$, where σ_e^2 is the residual variance. Orthogonal contrasts were estimated between the least square means (LSM) of traits for: (i) the effect of DIM: (a) the linear component and (b) the quadratic component; between LSM of traits for (ii) the effect of parity: (a) first parity *v.* second and subsequent parities and (b) second parity *v.* third and subsequent parities; and (iii) the effect of breed combination: (a) the effect of crossbreeding (HO *v.* all crossbred cows), (b) the effect of generation (first-generation *v.* second-generation crosses), (c) the effect of SR sires in the first-generation cross (SR \times HO *v.* MO \times HO + BS \times HO), (d) comparison of the two paternal Alpine breeds in the first-generation crosses (MO \times HO *v.* BS \times HO), (e) comparison of SR sires and MO sires in the second-generation crosses (SR \times (MO \times HO) *v.* MO \times (SR \times HO) + MO \times (BS \times HO)) and (f) the effect of maternal grand-sire breed in second-generation crosses with MO sire MO \times (SR \times HO) *v.* MO \times (BS \times HO).

Results and discussion

In the present work, we investigated the effect of animal factors, such as parity, stage of lactation and breed combination, and the effects of HTD on bovine milk protein fractions. Descriptive statistics for the investigated traits are reported in Table 1.

Table 1 Descriptive statistics for protein composition and quality traits of individual milk samples from purebred and crossbred cows ($n=505$)

Trait ¹	Mean	SD	P1	P99
Caseins				
g/l	32.66	4.50	22.18	42.55
%	82.29	3.18	74.28	88.96
κ -CN				
g/l	4.50	0.84	2.52	6.62
%	11.34	1.58	7.28	15.02
β -CN				
g/l	13.10	2.14	7.15	17.36
%	32.96	3.02	21.902	38.72
α_{S1} -CN				
g/l	11.64	1.55	8.00	15.39
%	29.40	1.98	24.25	34.05
α_{S2} -CN				
g/l	3.42	0.81	1.96	5.49
%	8.59	1.50	5.52	12.06
Whey proteins				
g/l	6.99	1.34	4.21	9.89
%	17.71	3.18	11.44	27.60
α -LG				
g/l	1.01	0.21	0.39	1.45
%	2.56	0.48	1.20	3.61
β -LG				
g/l	5.98	1.28	3.37	8.90
%	15.15	3.13	9.26	24.00
Milk quality traits				
Fat (%)	4.09	0.86	2.09	6.36
Protein (%)	3.71	0.30	2.94	4.41
SCS (units)	2.56	1.84	-0.47	7.43
pH	6.47	0.08	6.29	6.68
Milk yield (kg/day)	31.82	9.95	11.78	55.20

P1 = first percentile; P99 = 99th percentile; κ -CN = κ -casein; β -CN = β -casein; α_{S1} -CN = α_{S1} -casein; α_{S2} -CN = α_{S2} -casein; α -LA = α -lactalbumin; β -LG = β -lactoglobulin; SCS = somatic cell score.

¹Contents of all protein fractions were measured by reversed-phase HPLC on skim milk. The proportion of protein fractions were expressed as percentage (%) on total true protein content (g/l). Caseins: sum of total casein fraction; whey proteins: sum of total whey fraction.

Effects of herd-test-day

We found that HTD significantly affected all milk protein fraction contents and proportions ($P < 0.001$), in particular α_{S1} -CN and α -LA, which might reflect the effect of differences in the environmental, management and feeding conditions across herds, and variations in the test records collected on different days. However, HTD explained ~11% of variance for milk protein fractions on average, in line with previous studies which found that the proportion of phenotypic variation for milk proteins explained by herd was relatively small (Schopen *et al.*, 2009), especially compared with fat, while genetic factors play the most important role.

Effects of days in milk and parity

Lactation stage was confirmed as an important source of variation in milk protein fractions (Table 2), as previously reported (Jöduu *et al.*, 2008). In this study, the total amount of protein, which averaged 39.65 g/l, increased linearly over

Table 2 Least squares means of milk protein fraction and milk protein composition across days in milk (DIM)

Trait ¹	DIM					Contrast	
	<60 day	60 to 120 days	121 to 180 days	181 to 240 days	>240 days	L	Q
Caseins							
g/l	31.04	31.61	33.32	33.64	33.80	***	Ns
%	82.74	82.62	82.51	82.21	81.43	**	Ns
κ-CN							
g/l	4.41	4.51	4.67	4.60	4.42	Ns	**
%	11.70	11.79	11.58	11.29	10.67	***	**
β-CN							
g/l	12.09	12.46	13.36	13.56	13.79	***	Ns
%	32.24	32.52	33.06	33.11	33.17	*	Ns
α_{S1}-CN							
g/l	11.18	11.27	11.81	11.93	12.03	***	Ns
%	29.88	29.54	29.30	29.18	29.06	***	Ns
α_{S2}-CN							
g/l	3.36	3.37	3.48	3.55	3.56	*	Ns
%	8.92	8.77	8.57	8.63	8.53	*	Ns
Whey proteins							
g/l	6.44	6.60	6.96	7.20	7.67	***	Ns
%	17.26	17.38	17.49	17.79	18.57	**	Ns
α-LA							
g/l	0.97	1.01	1.03	1.06	1.03	**	Ns
%	2.57	2.64	2.56	2.62	2.53	Ns	Ns
β-LG							
g/l	5.47	5.59	5.93	6.14	6.64	***	Ns
%	14.69	14.74	14.93	15.17	16.04	**	Ns

L = linear; Q = quadratic; κ -CN = κ -casein; β -CN = β -casein; α_{S1} -CN = α_{S1} -casein; α_{S2} -CN = α_{S2} -casein; α -LA = α -lactalbumin; β -LG = β -lactoglobulin.

¹Contents of all protein fractions were measured by reversed-phase HPLC on skim milk. The proportion of protein fractions were expressed as percentage (%) on total true protein content (g/L). Caseins: sum of total casein fraction; whey protein: sum of total whey fractions.

* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

the 5-bimonthly periods comprising lactation ($P < 0.001$), whereas total CN and whey contents (g/l) followed the same trend ($P < 0.001$; Table 2), consistent with previous data (Coulon *et al.*, 1998). Linear increases were also found for the individual protein fraction contents, except for κ -CN, which showed a quadratic pattern ($P < 0.01$; Table 2). Indeed, after the peak of lactation and just until the dry period, milk production starts to decrease, but at the same time fat, protein and body weight increase, as result of the concentrating effect of decreasing milk volumes and the energy balance change. Stage of lactation also significantly influenced CN and whey protein proportions (%), except for α -LA (Table 2), results that were only partially consistent with previous data (Ostensen *et al.*, 1997). In particular, in the present study the total casein proportion ($P < 0.01$) followed a decreasing trend, as did the κ -CN ($P < 0.001$), α_{S1} -CN ($P < 0.001$) and α_{S2} -CN ($P < 0.05$) proportions, which decreased linearly across lactation, while β -CN exhibited the opposite trend (linear increase; $P < 0.05$). On the other hand, the linear increase of the whey protein proportion ($P < 0.01$) was essentially due to the β -LG proportion being higher in late than in early lactation ($P < 0.01$). Along with the lactation an increase in milk SCS was detected (Malchiodi *et al.*, 2014b), which might be related to an increase in plasmin

activity (Politis *et al.*, 1989). The increasing proportion of total whey at the expense of casein might be likely due to the possible increase in plasmin hydrolysis of caseins, which makes late-lactation milk less suitable for cheese making (Akers, 2016). During lactation, mammary epithelial cells exhibit an increase in sensitivity to insulin which is an important inducer of milk protein gene expression (Bionaz and Looor, 2011). The different behavior of β -CN respect to the other casein proportions, which is in line with previous findings (Ostensen *et al.*, 1997; Barber *et al.*, 2005), might be explained by possible differential effects of insulin on individual casein fractions (Menziez *et al.*, 2009). Another possible explanation might rely, at least partially, on micelles characteristics; indeed, it was reported that micelle size increases in late lactation in the cow (Holt and Baird, 1978) and the relative proportion of β -CN increases with increasing micelle size while κ -CN showing the opposite trend (Ekstrand and Larsson-Raźnikiewicz, 1978).

Parity had significant effects on total CN content (g/l) ($P < 0.001$; Table 3), the milk of first-parity cows, in particular, having significantly higher contents of all individual CN fractions ($P < 0.001$). Significantly higher contents of α -LA were also found in primiparous cows ($P < 0.05$), while total whey and β -LG contents were unaffected by parity (Table 3),

Table 3 Least squares means of milk protein fraction and milk protein composition across parities

Trait ¹	Parity (P)			Contrasts	
	1	2	≥3	P1 v. P2 + ≥P3	P2 v. ≥P3
Caseins					
g/l	34.60	32.13	31.32	***	Ns
%	83.03	82.00	81.87	***	Ns
κ-CN					
g/l	4.86	4.35	4.37	***	Ns
%	11.65	11.11	11.46	*	Ns
β-CN					
g/l	13.94	12.83	12.38	***	Ns
%	33.46	32.74	32.26	**	Ns
α _{S1} -CN					
g/l	12.13	11.53	11.28	***	Ns
%	29.15	29.49	29.54	Ns	Ns
α _{S2} -CN					
g/l	3.67	3.42	3.29	***	Ns
%	8.77	8.66	8.61	Ns	Ns
Whey proteins					
g/l	7.06	7.00	6.87	Ns	Ns
%	16.97	18.00	18.13	***	Ns
α-LA					
g/l	1.05	1.00	1.00	*	Ns
%	2.53	2.58	2.64	Ns	Ns
β-LG					
g/l	6.01	6.00	5.87	Ns	Ns
%	14.44	15.42	15.49	***	Ns

κ-CN = κ-casein; β-CN = β-casein; α_{S1}-CN = α_{S1}-casein; α_{S2}-CN = α_{S2}-casein; α-LA = α-lactalbumin; β-LG = β-lactoglobulin.

¹Contents of all protein fractions were measured by reversed-phase HPLC on skim milk. The proportion of protein fractions were expressed as percentage (%) on total true protein content (g/l). Caseins: sum of total casein fraction; whey protein: sum of total whey fractions.

* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

a finding at variance with Jõudu *et al.* (2008), who reported a significant effect of parity on β-LG content. The proportion of total CN was also significantly higher in primiparous cows than in multiparous cows ($P < 0.001$; Table 3), in agreement with previous results (Ikonen *et al.*, 2004), while the percentages of total whey were significantly lower ($P < 0.001$; Table 3). Looking at the individual fractions as proportions of total protein content, those of κ-CN and β-CN were significantly higher ($P < 0.05$ and $P < 0.01$, respectively), and β-LG significantly lower ($P < 0.001$) in the milk of first-parity cows (Table 3). Remarkably, multiparous cows of this study were characterized by higher SCS respect to first-parity cows (Malchiodi *et al.*, 2014b). β-Casein was reported to be the most susceptible casein isoform to degradation by plasmin activity (Cruden *et al.*, 2005). Accordingly, the largest difference between primiparous and multiparous cows among caseins proportions was detected for β-CN. Overall, these results add weight to the evidence that, despite a higher yield, the milk from multiparous cows tends to be of lower quality due to a reduction in the protein content (Hansen *et al.*, 2006) and particularly of those caseins having a greater involvement in milk coagulation and curd firming.

Effects of breed

Results regarding the effects of crossbreeding on milk protein fractions are shown in Table 4. Significant effects were found for the proportions of total CN and total whey, and the contents and proportions of κ-CN, α-LA and β-LG. In particular, the milk of crossbred animals had significantly higher proportions of total CN and lower proportions of total whey proteins than purebred HO ($P < 0.05$). With respect to individual protein fractions, the milk from crossbred cows was characterized by higher contents and proportions of κ-CN ($P < 0.01$ and $P < 0.001$, respectively) and α-LA ($P < 0.05$), and lower contents and proportions of β-LG ($P < 0.05$ and $P < 0.01$, respectively). The higher proportion of total caseins in crossbred cows was essentially due to the higher content and proportion of κ-CN content which is related to the higher frequency of B allele in the crossbred populations of this study (38.4% on average) respect to HO (28.9%), with the only exception of SR × HO (18.9%). Selective breeding is able to alter the milk protein composition to improve milk quality and milk technological properties; for instance, a higher amount of κ-CN is associated with better milk coagulation properties (Bittante *et al.*, 2012). Several studies have focused on the relationship between protein composition, milk production and milk protein variants (e.g., Heck *et al.*, 2009; Demeter *et al.*, 2010). Furthermore, cheese yield increases with a greater CN concentration, and cheese-making properties, like milk coagulation time and curd firmness, depend on casein composition (Wedholm *et al.*, 2008). Therefore, the high content and proportion of κ-CN observed in crossbred cows may help to explain the more favorable curd firmness characteristics of their milk (Malchiodi *et al.*, 2014b).

Significant differences between the F1 and F2 generations were observed only for α-LA, which was found in higher proportions in the F2 generation ($P < 0.05$). This could be partially due to recombination effects between the loci, differences in the gene frequencies, or additive by additive and dominance by dominance epistatic effects (William and Pollak, 1985). Looking specifically at the various crossbred combinations, the only significant difference was SR × (MO × HO) v. MO × (SR × HO) + MO × (BS × HO) for total CN and total whey % ($P < 0.05$), and for both the content and proportion of β-LG ($P < 0.05$) (Table 4). This result might be explained by differences between these crossbreds due to genotype interactions and cross-combinations among MO, BS and HO.

Conclusions

In conclusion, our results show that milk protein fraction contents and proportions vary with HTD, stage of lactation and number of lactations, albeit to different extents. Crossbreeding programs can be a very useful strategy in animal breeding. Besides the well-known effects on fertility, longevity and robustness traits, crossbreeding may be a viable option for improving milk quality, in particular milk protein composition. Indeed, the crossbred animals included in this study produced milk with higher amounts of κ-CN, which is desirable in cheese production.

Table 4 Least squares means of milk protein fraction and milk protein composition across breed combinations

Trait ¹	Purebreds		F1 crossbreds			F2 crossbreds		Contrasts	
	HO × HO	SR × HO	MO × HO	BS × OH	SR × (MO × HO)	MO × (SR × HO)	MO × (BS × HO)	Purebreds v. Crossbreds	F1 v. F2 crossbreds
Caseins									
g/l	32.30	33.37	33.10	32.24	33.47	32.69	31.59	Ns	Ns
%	81.71	82.17	82.97	81.69	83.73 ^a	82.40	81.45	*	Ns
κ-CN									
g/l	4.31	4.54	4.68	4.56	4.72	4.57	4.30	**	Ns
%	10.89	11.16	11.74	11.56	11.84	11.59	11.06	***	Ns
β-CN									
g/l	13.10	13.49	13.03	12.67	13.44	13.03	12.58	Ns	Ns
%	33.11	33.19	32.65	32.09	33.57	32.76	32.40	Ns	Ns
α₅₁-CN									
g/l	11.54	11.93	11.72	11.56	11.89	11.62	11.25	Ns	Ns
%	29.27	29.43	29.45	29.41	29.77	29.34	29.07	Ns	Ns
α₅₂-CN									
g/l	3.35	3.41	3.67	3.45	3.42	3.47	3.46	Ns	Ns
%	8.44	8.39	9.13	8.63	8.55	8.71	8.92	Ns	Ns
Whey proteins									
g/l	7.19	7.22	6.78	7.15	6.45	6.93	7.11	Ns	Ns
%	18.29	17.83	17.03	18.31	16.27 ^a	17.60	18.55	*	Ns
α-LA									
g/l	0.98	1.02	1.00	0.97	1.08	1.07	1.02	*	Ns
%	2.48	2.52	2.51	2.49	2.70	2.71	2.68	*	*
β-LG									
g/l	6.21	6.20	5.78	6.18	5.37 ^a	5.86	6.09	*	Ns
%	15.81	15.31	14.52	15.82	13.57 ^a	14.89	15.87	**	Ns

HO = Holstein; SR = Swedish Red; BS = Brown Swiss; MO = Montbeliarde; κ-CN = κ-casein; β-CN = β-casein; α₅₂-CN = α₅₂-casein; α₅₁-CN = α₅₁-casein; α-LA = α-lactalbumin; β-LG = β-lactoglobulin.

^a SR × (MO × HO) significantly different from MO × (SR × HO) + MO × (BS × HO) at $P < 0.05$.

¹ Contents of all protein fractions were measured by reversed-phase HPLC on skim milk. The proportion of protein fractions were expressed as percentage (%) on total true protein content (g/l). Caseins: sum of total casein fraction; whey protein: sum of total whey fractions.

* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

Acknowledgments

The authors thank the Genesis Project SRL (Castelnuovo Sotto, Reggio Emilia, Italy), and the managers of the three dairy farms for their help with the study and for providing data on their cows.

Declaration of interest

No conflict of interest is associated with this publication.

Ethics statement

The cows included in this study were not subjected to any invasive procedures. Milk samples were collected during the routine milk recording coordinated by authorized technicians.

Software and data repository resources

No data repository resources are available for this publication.

References

Ali AKA, Shook GE, Gabler FR and Peters J 1980. An optimum transformation for somatic cell concentration in milk. *Journal of Dairy Science* 63, 487–490.

Akers RM 2016. Lactation and the mammary gland. Blackwell Publishing Professional, Ames, IA, USA.

Barber DG, Houlihan AV, Lynch FC and Poppi DP 2005. The influence of nutrition, genotype and stage of lactation on milk casein composition. In *Indicators of milk and beef quality* (ed. JF Hocquette and S Gigli), pp. 203–216. Wageningen Academic Publishers, Wageningen, The Netherlands.

Bionaz M and Looor JJ 2011. Gene networks driving bovine mammary protein synthesis during the lactation cycle. *Bioinformatics and Biology Insights* 5, 83–98.

Bittante G, Penasa M and Cecchinato A 2012. Invited review: genetics and modeling of milk coagulation properties. *Journal of Dairy Science* 95, 6843–6870.

Bonfatti V, Di Martino G, Cecchinato A, Degano L and Carnier P 2010. Effects of β-κ-casein (CSN2-CSN3) haplotypes, β-lactoglobulin (BLG) genotypes, and detailed protein composition on coagulation properties of individual milk of Simmental cows. *Journal of Dairy Science* 93, 3809–3817.

Coulon JB, Verdier I, Pradel P and Almerna M 1998. Effect of lactation stage on the cheesemaking properties of milk and the quality of Saint-Nectaire-type cheese. *Journal of Dairy Research* 65, 295–305.

Creamer LK, Plowman JE, Liddell MJ, Smith MH and Hill JP 1998. Micelle stability: k-casein structure and function. *Journal of Dairy Science* 81, 3004–3012.

Crudden A, Fox PF and Kelly AL 2005. Factors affecting the hydrolytic action of plasmin in milk. *International Dairy Journal* 15, 305–313.

Dechow CD, Rogers GW, Cooper JB, Phelps MI and Mosholder AL 2007. Milk, fat, protein, somatic cell score, and days open among Holstein, Brown Swiss, and their crosses. *Journal of Dairy Science* 90, 3542–3549.

Demeter RM, Markiewicz K, van Arendonk JAM, Bovenhuis H, Andren A, Bovenhuis H, van Valenberg HJF, van Arendonk JAM, Wauthy JM and Winter KA

2010. Relationships between milk protein composition, milk protein variants, and cow fertility traits in Dutch Holstein-Friesian cattle. *Journal of Dairy Science* 93, 5495–5502.
- Dezetter C, Leclerc H, Mattalia S, Barbat A, Boichard D and Ducrocq V 2015. Inbreeding and crossbreeding parameters for production and fertility traits in Holstein, Montbéliarde, and Normande cows. *Journal of Dairy Science* 98, 4904–4913.
- de Haas Y, Smolders EAA, Hoorneman JN, Nauta WJ and Veerkamp RF 2013. Suitability of cross-bred cows for organic farms based on cross-breeding effects on production and functional traits. *Animal* 7, 655–665.
- Ekstrand B and Larsson-Raźnikiewicz M 1978. The monomeric casein composition of different size bovine casein micelles. *Biochimica et Biophysica Acta* 536, 1–9.
- Hansen JV, Friggens NC, Højsgaard S, Kolver E, Pleasants A, Rook A and Beever D 2006. The influence of breed and parity on milk yield, and milk yield acceleration curves. *Livestock Science* 104, 53–62.
- Hazel AR, Heins BJ and Hansen LB 2017. Production and calving traits of Montbéliarde × Holstein and Viking Red × Holstein cows compared with pure Holstein cows during first lactation in 8 commercial dairy herds. *Journal of Dairy Science* 100, 4139–4149.
- Heck JML, Schennink A, van Valenberg HJF, Bovenhuis H, Visker MHPW, van Arendonk JAM and van Hooijdonk ACM 2009. Effects of milk protein variants on the protein composition of bovine milk. *Journal of Dairy Science* 92, 1192–1202.
- Heins BJ and Hansen LB 2012. Short communication: fertility, somatic cell score, and production of Normande × Holstein, Montbéliarde × Holstein, and Scandinavian Red × Holstein crossbreds versus pure Holsteins during their first 5 lactations. *Journal of Dairy Science* 95, 918–924.
- Heins BJ, Hansen LB and Seykora AJ 2006. Production of pure Holsteins versus crossbreds of Holstein with Normande, Montbéliarde, and Scandinavian Red. *Journal of Dairy Science* 89, 2799–2804.
- Holland B, Rahimi Yazdi S, Ion Titapiccolo G and Corredig M 2010. Short communication: separation and quantification of caseins and casein macropeptides using ion-exchange chromatography. *Journal of Dairy Science* 93, 893–900.
- Holt C and Baird L 1978. Natural variations in the average size of bovine casein micelles: I. Milks from individual Ayrshire cows. *Journal of Dairy Research* 45, 339–345.
- Ikonen T, Morri S, Tyrisevä AM, Ruottinen O and Ojala M 2004. Genetic and phenotypic correlations between milk coagulation properties, milk production traits, somatic cell count, casein content, and pH of milk. *Journal of Dairy Science* 87, 458–467.
- Jensen HB, Holland JW, Poulsen NA and Larsen LB 2012. Milk protein genetic variants and isoforms identified in bovine milk representing extremes in coagulation properties. *Journal of Dairy Science* 95, 2891–2903.
- Jöudu I, Henno M, Kaart T, Püssa T and Kärt O 2008. The effect of milk protein contents on the rennet coagulation properties of milk from individual dairy cows. *International Dairy Journal* 18, 964–967.
- Lopez-Villalobos N, Garrick DJ, Holmes CW, Blair HT and Spelman RJ 2000. Profitabilities of some mating systems for dairy herds in New Zealand. *Journal of Dairy Science* 83, 144–153.
- Malchiodi F, Cecchinato A and Bittante G 2014a. Fertility traits of purebred Holsteins and 2- and 3-breed crossbred heifers and cows obtained from Swedish Red, Montbéliarde, and Brown Swiss sires. *Journal of Dairy Science* 97, 7916–7926.
- Malchiodi F, Cecchinato A, Penasa M, Cipolat-Gotet C and Bittante G 2014b. Milk quality, coagulation properties, and curd firmness modeling of purebred Holsteins and first- and second-generation crossbred cows from Swedish Red, Montbéliarde, and Brown Swiss bulls. *Journal of Dairy Science* 97, 4530–4541.
- Malchiodi F, Penasa M, Tiezzi F and Bittante G 2011. Milk yield traits, somatic cell score, milking time and age at calving of pure Holstein versus crossbred cow. *Agriculturae Conspectus Scientificus* 76, 259–261.
- Maurmayr A, Cecchinato A, Grigoletto L and Bittante G 2013. Detection and Quantification of α S1-, α S2-, β -, κ -casein, α -lactalbumin, β -lactoglobulin and lactoferrin in bovine milk by reverse-phase high-performance liquid chromatography. *Agriculturae Conspectus Scientificus* 78, 201–205.
- McSweeney PLH and Fox PF 2013. *Advanced dairy chemistry. Volume 1A, Proteins: basic aspects*, 524pp. Springer. New York, NY, USA.
- Menzies KK, Lefevre C, Macmillan KL and Nicholas KR 2009. Insulin regulates milk protein synthesis at multiple levels in the bovine mammary gland. *Functional & Integrative Genomics* 9, 197–217.
- Ostensen S, Foldager J and Hermansen JE 1997. Effects of stage of lactation, milk protein genotype and body condition at calving on protein composition and renneting properties of bovine milk. *Journal of Dairy Research* 64, 207–219.
- Politis I, Ng Kwai Hang KF and Giroux RN 1989. Environmental factors affecting plasmin activity in milk. *Journal of Dairy Science* 72, 1713–1718.
- Schopen GCB, Heck JML, Bovenhuis H, Visker MHPW, van Valenberg HJF and van Arendonk JAM 2009. Genetic parameters for major milk proteins in Dutch Holstein-Friesians. *Journal of Dairy Science* 92, 1182–1191.
- Sørensen MK, Norberg E, Pedersen J and Christensen LG 2008. Invited review: crossbreeding in dairy cattle: a Danish perspective. *Journal of Dairy Science* 91, 4116–4128.
- Wedholm A, Larsen LB, Lindmark-Månsson H, Karlsson AH and Andrén A 2006. Effect of protein composition on the cheese-making properties of milk from individual dairy cows. *Journal of Dairy Science* 89, 3296–3305.
- Wedholm A, Møller HS, Stensballe A, Lindmark-Månsson H, Karlsson AH, Andersson R, Andrén A and Larsen LB 2008. Effect of minor milk proteins in Chymosin separated whey and casein fractions on cheese yield as determined by proteomics and multivariate data analysis. *Journal of Dairy Science* 91, 3787–3797.
- William RL and Pollak E 1985. Theory of heterosis. *Journal of Dairy Science* 68, 2411–2417.