

Black soldier fly as dietary protein source for broiler quails: meat proximate composition, fatty acid and amino acid profile, oxidative status and sensory traits

M. Cullere¹, G. Tasoniero¹, V. Giaccone¹, G. Acuti², A. Marangon³ and A. Dalle Zotte^{1†}

¹Department of Animal Medicine, Production and Health, University of Padova, Agripolis, Viale dell'Università 16, 35020 Legnaro, Padova, Italy; ²Department of Veterinary Medicine, University of Perugia, Via San Costanzo 4, 06126 Perugia, Italy; ³Laboratorio Analisi Sensoriale, Veneto Agricoltura – Istituto per la Qualità e le Tecnologie Agroalimentari, Via S. Gaetano 74, 36016 Thiene, Vicenza, Italy

(Received 4 April 2017; Accepted 29 June 2017; First published online 24 July 2017)

*In the perspective of improving the sustainability of meat production, insects have been rapidly emerging as innovative feed ingredient for some livestock species, including poultry. However, at present, there is still limited knowledge regarding the quality and sensory traits of the derived meat. Therefore, the present study tested the effect of a partial substitution of soya bean meal and oil with defatted black soldier fly (*Hermetia illucens*) larvae meal (H) in the diet for growing broiler quails (*Coturnix coturnix japonica*) on meat proximate composition, cholesterol, amino acid and mineral contents, fatty acid profile, oxidative status and sensory characteristics. To this purpose, three dietary treatments were designed: a control diet (C) and two diets (H1 and H2) corresponding to 10% and 15% H inclusion levels, respectively, were fed to growing quails from 10 to 28 days of age. At 28 days of age, quails were slaughtered and breast meat was used for meat quality evaluations. Meat proximate composition, cholesterol content and oxidative status remained unaffected by H supplementation as well as its sensory characteristics and off-flavours perception. Differently, with increasing the dietary H inclusion, the total saturated fatty acid and total monounsaturated fatty acid proportions raised to the detriment of the polyunsaturated fatty acid fraction thus lowering the healthiness of the breast meat. The H2 diet increased the contents of aspartic acid, glutamic acid, alanine, serine, tyrosine and threonine thus further enhancing the biological value of the meat protein. As a direct result of the dietary content of Ca and P, the meat of quails fed with the highest H level, displayed the highest Ca and the lowest P values. Therefore, meat quality evaluations confirmed H to be a promising insect protein source for quails. The only potential drawback from feeding H to broiler quails regarded the fatty acid profile of the meat, therefore requiring further research efforts to understand to what extent the fatty acid profile of H can be improved.*

Keywords: insect meal, quail, meat quality, fatty acids, amino acids

Implications

Reducing the feed-food competition is a critical aspect for modern livestock production and will be further stressed in the near future due to the growing world population. Therefore, the search for alternatives to the common feed ingredients represent a key aspect for the future sustainability of the livestock sector and, in this context, insects are candidates with great potential. This study showed that insect meal from *Hermetia illucens* larvae can partly substitute conventional ingredients in the diet for broiler quails as their meat showed satisfactory nutritional quality. However, research to improve the fatty acid profile of larvae is required.

Introduction

In the perspective of a rising world population with growing meat requirements, insects are considered an alternative feed source for livestock, which would offer a possible solution to improve the sustainability of the livestock supply chain and thus meat production (Sánchez-Muros *et al.*, 2014). Insects constitute a raw material that is included in the European Union feed material register and, although they are currently authorized only for pets and aquaculture animals, insect-derived feeds could also represent a suitable ingredient for pigs and poultry feed manufacturing. To allow a clear legislation and possible rapid industrial development of the sector, research efforts are currently being performed.

One of the most promising insect species is the black soldier fly (*H. illucens*), whose larvae can successfully grow

† E-mail: antonella.dallezotte@unipd.it

on substrates made of various by-products such as digestate, vegetable and restaurant leftover (Spranghers *et al.*, 2017). This aspect is interesting as it allows to transform potential wastes with an environmental impact into a nutrient rich product, and the final substrate can be successfully used as organic fertilizer (Lalander *et al.*, 2016). In addition, this new feed source would help in reducing the feed-food competition which is already a critical aspect for modern livestock production and which will be further stressed in the near future due to the growing world population. Regarding the poultry sector, soy is one of the main dietary ingredients and it is used both as a protein and a fat source but it is also a food for humans. Soy requires a lot of land and water to grow and therefore has a relevant environmental impact (Charlton *et al.*, 2015). Due to this, soy supplies represent a key aspect now and even more for the future. In fact, current poultry diets of the European Union use soy which is mainly imported from United States, Argentina and Brazil (Boerema *et al.*, 2016), highlighting once more that the search for alternative feed ingredients is imperative.

Black soldier fly has recently been successfully tested as feed ingredient in conventional poultry diets as fat source (Schiavone *et al.*, 2017) and in laying hens as a protein meal (Al-Qazzaz *et al.*, 2016; Maurer *et al.*, 2016). In addition, in the only study on meat quails, defatted black soldier fly larvae meal provided satisfactory results in terms of nutrient digestibility, carcass composition and meat physical traits (Cullere *et al.*, 2016).

However, the impact of this new feed ingredient on the meat quality is a key aspect for consumers that will directly influence the concrete interest by the industry in this new feed source.

Based on these considerations, the present research studied the effect of a dietary inclusion of defatted insect meal from *H. illucens* larvae on several meat quality aspects of broiler meat quails: meat proximate composition, cholesterol content, oxidative status, fatty acid profile, amino acid and mineral contents, sensory profile and off-odours and flavours perception.

Material and methods

Animals and diets

The experiment was carried out in a private quail farm ("La Colombara" Società Agricola, Castelnuovo di Isola Vicentina, VI, Italy), after the approval by the veterinary authority and in accordance with the article 2, DL 4 March 2014, No. 26 of the Official Journal of the Italian Republic (<http://www.gazzettaufficiale.it/eli/id/2014/03/14/14G00036/sg>), implementing the EC Directive 86/609/2010 EU regarding the protection of animals used for experimental and other scientific purposes.

A total of 450 quails, 10-day-old (*Coturnix coturnix japonica*), of both sexes were housed in batteries in an environmentally controlled room. The chicks were allocated by 30 in 15 cages and received three dietary treatments (five replicates per treatment) until slaughtering: a control diet (C) which was formulated referring to the common grower diet

used in the farm, H1 and H2 diets in which conventional protein/fat sources were partly substituted with defatted black soldier fly (*H. illucens*) larvae meal (H): 10% H for H1 and 15% H for H2. In H1, H replaced 28.4% of soya bean oil and 16.1% of soya bean meal, whereas in H2, H substituted 100% of soya bean oil and 24.8% of soya bean meal. Mashed feeds and water were provided *ad libitum*. All diets were formulated to meet the minimum requirements for Japanese quails (National Research Council, 1994). A detailed description of the origin, the chemical composition and the energy content of H, as well as the ingredients, the chemical composition and the energy content of experimental diets are reported elsewhere (Cullere *et al.*, 2016).

Slaughtering, carcass dissection and samples preparation

At 28 days of age, quails were deprived from feed and transported to the commercial slaughterhouse 'Quaja Veneta' Società Agricola (Malo, VI, Italy), which is located 8 km far from the farm. Fasting lasted 6 h, from feed withdrawal until slaughtering. Quails were electrically stunned, slaughtered and processed following commercial conditions. Stunned quails were bled, plucked, eviscerated and freed from head, neck, shanks and abdominal fat. Afterwards, they were refrigerated for 1 h in the refrigeration tunnel at +2°C, transported in chilled conditions to the Department of Animal Medicine, Production and Health (MAPS) of the University of Padova and stored at +2°C. The following day, breast muscles (right and left side) were excised from 96 quails/treatment and divided in accordance with the subsequent analytical step.

A total of 60 breasts/treatment ($n = 12$ breasts/replicated cage) were vacuum-packaged by 10 using a CSV-41n ORVED machine (99% vacuum level) in polyethylene bags (water vapour transmission rate: $3.5 \pm 1 \text{ g/m}^2 \text{ day}$ at 23°C and $85 \pm 2\%$ relative humidity), and stored at -40°C for subsequent analysis: lipid oxidation ($n = 10$ breasts/treatment), amino acid profile ($n = 10$ breasts/treatment) and sensory analysis ($n = 40$ breasts/treatment). The other $n = 36$ quail breasts/treatment ($n = 7, 7, 7, 7$ and 8 breasts/replicated cage) were randomly ground by three (to have enough sample to perform all the scheduled analysis) with a Retsch Grindomix GM 200 (7000 g for 10 s) for a total of 12 samples/treatment. Subsequently, meat samples were frozen at -40°C, freeze-dried and ground again (7000 g for 5 s) to obtain a fine powder which was used to determine proximate and fatty acid composition, cholesterol and mineral contents.

Meat proximate composition, cholesterol content and oxidative status

The proximate composition of meat was analysed in accordance with the AOAC (1995), whereas protein content, including glucidic molecules and their catabolites (0.25%), was calculated by difference (Ouhayoun *et al.*, 1990).

The cholesterol content was determined through absolute quantitative analysis using HPLC following the method described by Casiraghi *et al.* (1994).

After 2 weeks of storage, 10 quail breasts/treatment were allowed thawing for 12 h at +4°C, freed from polyethylene bags and individually ground with a Retsch Grindomix GM 200 (7000 g for 10 s). The extent of muscle lipid oxidation was evaluated with a spectrophotometer (Hitachi U-2000; Hitachi, Mannheim, Germany) set at 532 nm, that measured the absorbance of thiobarbituric acid-reactive substances and a 1,1,3,3-tetraethoxypropane calibration curve (Botsoglou *et al.*, 1994). Oxidation products were quantified as malondialdehyde (MDA) equivalents (mg MDA/kg muscle).

Mineral and amino acid contents of defatted *Hermetia illucens* meal, diets and meat

Aliquots of H, experimental diets and 12 freeze-dried meat samples/treatment were sent to the Department of Veterinary Medicine, University of Perugia to analyse the mineral content (AOAC, 2012). The methods to analyse the mineral content of the H and the experimental diets were: method 968.08 (Ca and Mg), method 995.11 (P), method 956.01 (Na), method 975.03 (K), method 968.08 (Cu, Fe, Zn and Mn). Differently, the methods used for meat samples were: method 991.25 (Ca and Mg), method 991.27 (P), method 956.01 (Na), method 975.03 (K, Cu, Fe, Zn) and method 921.02 (Mn).

The amino acid contents of the H, the experimental diets and the meat were analysed by EPTA NORD srl (Via Padova, Conselve, Italy, Internal method 693 rev 0 2007).

The mineral and the amino acid contents of the H and the experimental diets are presented in Table 1.

Fatty acid profile of defatted *Hermetia illucens* meal, diets and meat

Lipid extraction was performed combining the traditional Folch method (Folch *et al.*, 1957) with that provided by Lee *et al.* (1996) and the Accelerated Solvent Extraction (M-ASE), in which petroleum ether (H and experimental diets) or chloroform/methanol 1 : 2 (meat) were the solvents used for the extraction. Total lipid content was determined gravimetrically after the removal of the solvent by evaporation under nitrogen stream at 50°C. Samples were then transmethylated using a methanolic solution of H₂SO₄ (4%) to determine fatty acid methyl esters (FAME). Biphasic separation was obtained by adding 0.5 ml of distilled water and 1.5 ml of *N*-heptane to each sample. Fatty acid methyl esters were quantified by gas chromatography (Shimadzu GC17A, equipped with an Omegawax (Sigma-Aldrich Co. LLC., Saint Louis, USA) 250 column (30 m × 0.25 µm × 0.25 µm)) and flame ionization detector. Helium was used as carrier gas at a constant flow of 0.8 ml/min. Injector and detector temperatures were 260°C. Peaks were identified based on commercially available FAME mixtures (37-Component FAME Mix; Supelco Inc., Bellefonte, PA, USA) and the data obtained were expressed as % of total detected FAME.

The fatty acid profiles of the H and the experimental diets are shown in Table 2.

Meat sensory analysis

Quail breasts were subjected to a descriptive sensory analysis, to detect possible differences among the dietary treatments

Table 1 Mineral (mg/kg) and amino acid contents (g/100 g as fed) of the defatted *Hermetia illucens* larvae meal (H) and experimental diets for growing quails

	<i>Hermetia illucens</i>	Experimental diets		
	meal	Control	H1	H2
	H			
Minerals				
Ca	9770	4800	5200	5500
P	8329	6070	5960	5930
Na	1812	1740	1640	1720
Mg	2507	2280	2120	2160
K	11 541	10 200	9600	8400
Cu	25	36	34	35
Fe	297	208	204	206
Zn	96	82	78	74
Mn	165	80	78	82
Indispensable amino acids				
Arginine	1.64	1.02	0.91	0.86
Histidine	0.47	0.18	0.18	0.18
Isoleucine	2.24	0.95	0.92	0.92
Leucine	3.30	1.76	1.69	1.71
Lysine	1.96	0.33	0.37	0.46
Methionine	0.62	0.15	0.17	0.21
Phenylalanine	1.69	0.84	0.79	0.80
Threonine	1.93	0.91	0.89	0.91
Valine	3.58	1.22	1.26	1.32
Dispensable amino acids				
Alanine	4.64	1.51	1.63	1.80
Aspartic acid	4.49	2.37	2.23	2.16
Cysteine	0.09	0.11	0.09	0.10
Glutamic acid	4.52	3.62	3.35	3.09
Glycine	4.43	1.64	1.61	1.73
Proline	2.91	1.35	1.37	1.40
Serine	2.13	1.36	1.25	1.23
Tryptophan	0.02	1.40	1.16	1.30
Tyrosine	2.70	1.11	1.11	1.19
Total	43.4	21.8	21.0	21.4

H1 and H2 are diets corresponding to 10% and 15% H inclusion levels, respectively.

(C v. H1 v. H2). The sensory analysis was performed by an eight-member trained panel, which were qualified as experts according to ISO 8586 and had experience with descriptive tests (ISO 13299) on various food matrixes. All judges who perform tests with accredited methods undergo training every 3 years. Panelists underwent two pre-test training sessions of 1 h each to familiarize with the matrix and select appropriate descriptors, possible off-odours and off-flavours, also drawn from the literature. The panel received a list of descriptors to score on numerical and continuous scales from 0 (the lowest score for each attribute) to 10 (the highest score for each attribute).

Olfactory, gustative and textural aspects were evaluated. The descriptors were: odour intensity, off-odour intensity, flavour intensity, off-flavour intensity, juiciness, toughness,

chewiness and fibrousness. The chosen off-odours and off-flavours were: game meat, liver, oil/fat, peanut/hazelnut. The meat used for the training sessions was that of the C group ($n = 8$ quail breasts) and was processed, stored, handled and cooked in the same manner of the samples which were used for the subsequent sensory analysis.

For the experiment, a total of 32 quail breasts/treatment were used and 2 days of analysis were scheduled (16 quail breasts/treatment per session). After 1 month of frozen storage at -40°C , quail breasts were allowed thawing for 16 h at $+4^{\circ}\text{C}$. Each sample was then placed in a cooking plate (model GR6010 XL Health Comfort, 2400 Watt; Rowenta, Erbach, Germany) set at thermostat position '2' and cooked 5 min/side, until core temperature reached 74°C . Subsequently, samples were put in aluminium trays and served to the panel in random sequence. Samples were identified by a random three-digit code.

The evaluation sheet, distribution of samples to the judges and data acquisition were performed using FIZZ software

Table 2 Fatty acids (FA) profile (% of total fatty acids) of the defatted *Hermetia illucens* larvae meal (H) and experimental diets

	Insect meal	Experimental diets		
	H	Control	H1	H2
C10:0	1.14	0.00	0.46	0.63
C12:0	42.8	0.11	16.9	23.6
C14:0	8.12	0.23	3.44	4.81
C15:0	–	0.14	0.00	0.00
C16:0	13.9	14.7	13.7	13.9
C17:0	0.16	0.12	0.12	0.13
C18:0	2.86	3.93	3.21	2.90
C20:0	0.34	0.31	0.25	0.28
C22:0	–	0.00	0.09	0.00
C24:0	–	0.19	0.12	0.10
SFA	69.4	19.7	38.3	46.3
C14:1	0.15	0.00	0.00	0.10
C16:1	2.06	0.26	0.99	1.32
C17:1	–	0.15	0.13	0.17
C18:1n-9	10.4	23.6	19.4	17.9
C18:1n-11	0.31	1.24	0.84	0.64
C20:1n-9	0.26	0.00	0.00	0.00
C22:1n-9	–	0.00	0.10	0.00
MUFA	13.2	25.3	21.5	20.1
C18:2n-6	12.6	47.3	35.3	30.4
C20:4n-6	0.17	0.00	0.00	0.00
PUFAn-6	12.8	47.3	35.3	30.4
C18:3n-3	1.16	4.05	2.66	1.80
C20:5n-3	0.21	0.30	0.20	0.10
PUFAn-3	1.37	4.35	2.86	1.90
PUFA	14.1	51.7	38.2	32.3
PUFAn-6/n-3	9.31	10.9	12.4	16.0
UFA/SFA	0.39	3.91	1.56	1.13
Identified FA (%)	96.7	96.6	98.0	98.7

SFA = saturated fatty acids; MUFA = monounsaturated fatty acids; PUFA = polyunsaturated fatty acids; UFA = unsaturated fatty acids. H1 and H2 are diets corresponding to 10% and 15% H inclusion levels, respectively.

(BIOSYSTEMES FRANCE, St-Ouen l'Aumône, France) installed in eight terminals in the tasting booths of the laboratory. In addition, for each sample, panelists were asked to indicate if and which of the listed off-odours and off-flavours they could recognize. Still water at room temperature and unsalted crackers were available to panelists throughout each sensory session.

Statistical analysis

Meat proximate composition, oxidative status, cholesterol, mineral and amino acid contents and fatty acid profile data were subjected to a one-way ANOVA with experimental diet (C, H1 and H2) as fixed effect, following the GLM procedure of the SAS 9.1.3 Statistical Analysis Software for Windows (SAS Institute, 2008). A mixed model (PROC MIXED) was used to detect any dietary influence on sensory analysis scores, therefore considering experimental diet and the eight panelists as fixed and random effects, respectively. Least square means were obtained using Bonferroni test and the significance was calculated at a 5% confidence level. A χ^2 test with Marascuilo (1966) procedure was performed on off-odours and off-flavours characterization to detect the differences among treatments.

Results

Meat chemical composition and oxidative status

Independently to the inclusion level, H did not affect the proximate composition, cholesterol content and oxidative status of quail breast meat (Table 3) which exhibited an average 18.5% protein, 4.6% lipids, 73.3 mg/100 g meat cholesterol, and 0.36 mg MDA/kg meat.

The mineral profile of the meat (Table 4) was comparable among treatments with the exceptions of Ca and K contents which followed the same trend displayed by the diets. The meat content of Ca linearly increased from C to H2 meat (515, 554 and 586 mg/kg for C, H1 and H2 meat,

Table 3 Effect of the dietary inclusion of the defatted *Hermetia illucens* larvae meal (H) on the proximate composition (%), cholesterol content (mg/100 g meat) and oxidative status (mg MDA/kg meat) of Japanese quail breast meat

	Experimental groups			P-value	rSD
	Control	H1	H2		
No. ¹	12	12	12		
Water	75.3	75.4	75.4	0.8101	0.30
Protein	18.4	18.5	18.5	0.3810	0.32
Lipids	4.62	4.52	4.56	0.7035	0.28
Ash	1.71	1.54	1.62	0.2272	0.23
Cholesterol	71.6	73.3	74.9	0.1954	4.34
TBARs ²	0.36	0.35	0.36	0.2357	0.04

MDA = malondialdehyde; TBARs = thiobarbituric acid-reactive substances. H1 and H2 are diets corresponding to 10% and 15% H inclusion levels, respectively.

¹Each sample is made from three quail breasts.

²TBARs analysis was conducted on No. = 10 samples/treatment.

Table 4 Effect of the dietary inclusion of the defatted *Hermetia illucens* larvae meal (H) on the mineral content (mg/kg) of Japanese quail breast meat

	Experimental groups			P-value	rSD
	Control	H1	H2		
No. ¹	12	12	12		
Ca	515 ^b	554 ^{ab}	586 ^a	0.0297	13.7
P	8930	8881	8997	0.2586	112
Mg	1092	1079	1083	0.7513	11.9
Na	2488	2319	2400	0.4802	75.6
K	14 925 ^A	14 473 ^{AB}	13 958 ^B	0.0032	184
Cu	4.71	4.44	4.64	0.7069	0.24
Fe	47.2	50.7	47.2	0.3877	2.05
Zn	23.1	23.2	22.2	0.3342	0.54
Mn	Nd	Nd	Nd	–	–

Nd = not detected.

H1 and H2 are diets corresponding to 10% and 15% H inclusion levels, respectively.

^{a,b}Values within a row with different superscripts differ significantly at $P < 0.05$.

^{A,B}Values within a row with different superscripts differ significantly at $P < 0.01$.

¹Each sample is made by three quail breasts.

respectively; $P < 0.05$), whereas K content followed an opposite trend (14 925, 14 473 and 13 958 mg/kg for C, H1 and H2 meat, respectively; $P < 0.01$).

From the results displayed in Table 5, it was observed that the 10% and 15% inclusion of H in quail diets provided meat with satisfactory amino acid profile, which resulted improved ($P < 0.05$) for aspartic acid, glutamic acid, alanine, serine, tyrosine and threonine in H2 group compared to C.

Fatty acid profile of *Hermetia illucens* meal and quail meat

The fatty acid profile of H was mostly represented by saturated fatty acids (SFA) which accounted for more than 69% of total fatty acids with lauric (C12:0, 42.8%), palmitic (C16:0, 13.9%) and myristic (C14:0, 8.1%) fatty acids being the most representative. Oleic (C18:1n-9, 10.4%) and linoleic (C18:2n-6, 12.6%) fatty acids were the most abundant monounsaturated fatty acids (MUFA) and polyunsaturated fatty acids (PUFA), respectively. Consequently, the dietary inclusion of H greatly changed the fatty acid proportions of quail breast meat (Table 6). With increasing H level, SFA increased (34.6 v. 37.3 and 37.9% for C, H1 and H2, respectively; $P < 0.0001$) because of raising C10:0, C12:0, C14:0, C16:0 and C20:0 fatty acids. A similar trend was observed for MUFA, which increased from C to H2 meat (13.3 v. 14.6 v. 16.3% for C, H1 and H2, respectively; $P < 0.0001$). On the opposite, PUFA significantly decreased from C to H2 meat (42.7 v. 40.3 v. 38.0% for C, H1 and H2, respectively; $P < 0.0001$) with the PUFA_{n-3} fraction showing the highest decrease. In fact, H2 meat had 41% less n-3 than C. The latter determined an increasing trend in the PUFA_{n-6/n-3} ratio from C to H2 meat (9.34 v. 11.5 v. 14.7, for C, H1 and H2 meat; $P < 0.0001$).

Table 5 Effect of the dietary inclusion of the defatted *Hermetia illucens* larvae meal (H) on the amino acids content (g/100 g) of Japanese quail breast meat

	Experimental groups				
	Control	H1	H2	P-value	rSD
No.	10	10	10		
Indispensable amino acids					
Arginine	1.11	1.12	1.18	0.0700	0.032
Histidine	0.03	0.04	0.04	0.0787	0.003
Isoleucine	1.09	1.10	1.14	0.1761	0.034
Leucine	1.77	1.79	1.88	0.1214	0.058
Lysine	0.46	0.48	0.52	0.0783	0.029
Methionine	0.49	0.49	0.52	0.4699	0.031
Phenylalanine	0.53	0.55	0.55	0.8689	0.040
Threonine	1.12 ^b	1.13 ^{ab}	1.20 ^a	0.0173	0.026
Valine	1.34	1.36	1.42	0.0841	0.037
Dispensable amino acids					
Alanine	1.89 ^b	1.90 ^{ab}	2.03 ^a	0.0309	0.052
Aspartic acid	2.21 ^b	2.23 ^b	2.37 ^a	0.0118	0.048
Cystine	0.12	0.12	0.13	0.8503	0.008
Glutamic acid	3.54 ^b	3.57 ^{ab}	3.78 ^a	0.0343	0.091
Glycine	1.73	1.71	1.85	0.1314	0.080
Proline	0.92	0.91	0.83	0.7157	0.156
Serine	1.18 ^{ab}	1.16 ^b	1.23 ^a	0.0422	0.028
Tryptophan	0.10	0.10	0.09	0.7290	0.015
Tyrosine	1.85 ^b	1.88 ^{ab}	1.99 ^a	0.0279	0.049
Total	21.5	21.6	22.8	–	–

H1 and H2 are diets corresponding to 10% and 15% H inclusion levels, respectively.

^{a,b}Values within a row with different superscripts differ significantly at $P < 0.05$.

Meat sensory profile, off-odours and off-flavours

Results depicted in Table 7, showed that odour, flavour and texture attributes of quail meat deriving from H1 and H2 groups was not affected by the H inclusion level and resulted like the scores of C group. In addition, the average intensity perception of off-odour (score: 1.24 out of 10) and off-flavour (score: 1.22 out of 10) was very low. Regarding specific off-odours and off-flavours (Table 8), liver was the most frequently perceived (41% and 31% of the tested samples for off-odours and off-flavours, respectively) but this was independent to the treatment, thus suggesting it as a typical sensory characteristic of quail meat.

Discussion

The proximate composition and cholesterol content of quail meat in the present experiment corresponded to values found in literature (Güven *et al.*, 2015), with cholesterol being quite high if compared with data reported for some common meat species (Dalle Zotte and Szendrő, 2011) namely beef (48.7 mg/100 g *Longissimus dorsi* meat), pork (62 mg/100 g *Longissimus dorsi* meat) and chicken (55.3 mg/100 g breast meat). Even if some recent research papers dealt with the effect of feeding insects to poultry species, knowledge on the quality of the derived meat is still scarce

Table 6 Effect of the dietary inclusion of the defatted *Hermetia illucens* larvae meal (H) on the fatty acid (FA) profile (% of total fatty acids) of Japanese quail breast meat

	Experimental groups			P-value	rSD
	Control	H1	H2		
No. ¹	12	12	12		
C10:0	0.00 ^B	0.14 ^A	0.13 ^A	<0.0001	0.017
C12:0	0.00 ^C	1.26 ^B	2.10 ^A	<0.0001	0.272
C14:0	0.25 ^C	0.92 ^B	1.34 ^A	<0.0001	0.133
C16:0	17.8 ^B	18.8 ^A	19.0 ^A	0.0006	0.731
C17:0	0.19	0.18	0.17	0.6217	0.038
C18:0	14.3	14.1	13.5	0.0757	0.877
C20:0	0.03 ^B	0.07 ^{AB}	0.13 ^A	0.0024	0.069
C22:0	0.36	0.41	0.33	0.7725	0.250
C24:0	1.66 ^{Aa}	1.50 ^{Ab}	1.17 ^{Bab}	<0.0001	0.131
SFA	34.6 ^B	37.3 ^A	37.9 ^A	<0.0001	1.034
C14:1	0.00 ^C	0.08 ^B	0.18 ^A	<0.0001	0.050
C16:1	1.25 ^{Bc}	1.78 ^{ABb}	2.22 ^{Aa}	<0.0001	0.416
C17:1	0.16	0.16	0.14	0.4839	0.042
C18:1n-9	10.2 ^{Bab}	11.0 ^{ABb}	12.1 ^{Aa}	<0.0001	0.928
C18:1n-11	1.73	1.65	1.65	0.5001	0.200
MUFA	13.3 ^{Bab}	14.6 ^{ABb}	16.3 ^{Aa}	<0.0001	1.382
C18:2n-6	26.5	25.4	25.1	0.0608	1.468
C20:2n-6	0.53 ^A	0.17 ^B	0.30 ^B	<0.0001	0.174
C18:3n-6	0.03 ^B	0.20 ^A	0.24 ^A	<0.0001	0.049
C20:3n-6	0.65 ^C	0.81 ^B	1.00 ^A	<0.0001	0.116
C20:4n-6	10.9 ^A	10.5 ^A	9.0 ^B	<0.0001	0.893
PUFAn-6	38.0 ^A	36.9 ^{AB}	35.3 ^B	0.0035	1.818
C20:3n-3	0.03	0.01	0.00	0.5259	0.051
C18:3n-3	1.20 ^A	0.87 ^B	0.67 ^C	<0.0001	0.110
C20:5n-3	0.24	0.18	0.18	0.0617	0.066
C22:6n-3	2.63 ^A	2.17 ^B	1.56 ^C	<0.0001	0.258
PUFAn-3	4.09 ^A	3.22 ^B	2.42 ^C	<0.0001	0.290
PUFA	42.7 ^{Aa}	40.3 ^{ABb}	38.0 ^{Bc}	<0.0001	1.931
PUFAn-6/n-3	9.34 ^C	11.5 ^B	14.7 ^A	<0.0001	0.889
UFA/SFA	1.62 ^A	1.47 ^B	1.43 ^B	<0.0001	0.054
Identified FA (%)	90.6	92.2	92.2		

SFA = saturated fatty acids; MUFA = monounsaturated fatty acids; PUFA = polyunsaturated fatty acids; UFA = unsaturated fatty acids.

H1 and H2 are diets corresponding to 10% and 15% H inclusion levels, respectively.

^{a,b,c}Values within a row with different superscripts differ significantly at $P < 0.05$.

^{A,B,C}Values within a row with different superscripts differ significantly at $P < 0.01$.

¹Each sample is made by three quail breasts.

(Maurer *et al.*, 2016). In the present study, it was demonstrated that the proximate composition, cholesterol content and oxidative status of insects-fed quails is comparable with birds fed on conventional diets, which is a key aspect from the nutritional point of view.

In general, the mineral content of H was in line to data reported by Makkar *et al.* (2014), but slightly higher than literature values reported for whole insects (Finke, 2013), which was probably due to the defatting procedure which presumably concentrated the minerals. *Hermetia illucens* is a good source of Ca and P, which is in accordance with the present experiment, especially for Ca. Despite K content of H

Table 7 Effect of the dietary inclusion the defatted *Hermetia illucens* larvae meal (H) on the sensory profile scores of Japanese quail breast meat

	Experimental groups				
	Control	H1	H2	P-value	rSD
No.	32	32	32		
Odour					
Odour intensity	5.73	5.61	5.45	0.5933	1.08
Off-odour intensity	1.36	1.12	1.25	0.7147	1.19
Flavour					
Flavour intensity	5.68	5.57	5.41	0.5077	0.94
Off-flavour intensity	1.33	1.19	1.13	0.6967	1.00
Texture					
Juiciness	4.42	4.29	4.34	0.9143	1.28
Toughness	4.47	4.69	4.63	0.6390	0.97
Chewiness	5.38	5.19	5.33	0.8727	1.50
Fibrousness	5.00	5.00	5.00	0.9999	1.11

$n = 8$ breasts of the C group were used for the training sessions; H1 and H2 are diets corresponding to 10% and 15% H inclusion levels, respectively.

Table 8 Effect of the dietary inclusion of the defatted *Hermetia illucens* larvae meal (H) on the off-odours and flavours perception (% on total tested samples) of Japanese quail meat

	Experimental groups				P-value	χ^2
	Control	H1	H2			
No.	32	32	32			
Off-odours						
Game meat	21.9	15.6	25.0	0.6427	0.88	
Liver	37.5	40.6	43.8	0.8785	0.26	
Oil/fat	12.5	9.38	6.25	0.6922	0.74	
Off-flavours						
Game meat	12.5	9.4	12.5	0.9024	0.21	
Liver	37.5	28.1	28.1	0.6464	0.87	
Oil/fat	12.5	9.4	9.4	0.8944	0.22	
Peanut/hazelnut	0.00	3.13	3.13	0.3120	1.02	

$n = 8$ breasts of the C group were used for the training sessions; H1 and H2 are diets corresponding to 10% and 15% H inclusion levels, respectively.

showed to be about 67% higher than values showed by Makkar *et al.* (2014), it was not enough to provide a comparable K content in the three experimental diets, which reflected also in the meat content of this mineral.

The result of the amino acid profile of H, confirmed that this insect provides protein with high biological value, in line with literature data (Józefiak *et al.*, 2016), but which is, once again, highly dependent on the chemical composition of the growing substrate (Tschirner and Simon, 2015). The fact that the inclusion of H in the diet for growing quails increased the breast meat content of threonine, was not dependent on the content of this amino acid in the experimental diets. Except for alanine, also for the dispensable amino acids aspartic acid, glutamic acid, serine and tyrosine, their meat content increased with increasing the dietary H inclusion, but this was independent to the absolute amount of these amino acid

in the experimental diets. This was not surprising as growing evidence suggests that the content and proportion of the amino acids of the body does not directly correspond to the dietary ones, with discrepancies being particularly high for arginine, cysteine, glutamic acid, glycine, histidine, methionine, proline and serine. This is due to several factors as the different rates at which, in the small intestine, individual amino acids are catabolized and transformed. In addition, the amino acid composition in the diet does not correspond to their concentration in the blood circulation, individual amino acids in the plasma have a different metabolic destiny in different tissues and the pattern of dietary amino acids differs to that of amino acids in tissue proteins (Guoyao *et al.*, 2014). In general, however, it could be observed that breast meat of H2 quails showed an enhanced amino acid profile compared with that of C quails.

Black soldier fly was tested also in another recent trial on poultry but only as a fat substitute, where it replaced the soya bean oil (Schivone *et al.*, 2017). Results were overall comparable with those of the conventional diet with the only drawback related to the fatty acid profile of the derived meat which is exactly what it was observed in the present study.

In fact, increasing the content of H lowered the healthiness of the meat as SFA increased to the detriment of PUFA. Both the defatted larvae meal tested in the present trial and the black soldier fat used in the trial by Schivone *et al.* (2017) originated from larvae which were reared on a conventional substrate which was layer mash, therefore the two similar results on the fatty acid profile of quail and chicken breast meat, respectively, were expected. Interestingly, both the present and the above cited studies showed that a decrease in the dietary content of MUFA did not correspond to the same trend when considering meat. In fact, in the present experiment an increase in meat MUFA content with H inclusion was observed. The latter can be ascribable to elongase and $\Delta 9$ desaturase activities mainly from C12:0, C14:0 and C16:0 SFA whose content increased with increasing dietary H. The $\Delta 9$ desaturase is a lipogenic enzyme which was reported to be up-regulated by low-fat high-carbohydrate diets and down-regulated by the dietary addition of PUFA (Poureslami *et al.*, 2010). As C, H1 and H2 diets showed a decrease in crude fat content, an increase in nitrogen-free extracts (Cullere *et al.*, 2016) and a reduction in PUFA, a higher $\Delta 9$ desaturase activity was hypothesized in H supplemented diets compared to C.

Based on results of the present trial, *H. illucens* can therefore not be considered a feedstuff with a healthy fatty acid profile namely low in SFA and high in PUFA, and with a favourably low n-6/n-3 ratio (Karami *et al.*, 2013). However, as larvae are monogastric animals, a possible solution to solve this key issue could be the modulation of the rearing substrate aiming to modify the fatty acid profile of black soldier fly. In fact, a recent study by Spranghers *et al.* (2017) showed that the ether extract content of the larvae can vary greatly depending to the substrate, as well as the fatty acid profile. Similarly, also the study by Tschirner and Simon

(2015) observed that mixture of middlings, dried distillers' grains with solubles and dried sugar beet pulp, greatly changed the chemical composition of the larvae. Furthermore, farming black soldier fly larvae on fish offal was shown to provide a feedstuff substantially enriched in n-3 PUFA (St-Hilaire *et al.*, 2007). Therefore, it should be stressed that the research of the optimal feeding formulation for growing larvae is a key point for both a successful larvae production in terms of quantity and quality, as well as for a satisfactory nutritional quality of the meat derived from insects' fed animals.

In the perspective of a potential successful application of insects-based feed to livestock animals and thus to be fruitfully introduced in the meat sector, consumers must be willing to accept insects as feed ingredients, and the sensory characteristics of meat deriving from insects-fed animals must be comparable with the conventional one. A recent study by Laureati *et al.* (2016) showed that only 21.7% of Italian consumers recruited for the test were unwilling to accept insects as feed for livestock farming and that information plays a crucial role in driving the change towards acceptance. The present research is the first one in assessing the sensory attributes of meat coming from quails fed with black soldier fly larvae meal. Our findings agreed to those of Sealey *et al.* (2011) on rainbow trout fed with increasing levels of black soldier fly larvae as a replacement of fish meal, in which untrained panelists did not detect any significant difference in a blind comparison among the treatments. Differently, a recent study by Borgogno *et al.* (2017) found that a dietary inclusion of *H. illucens* larvae meal induced significant differences in the sensory description of rainbow trout fish samples. However, similarly to our research, no defects or off-flavours were perceived in samples deriving from fish fed with *H. illucens*. Finally, Pieterse *et al.* (2014) found that the sensory profile of meat derived from chickens fed with a 10% *Musca domestica* larvae meal was slightly different from the control group for a higher perception of metallic aroma and aftertaste but a higher sustained juiciness and a lower mealiness (dry sensation) in the mouth.

In conclusion, the present research showed that a partial replacement with a defatted *H. illucens* larvae meal to the common soya bean meal and oil in the diet for growing broiler quails (up to 15% inclusion level) is technically feasible and provide meat of comparable quality to that of quails fed a conventional diet. The fact that the sensory profile of meat was not affected by H dietary inclusion is a fundamental aspect for the potential of this new feedstuff to be successfully included in commercial diets for growing quails. Differently, the fatty acid profile of the larvae must be surely improved to provide healthier meat for the modern consumer. Therefore, further studies should be planned to assess to what extent the fatty acid profile of *H. illucens* can be improved, thus allowing to design an optimal growth substrate to produce larvae with a nutritional composition in line with the required health standards.

Acknowledgements

Authors would like to thank Rino Cailotto ("La Colombara" Società Agricola, Castelnovo di Isola Vicentina, VI, Italy) for providing the quails, for the technical support and for allowing us to conduct the study in his farm. Special thanks to Quaja Veneta® Società Cooperativa Agricola (Malo, VI, Italy) for slaughtering and for the technical help which represented a fundamental basis to conduct a successful experiment. Finally, authors acknowledge the feed company FANIN Srl (San Tomio di Malo, VI, Italy) for supplying the raw materials for the diets. The research was supported by the University of Padova funds (ex 60%, 2016 – DOR1603318) and Senior Researcher Scholarship (Prot. n. 1098, 2015).

References

- Al-Qazzaz MFA, Dahlan I, Akit H and Lokman HI 2016. Effect of using insect larvae meal as a complete protein source on quality and productivity characteristics of laying hens. *Revista Brasileira de Zootecnia* 45, 518–523.
- AOAC 1995. Official methods of analysis, 15th edition. Association of Official Analytical Chemists, Arlington, VA, USA.
- AOAC 2012. Official methods of analysis, 19th edition. Association of Official Analytical Chemists, Arlington, VA, USA.
- Boerema A, Peeters A, Swolfs S, Vandevenne F, Jacobs S, Staes J and Meire P 2016. Soybean trade: balancing environmental and socio-economic impacts of an intercontinental market. *PLoS One* 11, e0155222.
- Borgogno M, Dinnella C, Iaconisi V, Fusi R, Scarpaleggia C, Schiavone A, Monteleone E, Gasco L and Parisi G 2017. Inclusion of *Hermetia illucens* larvae meal on rainbow trout (*Oncorhynchus mykiss*) feed: effect on sensory profile according to static and dynamic evaluations. *Journal of the Science of Food and Agriculture* 97, 3402–3411.
- Botsoglou NA, Fletouris DJ, Papageorgiou GE, Vassilopoulos VN, Mantis AJ and Trakatellis AG 1994. Rapid, sensitive, and specific thiobarbituric acid method for measuring lipid peroxidation in animal tissue, food, and feedstuff samples. *Journal of Agriculture and Food Chemistry* 42, 1931–1937.
- Casiraghi E, Lucisano M, Pompei C and Dellea C 1994. Cholesterol determination in butter by high performance chromatography. *Milchwissenschaft* 49, 194–196.
- Charlton AJ, Dickinson M, Wakefield ME, Fitches E, Kenis M, Han R., Zhu F, Kone N, Grant M, Devic E, Bruggeman G, Prior R and Smith R 2015. Exploring the chemical safety of fly larvae as a source of protein for animal feed. *Journal of Insects as Food and Feed* 1, 7–16.
- Cullere M, Tasoniero G, Giaccone V, Miotti-Scapin R, Claeys E, De Smet S and Dalle Zotte A 2016. Black soldier fly as dietary protein source for boiler quails: apparent digestibility, excreta microbial load, feed choice, performance, carcass and meat traits. *Animal* 10, 1923–1930.
- Dalle Zotte A and Szendrő ZS 2011. The role of rabbit meat as functional food. *Meat Science* 88, 319–331.
- Finke MD 2013. Complete nutrient content of four species of feeder insects. *Zoo biology* 32, 27–36.
- Folch J, Lees M and Stanley GHS 1957. A simple method for the isolation and purification of total lipids from animal tissues. *Journal of Biological Chemistry* 226, 497–509.
- Guoyao W, Fuller WB, Zhaolai D, Defa L, Junjun W and Zhenlong W 2014. Amino acid nutrition in animals: protein synthesis and beyond. *Annual Review of Animal Biosciences* 2, 387–417.
- Güven R, Kiliç B and Özer CO 2015. Influence of using different oil sources in quail nutrition on meat composition and quality parameters. *Scientific Papers. Series D. Animal Science* 58, 115–118.
- Józefiak D, Józefiak A, Kierończyk B, Rawski M, Świątkiewicz S, Długosz J and Engberg RM 2016. Insects – a natural nutrient source for poultry – a review. *Annals of Animal Science* 16, 297–313.
- Karami M, Ponnampalam EN and Hopkins DL 2013. The effect of palm oil or canola oil on feedlot performance and tissue fatty acid profile and meat quality in goats. *Meat Science* 94, 165–169.
- Lalander C, Senecal J, Gros Calvo M, Ahrens L, Josefsson S, Wiberg K and Vinneras B 2016. Fate of pharmaceuticals and pesticides in fly larvae composting. *Science of the Total Environment* 565, 279–286.
- Laureati M, Proserpio C, Jucker C and Savoldelli S 2016. New sustainable protein sources: consumers' willingness to adopt insects as feed and food. *Italian Journal of Food Science* 28, 652–668.
- Lee CM, Trevino B and Chaipayat M 1996. A simple and rapid solvent extraction method for determining total lipids in fish tissue. *Journal of AOAC International* 79, 487–492.
- Makkar HPS, Tran G, Heuzé V and Ankers P 2014. State-of-the-art on use of insects as animal feed. *Animal Feed Science and Technology* 197, 1–33.
- Marascuilo LA 1966. Large-sample multiple comparisons. *Psychological Bulletin* 65, 280–290.
- Maurer V, Holinger M, Amsler Z, Früh B, Wohlfahrt J, Stamer A and Leiber F 2016. Replacement of soybean cake by *Hermetia illucens* meal in diets for layers. *Journal of Insects as Food and Feed* 2, 83–90.
- National Research Council, Subcommittee on Poultry Nutrition 1994. Nutrient requirements of ring-necked pheasants, Japanese quail, and bobwhite quail. In *Nutrient requirements of poultry*, 9th revised edition (ed.) National Academy Press, pp. 44–45. National Academy of Sciences, Washington, DC, USA.
- Ouhayoun J, Delmas D, Monin G and Roubiscoul P 1990. Abattage du Lapin. 2: Effet du mode de réfrigération sur la biochimie et la contraction des muscles. *Proceedings of the 5^{èmes} Journées de la Recherche Cunicole* 2, 45–56.
- Pieterse E, Pretorius Q, Hoffman LC and Drew DW 2014. The carcass quality, meat quality and sensory characteristics of broilers raised on diets containing either *Musca domestica*, larvae meal, fish meal or soya bean meal as the main protein source. *Animal Production Science* 54, 622–628.
- Poureslami R, Turchini GM, Raes K, Huyghebaert G and De Smet S 2010. Effect of diet, sex and age on fatty acid metabolism in broiler chickens: SFA and MUFA. *British Journal of Nutrition* 104, 204–213.
- Sánchez-Muros MJ, Barroso FG and Manzano-Agugliaro F 2014. Insect meal as renewable source of food for animal feeding: a review. *Journal of Cleaner Production* 65, 16–27.
- Schiavone A, Cullere M, De Marco M, Meneguz M, Biasato I, Bergagna S, Dezzutto D, Gai F, Dabbou S, Gasco L and Dalle Zotte A 2017. Partial or total replacement of soybean oil by black soldier fly larvae (*Hermetia illucens* L.) fat in broiler diets: effect on growth performances, feed-choice, blood traits, carcass characteristics and meat quality. *Italian Journal of Animal Science* 16, 93–100.
- Sealey WM, Gaylord TG, Barrows FT, Tomberlin JK, McGuire MA and Ross C 2011. Sensory analysis of rainbow trout, *Oncorhynchus mykiss*, fed enriched black soldier fly prepupae, *Hermetia illucens*. *Journal of the World Aquaculture Society* 42, 34–45.
- Spranghers T, Ottoboni M, Klootwijk C, Ovin A, Deboosere S, De Meulenaer B, Michielis J, Eeckhout M, De Clercq P and De Smet S 2017. Nutritional composition of black soldier fly (*Hermetia illucens*) prepupae reared on different organic waste substrates. *Journal of the Science of Food and Agriculture* 97, 2594–2600.
- Statistical Analysis Software for Windows 2008. Statistics version 9.1.3. SAS Institute, Cary, NC, USA.
- St-Hilaire S, Cranfill K, McGuire MA, Mosley EE, Tomberlin JK, Newton L, Sealey W, Sheppard C and Irving S 2007. Fish offal recycling by the black soldier fly produces a foodstuff high in omega-3 fatty acids. *Journal of the World Aquaculture Society* 38, 309–313.
- Tschirner M and Simon A 2015. Influence of different growing substrates and processing on the nutrient composition of black soldier fly larvae destined for animal feed. *Journal of Insects as Food and Feed* 1, 249–259.