

# TOWARDS A BIHAMILTONIAN STRUCTURE FOR THE DOUBLE RAMIFICATION HIERARCHY

ALEXANDR BURYAK, PAOLO ROSSI, AND SERGEY SHADRIN

*To the memory of Boris Dubrovin, our teacher and friend*

ABSTRACT. We propose a remarkably simple and explicit conjectural formula for a bihamiltonian structure of the double ramification hierarchy corresponding to an arbitrary homogeneous cohomological field theory. Various checks are presented to support the conjecture.

## CONTENTS

Introduction	1
Organization of the paper	3
Acknowledgements	3
1. Double ramification hierarchy and the main conjecture	3
1.1. Cohomological field theories	3
1.2. Double ramification hierarchy	5
1.3. Bihamiltonian structure for the double ramification hierarchy	9
2. Evidence I: checks for a general CohFT	10
2.1. Genus 0	10
2.2. The first equation of the bihamiltonian recursion	12
2.3. The Schouten–Nijenhuis bracket of two skew-symmetric operators	12
3. Evidence II: central invariants	15
3.1. Miura transformations	15
3.2. The central invariants of a pair of compatible Poisson operators	16
3.3. The central invariants of the pair $(K_1, K_2)$	17
3.4. Relation with the Dubrovin–Zhang hierarchy	18
4. Evidence III: explicit examples	18
4.1. The trivial cohomological field theory	18
4.2. Higher $r$ -spin theories	19
4.3. The Gromov–Witten theory of $\mathbb{C}\mathbb{P}^1$	24
References	26

## INTRODUCTION

Cohomological field theories (or CohFTs for brevity) are systems of cohomology classes on the moduli space  $\overline{\mathcal{M}}_{g,n}$  of stable algebraic curves of genus  $g$  with  $n$  marked points. They were introduced by Kontsevich and Manin in [KM94] to axiomatize the properties of Gromov–Witten classes of a given target variety. Their compatibility with the strata structure and natural morphisms between moduli spaces makes them powerful tools for probing the cohomology of  $\overline{\mathcal{M}}_{g,n}$  and its tautological ring in particular.

Since the Kontsevich–Witten theorem [Wit91, Kon92] stating that the generating series of integrals over  $\overline{\mathcal{M}}_{g,n}$  of monomials in psi classes (the Chern classes of tautological line bundles)

is the logarithm of the tau function of a special solution to the Korteweg–de Vries (KdV) hierarchy, it is well known that integrable hierarchies of evolutionary, Hamiltonian, tau-symmetric PDEs control the intersection theory of CohFTs.

Dubrovin and Zhang [DZ01] give a systematic construction of such integrable hierarchy starting from a semisimple cohomological field theory. Their framework gives, among other things, the language for stating the analogue of the Kontsevich–Witten theorem for any semisimple CohFT, where the KdV hierarchy is replaced by the relevant Dubrovin–Zhang (DZ) hierarchy.

In fact, Dubrovin and Zhang’s method postulates the existence of a bihamiltonian structure (a pair of compatible Poisson structure producing the flows by a recursive procedure) for the DZ hierarchy of a homogeneous semisimple CohFTs. While one of the two Poisson structures was proved to exist (in the differential polynomial class) in [BPS12a, BPS12b], the existence of the second Hamiltonian structure is still an open problem. Notice that the approach of [BPS12a, BPS12b] also clarifies how the construction of the DZ hierarchy (as a Hamiltonian system with respect to the first Poisson bracket) can be based on the axioms and properties of CohFTs together with semisimplicity, without requiring the existence of the second Poisson structure or homogeneity. Nonetheless, the existence of this second Hamiltonian structure in the homogeneous case remains an important unproven feature of the DZ hierarchy.

In [Bur15] a novel approach to constructing integrable hierarchies starting from CohFTs was introduced. This approach does not require semisimplicity of the CohFT and, although based again on the intersection theory on  $\overline{\mathcal{M}}_{g,n}$ , it employs different tautological classes, notably the double ramification cycle (an appropriate compactification of the locus of smooth curves whose marked points support a principal divisor), which explains why this hierarchy was called the double ramification (DR) hierarchy.

The DR hierarchy is always Hamiltonian with respect to a very simple Poisson structure, which, as opposed to the one for the DZ hierarchy, does not essentially depend on the underlying CohFT. The two hierarchies coincide by definition in the dispersionless (genus 0) limit and, by a conjecture in [Bur15] called the *DR/DZ equivalence conjecture*, in the semisimple case they are related by a Miura transformation, which was completely identified in [BDGR18]. Although still unproven, the DR/DZ equivalence conjecture has accumulated a remarkable amount of evidence and verifications (see e.g. [BG16, BDGR18, BDGR19, BGR19, DR19]).

In this paper we propose a very simple formula for a second Poisson structure and collect some evidence for it to give a bihamiltonian structure for the DR hierarchy. These Poisson brackets do depend on the homogeneous CohFT under consideration in a remarkably explicit way. We also compute the central invariants of the resulting bihamiltonian structure, finding that it is Miura equivalent to the conjectured DZ bihamiltonian structure. Finally, we confirm that in several well-known examples of CohFTs our formula does give the expected bihamiltonian structure of the corresponding hierarchy.

Notice that, assuming that our conjectural second Poisson bracket gives a bihamiltonian structure for the DR hierarchy, the existence of a bihamiltonian structure for the DZ hierarchy follows from the DR/DZ equivalence conjecture. In fact, this means that the Miura transformation mapping the DZ hierarchy to the DR hierarchy would simplify not only the first Poisson structure (making it virtually independent of the CohFT), but also the second one, for which no explicit formula was previously known.

**Organization of the paper.** In Section 1 we present our explicit formula and the main conjecture that it gives a bihamiltonian structure for the DR hierarchy of a homogeneous CohFT.

In Section 2 we verify our conjecture in genus 0. Then we prove that the level 0 (primary) flows of the hierarchy can be obtained by bihamiltonian recursion from the level  $-1$  integrals of motion (the Casimir functionals of the first Poisson bracket). Moreover, assuming our conjectural second Poisson bracket satisfies the Jacobi identity, we prove that it is compatible with the first one (i.e., their Schouten–Nijenhuis bracket vanishes).

In Section 3, assuming again that our conjectural second Poisson bracket satisfies the Jacobi identity, we compute the central invariants of the Poisson pencil formed by the two Hamiltonian structures. Under mild hypotheses these central invariants classify bihamiltonian structures of the type relevant for the DR and DZ hierarchies up to Miura transformations. For the Dubrovin–Zhang bihamiltonian structure these invariants have been computed (see e.g. [Liu18]) and in this paper we show that they coincide with the ones for the DR bihamiltonian structure, as expected in light of the DR/DZ equivalence conjecture.

Finally, in Section 4 we prove our main conjecture for several important examples of CohFTs, including the trivial and  $r$ -spin CohFTs (for  $r \leq 5$ ) and the Gromov–Witten theory of the projective line.

*Notation and conventions.* Throughout the text we use the Einstein summation convention for repeated upper and lower Greek indices.

When it doesn't lead to a confusion, we use the symbol  $*$  to indicate any value, in the appropriate range, of a sub- or superscript.

For a topological space  $X$  let  $H^*(X)$  denote the cohomology ring of  $X$  with the coefficients in  $\mathbb{C}$ .

**Acknowledgements.** We are grateful to A. Arsie and P. Lorenzoni for valuable remarks about the preliminary version of the paper. We would like to thank G. Carlet and F. Hernández Iglesias for useful discussions on closely related topics.

The work of A. B. (Sections 1 and 4) was supported by the grant no. 20-11-20214 of the Russian Science Foundation. S. S. was supported by the Netherlands Organization for Scientific Research.

## 1. DOUBLE RAMIFICATION HIERARCHY AND THE MAIN CONJECTURE

In this section, after recalling the notion of cohomological field theory and the construction of the double ramification hierarchy, we present our conjectural formula for a bihamiltonian structure of the double ramification hierarchy.

**1.1. Cohomological field theories.** Let  $\overline{\mathcal{M}}_{g,n}$  be the Deligne–Mumford moduli space of stable curves of genus  $g$  with  $n$  marked points,  $g \geq 0$ ,  $n \geq 0$ ,  $2g - 2 + n > 0$ . Note that  $\overline{\mathcal{M}}_{0,3} = \text{pt}$ , and throughout the text we silently use the identification  $H^*(\overline{\mathcal{M}}_{0,3}) \cong \mathbb{C}$ . Recall the following system of standard maps between these spaces:

- $\pi_{g,n+1}: \overline{\mathcal{M}}_{g,n+1} \rightarrow \overline{\mathcal{M}}_{g,n}$  is the map that forgets the last marked point.
- $\text{gl}_{g_1, I_1; g_2, I_2}: \overline{\mathcal{M}}_{g_1, n_1+1} \times \overline{\mathcal{M}}_{g_2, n_2+1} \rightarrow \overline{\mathcal{M}}_{g_1+g_2, n_1+n_2}$ , is the gluing map that identifies the last marked points of curves of genus  $g_1$  and  $g_2$  and turns them into a node. The sets  $I_1$  and  $I_2$  of cardinality  $n_1$  and  $n_2$ ,  $I_1 \sqcup I_2 = \{1, \dots, n_1 + n_2\}$ , keep track of the relabelling of the remaining marked points.
- $\text{gl}_{g,n+2}^{\text{irr}}: \overline{\mathcal{M}}_{g,n+2} \rightarrow \overline{\mathcal{M}}_{g+1,n}$  is the gluing map that identifies the last two marked points and turns them into a node.

Abusing notation we denote these maps by  $\pi$ ,  $\mathfrak{gl}_2$ , and  $\mathfrak{gl}_1$ , respectively.

Let  $V$  be a finite dimensional vector space of dimension  $N$  with a distinguished vector  $e \in V$ , called the *unit*, and a symmetric nondegenerate bilinear form  $(\cdot, \cdot)$  on  $V$ , called the *metric*. We fix a basis  $e_1, \dots, e_N$  in  $V$  and let  $(\eta_{\alpha\beta})$  denote the matrix of the metric in this basis,  $\eta_{\alpha\beta} := (e_\alpha, e_\beta)$ , and  $A_\alpha$  the coordinates of  $e$  in this basis,  $e = A^\alpha e_\alpha$ . As usual,  $\eta^{\alpha\beta}$  denotes the entries of the inverse matrix,  $(\eta^{\alpha\beta}) := (\eta_{\alpha\beta})^{-1}$ .

**Definition 1.1** ([KM94]). A *cohomological field theory* (CohFT) is a system of linear maps

$$c_{g,n}: V^{\otimes n} \rightarrow H^{\text{even}}(\overline{\mathcal{M}}_{g,n}), \quad 2g - 2 + n > 0,$$

such that the following axioms are satisfied:

- (i) The maps  $c_{g,n}$  are equivariant with respect to the  $S_n$ -action permuting the  $n$  copies of  $V$  in  $V^{\otimes n}$  and the  $n$  marked points in  $\overline{\mathcal{M}}_{g,n}$ , respectively;
- (ii)  $\pi^* c_{g,n}(\otimes_{i=1}^n e_{\alpha_i}) = c_{g,n+1}(\otimes_{i=1}^n e_{\alpha_i} \otimes e)$  and  $c_{0,3}(e_{\alpha_1} \otimes e_{\alpha_2} \otimes e) = \eta_{\alpha_1 \alpha_2}$ ;
- (iii)  $\mathfrak{gl}_2^* c_{g_1+g_2, n_1+n_2}(\otimes_{i=1}^{n_1+n_2} e_{\alpha_i}) = c_{g_1, n_1+1}(\otimes_{i \in I_1} e_{\alpha_i} \otimes e_\mu) \otimes c_{g_2, n_2+1}(\otimes_{i \in I_2} e_{\alpha_i} \otimes e_\nu) \eta^{\mu\nu}$ ;
- (iv)  $\mathfrak{gl}_1^* c_{g+1, n}(\otimes_{i=1}^n e_{\alpha_i}) = c_{g, n+2}(\otimes_{i=1}^n e_{\alpha_i} \otimes e_\mu \otimes e_\nu) \eta^{\mu\nu}$ .

In all axioms above we assume  $\alpha_i \in \{1, \dots, N\}$  for any  $i = 1, 2, \dots$ .

For an arbitrary CohFT the formal power series

$$F = F(t^1, \dots, t^N) := \sum_{n \geq 3} \frac{1}{n!} \sum_{1 \leq \alpha_1, \dots, \alpha_n \leq N} \left( \int_{\overline{\mathcal{M}}_{0,n}} c_{0,n}(\otimes_{i=1}^n e_{\alpha_i}) \right) \prod_{i=1}^n t^{\alpha_i}$$

satisfies the equations

$$\begin{aligned} A^\mu \frac{\partial^3 F}{\partial t^\mu \partial t^\alpha \partial t^\beta} &= \eta_{\alpha\beta}, & 1 \leq \alpha, \beta \leq N, \\ \frac{\partial^3 F}{\partial t^\alpha \partial t^\beta \partial t^\mu} \eta^{\mu\nu} \frac{\partial^3 F}{\partial t^\nu \partial t^\gamma \partial t^\delta} &= \frac{\partial^3 F}{\partial t^\alpha \partial t^\gamma \partial t^\mu} \eta^{\mu\nu} \frac{\partial^3 F}{\partial t^\nu \partial t^\beta \partial t^\delta}, & 1 \leq \alpha, \beta, \gamma, \delta \leq N. \end{aligned}$$

Thus, the formal power series  $F$  defines a Dubrovin–Frobenius manifold structure on a formal neighbourhood of 0 in  $V$ .

**Definition 1.2.** A CohFT  $\{c_{g,n}\}$  is called *semisimple* (at the origin) if the algebra defined by the structure constants  $c_{\beta\gamma}^\alpha|_{t^*=0}$ , where  $c_{\beta\gamma}^\alpha := \eta^{\alpha\mu} \frac{\partial^3 F}{\partial t^\mu \partial t^\beta \partial t^\gamma}$ , doesn't have nilpotents.

1.1.1. *Homogeneity.* Let  $V$  be a graded vector space and assume the basis  $e_1, \dots, e_N$  is homogeneous with  $\deg e_\alpha = q_\alpha$ ,  $\alpha = 1, \dots, N$ . Assume also that  $\deg e = 0$ . By  $\text{Deg}: H^*(\overline{\mathcal{M}}_{g,n}) \rightarrow H^*(\overline{\mathcal{M}}_{g,n})$  we denote the operator that acts on  $H^i(\overline{\mathcal{M}}_{g,n})$  by multiplication by  $\frac{i}{2}$ .

**Definition 1.3.** A CohFT  $\{c_{g,n}\}$  is called *homogeneous*, or *conformal*, if there exist complex constants  $r^\alpha$ ,  $\alpha = 1, \dots, N$ , and  $\delta$  such that

$$(1.1) \quad \text{Deg } c_{g,n}(\otimes_{i=1}^n e_{\alpha_i}) + \pi_* c_{g,n+1}(\otimes_{i=1}^n e_{\alpha_i} \otimes r^\gamma e_\gamma) = \left( \sum_{i=1}^n q_{\alpha_i} + \delta(g-1) \right) c_{g,n}(\otimes_{i=1}^n e_{\alpha_i}).$$

The constant  $\delta$  is called the *conformal dimension* of CohFT.

For a homogeneous CohFT the formal power series  $F(t^1, \dots, t^N)$  satisfies the property

$$((1 - q_\alpha)t^\alpha + r^\alpha) \frac{\partial F}{\partial t^\alpha} = (3 - \delta)F + \frac{1}{2} A_{\alpha\beta} t^\alpha t^\beta,$$

where  $A_{\alpha\beta} := r^\mu c_{0,3}(e_\alpha \otimes e_\beta \otimes e_\mu)$ . Thus, the associated Dubrovin–Frobenius manifold on a formal neighbourhood of 0 in  $V$  is also homogeneous, with the Euler vector field given by

$$E = E^\alpha \frac{\partial}{\partial t^\alpha} := ((1 - q_\alpha)t^\alpha + r^\alpha) \frac{\partial}{\partial t^\alpha}.$$

## 1.2. Double ramification hierarchy.

1.2.1. *Formal loop space, differential polynomials, and local functionals.* Introduce formal variables  $u_i^\alpha$ ,  $\alpha = 1, \dots, N$ ,  $i = 0, 1, \dots$ . Following [DZ01] (see also [Ros17]) we define the ring of *differential polynomials*  $\mathcal{A}^0$  in the variables  $u^1, \dots, u^N$  as the ring of polynomials  $f(u^*, u_x^*, u_{xx}^*, \dots)$  in the variables  $u_i^\alpha$ ,  $i > 0$ , with coefficients in the ring of formal power series in the variables  $u^\alpha = u_0^\alpha$ :

$$\mathcal{A}^0 := \mathbb{C}[[u^*]][u_{\geq 1}^*].$$

*Remark 1.4.* This way we define a model of the loop space of the vector space  $V$  by describing its ring of functions. In particular, it is useful to think of the variables  $u^\alpha := u_0^\alpha$  as the components  $u^\alpha(x)$  of a formal loop  $u: S^1 \rightarrow V$  in the basis  $e_1, \dots, e_N$ . Then the variables  $u_1^\alpha := u_x^\alpha$ ,  $u_2^\alpha := u_{xx}^\alpha, \dots$  are the components of the iterated  $x$ -derivatives of a formal loop.

The *standard gradation* on  $\mathcal{A}^0$ , which we denote by  $\deg$ , is introduced by  $\deg u_i^\alpha := i$ . The homogeneous component of  $\mathcal{A}^0$  of standard degree  $d$  is denoted by  $\mathcal{A}_d^0$ . The operator

$$\partial_x := \sum_{i \geq 0} u_{i+1}^\alpha \frac{\partial}{\partial u_i^\alpha}$$

increases the standard degree by 1. Therefore, the quotient

$$\Lambda^0 := \mathcal{A}^0 / (\mathbb{C} \oplus \text{Im } \partial_x),$$

called the space of *local functionals*, inherits the standard gradation. The homogeneous component of  $\Lambda^0$  of standard degree  $d$  is denoted by  $\Lambda_d^0$ . The natural projection  $\mathcal{A}^0 \rightarrow \Lambda^0$  assigns to a differential polynomial  $f$  the local functional  $\bar{f} = \int f dx$ .

The *variational derivative*  $\frac{\delta}{\delta u^\alpha}: \mathcal{A}^0 \rightarrow \mathcal{A}^0$ ,  $\alpha = 1, \dots, N$ , is defined by

$$\frac{\delta}{\delta u^\alpha} := \sum_{i \geq 0} (-\partial_x)^i \circ \frac{\partial}{\partial u_i^\alpha}.$$

Since it vanishes on  $\mathbb{C} \oplus \text{Im } \partial_x$ , it is well defined on the space of local functionals and abusing notation we denote it by the same symbol,  $\frac{\delta}{\delta u^\alpha}: \Lambda^0 \rightarrow \mathcal{A}^0$ .

We associate with a differential polynomial  $f \in \mathcal{A}^0$  a sequence of differential operators indexed by  $\alpha = 1, \dots, N$  and  $k \geq 0$ :

$$L_\alpha^k(f) := \sum_{i \geq k} \binom{i}{k} \frac{\partial f}{\partial u_i^\alpha} \partial_x^{i-k}.$$

We often use the notation  $L_\alpha(f) := L_\alpha^0(f)$ . These operators satisfy the property

$$L_\alpha^k(\partial_x f) = \partial_x \circ L_\alpha^k(f) + L_\alpha^{k-1}(f), \quad k \geq 0,$$

where we adopt the convention  $L_\alpha^l(f) := 0$  for  $l < 0$ .

We associate with a local functional  $\bar{h} = \int h dx \in \Lambda^0$  a sequence of  $N \times N$  matrices  $\widehat{\Omega}^k(\bar{h}) = (\widehat{\Omega}^k(\bar{h})^{\alpha\beta})$  of differential operators, indexed by  $k \geq 0$ , defined as

$$\widehat{\Omega}^k(\bar{h})^{\alpha\beta} := \eta^{\alpha\mu} \eta^{\beta\nu} L_\nu^k \left( \frac{\delta \bar{h}}{\delta u^\mu} \right).$$

We often use the notation  $\widehat{\Omega}(\bar{h}) := \widehat{\Omega}^0(\bar{h})$ .

Consider an  $N \times N$  matrix  $K = (K^{\mu\nu})$  of differential operators of the form  $K^{\mu\nu} = \sum_{j \geq 0} K_j^{\mu\nu} \partial_x^j$ , where  $K_j^{\mu\nu} \in \mathcal{A}^0$  and the sum is finite. The conjugate matrix of differential operators  $K^* = ((K^*)^{\mu\nu})$  is defined by

$$(K^*)^{\mu\nu} := \sum_{j \geq 0} (-\partial_x)^j \circ K_j^{\nu\mu}.$$

**Lemma 1.5.** *For any  $\bar{h} \in \Lambda^0$  we have  $\widehat{\Omega}^k(\bar{h})^* = (-1)^k \widehat{\Omega}^k(\bar{h})$ .*

*Proof.* It is a straightforward computation. We have to check that

$$\left( L_\mu^k \left( \frac{\delta \bar{h}}{\delta u^\nu} \right) \right)^* = (-1)^k L_\nu^k \left( \frac{\delta \bar{h}}{\delta u^\mu} \right).$$

To this end, we compute

$$\begin{aligned} \left( L_\mu^k \left( \frac{\delta \bar{h}}{\delta u^\nu} \right) \right)^* &= \sum_{i=0}^{\infty} \binom{k+i}{i} (-\partial_x)^i \circ \frac{\partial}{u_{k+i}^\mu} \frac{\delta \bar{h}}{\delta u^\nu} = \sum_{i,j=0}^{\infty} \binom{k+i+j}{i,j} (-\partial_x)^i \frac{\partial}{u_{k+i+j}^\mu} \frac{\delta \bar{h}}{\delta u^\nu} (-\partial_x)^j \\ &= \sum_{i,j,l=0}^{\infty} (-1)^{k+i+j} \binom{k+i+j+l}{i,j,l} (-\partial_x)^{i+l} \frac{\partial}{u_{k+i+j+l}^\nu} \frac{\delta \bar{h}}{\delta u^\mu} (-\partial_x)^j \\ &= (-1)^k \sum_{j=0}^{\infty} \binom{k+j}{j} \frac{\partial}{u_{k+j}^\nu} \frac{\delta \bar{h}}{\delta u^\mu} \partial_x^j = (-1)^k L_\nu^k \left( \frac{\delta \bar{h}}{\delta u^\mu} \right). \end{aligned}$$

Here, in order to pass to the second line in the computation we use the identity

$$\frac{\partial}{u_t^\mu} \frac{\delta}{\delta u^\nu} = (-1)^t \sum_{l=0}^{\infty} \binom{l+t}{t} (-\partial_x)^l \frac{\partial}{u_{t+l}^\nu} \frac{\delta}{\delta u^\mu}$$

(cf. [LZ11, Lemma 2.1.5(i)]), and in the third line we use that for any  $t \geq 1$

$$\sum_{i+l=t} \frac{1}{i!l!} (-1)^i (-\partial_x)^{i+l} = 0.$$

□

Differential polynomials and local functionals can also be described using another set of formal variables, corresponding heuristically to the Fourier components  $p_k^\alpha$ ,  $k \in \mathbb{Z}$ , of the functions  $u^\alpha = u^\alpha(x)$ . We define a change of variables

$$(1.2) \quad u_j^\alpha = \sum_{k \in \mathbb{Z}} (ik)^j p_k^\alpha e^{ikx},$$

which allows us to express a differential polynomial  $f(u, u_x, u_{xx}, \dots) \in \mathcal{A}^0$  as a formal Fourier series in  $x$ . In the latter expression the coefficient of  $e^{ikx}$  is a power series in the variables  $p_j^\alpha$  with the sum of the subscripts in each monomial in  $p_j^\alpha$  equal to  $k$ . Moreover, the local functional  $\bar{f}$  corresponds to the constant term of the Fourier series of  $f$ .

**1.2.2. Poisson brackets and Hamiltonian hierarchies.** Let us describe a natural class of Poisson brackets on the space of local functionals. Given an  $N \times N$  matrix  $K = (K^{\mu\nu})$  of differential operators of the form  $K^{\mu\nu} = \sum_{j \geq 0} K_j^{\mu\nu} \partial_x^j$ , where  $K_j^{\mu\nu} \in \mathcal{A}^0$  and the sum is finite, we define

$$\{\bar{f}, \bar{g}\}_K := \int \left( \frac{\delta \bar{f}}{\delta u^\mu} K^{\mu\nu} \frac{\delta \bar{g}}{\delta u^\nu} \right) dx.$$

The bracket  $\{\cdot, \cdot\}_K$  is skew-symmetric if and only if  $K^* = -K$ . The bracket  $\{\cdot, \cdot\}_K$  satisfies the Jacobi identity if and only if the Schouten–Nijenhuis bracket of the bracket with itself vanishes,

$$[\{\cdot, \cdot\}_K, \{\cdot, \cdot\}_K] = 0,$$

which is discussed in details in Section 2.3.

**Definition 1.6.** An operator  $K$  is called *Poisson*, if the bracket  $\{\cdot, \cdot\}_K$  is skew-symmetric and satisfies the Jacobi identity.

*Example 1.7.* A standard example of a Poisson operator is given by the operator  $\eta^{-1} \partial_x$ . The corresponding Poisson bracket has a nice expression in terms of the variables  $p_k^\alpha$ :

$$\{p_k^\alpha, p_j^\beta\}_{\eta^{-1} \partial_x} = ik \eta^{\alpha\beta} \delta_{k+j,0}.$$

Consider the extensions  $\widehat{\mathcal{A}}^0 := \mathcal{A}^0 \otimes \mathbb{C}[[\varepsilon]]$  and  $\widehat{\Lambda}^0 := \Lambda^0 \otimes \mathbb{C}[[\varepsilon]]$  of the spaces  $\mathcal{A}^0$  and  $\Lambda^0$  with a new variable  $\varepsilon$  of standard gradation  $\deg \varepsilon := -1$ . Let  $\widehat{\mathcal{A}}_k^0$  and  $\widehat{\Lambda}_k^0$  denote the subspaces of degree  $k$  of  $\widehat{\mathcal{A}}^0$  and  $\widehat{\Lambda}^0$ , respectively. Abusing the terminology we still call their elements *differential polynomials* and *local functionals*.

We can also define a bracket  $\{\cdot, \cdot\}_K: \widehat{\Lambda}^0 \times \widehat{\Lambda}^0 \rightarrow \widehat{\Lambda}^0$  as above starting from an operator  $K = (K^{\mu\nu})$ ,  $K^{\mu\nu} = \sum_{i,j \geq 0} K_j^{[i],\mu\nu} \varepsilon^i \partial_x^j$ ,  $K_j^{[i],\mu\nu} \in \mathcal{A}^0$ , where for each  $i \geq 0$  the sum  $\sum_{j \geq 0} K_j^{[i],\mu\nu} \partial_x^j$  is finite. An operator  $K = (K^{\alpha\beta})$  and the corresponding bracket  $\{\cdot, \cdot\}_K$  have degree  $d$  if  $K^{\alpha\beta} = \sum_{s \geq 0} K_s^{\alpha\beta} \partial_x^s$ , where  $K_s^{\alpha\beta} \in \widehat{\mathcal{A}}_{-s+d}^0$ .

**Definition 1.8.** A *Hamiltonian hierarchy* of PDEs is a system of the form

$$(1.3) \quad \frac{\partial u^\alpha}{\partial \tau_i} = K^{\alpha\mu} \frac{\delta \bar{h}_i}{\delta u^\mu}, \quad 1 \leq \alpha \leq N, \quad i \geq 1,$$

where  $\bar{h}_i \in \widehat{\Lambda}_0^0$ ,  $K = (K^{\mu\nu})$  is a Poisson operator of degree 1, and the compatibility condition  $\{\bar{h}_i, \bar{h}_j\}_K = 0$  for  $i, j \geq 1$  is satisfied.

The local functionals  $\bar{h}_i$  are called the *Hamiltonians* of the system (1.3).

Consider Hamiltonian hierarchies of the form

$$(1.4) \quad \frac{\partial u^\alpha}{\partial t_q^\beta} = K_1^{\alpha\mu} \frac{\delta \bar{h}_{\beta,q}}{\delta u^\mu}, \quad 1 \leq \alpha, \beta \leq N, \quad q \geq 0,$$

equipped with  $N$  linearly independent Casimirs  $\bar{h}_{\alpha,-1}$ ,  $1 \leq \alpha \leq N$ , of the Poisson bracket  $\{\cdot, \cdot\}_{K_1}$ . Two Poisson operators  $K_1$  and  $K_2$  are said to be *compatible* if the linear combination  $K_2 - \lambda K_1$  is a Poisson operator for any  $\lambda \in \mathbb{C}$ .

**Definition 1.9.** A Hamiltonian hierarchy (1.4) is said to be *bihamiltonian* if it is endowed with a Poisson operator  $K_2$  of degree 1 compatible with the operator  $K_1$  and such that

$$(1.5) \quad \{\cdot, \bar{h}_{\alpha,i-1}\}_{K_2} = \sum_{j=0}^i R_{i,\alpha}^{j,\beta} \{\cdot, \bar{h}_{\beta,i-j}\}_{K_1}, \quad 1 \leq \alpha \leq N, \quad i \geq 0,$$

where  $R_i^j = (R_{i,\alpha}^{j,\beta})$ ,  $0 \leq j \leq i$ , are constant  $N \times N$  matrices.

The relation (1.5) is called a *bihamiltonian recursion*.

*Remark 1.10.* Note that if the matrices  $R_i^0$ ,  $i \geq 0$ , are invertible, then the bihamiltonian recursion (1.5) determines the local functionals  $\bar{h}_{\alpha,i}$ ,  $i \geq 0$ , uniquely up to a triangular transformation

$$\bar{h}_{\alpha,i} \mapsto \bar{h}_{\alpha,i} + \sum_{j=1}^{i+1} A_\alpha^{j,\beta} \bar{h}_{\beta,i-j}, \quad 1 \leq \alpha \leq N, \quad i \geq 0, \quad A_\alpha^{j,\beta} \in \mathbb{C}.$$

In the bihamiltonian hierarchies considered below all but finitely many matrices  $R_i^0$  are invertible.

**1.2.3. Construction of the double ramification hierarchy.** Denote by  $\psi_i \in H^2(\overline{\mathcal{M}}_{g,n})$  the first Chern class of the line bundle over  $\overline{\mathcal{M}}_{g,n}$  formed by the cotangent lines at the  $i$ -th marked point. Denote by  $\mathbb{E}$  the rank  $g$  Hodge vector bundle over  $\overline{\mathcal{M}}_{g,n}$  whose fibers are the spaces of holomorphic one-forms on stable curves. Let  $\lambda_j := c_j(\mathbb{E}) \in H^{2j}(\overline{\mathcal{M}}_{g,n})$ .

For any  $a_1, \dots, a_n \in \mathbb{Z}$ ,  $\sum_{i=1}^n a_i = 0$ , denote by  $\text{DR}_g(a_1, \dots, a_n) \in H^{2g}(\overline{\mathcal{M}}_{g,n})$  the *double ramification (DR) cycle*. We refer the reader, for example, to [BSSZ15] for the definition of the DR cycle on  $\overline{\mathcal{M}}_{g,n}$ , which is based on the notion of a stable map to  $\mathbb{C}\mathbb{P}^1$  relative to 0 and  $\infty$ . If not all the multiplicities  $a_i$  are equal to zero, then one can think of the class  $\text{DR}_g(a_1, \dots, a_n)$  as the Poincaré dual to a compactification in  $\overline{\mathcal{M}}_{g,n}$  of the locus of pointed smooth curves  $(C; p_1, \dots, p_n)$  satisfying  $\mathcal{O}_C(\sum_{i=1}^n a_i p_i) \cong \mathcal{O}_C$ . Consider the Poincaré dual to the double ramification cycle  $\text{DR}_g(a_1, \dots, a_n)$  in the space  $\overline{\mathcal{M}}_{g,n}$ . It is an element of  $H_{2(2g-3+n)}(\overline{\mathcal{M}}_{g,n})$ , and abusing notation it is also denoted by  $\text{DR}_g(a_1, \dots, a_n)$ .

Consider a CohFT  $\{c_{g,n}: V^{\otimes n} \rightarrow H^{\text{even}}(\overline{\mathcal{M}}_{g,n})\}$ . The Hamiltonians of the *double ramification hierarchy* are defined as follows:

$$(1.6) \quad \bar{g}_{\alpha,d} := \sum_{\substack{g \geq 0 \\ n \geq 2}} \frac{(-\varepsilon^2)^g}{n!} \sum_{\substack{a_1, \dots, a_n \in \mathbb{Z} \\ \sum a_i = 0}} \left( \int_{\text{DR}_g(0, a_1, \dots, a_n)} \lambda_g \psi_1^d c_{g,n+1}(e_\alpha \otimes \otimes_{i=1}^n e_{\alpha_i}) \right) \prod_{i=1}^n p_{a_i}^{\alpha_i},$$

for  $\alpha = 1, \dots, N$  and  $d = 0, 1, 2, \dots$

The expression on the right-hand side of (1.6) can be uniquely written as a local functional from  $\widehat{\Lambda}_0^0$  using the change of variables (1.2). Concretely, it can be done in the following way. The restriction  $\text{DR}_g(a_1, \dots, a_n)|_{\mathcal{M}_{g,n}^{\text{ct}}}$ , where  $\mathcal{M}_{g,n}^{\text{ct}}$  is the moduli space of stable curves of compact type, is a homogeneous polynomial in  $a_1, \dots, a_n$  of degree  $2g$  with the coefficients in  $H^{2g}(\mathcal{M}_{g,n}^{\text{ct}})$ . This follows from Hain's formula [Hai13] for the version of the DR cycle defined using the universal Jacobian over  $\mathcal{M}_{g,n}^{\text{ct}}$  and the result of the paper [MW13], where it is proved that the two versions of the DR cycle coincide on  $\mathcal{M}_{g,n}^{\text{ct}}$  (the polynomiality of the DR cycle on  $\overline{\mathcal{M}}_{g,n}$  is proved in [JPPZ17]). The polynomiality of the DR cycle on  $\mathcal{M}_{g,n}^{\text{ct}}$  together with the fact that  $\lambda_g$  vanishes on  $\overline{\mathcal{M}}_{g,n} \setminus \mathcal{M}_{g,n}^{\text{ct}}$  (see e.g. [FP00, Section 0.4]) imply that the integral

$$(1.7) \quad \int_{\text{DR}_g(0, a_1, \dots, a_n)} \lambda_g \psi_1^d c_{g,n+1}(e_\alpha \otimes \otimes_{i=1}^n e_{\alpha_i})$$

is a homogeneous polynomial in  $a_1, \dots, a_n$  of degree  $2g$ , which we denote by

$$P_{\alpha,d,g;\alpha_1, \dots, \alpha_n}(a_1, \dots, a_n) = \sum_{\substack{b_1, \dots, b_n \geq 0 \\ b_1 + \dots + b_n = 2g}} P_{\alpha,d,g;\alpha_1, \dots, \alpha_n}^{b_1, \dots, b_n} a_1^{b_1} \dots a_n^{b_n}.$$

Then we have

$$\bar{g}_{\alpha,d} = \int \sum_{\substack{g \geq 0 \\ n \geq 2}} \frac{\varepsilon^{2g}}{n!} \sum_{\substack{b_1, \dots, b_n \geq 0 \\ b_1 + \dots + b_n = 2g}} P_{\alpha,d,g;\alpha_1, \dots, \alpha_n}^{b_1, \dots, b_n} u_{b_1}^{\alpha_1} \dots u_{b_n}^{\alpha_n} dx.$$

Note that the integral (1.7) is defined only when  $a_1 + \dots + a_n = 0$ . Therefore, the polynomial  $P_{\alpha,d,g;\alpha_1, \dots, \alpha_n}$  is not uniquely defined. However, the resulting local functional  $\bar{g}_{\alpha,d} \in \widehat{\Lambda}_0^0$  doesn't depend on this ambiguity, see [Bur15]. In fact, in [BR16a] a special choice of the differential polynomial densities  $g_{\alpha,d} \in \widehat{\mathcal{A}}_0^0$  for  $\bar{g}_{\alpha,d} = \int g_{\alpha,d} dx$  is proposed. The densities are defined in terms of the  $p$ -variables as

$$g_{\alpha,d} := \sum_{\substack{g \geq 0, n \geq 1 \\ 2g-1+n > 0}} \frac{(-\varepsilon^2)^g}{n!} \sum_{\substack{a_0, \dots, a_n \in \mathbb{Z} \\ \sum a_i = 0}} \left( \int_{\text{DR}_g(a_0, a_1, \dots, a_n)} \lambda_g \psi_1^d c_{g,n+1}(e_\alpha \otimes \otimes_{i=1}^n e_{\alpha_i}) \right) \prod_{i=1}^n p_{a_i}^{\alpha_i} e^{-ia_0 x}$$

and converted unequivocally to differential polynomials using the change of variables (1.2).

It is proved in [Bur15] that the local functionals  $\bar{g}_{\alpha,d}$  mutually commute with respect to the standard bracket  $\{\cdot, \cdot\}_{\eta^{-1}\partial_x}$ .

**Definition 1.11.** The system of local functionals  $\bar{g}_{\alpha,d}$ , for  $\alpha = 1, \dots, N$ ,  $d = 0, 1, 2, \dots$ , and the corresponding system of Hamiltonian PDEs with respect to the Poisson bracket  $\{\cdot, \cdot\}_{\eta^{-1}\partial_x}$ ,

$$\frac{\partial u^\alpha}{\partial t_q^\beta} = \eta^{\alpha\mu} \partial_x \frac{\delta \bar{g}_{\beta,q}}{\delta u^\mu}, \quad 1 \leq \alpha, \beta \leq N, \quad q \geq 0,$$

is called the *double ramification hierarchy* (or the *DR hierarchy* for brevity).



1.2.4. *Extra structures for the double ramification hierarchy.* We can equip the DR hierarchy with the following  $N$  linearly independent Casimirs of its Poisson bracket  $\{\cdot, \cdot\}_{\eta^{-1}\partial_x}$ :

$$\bar{g}_{\alpha,-1} := \int \eta_{\alpha\beta} u^\beta dx, \quad 1 \leq \alpha \leq N.$$

Another important object related to the DR hierarchy is the local functional  $\bar{g}$  defined in terms of the  $p$ -variables as

$$\bar{g} := \sum_{\substack{g,n \geq 0 \\ 2g-2+n > 0}} \frac{(-\varepsilon^2)^g}{n!} \sum_{\substack{a_1, \dots, a_n \in \mathbb{Z} \\ \sum a_i = 0}} \left( \int_{\text{DR}_g(a_1, \dots, a_n)} \lambda_g c_{g,n} (\otimes_{i=1}^n e_{\alpha_i}) \right) \prod_{i=1}^n p_{\alpha_i}^{\alpha_i}.$$

We have the following relations [Bur15, Section 4.2.5]:

$$(1.8) \quad \bar{g}_{\mathbb{1},1} = (D-2)\bar{g} \quad \text{and} \quad \frac{\partial \bar{g}}{\partial u^\alpha} = \bar{g}_{\alpha,0},$$

where  $D := \sum_{n \geq 0} (n+1) u_n^\alpha \frac{\partial}{\partial u_n^\alpha}$  and  $\bar{g}_{\mathbb{1},1} := A^\alpha \bar{g}_{\alpha,1}$ .

1.3. **Bihamiltonian structure for the double ramification hierarchy.** Consider a homogeneous cohomological field theory and the associated DR hierarchy. Let

$$\mu_\alpha := q_\alpha - \frac{\delta}{2}$$

and define a diagonal matrix  $\mu$  by

$$\mu := \text{diag}(\mu_1, \dots, \mu_N).$$

Note that

$$(1.9) \quad \mu\eta + \eta\mu = 0.$$

Introduce an operator  $\widehat{E}$  by

$$\widehat{E} := \sum_{n \geq 0} ((1 - q_\alpha) u_n^\alpha + \delta_{n,0} r^\alpha) \frac{\partial}{\partial u_n^\alpha} + \frac{1 - \delta}{2} \varepsilon \frac{\partial}{\partial \varepsilon}.$$

**Definition 1.12.** Define an operator  $K_2$  of degree 1 by

$$K_2 := \widehat{E} \left( \widehat{\Omega}(\bar{g}) \right) \circ \partial_x + \widehat{\Omega}(\bar{g})_x \circ \left( \frac{1}{2} - \mu \right) + \partial_x \circ \widehat{\Omega}^1(\bar{g}) \circ \partial_x,$$

where the notation  $\widehat{E}(\widehat{\Omega}(\bar{g}))$  (respectively,  $\widehat{\Omega}(\bar{g})_x$ ) means that we apply the operator  $\widehat{E}$  (respectively,  $\partial_x$ ) to the coefficients of the operator  $\widehat{\Omega}(\bar{g})$ .

**Conjecture 1.13.**

- (1) The operator  $K_2$  is Poisson and compatible with the operator  $K_1 := \eta^{-1}\partial_x$ .
- (2) The Poisson brackets  $\{\cdot, \cdot\}_{K_2}$  and  $\{\cdot, \cdot\}_{K_1}$  give a bihamiltonian structure for the DR hierarchy with the following bihamiltonian recursion:

$$(1.10) \quad \{\cdot, \bar{g}_{\alpha,d}\}_{K_2} = \left( d + \frac{3}{2} + \mu_\alpha \right) \{\cdot, \bar{g}_{\alpha,d+1}\}_{K_1} + A_\alpha^\beta \{\cdot, \bar{g}_{\beta,d}\}_{K_1}, \quad d \geq -1,$$

where  $A_\alpha^\beta := \eta^{\beta\nu} A_{\nu\alpha}$ .

**Lemma 1.14.** The operator  $K_2$  has the following alternative expression:

$$(1.11) \quad K_2 = \partial_x \circ \widehat{\Omega}(\bar{g}) \circ \left( \frac{1}{2} - \mu \right) + \left( \frac{1}{2} - \mu \right) \circ \widehat{\Omega}(\bar{g}) \circ \partial_x + \eta^{-1} A \eta^{-1} \partial_x + \partial_x \circ \widehat{\Omega}^1(\bar{g}) \circ \partial_x,$$

where  $A = (A_{\alpha\beta})$ .

*Proof.* Equation (1.11) is equivalent to

$$\widehat{E} \left( \widehat{\Omega}(\overline{g}) \right) = \widehat{\Omega}(\overline{g}) \circ \left( \frac{1}{2} - \mu \right) + \left( \frac{1}{2} - \mu \right) \circ \widehat{\Omega}(\overline{g}) + \eta^{-1} A \eta^{-1},$$

which is equivalent to

$$\widehat{E} \left( L_\alpha \left( \frac{\delta \overline{g}}{\delta u^\beta} \right) \right) = (q_\alpha + q_\beta + 1 - \delta) L_\alpha \left( \frac{\delta \overline{g}}{\delta u^\beta} \right) + A_{\alpha\beta}.$$

Note that, since  $[\widehat{E}, \partial_x] = 0$ , the operator  $\widehat{E}$  acts on  $\widehat{\Lambda}^0$ . It is easy to see that Equation (1.1) implies that

$$\widehat{E}\overline{g} = (3 - \delta)\overline{g} + \int \frac{1}{2} A_{\alpha\beta} u^\alpha u^\beta dx.$$

Using this property we get

$$\begin{aligned} \widehat{E} \left( L_\alpha \left( \frac{\delta \overline{g}}{\delta u^\beta} \right) \right) &= L_\alpha \left( \frac{\delta(\widehat{E}\overline{g})}{\delta u^\beta} \right) - (2 - q_\alpha - q_\beta) L_\alpha \left( \frac{\delta \overline{g}}{\delta u^\beta} \right) \\ &= (q_\alpha + q_\beta + 1 - \delta) L_\alpha \left( \frac{\delta \overline{g}}{\delta u^\beta} \right) + A_{\alpha\beta}, \end{aligned}$$

as required.  $\square$

*Remark 1.15.* Formula (1.11) together with Lemma 1.5 immediately imply that the operator  $K_2$  is skew-symmetric.

## 2. EVIDENCE I: CHECKS FOR A GENERAL COHFT

In this section we check some parts of Conjecture 1.13 for an arbitrary homogeneous CohFT  $\{c_{g,n}\}$ : we prove it in genus 0, examine the first step of the bihamiltonian recursion, and compute the Schouten–Nijenhuis bracket of  $\{\cdot, \cdot\}_{K_2}$  and  $\{\cdot, \cdot\}_{K_1}$  (considered as bivector fields).

**2.1. Genus 0.** By reduction to genus 0 we mean the reduction to the dispersionless part of the hierarchy, that is, we set  $\varepsilon = 0$ .

**Proposition 2.1.** *Conjecture 1.13 is true in genus 0.*

*Proof.* Recall that  $F = F(t^1, \dots, t^N)$  denotes the Dubrovin–Frobenius manifold potential associated to a CohFT  $\{c_{g,n}\}$ . We have

$$\overline{g}|_{\varepsilon=0} = \int F(u^1, \dots, u^N) dx.$$

Therefore,

$$K_2^{[0], \alpha\beta} := K_2^{\alpha\beta} \Big|_{\varepsilon=0} = g^{\alpha\beta} \partial_x + \partial_x \Omega^{\alpha\beta} \left( \frac{1}{2} - \mu_\beta \right),$$

where

$$(2.1) \quad g^{\alpha\beta} := \eta^{\alpha\mu} \eta^{\beta\nu} E^\gamma \frac{\partial^3 F}{\partial t^\gamma \partial t^\mu \partial t^\nu} \Big|_{t^\theta = u^\theta} \quad \text{and} \quad \Omega^{\alpha\beta} := \eta^{\alpha\mu} \eta^{\beta\nu} \frac{\partial^2 F}{\partial t^\mu \partial t^\nu} \Big|_{t^\theta = u^\theta}.$$

The fact that the operators  $\eta^{-1} \partial_x$  and  $K_2^{[0]}$  form a pair of compatible Poisson operators is well known (see e.g. [DZ99, page 443]). This proves the first part of the conjecture in genus 0.

Now we check the second part of the conjecture. Let  $t_a^\alpha$ ,  $1 \leq \alpha \leq N$ ,  $a \geq 0$ , be formal variables, where we identify  $t_0^\alpha = t^\alpha$ . Consider the genus 0 potential of  $\{c_{g,n}\}$ :

$$\mathcal{F}_0(t_*^*) := \sum_{n \geq 3} \frac{1}{n!} \sum_{\substack{1 \leq \alpha_1, \dots, \alpha_n \leq N \\ d_1, \dots, d_n \geq 0}} \left( \int_{\overline{\mathcal{M}}_{0,n}} c_{0,n}(\otimes_{i=1}^n e_{\alpha_i}) \prod_{i=1}^n \psi_i^{d_i} \right) \prod_{i=1}^n t_{d_i}^{\alpha_i} \in \mathbb{C}[[t_*^*]].$$

It satisfies the string equation

$$(2.2) \quad \frac{\partial \mathcal{F}_0}{\partial t_0^{\mathbb{1}}} = \sum_{n \geq 0} t_{n+1}^\alpha \frac{\partial \mathcal{F}_0}{\partial t_n^\alpha} + \frac{1}{2} \eta_{\alpha\beta} t_0^\alpha t_0^\beta,$$

where we use the notation  $\frac{\partial}{\partial t_0^{\mathbb{1}}} := A^\alpha \frac{\partial}{\partial t_0^\alpha}$ , and the topological recursion relations

$$(2.3) \quad \frac{\partial^3 \mathcal{F}_0}{\partial t_{a+1}^\alpha \partial t_b^\beta \partial t_c^\gamma} = \frac{\partial^2 \mathcal{F}_0}{\partial t_a^\alpha \partial t_0^\mu} \eta^{\mu\nu} \frac{\partial^3 \mathcal{F}_0}{\partial t_0^\nu \partial t_b^\beta \partial t_c^\gamma}, \quad 1 \leq \alpha, \beta, \gamma \leq N, \quad a, b, c \geq 0.$$

Also, the homogeneity condition (1.1) implies that

$$(2.4) \quad \left( \sum_{n \geq 0} (1 - q_\alpha - n) t_n^\alpha \frac{\partial}{\partial t_n^\alpha} + r^\alpha \frac{\partial}{\partial t_0^\alpha} - \sum_{n \geq 0} A_\beta^\alpha t_{n+1}^\beta \frac{\partial}{\partial t_n^\alpha} \right) \mathcal{F}_0 = (3 - \delta) \mathcal{F}_0 + \frac{1}{2} A_{\alpha\beta} t_0^\alpha t_0^\beta.$$

Denote

$$\Omega_{\alpha,a;\beta,b} := \frac{\partial^2 \mathcal{F}_0}{\partial t_a^\alpha \partial t_b^\beta} \Big|_{t_c^\gamma = \delta_{c,0} u^\gamma}.$$

We obtain

$$\bar{h}_{\alpha,d} := \bar{g}_{\alpha,d}|_{\varepsilon=0} = \int \left( \frac{\partial \mathcal{F}_0}{\partial t_d^\alpha} \Big|_{t_c^\gamma = \delta_{c,0} u^\gamma} \right) dx \stackrel{\text{Eq. (2.2)}}{=} \int \Omega_{\mathbb{1},0;\alpha,d+1} dx.$$

Note that by Equation (2.2) we have

$$(2.5) \quad \frac{\partial \Omega_{\mathbb{1},0;\alpha,d+1}}{\partial u^\gamma} = \Omega_{\gamma,0;\alpha,d}, \quad d \geq -1,$$

where we adopt the convention  $\Omega_{\gamma,0;\alpha,-1} := \eta_{\gamma\alpha}$ . We need to check that for  $d \geq -1$  we have

$$(2.6) \quad \{\cdot, \bar{h}_{\alpha,d}\}_{K_2^{[0]}} = \left( d + \frac{3}{2} + \mu_\alpha \right) \{\cdot, \bar{h}_{\alpha,d+1}\}_{\eta^{-1}\partial_x} + A_\alpha^\beta \{\cdot, \bar{h}_{\beta,d}\}_{\eta^{-1}\partial_x}.$$

Equation (2.6) is equivalent to

$$(2.7) \quad K_2^{[0],\beta\gamma} \frac{\partial \Omega_{\mathbb{1},0;\alpha,d+1}}{\partial u^\gamma} = \left( d + \frac{3}{2} + \mu_\alpha \right) \eta^{\beta\gamma} \partial_x \frac{\partial \Omega_{\mathbb{1},0;\alpha,d+2}}{\partial u^\gamma} + \eta^{\beta\gamma} \partial_x \frac{\partial \Omega_{\mathbb{1},0;\mu,d+1}}{\partial u^\gamma} A_\alpha^\mu.$$

Using Equation (2.5) and the skew-symmetry of the operator  $K_2^{[0]}$  we have

$$K_2^{[0],\beta\gamma} \frac{\partial \Omega_{\mathbb{1},0;\alpha,d+1}}{\partial u^\gamma} = K_2^{[0],\beta\gamma} \Omega_{\gamma,0;\alpha,d} = \left( \partial_x \circ g^{\beta\gamma} + \left( \mu_\beta - \frac{1}{2} \right) \partial_x \Omega^{\beta\gamma} \right) \Omega_{\gamma,0;\alpha,d}.$$

By (2.3) we have  $g^{\beta\gamma} \Omega_{\gamma,0;\alpha,d} = \eta^{\beta\gamma} ((1 - q_\nu) u^\nu + r^\nu) \frac{\partial \Omega_{\gamma,0;\alpha,d+1}}{\partial u^\nu}$  and  $\partial_x \Omega^{\beta\gamma} \Omega_{\gamma,0;\alpha,d} = \eta^{\beta\gamma} \partial_x \Omega_{\gamma,0;\alpha,d+1}$ . Recall also that  $\mu\eta + \eta\mu = 0$ . Therefore, the left-hand side of Equation (2.7) is equal to

$$\eta^{\beta\gamma} \partial_x \left( ((1 - q_\nu) u^\nu + r^\nu) \frac{\partial \Omega_{\gamma,0;\alpha,d+1}}{\partial u^\nu} + \left( -\mu_\gamma - \frac{1}{2} \right) \partial_x \Omega_{\gamma,0;\alpha,d+1} \right).$$

Using Equation (2.5) we rewrite the right-hand side of Equation (2.7) as

$$\eta^{\beta\gamma} \partial_x \left( \left( d + \frac{3}{2} + \mu_\alpha \right) \Omega_{\gamma,0;\alpha,d+1} + \Omega_{\gamma,0;\mu,d} A_\alpha^\mu \right).$$

Therefore, Equation (2.7) would follow from

$$(2.8) \quad ((1 - q_\nu) u^\nu + r^\nu) \frac{\partial \Omega_{\gamma,0;\alpha,d+1}}{\partial u^\nu} = (d + 2 + \mu_\alpha + \mu_\gamma) \Omega_{\gamma,0;\alpha,d+1} + \Omega_{\gamma,0;\mu,d} A_\alpha^\mu.$$

Applying  $\frac{\partial^2}{\partial t_0^\gamma \partial t_{d+1}^\alpha}$  to both sides of Equation (2.4) and setting  $t_p^\nu = \delta_{p,0} u^\nu$  we prove Equation (2.8), hence Equation (2.7), hence Equation (2.6). This completes the proof of the proposition.  $\square$

*Remark 2.2.* The genus 0 part of the DR hierarchy coincides with a principal hierarchy (a genus 0 hierarchy in the language of [DZ98]) associated to the Dubrovin–Frobenius manifold given by the potential  $F$  [Bur15, Section 4.2.2]. The family of principal hierarchies associated to a Dubrovin–Frobenius manifold is parameterized by the choice of deformed flat coordinates  $\theta^\alpha(t^*, z)$  for a certain flat connection on a formal neighbourhood of  $0 \in \mathbb{C}^N$  (see e.g. [DZ98, Equation (2.38)]). In the literature the deformed flat coordinates are often chosen in such a way that they satisfy an additional quasihomogeneity constraint (see e.g. [DZ98, Equation (2.74)]). An analogue of relation (2.6) for the corresponding principal hierarchy is well known [DZ98, Proposition 2 and Equation (2.75)]. The genus 0 part of the DR hierarchy corresponds to a unique choice of the deformed flat coordinates  $\theta^\alpha(t^*, z) = t^\alpha + \sum_{p \geq 1} \theta_p^\alpha(t^*) z^p$  satisfying  $\frac{\partial \theta_p^\alpha}{\partial t^\beta} \Big|_{t^*=0} = 0$ .

## 2.2. The first equation of the bihamiltonian recursion.

**Proposition 2.3.** *Equation (1.10) is true for  $d = -1$ .*

*Proof.* For  $d = -1$  Equation (1.10) is equivalent to

$$K_2^{\beta\gamma} \frac{\delta}{\delta u^\gamma} \left( \int \eta_{\alpha\nu} u^\nu dx \right) = \left( \mu_\alpha + \frac{1}{2} \right) \eta^{\beta\gamma} \partial_x \frac{\delta \bar{g}_{\alpha,0}}{\delta u^\gamma} + A_\alpha^\nu \eta^{\beta\gamma} \partial_x \frac{\delta}{\delta u^\gamma} \left( \int \eta_{\nu\theta} u^\theta dx \right).$$

The variational derivative of  $\int u^* dx$  is constant, therefore, expanding the definition of  $K_2$  we see that the latter equation is equivalent to

$$\widehat{\Omega}(\bar{g})_x^{\beta\gamma} \left( \left( \frac{1}{2} - \mu_\gamma \right) \eta_{\gamma\alpha} \right) = \left( \mu_\alpha + \frac{1}{2} \right) \eta^{\beta\gamma} \partial_x \frac{\delta \bar{g}_{\alpha,0}}{\delta u^\gamma}.$$

Expanding the definition of  $\widehat{\Omega}$  and using the skew-symmetry of  $\mu$  with respect to  $\eta$  we see that the latter equation is equivalent to

$$\left( \mu_\alpha + \frac{1}{2} \right) \eta^{\beta\gamma} \partial_x \frac{\partial}{\partial u^\alpha} \left( \frac{\delta \bar{g}}{\delta u^\gamma} \right) = \left( \mu_\alpha + \frac{1}{2} \right) \eta^{\beta\gamma} \partial_x \frac{\delta \bar{g}_{\alpha,0}}{\delta u^\gamma}.$$

This last equation holds, since

$$\frac{\partial}{\partial u^\alpha} \left( \frac{\delta \bar{g}}{\delta u^\gamma} \right) = \frac{\delta}{\delta u^\gamma} \left( \frac{\partial \bar{g}}{\partial u^\alpha} \right) \stackrel{\text{Eq. (1.8)}}{=} \frac{\delta \bar{g}_{\alpha,0}}{\delta u^\gamma}.$$

□

**2.3. The Schouten–Nijenhuis bracket of two skew-symmetric operators.** In this section we prove that the Schouten–Nijenhuis bracket of the brackets  $\{\cdot, \cdot\}_{K_1}$  and  $\{\cdot, \cdot\}_{K_2}$  considered as bivector fields vanishes. In order to formulate and prove this result, we recall the necessary background regarding the Schouten–Nijenhuis bracket following [LZ13].

**2.3.1. Local multivector fields.** The space of *densities of local multivector fields* (on the formal loop space of  $V$ ) is the associative graded commutative algebra

$$\widehat{\mathcal{A}} := \mathbb{C}[[u^*]][u_{\geq 1}^*, \theta_{*,*}][[\varepsilon]],$$

where the range of the indexes of the new formal variables  $\theta_{\alpha,k}$  is  $1 \leq \alpha \leq N$ ,  $k \geq 0$ . The algebra  $\widehat{\mathcal{A}}$  is graded commutative with respect to the *super gradation*, denoted by  $\deg_\theta$ , which is defined by  $\deg_\theta \theta_{\alpha,k} := 1$  and  $\deg_\theta u_k^\alpha = \deg_\theta \varepsilon := 0$ . Note that the super degree 0 homogeneous component coincides with the space of differential polynomials  $\widehat{\mathcal{A}}^0$ . The super degree  $p$  homogeneous component of  $\widehat{\mathcal{A}}$  is denoted by  $\widehat{\mathcal{A}}^p$ .

The *standard gradation* is extended from  $\widehat{\mathcal{A}}^0 \subset \widehat{\mathcal{A}}$  to  $\widehat{\mathcal{A}}$  by assigning  $\deg \theta_{\alpha,k} := k$ . The homogeneous component of  $\widehat{\mathcal{A}}$  of standard degree  $d$  is denoted by  $\widehat{\mathcal{A}}_d$ , and we also denote

$$\widehat{\mathcal{A}}_d^p := \widehat{\mathcal{A}}_d \cap \widehat{\mathcal{A}}^p.$$

The operator  $\partial_x$  is extended from  $\widehat{\mathcal{A}}^0$  to  $\widehat{\mathcal{A}}$  by

$$\partial_x := \sum_{k \geq 0} u_{k+1}^\alpha \frac{\partial}{\partial u_k^\alpha} + \sum_{k \geq 0} \theta_{\alpha, k+1} \frac{\partial}{\partial \theta_{\alpha, k}}.$$

It increases the standard degree by 1 and preserves the super degree. Therefore, the space

$$\widehat{\Lambda} := \widehat{\mathcal{A}} / (\mathbb{C}[[\varepsilon]] \oplus \text{Im } \partial_x),$$

called the space of *local multivector fields*, still possesses both gradations. The corresponding homogeneous components are denoted by  $\widehat{\Lambda}_d^p$ . If  $f \in \widehat{\mathcal{A}}$ , its image in  $\widehat{\Lambda}$  is denoted by  $\overline{f} = \int f dx$ .

The *variational derivatives*  $\frac{\delta}{\delta u^\alpha}$  and  $\frac{\delta}{\delta \theta_\alpha}$  on  $\widehat{\mathcal{A}}$  are defined by

$$\frac{\delta}{\delta u^\alpha} := \sum_{i \geq 0} (-\partial_x)^i \circ \frac{\partial}{\partial u_i^\alpha} \quad \text{and} \quad \frac{\delta}{\delta \theta_\alpha} := \sum_{i \geq 0} (-\partial_x)^i \circ \frac{\partial}{\partial \theta_{\alpha, i}},$$

and, since they vanish on  $\mathbb{C}[[\varepsilon]] \oplus \text{Im } \partial_x$ , they restrict to  $\widehat{\Lambda}$ :

$$\frac{\delta}{\delta u^\alpha}, \frac{\delta}{\delta \theta_\alpha} : \widehat{\Lambda} \rightarrow \widehat{\mathcal{A}}.$$

There is a bijection between the space of  $N$ -tuples  $\overline{Q} = (Q^1, \dots, Q^N)$ ,  $Q^\alpha \in \widehat{\mathcal{A}}^0$ , and the space  $\widehat{\Lambda}^1$  (the space of *local vector fields*) given by

$$\overline{Q} \mapsto V_{\overline{Q}} := \int Q^\alpha \theta_\alpha dx \in \widehat{\Lambda}^1.$$

There is also a bijection between the space of skew-symmetric operators  $K = (K^{\alpha\beta})$ ,  $K^{\alpha\beta} = \sum_{s \geq 0} K_s^{\alpha\beta} \partial_x^s$ ,  $K^* = -K$ , and the space  $\widehat{\Lambda}^2$  (the space of *local bivector fields*) given by

$$K \mapsto B_K := \frac{1}{2} \int \sum_{s \geq 0} K_s^{\alpha\beta} \theta_{\alpha, 0} \theta_{\beta, s} dx \in \widehat{\Lambda}^2.$$

**Definition 2.4.** The *Schouten–Nijenhuis bracket*  $[\cdot, \cdot] : \widehat{\Lambda}^p \times \widehat{\Lambda}^q \rightarrow \widehat{\Lambda}^{p+q-1}$  is defined by

$$[P, Q] := \int \left( \frac{\delta P}{\delta \theta_\alpha} \frac{\delta Q}{\delta u^\alpha} + (-1)^p \frac{\delta P}{\delta u^\alpha} \frac{\delta Q}{\delta \theta_\alpha} \right) dx.$$

**Lemma 2.5** ([LZ11]). *For any  $P \in \widehat{\Lambda}^p$ ,  $Q \in \widehat{\Lambda}^q$ ,  $R \in \widehat{\Lambda}^r$  the Schouten–Nijenhuis bracket satisfies the properties*

$$\begin{aligned} [P, Q] &= (-1)^{pq} [Q, P], \\ (-1)^{pr} [[P, Q], R] + (-1)^{rq} [[R, P], Q] + (-1)^{qp} [[Q, R], P] &= 0. \end{aligned}$$

A skew-symmetric operator  $K$  is *Poisson* if and only if  $[B_K, B_K] = 0$ . For a Poisson operator  $K$  and any  $R \in \widehat{\Lambda}$  we have

$$(2.9) \quad [[R, B_K], B_K] = 0.$$

For a skew-symmetric operator  $K$ , an  $N$ -tuple  $\overline{Q} \in (\widehat{\mathcal{A}}^0)^N$ , and a local functional  $\overline{f} \in \widehat{\Lambda}^0$  we have the following properties:

$$\begin{aligned} \frac{\delta B_K}{\delta \theta_\alpha} &= \sum_{s \geq 0} K_s^{\alpha\beta} \theta_{\beta, s} \in \widehat{\mathcal{A}}^1; \\ [\overline{f}, B_K] &= -V_{\overline{f}}, \quad P^\alpha = \sum_{s \geq 0} K_s^{\alpha\beta} \partial_x^s \frac{\delta \overline{f}}{\delta u^\beta}; \end{aligned}$$

$$(2.10) \quad [V_{\overline{Q}}, B_K] = -B_{\widetilde{K}}, \quad \widetilde{K}^{\alpha\beta} = L_\mu(Q^\alpha) \circ K^{\mu\beta} + K^{\alpha\nu} \circ L_\nu^*(Q^\beta) - \sum_{p, s \geq 0} (\partial_x^p Q^\gamma) \frac{\partial K_s^{\alpha\beta}}{\partial u_p^\gamma} \partial_x^s.$$

Note that  $\{\bar{f}, \bar{g}\}_K = [[B_K, \bar{f}], \bar{g}]$  for any  $\bar{f}, \bar{g} \in \widehat{\Lambda}^0$ .

2.3.2. *The Schouten–Nijenhuis bracket of  $K_1$  and  $K_2$ .* The fact that the skew-symmetric operator  $K_2$  is Poisson and is compatible with the operator  $K_1$  is equivalent to the system of the following two equations:

$$[B_{K_2}, B_{K_2}] = 0 \quad \text{and} \quad [B_{K_2}, B_{K_1}] = 0.$$

**Proposition 2.6.** *We have  $[B_{K_2}, B_{K_1}] = 0$ .*

*Proof.* Equation (2.9) implies that it is sufficient to represent  $B_{K_2}$  as  $[V_{\bar{R}}, B_{K_1}]$  for some local vector field. We do it below in Lemma 2.7.  $\square$

Introduce the *higher Euler operators*  $\mathcal{T}_{\alpha,k}: \widehat{\mathcal{A}}^0 \rightarrow \widehat{\mathcal{A}}^0$ ,  $1 \leq \alpha \leq N$ ,  $k \geq 0$ , by

$$\mathcal{T}_{\alpha,k} := \sum_{n \geq k} \binom{n}{k} (-\partial_x)^{n-k} \circ \frac{\partial}{\partial u_n^\alpha}.$$

Clearly,  $\mathcal{T}_{\alpha,0} = \frac{\delta}{\delta u^\alpha}$ . The operators  $\mathcal{T}_{\alpha,k}$  satisfy the property  $\mathcal{T}_{\alpha,k+1} \circ \partial_x = \mathcal{T}_{\alpha,k}$ ,  $k \geq 0$  (see [KMGZ70], [BPS12b, Section 5.1], [LZ11, Section 2.1]).

Let us choose a representative  $g \in \widehat{\mathcal{A}}^0$  for the local functional  $\bar{g} \in \widehat{\Lambda}^0$  and define an  $N$ -tuple  $\bar{R} = (R^1, \dots, R^N) \in (\widehat{\mathcal{A}}^0)^N$  by

$$R^\alpha := \eta^{\alpha\beta} \left( \left( -\frac{1}{2} - \mu_\beta \right) \frac{\delta \bar{g}}{\delta u^\beta} - \frac{1}{2} A_{\beta\gamma} u^\gamma + \partial_x \mathcal{T}_{\beta,1}(g) \right).$$

**Lemma 2.7.** *We have  $B_{K_2} = [V_{\bar{R}}, B_{K_1}]$ .*

*Proof.* Split  $R^\alpha$  into the sum  $R^\alpha = R_1^\alpha + R_2^\alpha + R_3^\alpha$ , where

$$R_1^\alpha := -\eta^{\alpha\beta} \left( \frac{1}{2} + \mu_\beta \right) \frac{\delta \bar{g}}{\delta u^\beta}, \quad R_2^\alpha := -\frac{1}{2} A_\gamma^\alpha u^\gamma, \quad \text{and} \quad R_3^\alpha := \eta^{\alpha\beta} \partial_x \mathcal{T}_{\beta,1}(g).$$

Using Equation (1.11) we split  $K_2$  into the sum of skew-symmetric operators,  $K_2 = K_2^{(1)} + K_2^{(2)} + K_2^{(3)}$ , where

$$\begin{aligned} K_2^{(1)} &:= \partial_x \circ \widehat{\Omega}(\bar{g}) \circ \left( \frac{1}{2} - \mu \right) + \left( \frac{1}{2} - \mu \right) \circ \widehat{\Omega}(\bar{g}) \circ \partial_x, \\ K_2^{(2)} &:= \eta^{-1} A \eta^{-1} \partial_x, \\ K_2^{(3)} &:= \partial_x \circ \widehat{\Omega}^1(\bar{g}) \circ \partial_x. \end{aligned}$$

We prove below for  $i = 1, 2, 3$  that  $B_{K_2^{(i)}} = [V_{\bar{R}_i}, B_{K_1}]$ .

For  $i = 1$  we have

$$\begin{aligned} K_2^{(1);\alpha\beta} &= \left( \partial_x \circ \widehat{\Omega}(\bar{g})^* \circ \left( \frac{1}{2} - \mu \right) + \left( \frac{1}{2} - \mu \right) \circ \widehat{\Omega}(\bar{g}) \circ \partial_x \right)^{\alpha\beta} \\ &= \eta^{\alpha\nu} \partial_x \circ L_\nu^* \left( \eta^{\beta\gamma} \frac{\delta \bar{g}}{\delta u^\gamma} \left( \frac{1}{2} - \mu_\beta \right) \right) + L_\nu \left( \eta^{\alpha\gamma} \frac{\delta \bar{g}}{\delta u^\gamma} \left( \frac{1}{2} - \mu_\alpha \right) \right) \circ \eta^{\nu\beta} \partial_x \\ &= \eta^{\alpha\nu} \partial_x \circ L_\nu^* \left( \eta^{\beta\gamma} \frac{\delta \bar{g}}{\delta u^\gamma} \left( \frac{1}{2} + \mu_\gamma \right) \right) + L_\nu \left( \eta^{\alpha\gamma} \frac{\delta \bar{g}}{\delta u^\gamma} \left( \frac{1}{2} + \mu_\gamma \right) \right) \circ \eta^{\nu\beta} \partial_x. \end{aligned}$$

Then Equation (2.10) implies that  $B_{K_2^{(1)}} = [V_{\bar{R}_1}, B_{K_1}]$ .

For  $i = 2$  we have

$$K_2^{(2);\alpha\beta} = \frac{1}{2} (A_\nu^\alpha \circ \eta^{\nu\beta} \partial_x + \eta^{\alpha\nu} \partial_x \circ A_\nu^\beta),$$

which implies  $B_{K_2^{(2)}} = [V_{\bar{R}_2}, B_{K_1}]$ .

For  $i = 3$  we have (cf. the computation in the proof of Lemma 1.5)

$$\begin{aligned}
K_2^{(3);\alpha\beta} &= \partial_x \circ \eta^{\alpha\mu} \eta^{\beta\nu} \sum_{i=0}^{\infty} (i+1) \frac{\partial}{\partial u_{i+1}^\nu} \left( \frac{\delta \bar{g}}{\delta u^\mu} \right) \partial_x^{i+1} \\
&= \eta^{\alpha\mu} \eta^{\beta\nu} \sum_{i,k,m=0}^{\infty} \sum_{l=0}^{i+1} (i+1) (-1)^{l+1} \binom{m+k+l}{k,l} (-\partial_x)^{m+1} \circ \left( \frac{\partial^2 g}{\partial u_{i+1-l}^\nu \partial u_{m+k+l}^\mu} \right) \partial_x^{i+1+k} \\
&= \eta^{\alpha\mu} \eta^{\beta\nu} \sum_{m,n \geq 0} (-\partial_x)^{m+1} \circ \left( (m+1) \frac{\partial^2 g}{\partial u_{m+1}^\mu \partial u_n^\nu} - (n+1) \frac{\partial^2 g}{\partial u_m^\mu \partial u_{n+1}^\nu} \right) \circ \partial_x^{n+1} \\
&= \left( \sum_{m \geq 0} (m+1) (-\partial_x)^{m+1} \circ L_\nu \left( \eta^{\alpha\mu} \frac{\partial g}{\partial u_{m+1}^\mu} \right) \right) \circ \eta^{\nu\beta} \partial_x \\
&\quad + \eta^{\alpha\mu} \partial_x \circ \left( \sum_{n \geq 0} L_\mu^* \left( \eta^{\beta\nu} \frac{\partial g}{\partial u_{n+1}^\nu} \right) \circ (n+1) \partial_x^{n+1} \right) \\
&= L_\nu (-\eta^{\alpha\mu} \partial_x \mathcal{T}_{\mu,1}(g)) \circ \eta^{\nu\beta} \partial_x + \eta^{\alpha\mu} \partial_x \circ L_\mu^* (-\eta^{\beta\nu} \partial_x \mathcal{T}_{\nu,1}(g)),
\end{aligned}$$

which implies  $B_{K_2^{(3)}} = [V_{R_3}, B_{K_1}]$ . In this computation we pass from the second line to the third one using the following straightforward combinatorial identity:

$$\begin{aligned}
&\sum_{\substack{i+k=n \\ k+l=p}} (i+1) (-1)^{l+1} \binom{m+k+l}{k,l} \\
&= -(n+1) \sum_{k+l=p} (-1)^l \binom{m+k+l}{k,l} + \sum_{k+l=p} k (-1)^l \binom{m+k+l}{k,l} \\
&= -(n+1) \delta_{p,0} + (m+1) \delta_{p,1}.
\end{aligned}$$

□

### 3. EVIDENCE II: CENTRAL INVARIANTS

In this section, assuming that the first part of Conjecture 1.13, is true we compute the central invariants of the pair of operators  $(K_1, K_2)$ . We also discuss the relation between Conjecture 1.13, the conjecture on the equivalence of the DR hierarchy and the Dubrovin–Zhang hierarchy, and the Dubrovin–Zhang conjecture on existence of a bihamiltonian structure for the Dubrovin–Zhang hierarchy.

**3.1. Miura transformations.** Let us discuss changes of coordinates on the formal loop space. In order to distinguish rings of differential polynomials in different variables, let  $\mathcal{A}_w^0$  denote the ring of differential polynomials in variables  $w^1, \dots, w^N$ . The same notation is adopted for the extension  $\widehat{\mathcal{A}}_w^0$  and for the spaces of local functionals  $\Lambda_w^0$  and  $\widehat{\Lambda}_w^0$ .

**Definition 3.1.** Changes of variables of the form

(3.1)

$$u^\alpha \mapsto \tilde{u}^\alpha(u_*^*, \varepsilon) = \sum_{k \geq 0} \varepsilon^k f_k^\alpha(u_*^*), \quad 1 \leq \alpha \leq N, \quad \text{where } f_k^\alpha \in \mathcal{A}_{u;k}^0, \quad \det \left( \frac{\partial f_0^\alpha}{\partial u^\beta} \right) \Big|_{u^*=0} \neq 0,$$

are called *Miura transformations*.

We say that a Miura transformation is *close to identity* if  $f_0^\alpha = u^\alpha$ .

Under a Miura transformation any differential polynomial  $P(u_*^*, \varepsilon) \in \widehat{\mathcal{A}}_w^0$  can be rewritten as a formal power series in  $\varepsilon$  whose coefficients are polynomials in the variables  $\tilde{u}_i^\alpha$ ,  $i > 0$ ,

with coefficients in the ring of formal power series in the shifted variables  $\tilde{u}^\alpha - \tilde{u}_{\text{orig}}^\alpha$ , where  $\tilde{u}_{\text{orig}}^\alpha := \tilde{u}^\alpha(u^*, \varepsilon)|_{u^*=\varepsilon=0} = f_0^\alpha|_{u^*=0}$ . We introduce the notation

$$\widehat{\mathcal{A}}_{\tilde{u}}^{\tilde{u}_{\text{orig}};0} := \mathbb{C}[[\tilde{u}^\alpha - \tilde{u}_{\text{orig}}^\alpha][[\tilde{u}_{\geq 1}^\alpha][[\varepsilon]]], \quad \tilde{u}_{\text{orig}} = (\tilde{u}_{\text{orig}}^1, \dots, \tilde{u}_{\text{orig}}^N),$$

and still call the elements of the ring  $\widehat{\mathcal{A}}_{\tilde{u}}^{\tilde{u}_{\text{orig}};0}$  differential polynomials. Let  $\widehat{\Lambda}_{\tilde{u}}^{\tilde{u}_{\text{orig}};0}$  denote the corresponding space of local functionals. A differential polynomial  $P(u^*, \varepsilon) \in \widehat{\mathcal{A}}_{\tilde{u}}$  rewritten in the variables  $\tilde{u}^\alpha$  is denoted by  $P(\tilde{u}_*^*, \varepsilon) \in \widehat{\mathcal{A}}_{\tilde{u}}^{\tilde{u}_{\text{orig}};0}$ . The condition  $f_k^\alpha \in \mathcal{A}_{u;k}^0$  guarantees that if  $P(u^*, \varepsilon) \in \widehat{\mathcal{A}}_{u;d}^0$ , then  $P(\tilde{u}_*^*, \varepsilon) \in \widehat{\mathcal{A}}_{\tilde{u};d}^{\tilde{u}_{\text{orig}};0}$ . This means that a Miura transformation defines an isomorphism  $\widehat{\mathcal{A}}_{u;d}^0 \simeq \widehat{\mathcal{A}}_{\tilde{u};d}^{\tilde{u}_{\text{orig}};0}$ . It also induces an isomorphism  $\widehat{\Lambda}_{u;d}^0 \simeq \widehat{\Lambda}_{\tilde{u};d}^{\tilde{u}_{\text{orig}};0}$ . The image of a local functional  $\bar{h}[u] \in \widehat{\Lambda}_{u;d}^0$  under this isomorphism is denoted by  $\bar{h}[\tilde{u}] \in \widehat{\Lambda}_{\tilde{u};d}^{\tilde{u}_{\text{orig}};0}$ .

Let us describe the action of Miura transformations on Hamiltonian hierarchies. Consider a Hamiltonian hierarchy (1.3) and a Miura transformation (3.1). Then in the new variables  $\tilde{u}^\alpha$  the system (1.3) reads:

$$\frac{\partial \tilde{u}^\alpha}{\partial \tau_i} = K_{\tilde{u}}^{\alpha\mu} \frac{\delta \bar{h}_i[\tilde{u}]}{\delta \tilde{u}^\mu}, \quad K_{\tilde{u}}^{\alpha\beta} = L_\mu(\tilde{u}^\alpha(u^*, \varepsilon)) \circ K^{\mu\nu} \circ L_\nu^*(\tilde{u}^\beta(u^*, \varepsilon)).$$

**3.2. The central invariants of a pair of compatible Poisson operators.** Consider a pair of compatible Poisson operators  $(P_1, P_2)$  of degree 1:

$$P_a^{\alpha\beta} = g_a^{\alpha\beta} \partial_x + \Gamma_{a;\gamma}^{\alpha\beta} u_x^\gamma + O(\varepsilon), \quad g_a^{\alpha\beta}, \Gamma_{a;\gamma}^{\alpha\beta} \in \mathbb{C}[[u^*]], \quad a = 1, 2.$$

Assume that

$$\det(g_1^{\alpha\beta}) \Big|_{u^*=0} \neq 0$$

and that there exist formal power series  $\widehat{u}^i(u^*) \in \mathbb{C}[[u^*]]$ ,  $1 \leq i \leq N$ , such that

$$(3.2) \quad \det \left( g_2^{\alpha\beta} - \widehat{u}^i g_1^{\alpha\beta} \right) = 0, \quad 1 \leq i \leq N, \\ \det \left( \frac{\partial \widehat{u}^i}{\partial u^\alpha} \right) \Big|_{u^*=0} \neq 0.$$

Clearly, if such an  $N$ -tuple  $(\widehat{u}^1, \dots, \widehat{u}^N)$  exists, then it is unique up to permutations. Condition (3.2) implies that the functions  $\widehat{u}^i(u^*)$  can be used as a system of coordinates. These coordinates are called the *canonical coordinates* of the pair of operators  $(P_1, P_2)$ .

Let us use Latin superscripts to indicate components of the operators  $P_1, P_2$  in canonical coordinates,  $P_a^{ij} := P_{a;\widehat{u}}^{ij}$ . The operators  $P_1$  and  $P_2$  have the following form in canonical coordinates (see e.g. [DLZ06]):

$$(3.3) \quad P_a^{ij} = \sum_{k \geq 0} \varepsilon^k \sum_{l=0}^{k+1} P_{a;l}^{[k],ij} \partial_x^l, \quad \text{where}$$

$$(3.4) \quad P_{1;1}^{[0],ij} = \delta_{ij} f^i, \quad P_{2;1}^{[0],ij} = \delta_{ij} \widehat{u}^i f^i, \quad f^i \in \mathbb{C}[[\widehat{u}^j - \widehat{u}_{\text{orig}}^j]], \quad \widehat{u}_{\text{orig}}^j = \widehat{u}^j(u^*) \Big|_{u^*=0}.$$

The *central invariants* of the pair of operators  $(P_1, P_2)$  are the formal power series  $c_i(\widehat{u}^i) \in \mathbb{C}[[\widehat{u}^i - \widehat{u}_{\text{orig}}^i]]$ ,  $1 \leq i \leq N$ , defined by (see e.g. [DLZ06])

$$(3.5) \quad c_i(\widehat{u}^i) := \frac{1}{3(f^i)^2} \left( P_{2;3}^{[2],ii} - \widehat{u}^i P_{1;3}^{[2],ii} + \sum_{k \neq i} \frac{\left( P_{2;2}^{[1],ki} - \widehat{u}^i P_{1;2}^{[1],ki} \right)^2}{(\widehat{u}^k - \widehat{u}^i) f^k} \right).$$

*Remark 3.2.* In [DLZ06] the proof that the right-hand side of this formula is a formal power series that doesn't depend on  $\widehat{u}^j$  with  $j \neq i$  uses the assumption that  $\widehat{u}_{\text{orig}}^j \neq 0$  for all  $j$  and  $\widehat{u}_{\text{orig}}^j \neq \widehat{u}_{\text{orig}}^k$  for all  $j \neq k$ . Note, however, that the same argument allows to prove that  $\frac{\partial c_i}{\partial \widehat{u}^j} = 0$



for  $j \neq i$  without this assumption. Also, Equation (3.5) implies that  $c_i = \frac{g_i(\widehat{u}^1, \dots, \widehat{u}^N)}{\prod_{j \neq i} (\widehat{u}^j - \widehat{u}^i)}$  for some  $g_i(\widehat{u}^1, \dots, \widehat{u}^N) \in \mathbb{C}[[\widehat{u}^1 - \widehat{u}_{\text{orig}}^1, \dots, \widehat{u}^N - \widehat{u}_{\text{orig}}^N]]$ . For any fixed  $j \neq i$ , the fact that  $\frac{\partial c_i}{\partial \widehat{u}^j} = 0$  implies that  $g_i = (\widehat{u}^j - \widehat{u}^i) \frac{\partial g_i}{\partial \widehat{u}^j}$ . Therefore,  $g_i$  is divisible by  $\widehat{u}^j - \widehat{u}^i$  for any  $j \neq i$  in the ring  $\mathbb{C}[[\widehat{u}^1 - \widehat{u}_{\text{orig}}^1, \dots, \widehat{u}^N - \widehat{u}_{\text{orig}}^N]]$ . Thus, the quotient  $\frac{g_i(\widehat{u}^1, \dots, \widehat{u}^N)}{\prod_{j \neq i} (\widehat{u}^j - \widehat{u}^i)}$  belongs to the ring  $\mathbb{C}[[\widehat{u}^i - \widehat{u}_{\text{orig}}^i]]$ .

The central invariants of the pair  $(P_1, P_2)$  are invariant under Miura transformations of the variables  $\widehat{u}^i$  close to identity. Moreover, in case  $\widehat{u}_{\text{orig}}^i \neq 0$  and  $\widehat{u}_{\text{orig}}^i \neq \widehat{u}_{\text{orig}}^j$  for  $i \neq j$ , the central invariants classify pairs of compatible Poisson operators of the form (3.3)–(3.4) with fixed functions  $f^i$  under Miura transformations of the variables  $\widehat{u}^i$  close to identity (see [LZ05, DLZ06, LZ13, CPS18, CKS18]).

**3.3. The central invariants of the pair  $(K_1, K_2)$ .** Consider a homogeneous semisimple CohFT. There exist canonical coordinates  $\widehat{u}^i(t^*) \in \mathbb{C}[[t^*]]$ ,  $1 \leq i \leq N$ , for the associated Dubrovin–Frobenius manifold:

$$\frac{\partial}{\partial \widehat{u}^i} \circ \frac{\partial}{\partial \widehat{u}^j} = \delta_{ij} \frac{\partial}{\partial \widehat{u}^i},$$

where by  $\circ$  we denote here the multiplication in the tangent spaces of the Dubrovin–Frobenius manifold. After an appropriate shift  $\widehat{u}^i \mapsto \widehat{u}^i + a^i$ ,  $a^i \in \mathbb{C}$ , the Euler vector field  $E$  written in canonical coordinates is given by

$$E = \sum_{i=1}^N \widehat{u}^i \frac{\partial}{\partial \widehat{u}^i}.$$

Moreover,  $\det(g^{\alpha\beta} - \widehat{u}^i \eta^{\alpha\beta}) = 0$  for  $1 \leq i \leq N$ , where  $g^{\alpha\beta}$  is given by Equation (2.1) and we identify  $t^\gamma = u^\gamma$ .

Therefore, if we assume that the operator  $K_2$  is Poisson, then by Proposition 2.6 the operators  $K_1$  and  $K_2$  form a pair of compatible Poisson operators and we can compute the associated central invariants.

**Proposition 3.3.** *Suppose that the operator  $K_2$  corresponding to our homogeneous semisimple CohFT is Poisson. Then all the central invariants of the pair of operators  $(K_1, K_2)$  are equal to  $\frac{1}{24}$ .*

*Proof.* In canonical coordinates the operators  $K_1$  and  $K_2$  have the form

$$\begin{aligned} K_1^{ij} &= K_{1;1}^{[0],ij} \partial_x + K_{1;0}^{[0],ij}, & K_{1;1}^{[0],ij} &= \delta_{ij} f^i, & f^i &\in \mathbb{C}[[\widehat{u}^j - \widehat{u}_{\text{orig}}^j]], \\ K_2^{ij} &= K_{2;1}^{[0],ij} \partial_x + K_{2;0}^{[0],ij} + \sum_{k \geq 1} \varepsilon^{2k} \sum_{l=0}^{2k+1} K_{2;l}^{[2k],ij} \partial_x^l, & K_{2;1}^{[0],ij} &= \delta_{ij} \widehat{u}^i f^i. \end{aligned}$$

Therefore,

$$c_i = \frac{K_{2;3}^{[2],ii}}{3(f^i)^2}.$$

Consider the expansion

$$\bar{g} = \sum_{g \geq 0} \bar{g}^{[2g]} \varepsilon^{2g}, \quad \bar{g}^{[2g]} \in \Lambda_{2g}^0.$$

We have

$$\begin{aligned} K_2^{[2],\alpha\beta} &= \eta^{\alpha\mu} \eta^{\beta\nu} \left( \left( \frac{1}{2} - \mu_\beta \right) \partial_x \circ L_\nu \left( \frac{\delta \bar{g}^{[2]}}{\delta u^\mu} \right) + \left( \frac{1}{2} - \mu_\alpha \right) L_\nu \left( \frac{\delta \bar{g}^{[2]}}{\delta u^\mu} \right) \circ \partial_x \right. \\ &\quad \left. + \partial_x \circ L_\nu^1 \left( \frac{\delta \bar{g}^{[2]}}{\delta u^\mu} \right) \circ \partial_x \right), \end{aligned}$$

which implies that

$$K_{2;3}^{[2],\alpha\beta} = (3 - \mu_\alpha - \mu_\beta) \eta^{\alpha\mu} \eta^{\beta\nu} \frac{\partial}{\partial u_2^\nu} \frac{\delta \bar{g}^{[2]}}{\delta u^\mu}.$$

In [BDGR18, Lemma 8.1] it is proved that

$$\bar{g}^{[2]} = -\frac{1}{48} \int c_{\theta\xi}^\theta c_{\alpha\beta}^\xi u_x^\alpha u_x^\beta dx,$$

where we recall that  $c_{\beta\gamma}^\alpha = \eta^{\alpha\mu} \frac{\partial^3 F}{\partial t^\mu \partial t^\beta \partial t^\gamma}$  and we use the identification  $t^\theta = u^\theta$ . Therefore,  $\frac{\partial}{\partial u_\mu^\nu} \frac{\delta \bar{g}^{[2]}}{\partial u^\mu} = \frac{1}{24} c_{\theta\xi}^\theta c_{\mu\nu}^\xi$  and, hence,

$$K_{2;3}^{[2],\alpha\beta} = \frac{1}{24} (3 - \mu_\alpha - \mu_\beta) c_{\theta\xi}^\theta c^{\xi\alpha\beta},$$

where we raise the indices in the tensor  $c_{\beta\gamma}^\alpha$  using the metric  $\eta$ .

Note that

$$K_{2;3}^{[2],ij} = \frac{\partial \widehat{u}^i}{\partial u^\alpha} \frac{\partial \widehat{u}^j}{\partial u^\beta} K_{2;3}^{[2],\alpha\beta},$$

which means that the collection of functions  $K_{2;3}^{[2],\alpha\beta}$  is transformed to the canonical coordinates as a tensor. We use Latin indices for the expressions of the operator  $\mu$ , the metric  $\eta$ , and the tensor  $c_{\beta\gamma}^\alpha$  in canonical coordinates. We have  $\eta^{ij} = \delta_{ij} f^i$ ,  $c_{jk}^i = \delta_{ij} \delta_{jk}$ , and  $c^{ijk} = (f^i)^2 \delta_{ij} \delta_{jk}$ . Therefore,

$$K_{2;3}^{[2],ij} = \frac{1}{24} (3\delta_{ij} (f^i)^2 - \mu_j^i (f^j)^2 - (f^i)^2 \mu_i^j).$$

Equation (1.9) implies that  $\mu_j^i f^j + f^i \mu_i^j = 0$ , hence  $\mu_i^i = 0$ . We conclude that  $K_{2;3}^{[2],ii} = \frac{(f^i)^2}{8}$ , hence  $c_i = \frac{1}{24}$ .  $\square$

**3.4. Relation with the Dubrovin–Zhang hierarchy.** Beside the DR hierarchy, which we discuss in this paper, there is another Hamiltonian hierarchy associated to a semisimple homogeneous CohFT, called the *Dubrovin–Zhang (DZ) hierarchy* or the *hierarchy of topological type* [DZ01, BPS12b]. It has the same dispersionless part as the DR hierarchy. By a conjecture in [Bur15], called the *DR/DZ equivalence conjecture* (see also a stronger version in [BDGR18]), it is related to the DR hierarchy by a Miura transformation that is close to identity. A long-standing open conjecture proposed by Dubrovin and Zhang claims that the DZ hierarchy possesses a bihamiltonian structure.

We see that the Dubrovin–Zhang conjecture follows from the DR/DZ equivalence conjecture and Conjecture 1.13. The Dubrovin–Zhang conjecture is checked at the approximation up to  $\varepsilon^2$  [DZ98], and the central invariants of the resulting pair of compatible Poisson operators are all equal to  $\frac{1}{24}$  (see e.g. [Liu18]), which agrees with Proposition 3.3.

#### 4. EVIDENCE III: EXPLICIT EXAMPLES

In this section we prove Conjecture 1.13 in several special cases, for particular semisimple homogeneous CohFTs.

**4.1. The trivial cohomological field theory.** Consider the trivial CohFT with  $N = 1$ ,  $V = \langle e_1 \rangle$ ,  $e = e_1$ ,  $\eta_{1,1} = 1$ , and  $c_{g,n}^{\text{triv}}(e_1^{\otimes n}) = 1 \in H^*(\overline{\mathcal{M}}_{g,n})$ . Let  $u_n$  denote  $u_n^1$ . The corresponding DR hierarchy coincides with the KdV hierarchy,

$$\bar{g}_{1,d} = \bar{h}_d^{\text{KdV}}, \quad d \geq -1$$

(see [Bur15, Section 4.3.1]), where the sequence of Hamiltonians  $\bar{h}_d^{\text{KdV}} \in \widehat{\Lambda}_0^0$ ,  $d \geq -1$ , of the KdV hierarchy is uniquely determined by the properties

$$\begin{aligned} \bar{h}_1^{\text{KdV}} &= \int \left( \frac{u^3}{6} + \varepsilon^2 \frac{uu_{xx}}{24} \right) dx, \\ \bar{h}_d^{\text{KdV}} &= \int \left( \frac{u^{d+2}}{(d+2)!} + O(\varepsilon^2) \right) dx, \quad d \geq -1, \end{aligned}$$

and the commutation relations

$$\left\{ \overline{h}_d^{\text{KdV}}, \overline{h}_1^{\text{KdV}} \right\}_{\partial_x} = 0, \quad d \geq -1,$$

with respect to the Poisson bracket  $\{\cdot, \cdot\}_{\partial_x}$ .

The trivial CohFT is homogeneous with  $q_1 = 0$ ,  $\delta = 0$ ,  $A = 0$ ,  $\mu_1 = 0$ . We have

$$\overline{g} = \int \left( \frac{u^3}{6} + \varepsilon^2 \frac{uu_{xx}}{48} \right) dx, \quad \frac{\delta \overline{g}}{\delta u} = \frac{u^2}{2} + \varepsilon^2 \frac{u_{xx}}{24}, \quad \widehat{\Omega}(\overline{g}) = u + \frac{\varepsilon^2}{24} \partial_x^2, \quad \widehat{\Omega}^1(\overline{g}) = \frac{\varepsilon^2}{12} \partial_x,$$

which, using formula (1.11), gives

$$K_2 = u \partial_x + \frac{1}{2} u_x + \frac{\varepsilon^2}{8} \partial_x^3.$$

The fact that the pair  $(\partial_x, u \partial_x + \frac{1}{2} u_x + \frac{\varepsilon^2}{8} \partial_x^3)$  is a pair of compatible Poisson operators and that the Hamiltonians  $\overline{h}_d^{\text{KdV}}$  satisfy the relations

$$\left\{ \cdot, \overline{h}_d^{\text{KdV}} \right\}_{K_2} = \left( d + \frac{3}{2} \right) \left\{ \cdot, \overline{h}_{d+1}^{\text{KdV}} \right\}_{\partial_x}, \quad d \geq -1,$$

is a standard fact in the theory of the KdV hierarchy (see e.g. [Dic03]). Thus, Conjecture 1.13 holds for the trivial CohFT.

**4.2. Higher  $r$ -spin theories.** Let  $r \geq 2$  and consider an  $(r-1)$ -dimensional complex vector space  $V$  with a fixed basis  $e_1, \dots, e_{r-1}$ . Let  $\eta_{\alpha\beta} := \delta_{\alpha+\beta, r}$ . There exists a unique CohFT (see e.g. [PPZ15])  $c_{g,n}^{r\text{-spin}}: V^{\otimes n} \rightarrow H^{\text{even}}(\overline{\mathcal{M}}_{g,n})$  satisfying the following properties:

- (1)  $c_{g,n}^{r\text{-spin}}(\otimes_{i=1}^n e_{\alpha_i}) = 0$  if  $r \nmid (g-1)(r-2) + \sum_{i=1}^n (\alpha_i - 1)$ ;
- (2)  $\deg c_{g,n}^{r\text{-spin}}(\otimes_{i=1}^n e_{\alpha_i}) = 2 \frac{(g-1)(r-2) + \sum_{i=1}^n (\alpha_i - 1)}{r}$  if  $r \mid (g-1)(r-2) + \sum_{i=1}^n (\alpha_i - 1)$ ;
- (3)  $c_{0,3}^{r\text{-spin}}(e_\alpha \otimes e_\beta \otimes e_\gamma) = \delta_{\alpha+\beta+\gamma, r+1} \in H^0(\overline{\mathcal{M}}_{0,3}) \cong \mathbb{C}$ ;
- (4)  $c_{0,4}^{r\text{-spin}}(e_2^{\otimes 2} \otimes e_{r-1}^{\otimes 2}) = \frac{1}{r} [\text{pt}] \in H^2(\overline{\mathcal{M}}_{0,4})$  if  $r \geq 3$ .

Here [pt] denotes the cohomology class dual to a point in  $\overline{\mathcal{M}}_{0,4}$ . This CohFT is called the  $r$ -spin CohFT. It is homogeneous with  $q_\alpha = \frac{\alpha-1}{r}$ ,  $r^\alpha = 0$ ,  $\delta = \frac{r-2}{r}$ ,  $\mu_\alpha = \frac{2\alpha-r}{2r}$ ,  $A_{\alpha\beta} = 0$ . The 2-spin CohFT coincides with the trivial CohFT.

In this section we check Conjecture 1.13 for the  $r$ -spin CohFT for  $r = 3, 4, 5$ . To this end, we first describe the construction of the Gelfand–Dickey hierarchy, then we present its relation (through the Dubrovin–Zhang hierarchy) to the DR hierarchy for the  $r$ -spin theory for  $r = 3, 4, 5$  proved in [BG16], and then we finally check Conjecture 1.13 for the  $r$ -spin CohFT for these values of  $r$ .

*Remark 4.1.* Note that for any differential polynomial  $f \in \mathcal{A}^0$  there exists a unique differential polynomial  $\tilde{f} \in \widehat{\mathcal{A}}_0^0$  such that  $\tilde{f}|_{\varepsilon=0} = f$ . This gives a natural inclusion  $\mathcal{A}^0 \hookrightarrow \widehat{\mathcal{A}}_0^0$ . The same remark is true for local functionals. Regarding Poisson brackets, note that any Poisson operator  $K$ ,  $K^{\alpha\beta} = \sum_{i \geq 0} K_i^{\alpha\beta} \partial_x^i$ ,  $K_i^{\alpha\beta} \in \mathcal{A}^0$ , can be expressed in a unique way as the sum of homogeneous operators:  $K^{\alpha\beta} = \sum_{d \geq 0} \left( \sum_{i+j=d} K_{i,j}^{\alpha\beta} \partial_x^i \right)$ , where  $K_{i,j}^{\alpha\beta} \in \mathcal{A}_{d-i}^0$ . If its degree zero part vanishes, i.e., if  $K_{0,0}^{\alpha\beta} = 0$ , then we can naturally associate to such a Poisson operator  $K$  the Poisson operator  $\tilde{K}^{\alpha\beta} = \sum_{d \geq 1} \varepsilon^{d-1} \left( \sum_{i+j=d} K_{i,j}^{\alpha\beta} \partial_x^i \right)$  that gives a Poisson bracket of degree 1 on the space  $\widehat{\Lambda}^0$ . We silently use this correspondence throughout this section.

**4.2.1. Gelfand–Dickey hierarchy.** We review the construction of the Gelfand–Dickey hierarchy and its bihamiltonian structure following [Dic03].

Consider variables  $f_0, f_1, \dots, f_{r-2}$ . A pseudo-differential operator  $A$  is a Laurent series

$$A = \sum_{n=-\infty}^m a_n \partial_x^n,$$

where  $m$  is an arbitrary integer and  $a_n \in \mathcal{A}_{f_0, f_1, \dots, f_{r-2}}^0$ . Let

$$A_+ := \sum_{n=0}^m a_n \partial_x^n, \quad \text{res } A := a_{-1}.$$

The product of pseudo-differential operators is defined by the following commutation rule:

$$\partial_x^k \circ a := \sum_{l=0}^{\infty} \frac{k(k-1)\dots(k-l+1)}{l!} (\partial_x^l a) \partial_x^{k-l}, \quad a \in \mathcal{A}_{f_0, f_1, \dots, f_{r-2}}^0, \quad k \in \mathbb{Z}.$$

For any  $m \geq 2$  and a pseudo-differential operator  $A$  of the form

$$A = \partial_x^m + \sum_{n=1}^{\infty} a_n \partial_x^{m-n}$$

there exists a unique pseudo-differential operator  $A^{\frac{1}{m}}$  of the form

$$A^{\frac{1}{m}} = \partial_x + \sum_{n=0}^{\infty} \tilde{a}_n \partial_x^{-n}$$

such that  $(A^{\frac{1}{m}})^m = A$ . Let

$$L := \partial_x^r + f_{r-2} \partial_x^{r-2} + \dots + f_1 \partial_x + f_0.$$

The  $r$ -th Gelfand–Dickey (GD) hierarchy is the following system of partial differential equations:

$$(4.1) \quad \frac{\partial L}{\partial T_a^\alpha} = \left[ (L^{a+\frac{\alpha}{r}})_+, L \right], \quad 1 \leq \alpha \leq r-1, \quad a \geq 0.$$

Let us describe a Hamiltonian structure for the GD hierarchy. Let  $X_0, X_1, \dots, X_{r-2} \in \mathcal{A}_{f_0, \dots, f_{r-2}}^0$  be some differential polynomials. Consider a pseudo-differential operator

$$X := \partial_x^{-(r-1)} \circ X_{r-2} + \dots + \partial_x^{-1} \circ X_0.$$

It is easy to see that the operator  $[X, L]_+$  has the form

$$[X, L]_+ = \sum_{0 \leq \alpha, \beta \leq r-2} \left( K_1^{\text{GD}; \alpha\beta} X_\beta \right) \partial_x^\alpha,$$

where

$$K_1^{\text{GD}; \alpha\beta} = \sum_{i \geq 0} K_{1;i}^{\text{GD}; \alpha\beta} \partial_x^i, \quad K_{1;i}^{\text{GD}; \alpha\beta} \in \mathcal{A}_{f_0, \dots, f_{r-2}}^0,$$

and the sum is finite. The operator  $K_1^{\text{GD}} = \left( K_1^{\text{GD}; \alpha\beta} \right)_{0 \leq \alpha, \beta \leq r-2}$  is Poisson and its degree zero part vanishes.

Consider the local functionals

$$\bar{h}_{\alpha, a}^{\text{GD}} := -\frac{r}{(a+1)r + \alpha} \int \text{res } L^{a+1+\frac{\alpha}{r}} dx, \quad 1 \leq \alpha \leq r-1, \quad a \geq -1.$$

We have

$$\left\{ \bar{h}_{\alpha, a}^{\text{GD}}, \bar{h}_{\beta, b}^{\text{GD}} \right\}_{K_1^{\text{GD}}} = 0, \quad 1 \leq \alpha, \beta \leq r-1, \quad a, b \geq 0,$$

and the local functionals  $\bar{h}_{\alpha, -1}^{\text{GD}}$ ,  $1 \leq \alpha \leq N$ , are linearly independent Casimirs of the Poisson bracket  $\{\cdot, \cdot\}_{K_1^{\text{GD}}}$ . For a local functional  $\bar{h} \in \Lambda_{f_0, f_1, \dots, f_{r-2}}^0$  define a pseudo-differential operator  $\frac{\delta \bar{h}}{\delta L}$  by

$$\frac{\delta \bar{h}}{\delta L} := \partial_x^{-(r-1)} \circ \frac{\delta \bar{h}}{\delta f_{r-2}} + \dots + \partial_x^{-1} \circ \frac{\delta \bar{h}}{\delta f_0}.$$

Then the right-hand side of (4.1) can be written in the following way:

$$\left[ \left( L^{a+\frac{\alpha}{r}} \right)_+, L \right] = \left[ \frac{\delta \bar{h}_{\alpha,a}^{\text{GD}}}{\delta L}, L \right]_+ = \sum_{0 \leq \beta, \gamma \leq r-2} \left( K_1^{\text{GD}; \beta \gamma} \frac{\delta \bar{h}_{\alpha,a}^{\text{GD}}}{\delta f_\gamma} \right) \partial_x^\beta.$$

Therefore, the sequence of local functionals  $\bar{h}_{\alpha,a}^{\text{GD}}$  together with the Poisson operator  $K_1^{\text{GD}}$  define a Hamiltonian structure of the GD hierarchy (4.1).

The GD hierarchy is endowed with a bihamiltonian structure, which can be described in the following way. Let

$$\tilde{X} := \partial_x^{-r} \circ X_{r-1} + \partial_x^{-(r-1)} \circ X_{r-2} + \dots + \partial_x^{-1} \circ X_0.$$

It is easy to see that the operator  $\left( L \circ \tilde{X} \right)_+ \circ L - L \circ \left( \tilde{X} \circ L \right)_+$  has the form

$$\left( L \circ \tilde{X} \right)_+ \circ L - L \circ \left( \tilde{X} \circ L \right)_+ = \sum_{0 \leq \alpha, \beta \leq r-1} \left( \tilde{K}^{\alpha \beta} X_\beta \right) \partial_x^\alpha,$$

where

$$\tilde{K}^{\alpha \beta} = \sum_{i \geq 0} \tilde{K}_i^{\alpha \beta} \partial_x^i, \quad \tilde{K}_i^{\alpha \beta} \in \mathcal{A}_{f_0, \dots, f_{r-2}}^0,$$

and the sum is finite. There is the following result:

$$\sum_{\beta=0}^{r-1} \tilde{K}^{r-1, \beta} X_\beta = \text{res}[\tilde{X}, L] = \partial_x \left( \sum_{\substack{0 \leq j \leq r-1 \\ 1 \leq \alpha \leq r-j}} \binom{-j-1}{\alpha} \partial_x^{\alpha-1} (f_{j+\alpha} X_j) \right),$$

where we adopt the conventions  $f_r := 1$  and  $f_{r-1} := 0$ . Let

$$f(X_0, X_1, \dots, X_{r-2}) := \frac{1}{r} \left( \sum_{\substack{0 \leq j \leq r-2 \\ 1 \leq \alpha \leq r-j}} \binom{-j-1}{\alpha} \partial_x^{\alpha-1} (f_{j+\alpha} X_j) \right).$$

Define an operator  $K_2^{\text{GD}} = \left( K_2^{\text{GD}; \alpha \beta} \right)_{0 \leq \alpha, \beta \leq r-2}$  by the equation

$$\sum_{0 \leq \alpha, \beta \leq r-2} \left( \tilde{K}^{\alpha \beta} X_\beta \right) \partial_x^\alpha + \sum_{\alpha=0}^{r-2} \left( \tilde{K}^{\alpha, r-1} f(X_0, \dots, X_{r-2}) \right) \partial_x^\alpha = \sum_{0 \leq \alpha, \beta \leq r-2} \left( K_2^{\text{GD}; \alpha \beta} X_\beta \right) \partial_x^\alpha.$$

The operator  $K_2^{\text{GD}}$  is Poisson and compatible with the operator  $K_1^{\text{GD}}$ . It is easy to see that the degree zero part of the operator  $K_2^{\text{GD}}$  vanishes.

Consider the local functionals

$$\bar{g}_{\alpha,a}^{\text{GD}} := \frac{r}{ar + \alpha} \int \text{res} L^{a+\frac{\alpha}{r}} dx = -\bar{h}_{\alpha,a+1}^{\text{GD}}, \quad 1 \leq \alpha \leq r-1, \quad a \geq 0.$$

The right-hand side of (4.1) can be written in the following way using the local functionals  $\bar{g}_{\alpha,a}^{\text{GD}}$  and the Poisson operator  $K_2^{\text{GD}}$ :

$$\left[ \left( L^{a+\frac{\alpha}{r}} \right)_+, L \right] = \sum_{0 \leq \beta, \gamma \leq r-2} \left( K_2^{\text{GD}; \beta \gamma} \frac{\delta \bar{g}_{\alpha,a}^{\text{GD}}}{\delta f_\gamma} \right) \partial_x^\beta,$$

or, equivalently,

$$\left\{ \cdot, \bar{h}_{\alpha,a}^{\text{GD}} \right\}_{K_2^{\text{GD}}} = - \left\{ \cdot, \bar{h}_{\alpha,a+1}^{\text{GD}} \right\}_{K_1^{\text{GD}}}, \quad 1 \leq \alpha \leq r-1, \quad a \geq -1,$$

which endows the GD hierarchy with a bihamiltonian structure.

4.2.2. *Relation with the DR hierarchy.* As it is mentioned in Section 3.4, there is another bihamiltonian hierarchy associated, in particular, to the  $r$ -spin CohFT, the Dubrovin–Zhang (DZ) hierarchy. In this section, we recall the relation between the  $r$ -th GD hierarchy and the DZ hierarchy for the  $r$ -spin CohFT, and then present a relation between the DR hierarchy and the DZ hierarchy for  $r = 3, 4$ , and  $5$ .

Introduce new variables  $w^1, \dots, w^{r-1}$  and consider the following Miura transformation:

$$f_i \mapsto w^\alpha(f_{*,*}) = \frac{1}{(r-\alpha)(-r)^{\frac{r-\alpha-1}{2}}} \operatorname{res} L^{(r-\alpha)/r}.$$

Define Poisson operators  $K_1^{r\text{-spin}} = \left( K_1^{r\text{-spin};\alpha\beta} \right)_{1 \leq \alpha, \beta \leq r-1}$ ,  $K_2^{r\text{-spin}} = \left( K_2^{r\text{-spin};\alpha\beta} \right)_{1 \leq \alpha, \beta \leq r-1}$ , and local functionals  $\overline{h}_{\alpha,d}^{r\text{-spin}} \in \Lambda_{w^1, \dots, w^{r-1}}$ ,  $1 \leq \alpha \leq r-1$ ,  $d \geq -1$ , by

$$\begin{aligned} K_1^{r\text{-spin}} &:= (-r)^{\frac{r}{2}} K_{1;w}^{\text{GD}}, \\ K_2^{r\text{-spin}} &:= K_{2;w}^{\text{GD}}, \\ \overline{h}_{\alpha,d}^{r\text{-spin}} &:= \frac{1}{(-r)^{\frac{\alpha-1+r(d+1)}{2}-d} (\alpha+rd)!_r} \overline{h}_{\alpha,d}^{\text{GD}}[w], \end{aligned}$$

where

$$(\alpha+rd)!_r := \begin{cases} \prod_{i=0}^d (\alpha+ri), & \text{if } d \geq 0, \\ 1, & \text{if } d = -1. \end{cases}$$

Recall that  $K_{1;w}^{\text{GD}}$  denotes the Miura transform of the operator  $K_1^{\text{GD}}$ . The local functionals  $\overline{h}_{\alpha,d}^{r\text{-spin}}$  together with the Poisson operator  $K_1^{r\text{-spin}}$  define the Hamiltonian hierarchy

$$\frac{\partial w^\alpha}{\partial t_b^\beta} = K_1^{r\text{-spin};\alpha\mu} \frac{\delta \overline{h}_{\beta,b}^{r\text{-spin}}}{\delta w^\mu}, \quad 1 \leq \alpha, \beta \leq N, \quad b \geq 0,$$

which is bihamiltonian with the bihamiltonian recursion given by

$$(4.2) \quad \left\{ \cdot, \overline{h}_{\alpha,d}^{r\text{-spin}} \right\}_{K_2^{r\text{-spin}}} = \frac{\alpha + (d+1)r}{r} \left\{ \cdot, \overline{h}_{\alpha,d+1}^{r\text{-spin}} \right\}_{K_1^{r\text{-spin}}}, \quad 1 \leq \alpha \leq r-1, \quad d \geq -1.$$

This bihamiltonian hierarchy is the Dubrovin–Zhang hierarchy corresponding to the  $r$ -spin CohFT [DZ01, FSZ10].

In [BG16] it is proved that for  $r = 3, 4$ , and  $5$  the local functionals  $\overline{g}_{\alpha,d}$  and the Poisson operator  $K_1 = \eta^{-1} \partial_x$  of the DR hierarchy are related to the local functionals  $\overline{h}_{\alpha,d}^{r\text{-spin}}$  and the Poisson operator  $K_1^{r\text{-spin}}$  by the following Miura transformations:

$$(4.3) \quad \begin{cases} w^1 = u^1, \\ w^2 = u^2, \end{cases} \quad \text{for } r = 3;$$

$$(4.4) \quad \begin{cases} w^1 = u^1 + \frac{\varepsilon^2}{96} u_{xx}^3, \\ w^2 = u^2, \\ w^3 = u^3, \end{cases} \quad \text{for } r = 4;$$

$$(4.5) \quad \begin{cases} w^1 = u^1 + \frac{\varepsilon^2}{60} u_{xx}^3, \\ w^2 = u^2 + \frac{\varepsilon^2}{60} u_{xx}^4, \\ w^3 = u^3, \\ w^4 = u^4, \end{cases} \quad \text{for } r = 5.$$

4.2.3. *Proof of Conjecture 1.13 for  $r = 3, 4, 5$ .* For the first part of the conjecture it is sufficient to check that the operators  $K_2$  and  $K_2^{r\text{-spin}}$  are related by the Miura transformations given by equations (4.3), (4.4), and (4.5).

For  $r = 3$  we have the following formula for  $K_2^{3\text{-spin}}$ :

$$\left( \frac{2}{9}(w^2)^2\partial_x + \frac{2}{9}w^2w_x^2 + \varepsilon^2 \left( \frac{5}{54}w^2\partial_x^3 + \frac{5}{36}w_x^2\partial_x^2 + \frac{1}{12}w_{xx}^2\partial_x + \frac{1}{54}w_{xxx}^2 \right) + \frac{\varepsilon^4}{162}\partial_x^5 \quad w^1\partial_x + \frac{1}{3}w_x^1 \right. \\ \left. \frac{2}{3}w^2\partial_x + \frac{1}{3}w_x^2 + \frac{2}{9}\varepsilon^2\partial_x^3 \right).$$

On the other hand, from [BG16, Proposition 3.7]) and (1.8) we obtain

$$\bar{g} = \int \left( \frac{(u^1)^2u^2}{2} + \frac{(u^2)^4}{72} + \varepsilon^2 \left( \frac{(u^2)^2u_2^2}{144} + \frac{u^1u_2^1}{24} \right) + \frac{\varepsilon^4}{1728}u^2u_4^2 \right) dx.$$

Computing the operator  $K_2$  using this expression (and identifying  $u^\alpha = w^\alpha$ ) we get exactly the operator  $K_2^{3\text{-spin}}$ .

For  $r = 4$  the operator  $K_2^{4\text{-spin}}$  is given by

$$K_2^{4\text{-spin};1,1} = \left( \frac{(w^3)^3}{32} + \frac{3(w^2)^2}{16} \right) \partial_x + \frac{3w^2w_x^2}{16} + \frac{3(w^3)^2w_x^3}{64} \\ + \varepsilon^2 \left( \left( \frac{7(w^3)^2}{256} + \frac{w^1}{48} \right) \partial_x^3 + \left( \frac{w_x^1}{32} + \frac{21w^3w_x^3}{256} \right) \partial_x^2 + \left( \frac{5(w_x^3)^2}{128} + \frac{13w^3w_{xx}^3}{256} + \frac{w_{xx}^1}{24} \right) \partial_x \right. \\ \left. + \frac{3w_x^3w_{xx}^3}{128} + \frac{w_{xxx}^1}{64} + \frac{3w^3w_{xxx}^3}{256} \right) \\ + \varepsilon^4 \left( \frac{7w^3}{1152}\partial_x^5 + \frac{35w_x^3}{2304}\partial_x^4 + \frac{91w_{xx}^3}{4608}\partial_x^3 + \frac{133w_{xxx}^3}{9216}\partial_x^2 + \frac{47w_4^3}{9216}\partial_x + \frac{w_5^3}{1536} \right) \\ + \varepsilon^6 \frac{17}{36864}\partial_x^7,$$

$$K_2^{4\text{-spin};1,2} = - \left( K_2^{4\text{-spin};2,1} \right)^* \\ = \frac{5w^2w^3}{16}\partial_x + \frac{w^3w_x^2}{8} + \frac{w^2w_x^3}{8} + \varepsilon^2 \left( \frac{7w^2}{64}\partial_x^3 + \frac{7w_x^2}{48}\partial_x^2 + \frac{17w_{xx}^2}{192}\partial_x + \frac{w_{xxx}^2}{48} \right),$$

$$K_2^{4\text{-spin};1,3} = - \left( K_2^{4\text{-spin};3,1} \right)^* = w^1\partial_x + \frac{w_x^1}{4} + \varepsilon^2 \left( \frac{7w^3}{192}\partial_x^3 + \frac{7w_x^3}{192}\partial_x^2 \right) + \varepsilon^4 \frac{7}{768}\partial_x^5,$$

$$K_2^{4\text{-spin};2,2} = \left( \frac{(w^3)^2}{8} + w^1 \right) \partial_x + \frac{w_x^1}{2} + \frac{w^3w_x^3}{8} + \varepsilon^2 \left( \frac{w^3}{8}\partial_x^3 + \frac{3w_x^3}{16}\partial_x^2 + \frac{w_{xx}^3}{12}\partial_x + \frac{w_{xxx}^3}{96} \right) + \frac{\varepsilon^4}{64}\partial_x^5,$$

$$K_2^{4\text{-spin};2,3} = - \left( K_2^{4\text{-spin};3,2} \right)^* = \frac{3w^2}{4}\partial_x + \frac{w_x^2}{4},$$

$$K_2^{4\text{-spin};3,3} = \frac{w^3}{2}\partial_x + \frac{w_x^3}{4} + \varepsilon^2 \frac{5}{16}\partial_x^3.$$

On the other hand, from [BG16, Proposition 3.8]) and (1.8) we obtain

$$\bar{g} = \int \left[ \frac{(u^1)^2u^3}{2} + \frac{u^1(u^2)^2}{2} + \frac{(u^2)^2(u^3)^2}{16} + \frac{(u^3)^5}{960} + \right. \\ \left. + \varepsilon^2 \left( \frac{u^1u_2^1}{16} + \frac{u_2^3(u^2)^2}{192} + \frac{u^3u^2u_2^2}{48} + \frac{u_2^1(u^3)^2}{192} + \frac{(u^3)^3u_2^3}{768} \right) + \right. \\ \left. + \varepsilon^4 \left( \frac{u^2u_4^2}{640} + \frac{(u^3)^2u_4^3}{4096} + \frac{3u^1u_4^3}{2560} \right) + \varepsilon^6 \frac{u^3u_6^3}{49152} \right] dx.$$

Computing the operator  $K_2$  using this expression and performing the Miura transformation (4.4) we obtain the operator  $K_2^{4\text{-spin}}$ .

For  $r = 5$  the first part of Conjecture 1.13 is proved using the same scheme: we compute the operator  $K_2^{5\text{-spin}}$  explicitly, then we compute the operator  $K_2$  using the expression for the local functional  $\bar{g}_{1,1}$  from [BG16, Proposition 3.10] and the first equation in (1.8), and finally we check that the resulting two operators are related by the Miura transformation (4.5). This computation is performed in **Mathematica** and is not presented here, since the computations and even the expressions for the Poisson operators get too involved.

After we have proved the first part of the conjecture, the second part becomes easy in all three cases. Indeed, since  $\mu_\alpha = \frac{2\alpha-r}{2r}$ , one can easily check that Equations (1.10) and (4.2) agree.

**4.3. The Gromov–Witten theory of  $\mathbb{CP}^1$ .** Consider the CohFT controlling the Gromov–Witten theory of  $\mathbb{CP}^1$  (see e.g. [BR16a, Section 6]). For simplicity, we set the degree parameter (typically denoted by  $q$ ) equal to 1. This CohFT has rank 2 and is homogeneous with

$$(q_1, q_2) = (0, 1), \quad (r^1, r^2) = (0, 2), \quad \delta = 1, \quad \eta = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \mu = \begin{pmatrix} -\frac{1}{2} & 0 \\ 0 & \frac{1}{2} \end{pmatrix}, \quad A = \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}.$$

The corresponding DR hierarchy is studied in [BR16a]. In particular, it is related to the extended Toda hierarchy (see e.g. [CDZ04]).

**Notation 4.2.** In computations we use the following two formal power series:

$$S(z) := \frac{e^{z/2} - e^{-z/2}}{z} \quad \text{and} \quad \tilde{S}(z) := \frac{e^{z/2} + e^{-z/2}}{2}.$$

Let us compute the operator  $K_2$  for the DR hierarchy.

**Lemma 4.3.** *We have*

$$(4.6) \quad \bar{g} = \int \left( \frac{(u^1)^2 u^2}{2} + \sum_{g \geq 1} \varepsilon^{2g} \frac{B_{2g}}{(2g)(2g)!} u^1 u_{2g}^1 + e^{S(\varepsilon \partial_x) u^2} - u^2 - \frac{(u^2)^2}{2} \right) dx,$$

where  $B_{2g}$  are the Bernoulli numbers.

*Proof.* By [BR16b] we have

$$\bar{g}_{1,1} = \int \left( \frac{(u^1)^2 u^2}{2} + \sum_{g \geq 1} \varepsilon^{2g} \frac{B_{2g}}{(2g)!} u^1 u_{2g}^1 + \left( \tilde{S}(\varepsilon \partial_x) u^2 - 2 \right) e^{S(\varepsilon \partial_x) u^2} + u^2 \right) dx.$$

Denote by  $f$  the expression under the integral sign on the right-hand side of (4.6). Note that  $f|_{\varepsilon=0} \in \mathbb{C}[[u^1, u^2]]$  starts from cubic terms in the variables  $u^1, u^2$ . Therefore, it is sufficient to check that

$$(D - 2)f = \frac{(u^1)^2 u^2}{2} + \sum_{g \geq 1} \varepsilon^{2g} \frac{B_{2g}}{(2g)!} u^1 u_{2g}^1 + \left( \tilde{S}(\varepsilon \partial_x) u^2 - 2 \right) e^{S(\varepsilon \partial_x) u^2} + u^2,$$

where we recall that  $D = \sum_{n \geq 0} (n+1) u_n^\alpha \frac{\partial}{\partial u_n^\alpha}$ . Since

$$\begin{aligned} (D - 2) \frac{(u^1)^2 u^2}{2} &= \frac{(u^1)^2 u^2}{2}, & (D - 2) u^1 u_{2g}^1 &= 2g u^1 u_{2g}^1, \\ (D - 2) \frac{(u^2)^2}{2} &= 0, & (D - 2) u^2 &= -u^2, \end{aligned}$$

it is sufficient to check that

$$D e^{S(\varepsilon \partial_x) u^2} = \tilde{S}(\varepsilon \partial_x) u^2 \cdot e^{S(\varepsilon \partial_x) u^2},$$

which follows from

$$D e^{S(\varepsilon \partial_x) u^2} = (\varepsilon \partial_x \circ S'(\varepsilon \partial_x) + S(\varepsilon \partial_x)) u^2 \cdot e^{S(\varepsilon \partial_x) u^2} = \tilde{S}(\varepsilon \partial_x) u^2 \cdot e^{S(\varepsilon \partial_x) u^2}.$$

□



We further compute

$$\frac{\delta \bar{g}}{\delta u^1} = u^1 u^2 + \sum_{g \geq 1} \varepsilon^{2g} \frac{B_{2g}}{g(2g)!} u_{2g}^1, \quad \frac{\delta \bar{g}}{\delta u^2} = \frac{(u^1)^2}{2} + S(\varepsilon \partial_x) e^{S(\varepsilon \partial_x) u^2} - 1 - u^2.$$

Therefore,

$$\begin{aligned} \widehat{\Omega}(\bar{g}) &= \begin{pmatrix} S(\varepsilon \partial_x) \circ e^{S(\varepsilon \partial_x) u^2} S(\varepsilon \partial_x) - 1 & u^1 \\ u^1 & u^2 + \sum_{g \geq 1} \frac{B_{2g}}{g(2g)!} (\varepsilon \partial_x)^{2g} \end{pmatrix}, \\ \widehat{\Omega}^1(\bar{g}) &= \begin{pmatrix} \varepsilon \left( S(\varepsilon \partial_x) \circ e^{S(\varepsilon \partial_x) u^2} S'(\varepsilon \partial_x) + S'(\varepsilon \partial_x) \circ e^{S(\varepsilon \partial_x) u^2} S(\varepsilon \partial_x) \right) & 0 \\ 0 & \sum_{g \geq 1} \varepsilon^{2g} \frac{2B_{2g}}{(2g)!} \partial_x^{2g-1} \end{pmatrix}. \end{aligned}$$

Finally, we obtain

$$\begin{aligned} K_2 &= \begin{pmatrix} S(\varepsilon \partial_x) \partial_x \circ e^{S(\varepsilon \partial_x) u^2} \widetilde{S}(\varepsilon \partial_x) + \widetilde{S}(\varepsilon \partial_x) \circ e^{S(\varepsilon \partial_x) u^2} S(\varepsilon \partial_x) \partial_x & u^1 \partial_x \\ \partial_x \circ u^1 & \sum_{g \geq 0} \varepsilon^{2g} \frac{2B_{2g}}{(2g)!} \partial_x^{2g+1} \end{pmatrix} \\ &= \begin{pmatrix} \varepsilon^{-1} \left( e^{\frac{\varepsilon \partial_x - 1}{\varepsilon \partial_x} u^2} e^{\varepsilon \partial_x} - e^{\frac{1 - e^{-\varepsilon \partial_x}}{\varepsilon \partial_x} u^2} e^{-\varepsilon \partial_x} \right) & u^1 \partial_x \\ \partial_x \circ u^1 & \frac{e^{\varepsilon \partial_x} + 1}{e^{\varepsilon \partial_x} - 1} \varepsilon \partial_x^2 \end{pmatrix}. \end{aligned}$$

On the other hand, consider the extended Toda hierarchy from [CDZ04]. The dependent variables of the extended Toda hierarchy are denoted by  $v$  and  $u$  in [CDZ04]. Following [BR16a] we denote  $v^1 = v$  and  $v^2 = u$ . The extended Toda hierarchy is bihamiltonian with the pair of compatible Poisson operators  $(K_1^{\text{Td}}, K_2^{\text{Td}})$  given by

$$\begin{aligned} K_1^{\text{Td}} &= \begin{pmatrix} 0 & \varepsilon^{-1} (e^{\varepsilon \partial_x} - 1) \\ \varepsilon^{-1} (1 - e^{-\varepsilon \partial_x}) & 0 \end{pmatrix}, \\ K_2^{\text{Td}} &= \begin{pmatrix} \varepsilon^{-1} (e^{\varepsilon \partial_x} \circ e^{v^2} - e^{v^2} e^{-\varepsilon \partial_x}) & \varepsilon^{-1} v^1 (e^{\varepsilon \partial_x} - 1) \\ \varepsilon^{-1} (1 - e^{-\varepsilon \partial_x}) \circ v^1 & \varepsilon^{-1} (e^{\varepsilon \partial_x} - e^{-\varepsilon \partial_x}) \end{pmatrix}. \end{aligned}$$

It is straightforward to check that the Miura transformation

$$u^1 = e^{-\frac{\varepsilon \partial_x}{2}} v^1, \quad u^2 = \frac{\varepsilon \partial_x}{e^{\frac{\varepsilon \partial_x}{2}} - e^{-\frac{\varepsilon \partial_x}{2}}} v^2$$

takes the pair  $(K_1^{\text{Td}}, K_2^{\text{Td}})$  to the pair  $(K_1, K_2)$ . Thus, Part 1 of Conjecture 1.13 holds for the CohFT of the Gromov–Witten theory of  $\mathbb{C}P^1$ .

Let us check Part 2 of Conjecture 1.13. Let  $\bar{h}_{\alpha,d}^{\text{Td}}$  denote the Hamiltonians of the extended Toda hierarchy. In [BR16a] it is proved that

$$(4.7) \quad \bar{g}_{\alpha,d} = \sum_{i=0}^{d+1} (-1)^i (S_i^*)_{\alpha}^{\mu} \bar{h}_{\mu,d-i}^{\text{Td}}[u], \quad d \geq -1,$$

where  $S_i^*$ ,  $i \geq 0$ , are certain matrices described in [BR16a, Section 6]. The bihamiltonian recursion for the extended Toda hierarchy is given by (see [CDZ04, Theorem 3.1])

$$\left\{ \cdot, \bar{h}_{\alpha,d}^{\text{Td}} \right\}_{K_2^{\text{Td}}} = \left( d + \frac{3}{2} + \mu_{\alpha} \right) \left\{ \cdot, \bar{h}_{\alpha,d+1}^{\text{Td}} \right\}_{K_1^{\text{Td}}} + R_{\alpha}^{\beta} \left\{ \cdot, \bar{h}_{\beta,d}^{\text{Td}} \right\}_{K_1^{\text{Td}}}, \quad d \geq -1,$$

where

$$R = (R_{\alpha}^{\beta}) := \begin{pmatrix} 0 & 0 \\ 2 & 0 \end{pmatrix}.$$

Using (4.7) and the property

$$A_{\alpha}^{\gamma} (S_i^*)_{\gamma}^{\beta} = (S_i^*)_{\alpha}^{\gamma} R_{\gamma}^{\beta} + (S_{i+1}^*)_{\alpha}^{\beta} (i + 1 + \mu_{\alpha} - \mu_{\beta}), \quad i \geq 0,$$

which is a standard fact in Gromov–Witten theory (or can be checked using the explicit formula for the matrices  $S_i^*$  given in [BR16a, Section 6]), we obtain (1.10) in this case.

## REFERENCES

- [Bur15] A. Buryak. *Double ramification cycles and integrable hierarchies*. Communications in Mathematical Physics 336 (2015), no. 3, 1085–1107.
- [BDGR18] A. Buryak, B. Dubrovin, J. Guéré, P. Rossi. *Tau-structure for the double ramification hierarchies*. Communications in Mathematical Physics 363 (2018), no. 1, 191–260.
- [BDGR19] A. Buryak, B. Dubrovin, J. Guéré, P. Rossi. *Integrable systems of double ramification type*. International Mathematics Research Notices, rnz029, <https://doi.org/10.1093/imrn/rnz029>.
- [BG16] A. Buryak, J. Guere. *Towards a description of the double ramification hierarchy for Witten’s  $r$ -spin class*. Journal de Mathématiques Pures et Appliquées 106 (2016), no. 5, 837–865.
- [BGR19] A. Buryak, J. Guéré, P. Rossi. *DR/DZ equivalence conjecture and tautological relations*. Geometry & Topology 23 (2019), no. 7, 3537–3600.
- [BPS12a] A. Buryak, H. Posthuma, S. Shadrin. *On deformations of quasi-Miura transformations and the Dubrovin–Zhang bracket*. Journal of Geometry and Physics 62 (2012), no. 7, 1639–1651.
- [BPS12b] A. Buryak, H. Posthuma, S. Shadrin. *A polynomial bracket for the Dubrovin–Zhang hierarchies*. Journal of Differential Geometry 92 (2012), no. 1, 153–185.
- [BR16a] A. Buryak, P. Rossi. *Recursion relations for double ramification hierarchies*. Communications in Mathematical Physics 342 (2016), no. 2, 533–568.
- [BR16b] A. Buryak, P. Rossi. *Double ramification cycles and quantum integrable systems*. Letters in Mathematical Physics 106 (2016), no. 3, 289–317.
- [BSSZ15] A. Buryak, S. Shadrin, L. Spitz, D. Zvonkine. *Integrals of  $\psi$ -classes over double ramification cycles*. American Journal of Mathematics 137 (2015), no. 3, 699–737.
- [CKS18] G. Carlet, R. Kramer, S. Shadrin. *Central invariants revisited*. Journal de l’École polytechnique. Mathématiques 5 (2018), 149–175.
- [CPS18] G. Carlet, H. Posthuma, S. Shadrin. *Deformations of semisimple Poisson pencils of hydrodynamic type are unobstructed*. Journal of Differential Geometry 108 (2018), no. 1, 63–89.
- [Dic03] L. A. Dickey. *Soliton equations and Hamiltonian systems*. Second edition. World Scientific, 2003.
- [DR19] A. du Crest de Villeneuve, P. Rossi. *Quantum  $D_4$  Drinfeld–Sokolov hierarchy and quantum singularity theory*. Journal of Geometry and Physics 141 (2019), 29–44.
- [DLZ06] B. Dubrovin, S.-Q. Liu, Y. Zhang. *On Hamiltonian perturbations of hyperbolic systems of conservation laws I: Quasi-triviality of bihamiltonian perturbations*. Communications on Pure and Applied Mathematics 59 (2006), no. 4, 559–615.
- [DZ98] B. Dubrovin, Y. Zhang. *Bihamiltonian hierarchies in 2D topological field theory at one-loop approximation*. Communications in Mathematical Physics 198 (1998), no. 2, 311–361.
- [DZ99] B. Dubrovin, Y. Zhang. *Frobenius manifolds and Virasoro constraints*. Selecta Mathematica. New Series 5 (1999), no. 4, 423–466.
- [DZ01] B. Dubrovin, Y. Zhang. *Normal forms of hierarchies of integrable PDEs, Frobenius manifolds and Gromov–Witten invariants*. arXiv:math/0108160.
- [CDZ04] G. Carlet, B. Dubrovin, Y. Zhang. *The extended Toda hierarchy*. Moscow Mathematical Journal 4 (2004), no. 2, 313–332.
- [FP00] C. Faber, R. Pandharipande. *Logarithmic series and Hodge integrals in the tautological ring. With an appendix by Don Zagier*. Dedicated to William Fulton on the occasion of his 60th birthday. Michigan Mathematical Journal 48 (2000), 215–252.
- [FSZ10] C. Faber, S. Shadrin, D. Zvonkine. *Tautological relations and the  $r$ -spin Witten conjecture*. Annales Scientifiques de l’École Normale Supérieure (4) 43 (2010), no. 4, 621–658.
- [Hai13] R. Hain. *Normal functions and the geometry of moduli spaces of curves*. In *Handbook of moduli*, Vol. I, 527–578. International Press, 2013.
- [JPPZ17] F. Janda, R. Pandharipande, A. Pixton, D. Zvonkine. *Double ramification cycles on the moduli spaces of curves*. Publications Mathématiques. Institut de Hautes Études Scientifiques 125 (2017), 221–266.
- [Kon92] M. Kontsevich. *Intersection theory on the moduli space of curves and the matrix Airy function*. Communications in Mathematical Physics 147 (1992), no. 1, 1–23.
- [KM94] M. Kontsevich, Yu. Manin. *Gromov–Witten classes, quantum cohomology, and enumerative geometry*. Communications in Mathematical Physics 164 (1994), no. 3, 525–562.
- [KMGZ70] M. Kruskal, R. Miura, C. Gardner, N. Zabusky. *Korteweg–de Vries equation and generalizations. V. Uniqueness and nonexistence of polynomial conservation laws*. Journal of Mathematical Physics 11 (1970), no. 3, 952–960.

- [Liu18] S.-Q. Liu. *Lecture notes on bihamiltonian structures and their central invariants*. In *B-model Gromov–Witten theory*, 573–625. Birkhäuser, Cham, 2018.
- [LZ05] S.-Q. Liu, Y. Zhang. *Deformations of semisimple bihamiltonian structures of hydrodynamic type*. *Journal of Geometry and Physics* 54 (2005), no. 4, 427–453.
- [LZ11] S.-Q. Liu, Y. Zhang. *Jacobi structures of evolutionary partial differential equations*. *Advances in Mathematics* 227 (2011), no. 1, 73–130.
- [LZ13] S.-Q. Liu, Y. Zhang. *Bihamiltonian cohomologies and integrable hierarchies I: A special case*. *Communications in Mathematical Physics* 324 (2013), no. 3, 897–935.
- [MW13] S. Marcus, J. Wise. *Stable maps to rational curves and the relative Jacobian*. arXiv:1310.5981.
- [PPZ15] R. Pandharipande, A. Pixton, D. Zvonkine. *Relations on  $\overline{\mathcal{M}}_{g,n}$  via 3-spin structures*. *Journal of the American Mathematical Society* 28 (2015), no. 1, 279–309.
- [Ros17] P. Rossi. *Integrability, quantization and moduli spaces of curves*. *SIGMA. Symmetry, Integrability and Geometry: Methods and Applications* 13 (2017), paper no. 060, 29 pp.
- [Wit91] E. Witten. *Two-dimensional gravity and intersection theory on moduli space*. *Surveys in Differential Geometry* 1 (1991), 243–310.

A. BURYAK:

FACULTY OF MATHEMATICS, NATIONAL RESEARCH UNIVERSITY HIGHER SCHOOL OF ECONOMICS,  
6 USACHEVA STR., 119048 MOSCOW, RUSSIAN FEDERATION; AND  
FACULTY OF MECHANICS AND MATHEMATICS, LOMONOSOV MOSCOW STATE UNIVERSITY,  
GSP-1, 119991 MOSCOW, RUSSIAN FEDERATION

*E-mail address:* aburyak@hse.ru

P. ROSSI:

DIPARTIMENTO DI MATEMATICA “TULLIO LEVI-CIVITA”, UNIVERSITÀ DEGLI STUDI DI PADOVA,  
VIA TRIESTE 63, 35121 PADOVA, ITALY

*E-mail address:* paolo.rossi@math.unipd.it

S. SHADRIN:

KORTEWEG–DE VRIES INSTITUUT VOOR WISKUNDE, UNIVERSITEIT VAN AMSTERDAM,  
POSTBUS 94248, 1090GE AMSTERDAM, THE NETHERLANDS

*E-mail address:* s.shadrin@uva.nl