

# Journal Pre-proof

Travellers' Risk Behaviors and Health Problems: Post-Travel Follow Up in Two Travel Medicine Centers in Italy

Armando Stefanati Dr Alice Pierobon Valeria Baccello Emanuela DeStefani Diego Gamberoni Patrizia Furlan Federica Sandri Antonio Stano Paolo Coin Vincenzo Baldo Giovanni Gabutti



PII: S0399-077X(20)30736-8

DOI: <https://doi.org/doi:10.1016/j.medmal.2020.10.009>

Reference: MEDMAL 4366

To appear in: *Médecine et Maladies Infectieuses*

Received Date: 12 February 2020

Revised Date: 15 April 2020

Accepted Date: 8 October 2020

Please cite this article as: Stefanati A, Pierobon A, Baccello V, DeStefani E, Gamberoni D, Furlan P, Sandri F, Stano A, Coin P, Baldo V, Gabutti G, Travellers' Risk Behaviors and Health Problems: Post-Travel Follow Up in Two Travel Medicine Centers in Italy, *Médecine et Maladies Infectieuses* (2020), doi: <https://doi.org/10.1016/j.medmal.2020.10.009>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier.

**TITLE PAGE****Travellers' Risk Behaviors and Health Problems: Post-Travel Follow Up in Two Travel Medicine Centers in Italy.**

Armando Stefanati<sup>1</sup>, Alice Pierobon<sup>2</sup>, Valeria Baccello<sup>1</sup>, Emanuela DeStefani<sup>2</sup>, Diego Gamberoni<sup>1</sup>, Patrizia Furlan<sup>3</sup>, Federica Sandri<sup>1</sup>, Antonio Stano<sup>2</sup>, Paolo Coin<sup>2</sup>, Vincenzo Baldo<sup>3</sup>, Giovanni Gabutti<sup>1</sup>.

<sup>1</sup> *University of Ferrara, Department of Medical Sciences, Ferrara, Italy*

<sup>2</sup> *Local Health Unit 7 Pedemontana, Veneto Regione, Department of Prevention, Bassano del Grappa, Italy*

<sup>3</sup> *University of Padua, Department of Cardiac, Thoracic and Vascular Sciences, Padua, Italy.*

Corresponding author: Dr Armando Stefanati

University of Ferrara - Department of Medical Sciences

Via Fossato di Mortara, 64/B - 44121 – Ferrara (FE) ITALY

Tel: +39 3389940576

E-mail: sta@unife.it

Dr A. Stefanati will handle correspondence concerning this article, press proofs, and orders for off-prints

The study described in the manuscript was previously reported as abstract at 51 SITI National Congress of Riva del Garda 17-20 October 2018 (Valutazione dello stato di salute nei soggetti al rientro da un viaggio internazionale nell'azienda USL di Ferrara. Assessment of the state of health in subjects returning from an international trip to the Ferrara Local Health Authority. G. Gabutti, L. Bertoni, V. Baccello, F. Brosio, A. Formaglio, A. Stefanati).

This research did not receive any grant from funding agencies in the public, commercial, or not for-profit sectors.

**Highlights**

- International travel can expose one to health risks and represents an important Public Health Issue.
- The compliance to prophylactic measures and behavioral advice given during pre-travel consultation are fundamental preventive measures.
- Pre-travel consultation is the only way to make travellers aware of specific risks that could occur during their trip and recommend specific prophylaxis, including immunization.

## ABSTRACT

**Objectives:** We examined the association between travellers' characteristics, compliance with pre-travel recommendations and health problems.

**Methods:** Volunteer travellers were enrolled and data collected using a questionnaire between 30-60 days after returning home. We analyzed the associations through bivariate and multivariate models.

**Results:** Of the 468 enrolled travelers, 68% consumed raw food and 81% food containing milk and/or eggs. 32% consumed street vendor food and 30% drinks containing ice. 24% used the recommended mechanical prophylaxis measures. 46% got sick during and/or after travel (gastrointestinal symptoms most frequently). Factors predisposing to health problems were female gender, youth/middle age, intermediate travel duration and profession. The American continent and staying in hostels and tents were significantly associated with febrile illness. Street vendor food was significantly associated with skin reactions.

**Conclusions:** Adherence to behavioral recommendations remains low. Travellers must be informed of health risks during and after travel.

**Keywords:** dangerous behavior, travel-associated infectious diseases, travel medicine, travel-related health problems, travel-related illness.

## INTRODUCTION

### Background

Between 1950 and 2017 the world population increased 2.9 fold while the number of travellers increased 53 fold.[1,2] In 2017, the largest number of international travellers were from Europe (51%), followed by Asia and the Pacific (25%), the Americas (16%), Africa (5%) and the Middle East (4%) [2]. Travel can expose one to health risks due to infectious diseases, changes in altitude, climate, humidity and temperature as well as accidents. The risk of health

problems during international travel depends on different factors such as the destination, the age and health of the traveller, the duration of the trip and planned activities [3]. Traveller's diarrhea is the most frequent illness reported [4]. According to the GeoSentinel, in the five-year period between 2007 and 2011 most travellers acquired the disorder in Asia (32.6%), followed by sub-Saharan Africa (26.7%) and Latin America and the Caribbean (19.2%). Among the main reasons for travel, 55.7% of subjects reporting symptoms traveled for tourism, 13.6% for work and 15.5% were Visiting Friends and Relatives (VFR). The most frequently reported diseases were gastrointestinal infections (34%), febrile illness (23.3%) and dermatological diseases (19.5%). Only 40.5% of travellers reporting symptoms pointed to a pre-travel consultation by a health professional [5]. Pre-travel consultation is an important tool to make travellers aware of health risks they may incur and the ways to avoid illness. The consultation should preferably take place about six weeks before travel in time to receive the recommended vaccinations and chemoprophylaxis indications [6]. There are still a limited number of studies in which traveller behavior and health problems during and after travel are assessed. Similarly, there are few studies analyzing which risk factors are most associated with health disorders.

## **Objective**

The objective of this study was to follow-up international travellers attending pre-travel health consultation services in two Italian centers in Northern Italy. We evaluated compliance to prophylactic measures and behavioral advice given during pre-travel consultation, to describe the types and frequencies of health problems occurring during and after travel. Finally, we assessed possible associations between risk factors and travel-related health problems.

## **MATERIAL AND METHODS**

### **Study design**

The study is a descriptive observational study conducted in two centers in Northern Italy: AULSS 7 Pedemontana in Veneto Region and AUSL of Ferrara in Emilia-Romagna Region. International travellers aged  $\geq 14$  years were enrolled in travel clinics in the period between 01 January, 2017 and 31 January, 2018. The only exclusion criterion was the lack of ability to fully understand the study procedures and thus express consent to participate. The travelers received pre-travel advice although consultations mainly concerned tropical or intertropical countries. Data was collected through a post-travel questionnaire administered through telephone interviews in a period between 30 and 60 days after their return. Questionnaires were sent by e-mail after 5 unsuccessful phone contact attempts. Written consent was obtained from all participants and Ethics Committee approvals were obtained for both travel clinics. The questionnaire was based on the available literature [7]. It consisted of 40 questions mostly in multiple-choice form, divided into three sections:

demographic and travel characteristics, compliance with advice given during the pre-travel consultation, health problems during and after travel including necessary medical evaluations after the trip. The assessment of compliance with protection measures against vector-borne diseases was administered only to travellers visiting high-risk areas according to the Center for Disease Control and Prevention (CDC) and the Scottish and British Travel Health consultation sites [8,9,10]. Diarrhea was defined according to WHO criteria [11].

### **Statistical methods**

Collected data was processed for descriptive analysis through Microsoft Excel®, 365. The univariate model was used to process answers provided by international travellers. Finally, we assessed the association between health problems and traveller characteristics (gender, age and profession), some travel characteristics (destination, duration of travel and type of accommodation) as well as some behavioral risks. Bivariate and multivariate models were used. For the bivariate model, the  $p$ -value of the Pearson Chi-square test and Fisher's exact test  $<0.05$  was considered statistically significant. The Yates correction for the Pearson Chi-square test was used when needed. When bivariate  $p$ -values were less than 0.10, a multivariate logistic regression model was used, and adjusted odds ratios and 95% confidence intervals were calculated. Statistical analyzes were carried out with the SPSS Statistics Program (Version 24, IBM Corp., USA).

## **RESULTS**

### **Sample description**

579 international travellers were enrolled from the two Travel Medicine Centers. Of these, a total of 468 travellers (80.8%) were admitted to the study after the telephone questionnaire: 61.5% from Travel Center in Veneto Region and the others from Travel Center in Emilia-Romagna Region. 27 travellers canceled or postponed the trip, 77 were not available after the 5 predetermined telephone attempts and subsequent e-mail notice. 5 travellers were excluded because of the lack of demographic data and 2 were excluded because their travel destination was within Europe, with a risk comparable to Italy.

Demographic features, destinations and travel information are presented in Table 1. 65.8% of all travellers preferred hotels but only 42.5% chose hotels as the only type of accommodation for the entire duration of the trip. 9.6% chose

B&B or guest houses as the only type of accommodation, 10.7% stayed exclusively in a private home, 2.1% in a hostel and 1.7% in a tent.

### **Chemoprophylaxis and vaccination**

50% of travellers went to malaria endemic areas and were advised to take chemoprophylaxis. Of those, 67% went to Africa, 23% to Asia and 10% to America. Only 56% of travellers actually took chemoprophylaxis. Most of them (58%) took atovaquone/proguanil, 37% mefloquine and 5% doxycycline. 96% of travellers went to at-risk areas for vector-borne diseases. Only 24% of them followed all recommended mechanical prophylaxis measures to protect against insect bites (insect repellents, light and covering clothes, mosquito nets on the windows or above the bed).

Vaccines against typhoid fever and hepatitis-A were the most recommended vaccinations (84% and 71% of all international travellers respectively), but only 70% of travellers decided to get a typhoid fever vaccination and 61% took the hepatitis-A vaccination. TBE vaccine is among the most recommended vaccination for travelers but was not considered because the population of travelers who choose Europe as their destination has been excluded.

Similarly, for other vaccinations, the percentage of travellers who decided to be vaccinated was lower than the percentage of travellers who were advised to be vaccinated.

### **Compliance to behavioral advice**

68% of international travellers consumed raw food at least once during their trip. In particular, 55% consumed raw vegetables, 36% fruit not personally peeled, 12% uncooked fish and 4% uncooked meat. Most travellers consumed raw/uncooked food in Asia. Uncooked meat/fish was eaten more frequently by young travellers (16-30 years) than middle-aged (31-50 years) and older travellers ( $\geq 51$  years). The percentage of travellers who consumed food containing milk and/or eggs was 81%. In particular, 64% of travellers consumed eggs, 32% cream, 23% fresh and soft cheeses and 22% mayonnaise and/or sauces. 32% of travellers consumed street vendor food, mostly in Asia. 30% of travellers consumed drinks containing ice, most frequently in Asia. 32% of travellers reported that they always washed their hands before meals and 47% always brushed their teeth with bottled water. Regarding their profession, 100% of travellers working in a restaurant or in the catering field always washed their hands before meals. Hands were washed before meals and teeth brushed with bottled water most frequently by travellers in Asia.

### **Health problems**

Health problems during travel were reported by 39% of international travellers. The most frequent reported health problems were gastrointestinal (75% of travellers with health problems during travel), followed by dermatological problems (15%) and febrile diseases (14%). In particular, 22% of all international travellers had diarrhea during travel.

Health problems after travel were reported by 19% of international travellers. As during travel, the most reported health problems after travel were gastrointestinal (68% of travellers with health problems after travel), followed by febrile diseases (16%). Demographics, destinations and travel information of international travellers with health problems during and after travel are presented in Table 2. 37% of travellers with health problems after travel experienced the illness the day of return and 46% within the first 7 days. In the remaining 17% of travellers, the disorders occurred between 15 and 30 days. 6% of travellers saw a doctor when they come back from their trip and 8% took medication. Most of them took antibiotics (37%). Altogether 46% of travellers presented health problems during and/or after travel. Most of them were females (54%), young travellers (16-30 years, 49%), went to Asia (51%), for a medium-length trip (15-29 days, 53%) and for tourism reasons (63%). The most reported health problems were gastrointestinal (75% of all travellers with health problems). Figure 1 summarizes demographics of participants, travel information and travellers' health problems divided according to travel destination.

### **Risk factors**

As reported in Table 3 the risk factors for total health problems (during and/or after travel) were female gender (OR=1.62; IC95% 1.08 – 2.44), youth (16-30 years; OR=4.00; IC95% 2.29 – 7.00) and intermediate age (31-50 years; OR=2.77; IC95% 1.61- 4.78), intermediate travel duration (15-29 days; OR=2.00 IC95% 1.24 – 3,21) and profession other than health workers and those in the catering field (OR=2.49; IC95% 1.21-5.09). Only youth (16-30 years; OR=5.39; IC95% 2.88 – 10.08) and intermediate age (31-50 years; OR=2.54; IC95% 1.35 – 4.75) and the intermediate travel duration (15-29 days; OR=1.79 IC95% 1.08 – 2.96) were risk factors for total gastrointestinal problems during and/or after travel. The risk factors for total febrile diseases (during and/or after travel) were intermediate travel duration (15-29 days; OR=4.48; IC95% 1.60 – 14.67), American continent as a destination (OR=2.83; IC95% 1.06 – 7.57), accommodation in hostels (OR=2.55; IC95% 1.02 – 6.36) and in a tent (OR=3.47; IC95% 1.27-9.48). The longer travel duration (over 30 days; OR=15.49; IC95% 1.64 – 146.48), Asian (OR=9.90; IC95% 1.16 – 84.69) and American continent as destination (OR=12.45; IC95% 1.36 – 113.86) were risk factors for total respiratory problems during and/or after travel. Finally, only youth (16-30 years; OR=5.37; IC95% 1.15 – 25.06) and street vendor food (OR=2.99; IC95% 1.32 – 6.80) were risk factors for total dermatological problems.

## **DISCUSSION**

### **Compliance to behavioral advice**

Due to the continuous increase of international travel, a qualified pre-travel consultation is essential. However, adherence to the recommendations remains low. This situation is in line with other study results. In a previous Italian study, one third of travellers consumed food and at-risk drinks [7]. In a German study, 80% of travellers did not follow the traditional food recommendations [12]. In our study, the consumption of at-risk food occurred mostly in young people going to Asia. The same situation was described in an Israeli study: adherence to food recommendations is much more respected in elderly travellers (over 60 years) than in young people (20-30 years) [13]. Even in a Finnish study, elderly travellers paid more attention to behavioral recommendations, compared to young people traveling to Asia [14].

### **Health problems**

Almost 40% of our travellers presented a health problem during their trip. This percentage is clearly lower than previous international studies: 76% of Finnish travellers, 64% of American travellers, 64% of British travellers, 70% of Israeli travellers [14,15,16,17]. Instead, this is similar to the percentage in other international studies: 43% in Germany, 38% in Switzerland and 49% in Sweden [18,19,20]. The percentage is higher only if compared with two recent American studies [21,22]. However, Balaban's study was carried out with public health professionals, who probably were better prepared in terms of behavioral risks. In line with all the studies mentioned above, also in our study, the most frequently reported health problem during travel was diarrhea.

Almost a fifth of the travellers in our study (19%) presented health problems upon their return. This percentage is lower than reported in Hill's (26%) and Vilkmann's study (32%) [14,23]. As in Hill's study, diarrhea was the most frequent health problem in our study. In Vilkmann's study fever was the most frequent illness after travel.

### **Risk factors**

Gender, age, duration of travel and profession are risk factors for health problems during and/or after travel. In particular, similarly to the Finnish study, female gender (OR=1.62; IC95% 1.08-2.44) was a factor predisposing to health problems [14]. Youth (16-30 years; OR=4; IC95% 2.29-7) was a risk factor, as in previous studies [12,13,23]. In our study, as in the Finnish one, adults (31-50 years; OR=2.77; IC95% 1.61-4.78) were found to be at greater risk than >50 years subjects [14]. Age and duration of travel were the only risk factors for gastrointestinal problems during and/or after the trip. Young people (16-30 years; OR=5.39; IC95% 2.88-10.08) and adults (31-50 years; OR=2.54; IC95% 1.35-4.75) were more at risk of developing gastrointestinal problems, but no specific foods resulted as a risk factor. This could be due to the fact that during travel young people and <50-year-old adults have a lifestyle more similar to that of the local population and greater interaction with them. The intermediate duration of travel (15-29 days; OR=1.79; IC95% 1.08-2.96) was found to be a risk factor. When on short trips, travellers are probably more compliant with the



recommendations given during the pre-travel consultation, while after two weeks of travel, there is a progressive decrease in compliance with recommendations and travellers engage in more dangerous behaviors.

The intermediate duration of travel (15-29 days; OR=4.84; IC95% 1.60-14.67), the American continent as destination (OR=2.83; IC95% 1.06-7.57) and accommodation in hostels (OR=2.55; IC95% 1.02-6.36) and a tent (OR=3.47; IC95% 1.27-9.48) were risk factors for fever during and/or after travel. This differs from previous studies, in which fever is the most frequent disorder in travellers heading to Sub-Saharan Africa and South East Asia [24,25]. Our travellers probably had greater exposure to fever-causing agents in America. The fact that even the type of accommodation (hostel and tent) is a risk factor, suggests that travellers were possibly more exposed to vector-borne diseases in America.

The duration of travel over 30 days (OR=15.49; IC95% 1.64-146.48), the American continent (OR=12.45; IC95% 1.36-113.86) and the Asian one as destination (OR=9.90; IC95% 1.16-84.69) were risk factors for respiratory problems during and/or after travel. This result was also reported in a Scottish study [26]. One hypothesis to explain this could be the greater exposure to air pollution in these two continents and the longer time spent airborne to reach these two continents from Italy. Long plane travel is a known risk of respiratory infections [27,28].

Finally, youth (16-30 years; OR=5.37; IC95% 1.15-25.06) and consumption of street vendor food (OR=2.99; IC95% 1.32-6.80) were risk factors for dermatological problems. Identifying the presence of specific allergens in street vendor food is not always easy, so this could be the cause of rash or dermatitis in travellers.

### **Limitations**

There are some limitations in our study. Firstly, our travellers are not representative of all travelers, as indeed they were enrolled among those who spontaneously requested pre-travel consultations. A wide range of travellers is excluded, especially those who, for various reasons, do not know or underestimate risks associated with staying in tropical and subtropical areas. Secondly, the proportion of non-Italian travellers, especially VFR, was also reduced due to linguistic barriers. This fact does not permit assessing the presence of different risk factors in these two categories of travellers. Thirdly, the type of restaurant where travellers consumed food and drink has not been investigated. This information could help to better understand the risk factors of gastrointestinal problems. Finally, statistical analyses in some subgroups of travellers, such as those with respiratory and dermatological diseases, are difficult to assess due to small sample size. An increase of the sample would be required.

### **CONCLUSIONS**

Strict adherence to behavioral recommendations must be implemented especially in young people despite the non-seriousness of symptoms reported during the trip and upon return; indeed, no patient required hospitalization. About

19% of travellers reported symptoms within 60 days after their return; this highlights the importance of advising travellers upon their return on potential health risks, as during pre-trip counseling.

Gastrointestinal disorders, mainly diarrhea, are not associated with the intake of specific types of food or drink, but rather depend on more generic factors; this implies the need to alert the travellers not only to risk factors, but also on measures to be implemented in the event of disorder onset and which drugs to always have with them for self-medication.

Travellers must undertake journeys that require long flights, such as those traveling to America or Asia, must be warned of the potential risks of acquiring respiratory diseases and to the need to carry self-medication.

In travellers with pre-existing allergies/intolerances it is recommended they avoid the intake of foods in which it is not possible to identify the presence of possible allergens, such as street vendor food.

Finally, even adherence to chemoprophylaxis, mechanical prophylaxis and vaccination remains low. It is important to enhance awareness of the risk of vaccine preventable diseases during the consultation.

Pre-travel consultation is the only way to alert travellers to specific risks associated with their trip and to recommend specific prophylaxis. Regarding vaccinations, it is important to inform travellers on the best timing to go for pre-travel consultation, in order to be immunized on time.

### **Funding**

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

### **Conflict of interest statement**

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors. GG received grants from GlaxoSmithKline Biologicals SA, Sanofi Pasteur MSD, Novartis, Crucell/Janssen, Seqirus, Sanofi Pasteur, Merck Italy, PaxVax and Pfizer for consultant work or taking part in an Advisory Board, expert meetings, being a speaker or an organizer of congresses/conferences and acting as investigator in clinical trials. AP received grants from GlaxoSmithKline and Pfizer for taking part to expert meetings. EDS received grants from Pfizer for taking part in expert meetings. All remaining authors have declared no conflicts of interest.

The study was approved by Ethics Committee of Ferrara and Padova.

**Authors' contributions**

All Authors have made a substantial contribution to conception, design of the article, as well as data analysis and interpretation, drafting the article and revising it critically for intellectual content; all Authors approve the final version of the manuscript.

Journal Pre-proof

**References**


- [1] United States. Census Bureau n.d. <https://www.census.gov/> (accessed September 4, 2019).
- [2] United Nations World Tourism Organization (UNWTO) 2018. <http://www2.unwto.org/en> (accessed September 5, 2019).
- [3] Kozarsky PE, Henry R. Introduction to Travel Health & the CDC Yellow Book 2018;16. <https://wwwnc.cdc.gov/travel/yellowbook/2020/introduction/introduction-to-travel-health-and-the-cdc-yellow-book> (accessed September 5, 2019).
- [4] R S, S. G. Epidemiology: Morbidity and Mortality in Travellers. *Travel Med* 3rd Ed 2013:5–11.
- [5] Leder K, Torresi J, Libman MD, Cramer JP, Castelli F, Schlagenhauf P, et al. GeoSentinel surveillance of illness in returned travelers, 2007-2011. *Ann Intern Med* 2013;158:456–68. doi:10.7326/0003-4819-158-6-201303190-00005.
- [6] Chen LH, Hochberg NS. The Pretravel Consultation - Chapter 2 - 2020 Yellow Book. *Travelers' Health*. CDC n.d. <https://wwwnc.cdc.gov/travel/yellowbook/2020/preparing-international-travelers/the-pretravel-consultation> (accessed September 5, 2019).
- [7] Laverone E, Boccalini S, Bechini A, Belli S, Santini MG, Baretta S, et al. Travelers' compliance to prophylactic measures and behavior during stay abroad: results of a retrospective study of subjects returning to a travel medicine center in Italy. *J Travel Med* n.d.;13:338–44. doi:10.1111/j.1708-8305.2006.00068.x.
- [8] Travelers' Health. CDC n.d. <https://wwwnc.cdc.gov/travel> (accessed September 5, 2019).
- [9] Health Protection Scotland. Fit for travel n.d. <https://www.fitfortravel.nhs.uk/home> (accessed September 5, 2019).
- [10] NaTHNaC - Home n.d. <https://travelhealthpro.org.uk/> (accessed September 5, 2019).
- [11] Diarrhoeal disease n.d. <https://www.who.int/news-room/fact-sheets/detail/diarrhoeal-disease> (accessed September 5, 2019).
- [12] Rack J, Wichmann O, Kamara B, Günther M, Cramer J, Schönfeld C, et al. Risk and Spectrum of Diseases in Travelers to Popular Tourist Destinations. *J Travel Med* 2006;12:248–53. doi:10.2310/7060.2005.12502.
- [13] Alon D, Shitrit P, Chowder M. Risk behaviors and spectrum of diseases among elderly travelers: a comparison of younger and older adults. *J Travel Med* n.d.;17:250–5. doi:10.1111/j.1708-8305.2010.00425.x.
- [14] Vilkin K, Pakkanen SH, Lääveri T, Siikamäki H, Kantele A. Travelers' health problems and behavior: prospective study with post-travel follow-up. *BMC Infect Dis* 2016;16:328. doi:10.1186/s12879-016-1682-0.
- [15] Chen LH, Han P V., Wilson ME, Stoney RJ, Jentes ES, Benoit C, et al. Self-reported illness among Boston-area international travelers: A prospective study. *Travel Med Infect Dis* 2016;14:604–13.

doi:10.1016/j.tmaid.2016.09.009.

- [16] Evans MR, Shickle D, Morgan MZ. Travel illness in British package holiday tourists: prospective cohort study. *J Infect* 2001;43:140–7. doi:10.1053/jinf.2001.0876.
- [17] Winer L, Alkan M. Incidence and precipitating factors of morbidity among Israeli travelers abroad. *J Travel Med* n.d.;9:227–32. doi:10.2310/7060.2002.24202.
- [18] Fleck S, Jäger H, Zeeb H. Travel and health status: a survey follow-up study. *Eur J Public Health* 2006;16:96–100. doi:10.1093/eurpub/cki144.
- [19] Bruni M, Steffen R. Impact of travel-related health impairments. *J Travel Med* 1997;4:61–4. doi:10.1111/j.1708-8305.1997.tb00781.x.
- [20] Ahlm C, Lundberg S, Fessé K, Wiström J. Health problems and self-medication among Swedish travellers. *Scand J Infect Dis* 1994;26:711–7. doi:10.3109/00365549409008640.
- [21] Stoney RJ, Kozarsky P, Bostick RM, Sotir MJ. International travellers from New Jersey: piloting a travel health module in the 2011 Behavioral Risk Factor Surveillance System survey. *J Travel Med* 2016;23. doi:10.1093/jtm/tav015.
- [22] Balaban V, Warnock E, Ramana Dhara V, Jean-Louis LA, Sotir MJ, Kozarsky P. Health risks, travel preparation, and illness among public health professionals during international travel. *Travel Med Infect Dis* n.d.;12:349–54. doi:10.1016/j.tmaid.2014.01.007.
- [23] Hill DR. Health problems in a large cohort of Americans traveling to developing countries. *J Travel Med* n.d.;7:259–66. doi:10.2310/7060.2000.00075.
- [24] Freedman DO, Weld LH, Kozarsky PE, Fisk T, Robins R, von Sonnenburg F, et al. Spectrum of disease and relation to place of exposure among ill returned travelers. (Erratum appears in *N Engl J Med*. 2006 Aug 31;355(9):967). *N Engl J Med* 2006;354:119–30. doi:10.1056/NEJMoa051331.
- [25] Wilson ME, Freedman DO. Etiology of travel-related fever. *Curr Opin Infect Dis* 2007;20:449–53. doi:10.1097/QCO.0b013e3282a95e27.
- [26] Redman CA, Maclennan A, Wilson E, Walker E. Diarrhea and respiratory symptoms among travelers to Asia, Africa, and South and Central America from Scotland. *J Travel Med* n.d.;13:203–11. doi:10.1111/j.1708-8305.2006.00046.x.
- [27] Hertzberg VS, Weiss H, Elon L, Si W, Norris SL, FlyHealthy Research Team. Behaviors, movements, and transmission of droplet-mediated respiratory diseases during transcontinental airline flights. *Proc Natl Acad Sci U S A* 2018;115:3623–7. doi:10.1073/pnas.1711611115.
- [28] Mangili A, Vindenes T, Gendreau M. Infectious Risks of Air Travel. *Microbiol Spectr* 2015;3.

doi:10.1128/microbiolspec.IOL5-0009-2015.

Journal Pre-proof



**Figure 1.** Travellers' characteristics, travel and health problems during and / or after travel, according to the continent of destination.