

# A $p$ -adically entire function with integral values on $\mathbb{Q}_p$ and entire liftings of the $p$ -divisible group $\mathbb{Q}_p/\mathbb{Z}_p$

Francesco Baldassarri

With an appendix by Maurizio Candilera

(Communicated by Peter Schneider)

*Dedicated to Christopher Deninger on his 60th birthday*

**Abstract.** We give a self-contained proof of the fact, discovered in [1] and proven in [2] with the methods of [6], that, for any prime number  $p$ , there exists a power series

$$\Psi = \Psi_p(T) \in T + T^2\mathbb{Z}[[T]],$$

which trivializes the addition law of the formal group of Witt covectors [6], [15, II.4], is  $p$ -adically entire and assumes values in  $\mathbb{Z}_p$  all over  $\mathbb{Q}_p$ . We actually generalize, following a suggestion of M. Candilera, the previous facts to any fixed unramified extension  $\mathbb{Q}_q$  of  $\mathbb{Q}_p$  of degree  $f$ , where  $q = p^f$ . We show that  $\Psi = \Psi_q$  provides a quasi-finite covering of the Berkovich affine line  $\mathbb{A}_{\mathbb{Q}_p}^1$  by itself. We prove in Section 4 new strong estimates for the growth of  $\Psi$ , in view of the application [3] to  $p$ -adic Fourier expansions on  $\mathbb{Q}_p$ . We refer to [3] for the proof of a technical corollary (Proposition 4.13) which we apply here to locate the zeros of  $\Psi$  and to obtain its product expansion (Corollary 4.16). We reconcile the present discussion (for  $q = p$ ) with the formal group proof given in [2], which takes place in the Fréchet algebra  $\mathbb{Q}_p\{x\}$  of the analytic additive group  $\mathbb{G}_{a, \mathbb{Q}_p}$  over  $\mathbb{Q}_p$ . We show that, for any  $\lambda \in \mathbb{Q}_p^\times$ , the closure  $\mathcal{E}_\lambda^\circ$  of  $\mathbb{Z}_p[\Psi(p^i x/\lambda) \mid i = 0, 1, \dots]$  in  $\mathbb{Q}_p\{x\}$  is a Hopf algebra object in the category of Fréchet  $\mathbb{Z}_p$ -algebras. The special fiber of  $\mathcal{E}_\lambda^\circ$  is the affine algebra of the  $p$ -divisible group  $\mathbb{Q}_p/p\lambda\mathbb{Z}_p$  over  $\mathbb{F}_p$ , while  $\mathcal{E}_\lambda^\circ[1/p]$  is dense in  $\mathbb{Q}_p\{x\}$ . From  $\mathbb{Z}_p[\Psi(\lambda x) \mid \lambda \in \mathbb{Q}_p^\times]$  we construct a  $p$ -adic analog  $\mathcal{AP}_{\mathbb{Q}_p}(\Sigma_\rho)$  of the algebra of Dirichlet series holomorphic in a strip  $(-\rho, \rho) \times i\mathbb{R} \subset \mathbb{C}$ . We start developing this analogy. It turns out that the Banach algebra of almost periodic functions on  $\mathbb{Q}_p$  identifies with the topological ring of germs of holomorphic almost periodic functions on strips around  $\mathbb{Q}_p$ .

## CONTENTS

1. Introduction	268
2. Rings of $p$ -adic analytic functions	273
3. Elementary proofs of the main properties of $\Psi$	279

4. Valuation and Newton polygons of $\Psi_q$	283
5. Rings of continuous functions on $\mathbb{Q}_p$	292
6. $p$ -adically entire functions bounded on $\mathbb{Q}_p$	301
Appendix A. Non archimedean topological algebra	306
Appendix B. Classical theory of almost periodic functions	309
Appendix C. Numerical calculations by M. Candilera	312

1. INTRODUCTION

1.1. **Foreword.** An unfortunate feature of  $p$ -adic numbers is that there exists no character

$$\psi: (\mathbb{Q}_p, +) \rightarrow (\mathbb{C}_p^\times, \cdot), \quad \psi \neq 1,$$

which extends to an entire function  $\mathbb{C}_p \rightarrow \mathbb{C}_p$ . In fact, let  $\pi_p \in \mathbb{C}_p^{\circ\circ}$  be such that the radius of convergence of  $\exp(\pi_p x)$  equals 1, so that  $\exp$  and  $\log$  establish an isomorphism

$$(\pi_p \mathbb{C}_p^{\circ\circ}, +) \xrightarrow{\sim} (\exp(\pi_p \mathbb{C}_p^{\circ\circ}), \cdot) \subset (1 + \mathbb{C}_p^{\circ\circ}, \cdot).$$

Now assume a  $\psi$  as above exists, and let  $n$  be a positive integer such that  $\psi(p^n) \in \exp(\pi_p \mathbb{C}_p^{\circ\circ})$  so that  $\psi$  restricts to a character  $\psi: (p^n \mathbb{Z}_p, +) \rightarrow (\exp(\pi_p \mathbb{C}_p^{\circ\circ}), \cdot)$ . Let  $a := \log(\psi(p^n))$ . Then, for any  $x \in \mathbb{Z}$ ,  $\psi(p^n x) = \psi(p^n)^x = \exp(ax)$ . But  $x \mapsto \exp(ax)$  has a finite radius of convergence.

We partially remedy to the previous inconvenience by showing the existence, for any  $\lambda \in \mathbb{Q}_p^\times$ , of a representable formal group functor

$$(1) \quad \mathbb{E}_\lambda: \mathcal{ACLM}_{\mathbb{Z}_p}^u \rightarrow \mathcal{A}b$$

(see Appendix A.1 for the notation), whose generic (resp. special) fiber is the  $\mathbb{Q}_p$ -analytic group  $\mathbb{G}_a$  (resp. the constant  $p$ -divisible group  $\mathbb{Q}_p/\lambda\mathbb{Z}_p$  over  $\mathbb{F}_p$ ). The idea is the following. Over the complex numbers the formulas

$$e^{iz} = \cos z + i \sin z, \quad e^{-iz} = \cos z - i \sin z,$$

show that the two (Hopf) algebras  $\mathbb{Z}[i][e^{iz}, e^{-iz}]$  and  $\mathbb{Z}[i][\sin z, \cos z]$  coincide. The sequence of functions

$$\Psi(x) = \Psi_p(x), \Psi(px), \Psi(p^2x), \dots$$

plays here the role of the pair  $(\cos z, \sin z)$  in that the  $p$ -adically entire and integral addition law (7) holds, and  $x$  is a logarithm for that formal group. So, while it is improper to say that  $\Psi$  plays the role of an entire character of  $\mathbb{Q}_p$ , it is suggestive to consider a suitable  $p$ -adic completion of the algebra  $\mathbb{Z}_p[\Psi(\lambda x) \mid \lambda \in \mathbb{Q}_p^\times]$  and to compare it with the classical algebras of Bohr’s almost periodic functions  $APH_{\mathbb{R}}$  and  $AP_{\mathbb{R}}$ . We review for convenience the classical definitions of real and complex Fourier analysis in Appendix B.5. A closer  $p$ -adic analog of those classical constructions, and a generalization of Amice–Fourier theory to  $p$ -adic functions on  $\mathbb{Q}_p$  will appear in [3].

1.2. **The function  $\Psi$ .** In the paper [2] we introduced, for any prime number  $p$ , a power series

$$\Psi(T) = \Psi_p(T) = T + \sum_{i=2}^{\infty} a_i T^i \in \mathbb{Z}[[T]],$$

which represents an entire  $p$ -adic analytic function, i.e., is such that

$$\limsup_{i \rightarrow \infty} |a_i|_p^{1/i} = 0.$$

This function has the remarkable property that  $\Psi_p(\mathbb{Q}_p) \subset \mathbb{Z}_p$  and that, for any  $i \in \mathbb{Z}$  and  $x \in \mathbb{Q}_p$ , if we write  $x$  as in (12), with  $x_i$  defined by (13), (14), then

$$x_{-i} = \Psi_p(p^i x) \pmod{p} \in \mathbb{F}_p.$$

The power series  $\Psi(T)$  is defined by the functional relation

$$(2) \quad \sum_{j=0}^{\infty} p^{-j} \Psi(p^j T)^{p^j} = T.$$

Its inverse function  $\beta = \beta_p \in T + T^2\mathbb{Z}[[T]]$  was shown to converge exactly in the region

$$|T|_p < p, \quad \text{i.e.,} \quad v_p(T) > -1.$$

One property we had failed to notice in [2] is the following.

**Proposition 1.3.** *The restriction of the function  $\Psi_p$  to a map  $\mathbb{Q}_p \rightarrow \mathbb{Z}_p$  is uniformly continuous. More precisely, for any  $j = 0, 1, \dots$  and  $x \in \mathbb{Q}_p$ ,*

$$\Psi_p(x + p^j \mathbb{C}_p^\circ) \subset \Psi_p(x) + p^j \mathbb{C}_p^\circ.$$

This is proven in Corollary 4.5 below. See also the more general Theorem 4.14 whose proof depends on Proposition 4.13, proven in [3].

1.4. **Our previous approach** [2]. Proofs in [2] were based on Barsotti–Witt algorithms [6]. The most basic notion of topological algebra in [6] is the one of a *simultaneously admissible* family, indexed by  $\alpha \in A$ , of sequences  $i \mapsto x_{\alpha,-i}$  for  $i = 0, 1, \dots$  in a Fréchet algebra  $R$  over  $\mathbb{Z}_p$  (in particular, over  $\mathbb{F}_p$ ) [6, Ch. 1, §1]. In case  $R$  is a Fréchet algebra over  $\mathbb{Q}_p$  the definition of simultaneous admissibility is more restrictive, but the name used in *loc. cit.* is the same. For clarity, the more restrictive notion will be called here (simultaneous) *PD-admissibility*, while the general notion will maintain the name of (simultaneous) *admissibility*.

Using the previous refined terminology, our main technical tool in [2] was a criterion [2, Lemma 1] of *simultaneous PD-admissibility* for a family indexed by  $\alpha \in A$ , of sequences  $i \mapsto x_{\alpha,-i}$  for  $i = 0, 1, \dots$  in a Fréchet  $\mathbb{Q}_p$ -algebra. In Barsotti’s theory of  $p$ -divisible groups one regards an admissible sequence  $i \mapsto x_{-i}$  as a *Witt covector*  $(\dots, x_{-2}, x_{-1}, x_0)$  [6, 15] with components  $x_{-i} \in R$ .

We take here only a short detour on the group functor viewpoint and refer the reader to [15] for precisions. As abelian group functors on a suitable

category of topological  $\mathbb{Z}_p$ -algebras, the direct limit  $W_n \rightarrow W_{n+1}$  of the Witt vector groups of length  $n$  via the *Verschiebung map*

$$V : (x_{-n}, \dots, x_{-1}, x_0) \rightarrow (0, x_{-n}, \dots, x_{-1}, x_0)$$

indeed exists. It is the group functor CW of *Witt covectors*. For a topological  $\mathbb{Z}_p$ -algebra  $R$  on which CW( $R$ ) is defined, it is convenient to denote an element  $x \in \text{CW}(R)$  by an inverse sequence

$$x = (\dots, x_{-2}, x_{-1}, x_0)$$

of elements of  $R$ , that is, a Witt covector with components in  $R$ . Two Witt covectors  $x = (\dots, x_{-2}, x_{-1}, x_0)$  and  $y = (\dots, y_{-2}, y_{-1}, y_0)$  with components  $R$  can be summed by taking limits of sums of finite Witt vectors. Namely, let

$$(3) \quad \varphi_i(X_0, \dots, X_i; Y_0, \dots, Y_i) \in \mathbb{Z}[X_0, \dots, X_i, Y_0, \dots, Y_i]$$

be the  $i$ -th (= the last!) entry of the Witt vector  $(X_0, \dots, X_i) + (Y_0, \dots, Y_i)$ . Then

$$x + y = z = (\dots, z_{-2}, z_{-1}, z_0)$$

means that, for any  $i = 0, -1, \dots$ ,

$$(4) \quad z_i = \lim_{n \rightarrow +\infty} \varphi_n(x_{i-n}, x_{i-n+1}, \dots, x_i; y_{i-n}, y_{i-n+1}, \dots, y_i)$$

converges in  $R$ . The convergence properties on the Witt covectors  $x$  and  $y$  above for the expressions (4) to converge, are dictated by the following lemma.

**Lemma 1.5** ([6, Teorema 1.11]). *With the notation as above, for  $i = 0, 1, 2, \dots$ , let us attribute the weight  $p^i$  to the variables  $X_i, Y_i$ . Then, for any  $i \geq 0$ , the polynomial  $\varphi_i$  in (3) is isobaric of weight  $p^i$ . Moreover, for any  $i \geq 1$ ,*

$$\begin{aligned} &\varphi_i(X_0, X_1, \dots, X_i; Y_0, Y_1, \dots, Y_i) - \varphi_{i-1}(X_1, \dots, X_i; Y_1, \dots, Y_i) \\ &\in X_0 Y_0 \mathbb{Z}[X_0, X_1, \dots, X_i, Y_0, Y_1, \dots, Y_i]. \end{aligned}$$

So, we equip the polynomial ring

$$\mathbb{Z}[X_0, X_{-1}, \dots, X_{-i}, \dots; Y_0, Y_{-1}, \dots, Y_{-i}, \dots]$$

with the linear topology defined by the powers of the ideals

$$I_N := (X_{-N}, X_{-N-1}, \dots; Y_{-N}, Y_{-N-1}, \dots),$$

and set

$$\mathcal{P} := \varprojlim_{N, M \rightarrow +\infty} \mathbb{Z}[X_0, X_{-1}, \dots, X_{-i}, \dots; Y_0, Y_{-1}, \dots, Y_{-i}, \dots] / I_N^M.$$

Then the sequence

$$(5) \quad i \mapsto \varphi_i(X_{-i}, \dots, X_{-1}, X_0; Y_{-i}, \dots, Y_{-1}, Y_0)$$

converges to an element

$$\Phi(X_0, X_{-1}, \dots, X_{-i}, \dots; Y_0, Y_{-1}, \dots, Y_{-i}, \dots) \in \mathcal{P}.$$

So, (4) is expressed more compactly as

$$(6) \quad z_i = \Phi(x_i, x_{i-1}, \dots; y_i, y_{i-1}, \dots)$$

**Remark 1.6.** The projective limit

$$W_{n+1} \rightarrow W_n, \quad (x_0, x_1, \dots, x_{n+1}) \mapsto (x_0, x_1, \dots, x_n),$$

produces instead the algebraic group  $W$  of *Witt vectors*.

The approach of Barsotti [6] is more flexible and easier to apply to analytic categories. If  $R$  is complete, for two simultaneously admissible Witt covectors  $x = (\dots, x_{-2}, x_{-1}, x_0)$  and  $y = (\dots, y_{-2}, y_{-1}, y_0)$  with components  $R$ , the expressions (6) all converge in  $R$  and define  $(\dots, z_{-2}, z_{-1}, z_0) = z =: x + y$ , which is in turn simultaneously admissible with  $x$  and  $y$ . In the  $\mathbb{Q}_p$ -algebra case, a Witt covector  $x = (\dots, x_{-2}, x_{-1}, x_0)$  has *ghost components*  $(\dots, x^{(-2)}, x^{(-1)}, x^{(0)})$  defined by

$$x^{(i)} = x_i + p^{-1}x_{i-1}^p + p^{-2}x_{i-2}^{p^2} + \dots, \quad i = 0, -1, -2, \dots$$

Under very general assumptions, [6, Teorema 1.11], a finite family of sequences  $(x_{\alpha, -i})_{i=0,1,\dots}$ , for  $\alpha \in A$  in a  $\mathbb{Q}_p$ -Fréchet algebra, are simultaneously PD-admissible if and only if the same holds for the family of sequences of ghost components  $(x_{\alpha}^{(-i)})_{i=0,1,\dots}$ , for  $\alpha \in A$ . Under these assumptions, for simultaneously PD-admissible covectors  $x$  and  $y$ ,  $x + y = z$  is equivalent to

$$z^{(i)} = x^{(i)} + y^{(i)}, \quad i = 0, -1, -2, \dots$$

In the present case, which coincides with the case treated in [2], the sequences  $i \mapsto x_{-i} := p^i x$  and  $i \mapsto y_{-i} := p^i y$  are simultaneously PD-admissible in the standard  $\mathbb{C}_p$ -Fréchet algebra  $\mathbb{C}_p\{x, y\}$  of entire functions on  $\mathbb{C}_p^2$  [2, Lemma 1 and Lemma 3]. It follows from relation (2) that  $i \mapsto x_{-i} := p^i x$ , for  $i = 0, 1, 2, \dots$ , is the sequence of ghost components of  $x \mapsto x^{(-i)} := \Psi(p^i x)$ . Therefore, from [6, *loc. cit.*], we conclude that the two sequences  $i \mapsto \Psi(p^i x)$  and  $i \mapsto \Psi(p^i y)$  are simultaneously admissible in  $\mathbb{C}_p\{x, y\}$  as well. Moreover, by [6, *loc. cit.*] and the definition of the addition law of Witt covectors with coefficients in  $\mathbb{C}_p\{x, y\}$ , we have

$$\begin{aligned} & (\dots, \Psi(p^2(x+y)), \Psi(p(x+y)), \Psi(x+y)) \\ &= (\dots, \Psi(p^2x), \Psi(px), \Psi(x)) + (\dots, \Psi(p^2y), \Psi(py), \Psi(y)). \end{aligned}$$

Equivalently,  $\Psi$  satisfies the addition law [2, (11)]

$$(7) \quad \Psi(x+y) = \Phi(\Psi(x), \Psi(px), \dots; \Psi(y), \Psi(py), \dots),$$

where

$$(8) \quad \begin{aligned} & \Phi(\Psi(x), \Psi(px), \dots; \Psi(y), \Psi(py), \dots) \\ &= \lim_{i \rightarrow \infty} \varphi_i(\Psi(p^i x), \dots, \Psi(px), \Psi(x); \Psi(p^i y), \dots, \Psi(py), \Psi(y)), \end{aligned}$$

for the polynomials  $\varphi_i$  of (3) and (5). Notice that (12) may be restated to say that, for any  $x \in \mathbb{Q}_p$ ,

$$x = (\dots, x_{-2}, x_{-1}; x_0, x_1, \dots),$$

where  $x_i \in \mathbb{F}_p$  is given by (13), as a *Witt bivector* [6] with coefficients in  $\mathbb{F}_p$ .

**1.7. Our present approach.** We present in Section 3 direct elementary proofs of the main properties of  $\Psi$ , which make no use of the Barsotti–Witt algorithms of [6]. Actually, following a suggestion of M. Candilera, we consider rather than (2), the more general functional relation for  $\Psi = \Psi_q$ ,  $q = p^f$ ,

$$(9) \quad \sum_{j=0}^{\infty} p^{-j} \Psi(p^j T)^{q^j} = T.$$

The result, at no extra work, will then be that (9) admits a unique solution  $\Psi_q(T) \in T + T^2\mathbb{Z}[[T]]$ . The series  $\Psi_q(T)$  represents a  $p$ -adically entire function such that  $\Psi_q(\mathbb{Q}_q) \subset \mathbb{Z}_q$ . In Section 4 we describe in the same elementary style the Newton and valuation polygons of the entire function  $\Psi_q$ , and obtain new estimates on the growth of  $|\Psi_q(x)|$  as  $|x| \rightarrow \infty$ , which will be crucial for the sequel [3]. From these estimates we also deduce, modulo the self-contained technical Proposition 4.13 whose proof appears in [3], the location of the zeros of  $\Psi_p$  (Theorem 4.16). Namely, any ball of radius 1,  $a + \mathbb{Z}_p \in \mathbb{Q}_p/\mathbb{Z}_p$ , contains precisely one (simple) zero of  $\Psi_p$ .

We present in Appendix C some numerical calculations due to M. Candilera, which exhibit the first coefficients of  $\Psi_p$  for small values of  $p$ . These calculations have been useful to us and we believe they may be quite convincing for the reader.

The function  $\Psi_q: \mathbb{A}_{\mathbb{Q}_p}^1 \rightarrow \mathbb{A}_{\mathbb{Q}_p}^1$  is a quasi-finite covering of the Berkovich affine line over  $\mathbb{Q}_p$  by itself. We do not know whether the previous covering is Galois.

**1.8. Convergence of Fourier-type expansions.** Section 2.1 describes some Hopf algebras whose existence follows from the addition properties of  $\Psi_p$ . Section 2.13 suggests an interpretation of the functions  $\Psi_p(x/\lambda)$ , for  $\lambda \in \mathbb{Q}_p^\times$ , as  $p$ -adic analogs of  $\exp(\frac{2\pi i}{\lambda}z)$ , for  $\lambda \in \mathbb{R}^\times$ . We are naturally lead to the question of which functions can be expressed as uniform limits on  $\mathbb{Q}_p$  of the previous functions. By analogy to the classical case, we call these functions uniformly almost periodic on  $\mathbb{Q}_p$  and denote by  $AP_{\mathbb{Q}_p}$  the corresponding closed subalgebra of the Banach algebra  $\mathcal{C}_{\text{unif}}^{\text{bd}}(\mathbb{Q}_p, \mathbb{Q}_p)$  of bounded uniformly continuous functions  $\mathbb{Q}_p \rightarrow \mathbb{Q}_p$ . Although we do not have an intrinsic characterization of these functions, we can show that they may be seen as germs of holomorphic functions on a neighborhood of  $\mathbb{Q}_p$ . We point out that colimits for topological algebras are not in general supported by set-theoretic inductive limits (see Remark 6.8). Therefore, our uniform approximation Theorem 2.27 does not state that any uniformly almost periodic function on  $\mathbb{Q}_p$  necessarily extends to an analytic function on a  $p$ -adic strip around  $\mathbb{Q}_p$ . On the other hand,  $AP_{\mathbb{Q}_p}$  is dense in the Fréchet algebra  $\mathcal{C}(\mathbb{Q}_p, \mathbb{Q}_p)$  of continuous functions  $\mathbb{Q}_p \rightarrow \mathbb{Q}_p$ , equipped with the topology of uniform convergence on compact open subsets of  $\mathbb{Q}_p$ . The proofs of these facts are detailed in Sections 5 and 6. We spend some time in Section 5 to explain in categorical terms (clearly stated in Appendix A) the natural limit/colimit/tensor product formulas which justify the linear topologies of the previous function algebras. For example,  $\mathcal{C}(\mathbb{Q}_p, \mathbb{Z}_p)$

(but not  $\mathcal{C}_{\text{unif}}(\mathbb{Q}_p, \mathbb{Z}_p)$ ) is a Hopf algebra related to the constant  $p$ -divisible group  $\mathbb{Q}_p/\mathbb{Z}_p$  over  $\mathbb{Z}_p$  and to its “universal covering”  $\mathbb{Q}_p$ . A more complete discussion of these topological algebras and of their duality relation with the affine algebra of the universal covering of the  $p$ -divisible torus, interpreted as an algebra of measures, will appear in [5].

In Section 6 we prove the facts announced in Section 2.13, namely, Theorem 2.17, Theorem 2.19, Proposition 2.20, Proposition 2.21, Proposition 2.23, Proposition 2.24, and Theorem 2.27.

## 2. RINGS OF $p$ -ADIC ANALYTIC FUNCTIONS

**2.1. Entire functions bounded on  $p$ -adic strips.** (See Appendix A for notation of topological algebra and nonarchimedean functional analysis.) We describe here the Hopf algebra object  $\mathbb{Q}_p\{x\}$  in the category of Fréchet  $\mathbb{Q}_p$ -algebras equipped with the completed projective = inductive tensor product  $\widehat{\otimes}_{\pi, \mathbb{Q}_p} = \widehat{\otimes}_{\iota, \mathbb{Q}_p}$ , which consists of the global sections of the  $\mathbb{Q}_p$ -analytic group  $\mathbb{G}_a$ . We also consider boundedness conditions for the functions in  $\mathbb{Q}_p\{x\}$  on suitable neighborhoods of  $\mathbb{Q}_p$  in the Berkovich affine line  $\mathbb{A}_{\mathbb{Q}_p}^1$  over  $\mathbb{Q}_p$ .

Our notation for coproduct (resp. counit, resp. inversion) of a Hopf algebra object  $A$  in a symmetric monoidal category with monoidal product  $\otimes$  and unit object  $I$  is usually  $\mathbb{P} = \mathbb{P}_A: A \rightarrow A \otimes A$  (resp.  $\varepsilon = \varepsilon_A: A \rightarrow I$ , resp.  $\rho = \rho_A: A \rightarrow A$ ).

**Definition 2.2.** For any closed subfield  $K$  of  $\mathbb{C}_p$ , we denote by  $K\{x\} = K\{x_1, \dots, x_n\}$  the ring of entire functions on the  $K$ -analytic affine space  $(\mathbb{A}_K^n, \mathcal{O}_K)$ . The *standard Fréchet topology* on the  $K$ -algebra  $K\{x\}$  is induced by the family  $\{w_r\}_{r \in \mathbb{Z}}$  of valuations

$$w_r(f) := \inf_{x \in (p^{-r}\mathbb{C}_p^\circ)^n} v(f(x)),$$

for any  $f \in K\{x\}$ .

**Remark 2.3.** More generally, for bounded functions  $f: X \rightarrow (S, | \cdot |)$ , where  $X$  is a set and  $(S, | \cdot |)$  is a Banach ring in multiplicative notation,  $\|f\|_X = \sup_{x \in X} |f(x)|$  will denote the supnorm on  $X$ .

**Definition 2.4.** For any  $\rho > 0$  and any finite extension  $K/\mathbb{Q}_p$ , the  *$p$ -adic  $n$ -strip of width  $\rho$  around  $K^n$*  is the analytic domain which is the union  $\Sigma_\rho(K) = \Sigma_\rho^{(n)}(K)$  of all affinoid  $n$ -polydiscs of radius  $\rho$  centered at  $K$ -rational points. We denote by

$$\text{Res}_\rho: \mathbb{C}_p\{x\} \rightarrow \mathcal{O}(\Sigma_\rho), \quad f \mapsto f|_{\Sigma_\rho},$$

the restriction map. Clearly, the map  $\text{Res}_\rho$  is an injection. We let  $\mathcal{O}_K^{\text{bd}}(\Sigma_\rho(K))$  (resp.  $\mathcal{O}_K^\circ(\Sigma_\rho(K))$ ) denote the subring of  $\mathcal{O}_K(\Sigma_\rho(K))$ , consisting of functions bounded (resp. bounded by 1) on  $\Sigma_\rho(K)$ . We denote by  $\| \cdot \|_{K, \rho}$  the supnorm on  $\Sigma_\rho(K)$ . The Banach algebra structure on  $\mathcal{O}_K^{\text{bd}}(\Sigma_\rho(K))$  (resp.  $\mathcal{O}_K^\circ(\Sigma_\rho(K))$ ) induced by the norm  $\| \cdot \|_{K, \rho}$  will be called  *$K$ -uniform*. The Fréchet structure of  $\mathcal{O}_K(\Sigma_\rho(K))$  (resp.  $\mathcal{O}_K^\circ(\Sigma_\rho(K))$ ) induced by the family of seminorms of

Definition 2.2 will be called *standard*. We set, in particular,  $\Sigma_\rho = \Sigma_\rho^{(1)}(\mathbb{Q}_p)$  but will keep the notation  $\|\cdot\|_{\mathbb{Q}_p, \rho}$ . We also denote by  $\mathcal{H}_K^{(n), \text{bd}}(\rho)$  (resp.  $\mathcal{H}_K^{(n), \circ}(\rho)$ ) the subring of  $K\{x\}$  of functions which are bounded (resp. bounded by 1) on  $\Sigma_\rho(K)$ . We set

$$\mathcal{H}_K^{(n), \text{bd}} := \bigcap_{\rho} \mathcal{H}_K^{(n), \text{bd}}(\rho).$$

For any  $\rho > 0$  and any  $f \in \mathcal{H}_K^{(n), \text{bd}}(\rho)$ , we introduce one further valuation

$$w_{K, \infty}(f) := \inf_{x \in K^n} v(f(x)).$$

For  $n = 1$  and  $K = \mathbb{Q}_p$ , we shorten  $\mathcal{H}_K^{(n), \text{bd}}(\rho)$  (resp.  $\mathcal{H}_K^{(n), \circ}(\rho)$ , resp.  $\mathcal{H}_K^{(n), \text{bd}}$ , resp.  $w_{K, \infty}$ , resp.  $K$ -uniform) to  $\mathcal{H}^{\text{bd}}(\rho)$  (resp.  $\mathcal{H}^\circ(\rho)$ , resp.  $\mathcal{H}^{\text{bd}}$ , resp.  $w_\infty$ , resp. uniform).

**Remark 2.5.** It is not a priori clear that  $\mathcal{H}^{\text{bd}}$  contains non-constant functions. We will prove below (Theorem 4.14) that  $\Psi(x) \in \mathcal{H}^{\text{bd}}$ .

**Remark 2.6.** For any  $n$  and any  $\rho > 0$ ,  $\mathcal{H}_K^{(n), \circ}(\rho)$  is a closed  $K^\circ$ -subalgebra of  $K\{x_1, \dots, x_n\}$ ; the induced Fréchet  $K^\circ$ -algebra structure on  $\mathcal{H}_K^{(n), \circ}(\rho)$  will be called *standard*. Now it follows from formula (2) below that, by contrast,  $\mathcal{H}_K^{(n), \text{bd}}(\rho) = \mathcal{H}_K^{(n), \circ}(\rho)[1/p]$  is dense in  $K\{x_1, \dots, x_n\}$ .

**Remark 2.7.** The Fréchet structure on  $\mathcal{O}_K(\Sigma_\rho)$ , which we call “standard”, is the one of analytic geometry: it coincides with the topology of uniform convergence on rigid discs of radius  $\rho$ . Similarly for  $\mathcal{O}_K^\circ(\Sigma_\rho(K))$ . The standard Fréchet algebra  $K\{x\}$  identifies with

$$K\{x\} = \varprojlim_{\rho \rightarrow +\infty} (\mathcal{O}_K(\Sigma_\rho), \text{standard}).$$

**Definition 2.8.** The *strip topology* on  $\mathcal{H}_K^{(n), \text{bd}}$  is the projective limit topology of the uniform topologies of Definition 2.4. So,

$$(\mathcal{H}_K^{(n), \text{bd}}, \text{strip}) = \varprojlim_{\rho \rightarrow +\infty} (\mathcal{O}_K^{\text{bd}}(\Sigma_\rho(K)), \|\cdot\|_{K, \rho})$$

is a  $K$ -Fréchet space.

**Remark 2.9.** We have a dense embedding  $\mathcal{H}_K^{(n), \text{bd}} \subset K\{x_1, \dots, x_n\}$ . The strip topology on  $\mathcal{H}_K^{(n), \text{bd}}$ , for which this algebra is complete, is finer than its (non complete) standard topology.

The next lemma shows that, for any non archimedean field  $K$  and  $\mathbb{G}_a = \mathbb{G}_{a, K}$ ,

$$\mathcal{O}(\mathbb{G}_a \times \mathbb{G}_a) = \mathcal{O}(\mathbb{G}_a) \widehat{\otimes}_{\pi, K} \mathcal{O}(\mathbb{G}_a),$$

so that  $\mathcal{O}(\mathbb{G}_a)$  is a Hopf algebra object in the category of Fréchet  $K$ -algebras.

**Lemma 2.10.** *There are natural identifications*

$$(10) \quad K\{x_1, \dots, x_n\} \widehat{\otimes}_{\pi, K} K\{y_1, \dots, y_m\} = K\{x_1, \dots, x_n, y_1, \dots, y_m\}.$$

sending  $x_i \otimes 1 \mapsto x_i$  and  $1 \otimes y_j \mapsto y_j$ , for  $i = 1, \dots, n, j = 1, \dots, m$ .



*Proof.* For any  $s \in \mathbb{Z}$ , the map of the statement produces isomorphisms of  $K$ -Tate algebras [11, §6.1.1, Cor. 8]

$$\begin{aligned} K\langle p^{-s}x_1, \dots, p^{-s}x_n \rangle \widehat{\otimes}_{\pi, K} K\langle p^{-s}y_1, \dots, p^{-s}y_m \rangle \\ = K\langle p^{-s}x_1, \dots, p^{-s}x_n, p^{-s}y_1, \dots, p^{-s}y_m \rangle. \end{aligned}$$

We now apply Proposition A.9. □

**Corollary 2.11.** *Let  $K$  be a finite extension of  $\mathbb{Q}_p$  and  $\rho > 0$ . The identifications (10) induce identifications*

$$(\mathcal{H}_K^{(n), \circ}(\rho), \text{standard}) \widehat{\otimes}_{K^\circ}^u (\mathcal{H}_K^{(m), \circ}(\rho), \text{standard}) \xrightarrow{\sim} (\mathcal{H}_K^{(m+n), \circ}(\rho), \text{standard}).$$

*Similarly for  $(\mathcal{O}_K^\circ(\Sigma_\rho(K)), \text{standard})$ .*

**Corollary 2.12.** *The map  $\mathbb{P}: x_i \mapsto x_i \widehat{\otimes} 1 + 1 \widehat{\otimes} x_i$  makes  $K\{x_1, \dots, x_n\}$  into a Hopf algebra object in the category of Fréchet  $K$ -algebras. The restriction of  $\mathbb{P}$  to  $\mathcal{H}_K^{(n), \circ}(\rho)$  induces a map*

$$\mathbb{P}: (\mathcal{H}_K^{(n), \circ}(\rho), \text{standard}) \rightarrow (\mathcal{H}_K^{(n), \circ}(\rho), \text{standard}) \widehat{\otimes}_{K^\circ}^u (\mathcal{H}_K^{(n), \circ}(\rho), \text{standard})$$

*which makes  $(\mathcal{H}_K^{(n), \circ}(\rho), \text{standard})$  a Hopf algebra object in the category of Fréchet  $K^\circ$ -algebras. Similarly for  $(\mathcal{O}_K^\circ(\Sigma_\rho(K)), \text{standard})$ .*

**2.13.  $p$ -adic almost periodic functions.** We sketch here the main ideas and results on  $p$ -adic almost periodic functions. Proofs are given in Section 6. We freely use in this introduction the (quite self-explanatory) notation of Section 5 for continuous, uniformly continuous, bounded rings of  $p$ -adic functions  $\mathbb{Q}_p \rightarrow \mathbb{Q}_p$  and their topologies.

The following elementary lemma shows that a naive  $p$ -adic analog of real Bohr’s uniformly almost periodic functions (see Definition B.3 in Appendix B), where “an interval of length  $\ell_\varepsilon$  in  $\mathbb{R}$ ” is taken to mean a coset  $a + p^h\mathbb{Z}_p$ , for  $a \in \mathbb{Q}_p$  and  $p^{-h} = \ell_\varepsilon$ , does not lead to a meaningful definition.

**Lemma 2.14.** *A continuous function  $f: \mathbb{Q}_p \rightarrow \mathbb{Q}_p$  which has the property that for any  $\varepsilon > 0$ , there exists  $h \in \mathbb{Z}$  such that any coset  $a + p^h\mathbb{Z}_p$  in  $\mathbb{Q}_p/p^h\mathbb{Z}_p$  contains an element  $t_a$  such that*

$$(11) \quad |f(x + t_a) - f(x)| < \varepsilon \quad \text{for all } x \in \mathbb{Q}_p$$

*is constant.*

*Proof.* In fact, from condition (11), for any  $a \in \mathbb{Q}_p$ , it follows by iteration that  $t_a$  may be replaced by any  $t \in \mathbb{Z}t_a$ . By continuity, we may replace  $t_a$  by any  $t \in \mathbb{Z}_p t_a$ . For  $a \notin p^h\mathbb{Z}_p$ ,  $\mathbb{Z}_p t_a = \mathbb{Z}_p a$ . So, if we pick  $a = p^{-N}$ , for  $N \gg 0$ , (11) implies that the variation of  $f(x)$  in  $p^{-N}\mathbb{Z}_p$  is less than  $\varepsilon$ . So, the variation of  $f(x)$  in  $\mathbb{Q}_p$  is less than  $\varepsilon$  for any  $\varepsilon > 0$ , hence  $f$  is constant. □

We resort to an *ad hoc* definition. For  $x \in \mathbb{Q}_p$ , let us consider the classical Witt vector expression

$$(12) \quad x = \sum_{i \gg -\infty}^{\infty} [x_i]p^i \in W(\mathbb{F}_p)[1/p] = \mathbb{Q}_p,$$

where  $[t]$ , for  $t \in \mathbb{F}_p$ , is the Teichmüller representative of  $t$  in  $W(\mathbb{F}_p) = \mathbb{Z}_p$ . Notice that, for any  $i \in \mathbb{Z}$ , the function

$$(13) \quad x_i : \mathbb{Q}_p \rightarrow \mathbb{F}_p, \quad x \mapsto x_i,$$

factors through a function, still denoted by  $x_i$ ,

$$(14) \quad x_i : \mathbb{Q}_p/p^{i+1}\mathbb{Z}_p \rightarrow \mathbb{F}_p, \quad h \mapsto h_i.$$

We regard the function in (14) as an  $\mathbb{F}_p$ -valued *periodic function of period  $p^{i+1}$*  on  $\mathbb{Q}_p$ . In the following, for any  $i \in \mathbb{Z}$  and any  $\lambda \in \mathbb{Q}_p^\times$ , we denote by “ $[(\lambda x)_i]$ ” the uniformly continuous function  $\mathbb{Q}_p \rightarrow \mathbb{Z}_p$ ,  $x \mapsto [(\lambda x)_i]$ . We observe that

$$[(\lambda p^j x)_i] = [(\lambda x)_{i-j}]$$

for any  $i, j \in \mathbb{Z}$  and  $\lambda \in \mathbb{Q}_p^\times$ .

**Definition 2.15.** We define the  $\mathbb{Q}_p$ -algebra  $AP_{\mathbb{Q}_p}$  (resp. the  $\mathbb{Z}_p$ -algebra  $AP_{\mathbb{Z}_p}$ ) of (resp. *integral*) *uniformly almost periodic (u.a.p. for short) functions*  $\mathbb{Q}_p \rightarrow \mathbb{Q}_p$  (resp.  $\mathbb{Q}_p \rightarrow \mathbb{Z}_p$ ) as the closure of

$$\mathbb{Q}_p [ [(\lambda x)_i] \mid i \in \mathbb{Z}, \lambda \in \mathbb{Z}_p^\times ] \quad (\text{resp. of } \mathbb{Z}_p [ [(\lambda x)_i] \mid i \in \mathbb{Z}, \lambda \in \mathbb{Z}_p^\times ])$$

in the  $\mathbb{Q}_p$ -Banach algebra  $\mathcal{C}_{\text{unif}}^{\text{bd}}(\mathbb{Q}_p, \mathbb{Q}_p)$  (resp. in the  $\mathbb{Z}_p$ -Banach ring  $\mathcal{C}_{\text{unif}}(\mathbb{Q}_p, \mathbb{Z}_p)$ ), equipped with the induced valuation  $w_\infty$ .

**Remark 2.16.** This remark is made to partially justify Definition 2.15. For any  $N \in \mathbb{Z}$ , we denote by  $S_N : \mathbb{Q}_p \rightarrow p^N \mathbb{Z}_p$  the function  *$N$ -th order fractional part*, namely,

$$x = \sum_{i \gg -\infty}^{\infty} [x_i] p^i \mapsto S_N(x) = \sum_{i=N}^{\infty} [x_i] p^i.$$

It is clear that, for any  $N$  and  $\lambda \in \mathbb{Q}_p^\times$ ,  $x \mapsto S_N(\lambda x)$  is a bounded uniformly continuous function. The function  $S_3$ , certainly not periodic, is a  $p$ -adic analog of the function

$$\mathbb{R} \rightarrow [0, 1), \quad \dots 1234.56789 \dots \mapsto 0.789 \dots,$$

which is genuinely periodic of period 0.01.

We will prove the following partial analog to Bohr’s “approximation theorem” (Theorem B.4 in Appendix B), where in fact the functions  $\cos(\frac{2\pi}{\lambda}x)$  and  $\sin(\frac{2\pi}{\lambda}x)$ , for  $\lambda \in \mathbb{R}^\times$ , are replaced by the functions  $\Psi(\lambda x)$ , for  $\lambda \in \mathbb{Q}_p^\times$ .

**Theorem 2.17.**  $(AP_{\mathbb{Q}_p}, w_\infty)$  (resp.  $(AP_{\mathbb{Z}_p}, w_\infty)$ ) is the completion of the valued ring

$$(\mathbb{Q}_p[\Psi(\lambda x) \mid \lambda \in \mathbb{Q}_p^\times], w_\infty) \quad (\text{resp. } (\mathbb{Z}_p[\Psi(\lambda x) \mid \lambda \in \mathbb{Q}_p^\times], w_\infty)).$$

**Definition 2.18.** For any  $\lambda \in \mathbb{Q}_p^\times$ , the Fréchet  $\mathbb{Z}_p$ -algebra  $\mathcal{E}_\lambda^\circ$  (resp.  $\mathcal{T}_\lambda^\circ$ ) is the closure of

$$\mathbb{Z}_p[\Psi(\lambda^{-1} p^j x) \mid j = 0, 1, \dots]$$

in  $\mathbb{Q}_p\{x\}$  (resp. in  $\mathcal{O}(\Sigma_{|\lambda|})$ ) with the standard topology. We then set  $\mathcal{E}_\lambda^{\text{bd}} := \mathcal{E}_\lambda^\circ[1/p]$  (resp.  $\mathcal{T}_\lambda^{\text{bd}} := \mathcal{T}_\lambda^\circ[1/p]$ ).

Finally, we define the Fréchet  $\mathbb{Z}_p$ -algebra  $\mathcal{E}^\circ$  as the closure of  $\mathbb{Z}_p[\Psi(\lambda^{-1}p^jx) \mid j = 0, 1, \dots]$  in  $\mathbb{Q}_p\{x\}$ , and set  $\mathcal{E}^{\text{bd}} := \mathcal{E}^\circ[1/p]$ .

**Theorem 2.19** (Approximation theorem on compacts). *The completion of the multivalued ring*

$$(\mathcal{E}^{\text{bd}}, \{\|\cdot\|_{p^r\mathbb{Z}_p}\}_{r \in \mathbb{Z}}) \quad (\text{resp. } (\mathcal{E}^\circ, \{\|\cdot\|_{p^r\mathbb{Z}_p}\}_{r \in \mathbb{Z}}))$$

is the Fréchet  $\mathbb{Q}_p$ -algebra (resp.  $\mathbb{Z}_p$ -algebra)  $\mathcal{C}(\mathbb{Q}_p, \mathbb{Q}_p)$  (resp.  $\mathcal{C}(\mathbb{Q}_p, \mathbb{Z}_p)$ ).

The following proposition follows from the estimates of Proposition 1.3 (see Corollary 4.5 or Theorem 4.14 for the proof) together with the fact that the conditions listed below are closed for the standard Fréchet structure. The proof of the latter fact is given in Lemma 6.6.

**Proposition 2.20.** *For any  $f \in \mathcal{E}_\lambda^\circ$  (resp.  $f \in \mathcal{T}_\lambda^\circ$ ), we have*

- (1)  $f$  is bounded by 1 on the  $p$ -adic strip  $\Sigma_{|\lambda|}$ ;
- (2)  $f(\mathbb{Q}_p) \subset \mathbb{Z}_p$ ;
- (3) For any  $r \in \mathbb{Z}$ ,  $a, j \in \mathbb{Z}_{\geq 0}$ , the function  $g(x) := f(p^{-r}x)^{p^a}$  satisfies

$$g(x + p^{r+j}\lambda\mathbb{C}_p^\circ) \subset g(x) + p^{a+j}\mathbb{C}_p^\circ \quad \text{for all } x \in \mathbb{Q}_p.$$

**Proposition 2.21.** *For any  $\lambda \in \mathbb{Q}_p^\times$ ,  $(\mathcal{E}_\lambda^\circ, \text{standard})$  (resp.  $(\mathcal{T}_\lambda^\circ, \text{standard})$ ) is a Hopf algebra object in the monoidal category  $(\mathcal{C}\mathcal{L}\mathcal{M}_{\mathbb{Z}_p}^u, \widehat{\otimes}_{\mathbb{Z}_p}^u)$  for the coproduct  $\mathbb{P}$  and coidentity  $\varepsilon$  given by*

$$(15) \quad \mathbb{P}(\Psi(\lambda^{-1}p^jx)) \mapsto \Psi(\lambda^{-1}p^jx) \widehat{\otimes} 1 + 1 \widehat{\otimes} \lambda^{-1}p^jx, \quad \varepsilon(\Psi(\lambda^{-1}p^jx)) = 0,$$

for  $j = 0, 1, \dots$ . This structure only depends upon  $|\lambda|$ .

**Definition 2.22.** We define  $\mathbb{E}_\lambda$  in (1) (resp.  $\mathbb{T}_\lambda$ ) as the abelian group functor on  $\mathcal{A}\mathcal{C}\mathcal{L}\mathcal{M}_{\mathbb{Z}_p}^u$ , represented by the Hopf algebras  $(\mathcal{E}_\lambda^\circ, \text{standard})$  (resp. by  $(\mathcal{T}_\lambda^\circ, \text{standard})$ ).

A partial  $p$ -adic analog of Féjer’s theorem, or, more precisely, of Theorem B.2 in Appendix B, is the following proposition.

**Proposition 2.23.** *For any  $\lambda \in \mathbb{Q}_p^\times$ , the completion of the valued ring*

$$(\mathbb{Z}_p[\Psi(\lambda^{-1}p^jx) \mid j = 0, 1, \dots], w_\infty)$$

coincides with its closure in  $\mathcal{C}_{\text{unif}}(\mathbb{Q}_p, \mathbb{Z}_p) = \mathbb{W}(\mathcal{C}_{\text{unif}}(\mathbb{Q}_p, \mathbb{F}_p))$ , equipped with the  $p$ -adic topology, and identifies with  $\mathbb{W}(\mathbb{F}_p[[\lambda^{-1}x]_{-j} \mid j = 0, 1, \dots]])$ , also equipped with the  $p$ -adic topology.

For the standard topology, we have the following result.

**Proposition 2.24.** *For any  $\lambda \in \mathbb{Q}_p^\times$ , the completion of the valued ring  $(\mathcal{E}_\lambda^\circ, w_\infty)$  (resp.  $(\mathcal{T}_\lambda^\circ, w_\infty)$ ) coincides with its closure in  $\mathcal{C}(\mathbb{Q}_p, \mathbb{Z}_p) = \mathbb{W}(\mathcal{C}(\mathbb{Q}_p, \mathbb{F}_p))$ , equipped with the product topology of the prodiscrete topologies on the components (33), and identifies with  $\mathbb{W}(\mathbb{F}_p(v(\lambda), \infty))$  (see Proposition 5.16 below for notation), also equipped with the product topology of the prodiscrete topologies on the components.*

The ring  $\mathcal{E}_\lambda^\circ \otimes_{\mathbb{Z}_p} \mathbb{F}_p$  (resp.  $\mathcal{T}_\lambda^\circ \otimes_{\mathbb{Z}_p} \mathbb{F}_p$ ), equipped with the quotient topology coincides with  $\mathbb{F}_p(v(\lambda), \infty) = \mathcal{C}(\mathbb{Q}_p/\lambda p\mathbb{Z}_p, \mathbb{F}_p)$ .

We now introduce our  $p$ -adic analog of the sheaf  $\mathcal{APH}_{\mathbb{C}}$  of almost periodic analytic functions (see Section B.5 in Appendix B).

**Definition 2.25.**

- (1) For any  $\rho > 0$ , we define the algebra of (resp. *integral*) *almost periodic  $p$ -adic analytic functions on the strip*  $\Sigma_\rho$  as the closure  $\mathcal{APH}_{\mathbb{Q}_p}(\Sigma_\rho)$  (resp.  $\mathcal{APH}_{\mathbb{Z}_p}(\Sigma_\rho)$ ) of  $\mathbb{Q}_p[\Psi(\lambda x) \mid \lambda \in \mathbb{Q}_p^\times]$  (resp.  $\mathbb{Z}_p[\Psi(\lambda x) \mid \lambda \in \mathbb{Q}_p^\times]$ ) in  $(\mathcal{O}^{\text{bd}}(\Sigma_\rho), \text{uniform})$ , with the induced Banach ring structure.
- (2) The algebra of *germs at 0 of almost periodic  $p$ -adic analytic functions* is the locally convex inductive limit

$$(\mathcal{APH}_{0, \mathbb{Q}_p}, \text{strip}) := \varinjlim_{\rho \rightarrow 0} \mathcal{APH}_{\mathbb{Q}_p}(\Sigma_\rho).$$

- (3) The algebra of *germs at 0 of integral almost periodic  $p$ -adic analytic functions* is

$$(\mathcal{APH}_{0, \mathbb{Z}_p}, \text{strip}) := \varinjlim^u_{\rho \rightarrow 0} \mathcal{APH}_{\mathbb{Z}_p}(\Sigma_\rho).$$

- (4) The algebra of *almost periodic  $p$ -adic entire functions* is

$$(\mathcal{APH}_{\mathbb{Q}_p}, \text{strip}) := \varprojlim_{\rho \rightarrow +\infty} \mathcal{APH}_{\mathbb{Q}_p}(\Sigma_\rho).$$

- (5) The algebra of *integral almost periodic  $p$ -adic entire functions* is the closure  $(\mathcal{APH}_{\mathbb{Z}_p}, \text{strip})$  of  $\mathbb{Z}_p[\Psi(\lambda x) \mid \lambda \in \mathbb{Q}_p^\times]$  in  $(\mathcal{APH}_{\mathbb{Q}_p}, \text{strip})$  equipped with the induced Fréchet  $\mathbb{Z}_p$ -algebra structure.
- (6) The Fréchet  $\mathbb{Z}_p$ -algebra  $\mathcal{E}^\circ$  is a Hopf algebra object in the category  $\mathcal{CLM}_{\mathbb{Z}_p}^u$  for the laws (15). The corresponding group functor

$$\mathbb{E}: \mathcal{ACLM}_{\mathbb{Z}_p}^u \rightarrow \mathcal{Ab}$$

will be called the *universal covering* of  $\mathbb{E}_\lambda$ , for any  $\lambda \in \mathbb{Q}_p^\times$ .

**Remark 2.26.** The special fiber of  $\mathbb{E}$  is the constant group

$$\mathbb{Q}_p = \varprojlim_{|\lambda| \rightarrow 0} \mathbb{Q}_p/\lambda\mathbb{Z}_p$$

over  $\mathbb{F}_p$ . On the other hand, equation (2) shows that  $\mathcal{E}^\circ[1/p]$  is dense in  $\mathbb{Q}_p\{x\}$ , so that the generic fiber of  $\mathbb{E}$  is  $\mathbb{G}_{a, \mathbb{Q}_p}$ .

Our Definition 2.25 is designed as to make the analog of Theorem B.7 in Appendix B a true statement. In the  $p$ -adic case, we actually get the following more precise statement.

**Theorem 2.27** (Uniform approximation theorem). *The natural  $\mathcal{CLM}_{\mathbb{Q}_p}^u$ -morphism (resp.  $\mathcal{CLM}_{\mathbb{Z}_p}^u$ -morphism)*

$$(\mathcal{APH}_{0, \mathbb{Q}_p}, \text{strip}) \rightarrow (\mathcal{AP}_{\mathbb{Q}_p}, w_\infty) \quad (\text{resp. } (\mathcal{APH}_{0, \mathbb{Z}_p}, \text{strip}) \rightarrow (\mathcal{AP}_{\mathbb{Z}_p}, w_\infty)),$$

*is an isomorphism.*

The similarity with classical Fourier expansions will be made more stringent in [3], where the classical Mahler binomial expansions of continuous functions  $\mathbb{Z}_p \rightarrow \mathbb{Z}_p$  is generalized to an expansion of any uniformly continuous functions  $\mathbb{Q}_p \rightarrow \mathbb{Q}_p$  as a series with countably many terms of entire functions of exponential type. Such a  $p$ -adic Fourier theory on  $\mathbb{Q}_p$  presents the same power and limitations as the classical Fourier theory on  $\mathbb{R}$ . Functions in  $AP_{\mathbb{Q}_p}$  play the role of Bohr’s uniformly almost periodic functions and a variation of the Bochner–Fejér approximation theorem [9, §I.9] holds. On the other hand, a Fourier series  $\mathcal{F}(f)$  (with countably many terms) does exist for a much more general class of functions  $f: \mathbb{Q}_p \rightarrow \mathbb{Q}_p$  and the classical question as to what extent the series  $\mathcal{F}(f)$  approximates  $f$  makes perfect sense, precisely as in classical Harmonic Analysis.

We ask whether the classical Bohr compactification of  $\mathbb{Q}_p$  has a  $p$ -adic analytic description, as it has one in terms of classical (i.e., complex-valued) harmonic theory on the locally compact group  $(\mathbb{Q}_p, +)$ .

We expect that a completely analogous theory should exist for any finite extension  $K/\mathbb{Q}_p$ . To develop it properly it will be necessary to extend Barsotti covector’s construction to ramified Witt vectors modeled on  $K$  and to relate this construction to Lubin–Tate groups over  $K^\circ$  [19].

### 3. ELEMENTARY PROOFS OF THE MAIN PROPERTIES OF $\Psi$

We prove here the basic properties of the function  $\Psi$ . In contrast to [2], the proofs are here completely self-contained.

**Proposition 3.1.** *The equation (9) has a unique solution in  $\Psi = \Psi_q \in T + T^2\mathbb{Z}[[T]]$ .*

*Proof.* We endow  $\mathbb{Z}[[T]]$  of the  $T$ -adic topology. It is clear that, for any  $\varphi \in T\mathbb{Z}[[T]]$ , the series  $\sum_{j=1}^\infty p^{-j}\varphi(p^jT)^{q^j}$  converges in  $T\mathbb{Z}[[T]]$ . Moreover, the map

$$\mathcal{L}: \varphi \mapsto T - \sum_{j=1}^\infty p^{-j}\varphi(p^jT)^{q^j}$$

is a contraction of the complete metric space  $T + T^2\mathbb{Z}[[T]]$ . In fact, let  $\varepsilon(T) \in T^r\mathbb{Z}[[T]]$ , with  $r \geq 3$ . For any  $\varphi \in T + T^2\mathbb{Z}[[T]]$ , we see that

$$\mathcal{L}(\varphi + \varepsilon) - \mathcal{L}(\varphi) \in T^{r(q-1)+q}\mathbb{Z}[[T]].$$

Since  $r(q - 1) + q > r$ , this shows that  $\mathcal{L}$  is a contraction. So, this map has a unique fixed point which is  $\Psi_q(T)$ . □

The following proposition, due to M. Candilera, provides an alternative proof of Proposition 3.1 and finer information on  $\Psi_q(T)$ .

**Proposition 3.2** (M. Candilera). *The functional equation for the unknown function  $u$*

$$1 = \sum_{j=0}^\infty p^{j(q^j-1)} T^{\frac{q^j-1}{q-1}} u(p^{j(q-1)}T)^{q^j}$$

admits a unique solution  $u(T) = u_q(T) \in 1 + T\mathbb{Z}[[T]]$ . We have

$$\Psi_q(T) = Tu_q(T^{q-1}).$$

*Proof.* In this case we consider the  $T$ -adic metric space  $1 + T\mathbb{Z}[[T]]$  and the map

$$\mathcal{M}: 1 + T\mathbb{Z}[[T]] \rightarrow 1 + T\mathbb{Z}[[T]], \quad \varphi \mapsto 1 - \sum_{j=1}^{\infty} p^{j(q^j-1)} T^{\frac{q^j-1}{q-1}} \varphi(p^{j(q-1)} T)^{q^j}.$$

We endow  $\mathbb{Z}[[T]]$  with the  $T$ -adic topology. It is clear that, for any  $\varphi \in T\mathbb{Z}[[T]]$ , the series  $\sum_{j=1}^{\infty} p^{-j} \varphi(p^j T)^{q^j}$  converges in  $T\mathbb{Z}[[T]]$ . If  $\varepsilon(T) \in T^r\mathbb{Z}[[T]]$ , with  $r \geq 2$ . For any  $\varphi \in 1 + T\mathbb{Z}[[T]]$ , we see that

$$\mathcal{M}(\varphi + \varepsilon) - \mathcal{M}(\varphi) \in T^{r+1}\mathbb{Z}[[T]].$$

So, the map  $\mathcal{M}$  is a contraction and its unique fixed point has the properties stated for the series  $u$  in the statement.  $\square$

**Proposition 3.3.** *The series  $\Psi(T) = \Psi_q(T)$  is entire.*

*Proof.* Since  $\Psi \in T + T^2\mathbb{Z}[[T]] \subset T\mathbb{Z}[[T]]$ , we deduce that  $\Psi$  converges for  $v_p(T) > 0$ . Since the coefficient of  $T$  in  $\Psi(T)$  is 1, whenever  $v_p(T) > 0$ , we have  $v_p(\Psi(T)) = v_p(T)$ .

Suppose  $\Psi$  converges for  $v_p(T) > \rho$ , for  $\rho \leq 0$ . Then, for  $j \geq 1$ ,  $\Psi(p^j T)^{q^j}$  converges for  $v_p(T) > \rho - 1$ . Moreover, if  $j > -\rho + 1$  and  $v_p(T) > \rho - 1$ , we have

$$v_p(p^{-j} \Psi(p^j T)^{q^j}) = -j + q^j(v_p(p^j T)) > -j + q^j(j + \rho - 1),$$

and this last term  $\rightarrow +\infty$ , as  $j \rightarrow +\infty$ .

This shows that the series  $T - \sum_{j=1}^{\infty} p^{-j} \Psi(p^j T)^{q^j}$  converges uniformly for  $v_p(T) > \rho - 1$ , so that its sum, which is  $\Psi$ , is analytic for  $v_p(T) > \rho - 1$ . It follows immediately from this that  $\Psi$  is an entire function.  $\square$

**Remark 3.4.** We have proven that, for any  $j = 0, 1, \dots$  and for  $v_p(T) > -j$ ,

$$v_p(p^{-j} \Psi(p^j T)^{q^j}) = -j + q^j(j + v_p(T)).$$

In particular, for any  $a \in \mathbb{Z}_q$ , ( $\Psi(a) \in \mathbb{Z}_q$  and)  $\Psi_q(a) \equiv a$ , modulo  $p\mathbb{Z}_q$ .

**Proposition 3.5.** *For any  $a \in \mathbb{Q}_q$ ,  $\Psi_q(a) \in \mathbb{Z}_q$ .*

*Proof.* Let  $a \in \mathbb{Z}_q$ . We define by induction the sequence  $\{a_i\}_{i=0,1,\dots}$ :

$$(16) \quad a_0 = a, \quad a_i = \sum_{j=0}^{i-1} p^{j-i} (a_j^{q^{i-j-1}} - a_j^{q^{i-j}}).$$

Since, for any  $a, b \in \mathbb{Z}_q$ , if  $a \equiv b \pmod p$ , then  $a^{q^n} \equiv b^{q^n} \pmod{pq^n}$ , hence modulo  $p^{n+1}$ , while  $a \equiv a^q \pmod p$ , we see that  $a_i \in \mathbb{Z}_q$  for any  $i$ .

We then see by induction that, for any  $i$ ,

$$a_i = p^{-i} \left( a - \sum_{j=0}^{i-1} p^j a_j^{q^{i-j}} \right) \quad \text{or, equivalently, } a = \sum_{j=0}^i p^j a_j^{q^{i-j}}.$$

Explicitly, if we substitute in the formula which defines  $a_i$ , namely,

$$p^i a_i = \sum_{j=0}^{i-1} p^j a_j^{q^{i-j-1}} - \sum_{j=0}^{i-1} p^j a_j^{q^{i-j}},$$

the  $(i - 1)$ -st step of the induction, namely,  $a = \sum_{j=0}^{i-1} p^j a_j^{q^{i-j-1}}$ , we get

$$p^i a_i = a - \sum_{j=0}^{i-1} p^j a_j^{q^{i-j}},$$

which is precisely the  $i$ -th inductive step.

From the functional equation (9) and from Remark 3.4, we have, for  $a \in \mathbb{Z}_q$  and  $i = 0, 1, 2, \dots$ ,

$$\begin{aligned} (17) \quad \Psi(p^{-i}a) &\equiv p^{-i}a - \sum_{\ell=1}^i p^{-\ell} \Psi(p^\ell p^{-i}a)^{q^\ell} \\ &= p^{-i} \left( a - \sum_{j=0}^{i-1} p^j \Psi(p^{-j}a)^{q^{i-j}} \right) \pmod{p\mathbb{Z}_q}. \end{aligned}$$

Notice that  $\Psi(a) \in \mathbb{Z}_q$  and that, modulo  $p\mathbb{Z}_q$ ,  $\Psi_q(a) \equiv a = a_0$ , defined as in (16). We now show by induction on  $i$  that for  $a_1, \dots, a_i, \dots$  defined as in (16),

$$\Psi(p^{-i}a) \equiv a_i \pmod{p\mathbb{Z}_q},$$

which proves the statement. In fact, assume  $\Psi(p^{-j}a) \equiv a_j \pmod{p\mathbb{Z}_q}$ , for  $j = 0, 1, \dots, i - 1$ , and plug this information in (17). We get

$$\Psi(p^{-i}a) \equiv p^{-i}a - \sum_{\ell=1}^i p^{-\ell} a_{i-\ell}^{q^\ell} = p^{-i} \left( a - \sum_{j=0}^{i-1} p^j a_j^{q^{i-j}} \right) = a_i \pmod{p\mathbb{Z}_q},$$

which is the  $i$ -th inductive step. □

**Remark 3.6.** Notice that from (17) it follows that, for any  $a \in p^{-n}\mathbb{Z}_q$ ,

$$a \equiv \sum_{\ell=0}^n p^{-\ell} \Psi_q(p^\ell a)^{q^\ell} \pmod{p\mathbb{Z}_q}.$$

The formula can be more precise using the functional equation (9) and Remark 3.4. We get, for any  $a \in \mathbb{Q}_q$ ,

$$a \equiv \sum_{\ell=0}^{-v_p(a)+i} p^{-\ell} \Psi_q(p^\ell a)^{q^\ell} \pmod{p^{i+1}\mathbb{Z}_q} \quad \text{for all } i \in \mathbb{Z}_{\geq 0},$$

that is,

$$(18) \quad a \equiv \sum_{\ell=0}^i p^{-\ell} \Psi_q(p^\ell a)^{q^\ell} \pmod{p^{i+v_p(a)+1}\mathbb{Z}_q} \quad \text{for all } i \in \mathbb{Z}_{\geq -v_p(a)}.$$

We generalize (12) as follows.

**Corollary 3.7.** For any  $a \in \mathbb{Q}_q$ , let

$$a_i := \Psi_q(p^{-i}a) \bmod p\mathbb{Z}_q \in \mathbb{F}_q.$$

We have

$$a = \sum_{i \gg -\infty}^{\infty} [a_i]p^i \in W(\mathbb{F}_q)[1/p] = \mathbb{Q}_q.$$

*Proof.* Assume first that  $a \in \mathbb{Z}_q$ . In this case (18) implies

$$a \equiv \sum_{\ell=0}^i p^{-\ell} \Psi_q(p^\ell a)^{q^\ell} \bmod p^{i+1}\mathbb{Z}_q \quad \text{for all } i \in \mathbb{Z}_{\geq 0}.$$

So, the statement follows from the following lemma.

**Lemma 3.8.** Let  $i \mapsto b_i$  and  $i \mapsto c_i$ , for  $i = 0, 1, \dots$ , be two sequences in  $\mathbb{Z}_q$  such that

$$\sum_{j=0}^i p^j b_j^{q^{i-j}} \equiv \sum_{j=0}^i p^j c_j^{q^{i-j}} \bmod p^{i+1}\mathbb{Z}_q \quad \text{for all } i \in \mathbb{Z}_{\geq 0}.$$

Then

$$b_i \equiv c_i \bmod p\mathbb{Z}_q \quad \text{for all } i \in \mathbb{Z}_{\geq 0}.$$

*Proof.* Immediate by induction on  $i$ . □

In the general case, assume  $a \in p^{-n}\mathbb{Z}_q$ . Then

$$p^n a = \sum_{i=0}^{\infty} [\Psi_q(p^{n-i}a) \bmod p\mathbb{Z}_q] p^i \in W(\mathbb{F}_q),$$

hence

$$a = \sum_{i=0}^{\infty} [\Psi_q(p^{n-i}a) \bmod p\mathbb{Z}_q] p^{i-n} \in p^{-n}W(\mathbb{F}_q). \quad \square$$

From the previous corollary, it follows that  $a \in \mathbb{Q}_q$  has the following expression as a Witt bivector with coefficients in  $\mathbb{F}_q$ :

$$a = (\dots, a_{-i}^{(q/p)^i}, \dots, a_{-2}^{(q/p)^2}, a_{-1}^{q/p}; a_0, a_1^p, a_1^{p^2}, \dots),$$

which obviously equals  $(\dots, a_{-i}, \dots, a_{-2}, a_{-1}; a_0, a_1, a_1, \dots)$ , if  $q = p$ .

**Remark 3.9.** We have tried to provide a simple addition formula for  $\Psi_q$  of the form (7), in terms of the same power-series  $\Phi$ . We could not get one, nor were we able to establish the relation between  $\Psi_q$  and  $\Psi_p$  for  $q = p^f$  and  $f > 1$ . On the other hand, it is clear that Barsotti’s construction of Witt bivectors, based on classical Witt vectors, extends to the  $L$ -Witt vectors of [19, Ch. 1], where  $L/\mathbb{Q}_p$  denotes any fixed finite extension. In our case, we would only need the construction of *loc. cit.* in the case of the field  $L = \mathbb{Q}_q$ . We believe that the inductive limit of  $\mathbb{Z}_q$ -groups  $W_{\mathbb{Q}_q, n} \rightarrow W_{\mathbb{Q}_q, n+1}$  under Verschiebung

$$V: (x_{-n}, \dots, x_{-1}, x_0) \rightarrow (0, x_{-n}, \dots, x_{-1}, x_0)$$



is a  $\mathbb{Z}_q$ -formal groups whose addition law is expressed by a power-series  $\Phi_q$  analog to Barsotti's  $\Phi$ . We believe that equation (7) still holds true for  $\Psi_q$  if we replace  $\Phi$  by  $\Phi_q$ . We also believe that a generalized  $\Psi$  exists for any finite extension  $L/\mathbb{Q}_p$ , with analogous properties.

4. VALUATION AND NEWTON POLYGONS OF  $\Psi_q$

This section is dedicated to establishing the growth behavior of  $|\Psi_q(x)|$  as  $|x| \rightarrow \infty$ . These results will be essential to get the delicate estimates of [3].

4.1. **Valuation polygon of  $\Psi_q$ .** We recall from [17] that the valuation polygon of a Laurent series  $f = \sum_{i \in \mathbb{Z}} a_i T^i$  with coefficients  $a_i \in \mathbb{C}_p$ , converging in an annulus  $A := \alpha \leq v_p(T) \leq \beta$ , is the graph  $\text{Val}(f)$  of the function  $\mu \mapsto v(f, \mu) := \inf_i (v_p(a_i) + i\mu)$ , which is in fact finite along the segment  $\alpha \leq \mu \leq \beta$ . The function  $\mu \mapsto v(f, \mu)$  is continuous, piecewise affine, and concave on  $[\alpha, \beta]$ . For any  $\mu \in [\alpha, \beta]$ , we have  $v(f, \mu) = \inf \{v_p(\Psi(x)) \mid v_p(x) = \mu\}$ . In the case of  $\Psi$ ,  $A = \mathbb{C}_p$  and the segment  $[\alpha, \beta]$  is the entire  $\mu$ -line. For the convenience of the reader, we have recalled below the relation between the valuation polygon and the Newton polygon of  $f$ .

**Theorem 4.2.** *The valuation polygon of  $\Psi_q$  goes through the origin, has slope 1 for  $\mu > -1$ , and slope  $q^j$ , for  $-j - 1 < \mu < -j$ ,  $j = 1, 2, \dots$  (see Figure 1).*




FIGURE 1. The valuation polygon of  $\Psi_q$ .

*Proof.* We recall that if both  $f$  and  $g$  converge in the annulus  $A := \alpha \leq v_p(T) \leq \beta$ , then, for any  $\mu \in [\alpha, \beta]$ ,  $v(f + g, \mu) \geq \inf(v(f, \mu), v(g, \mu))$ , and that equality holds at  $\mu$  if  $v(f, \mu) \neq v(g, \mu)$ . Moreover, for any  $n \in \mathbb{N}$ ,  $v(f^n, \mu) = n v(f, \mu)$ .

In the polygon in Figure 1, for  $j = 1, 2, \dots$ , the side of projection  $[-j, 1 - j]$  on the  $\mu$ -axis is the graph of the function

$$\sigma_j(\mu) := q^{j-1}(\mu + j - 1) - q^{j-2} - \dots - q - 1.$$

Notice that

$$\sigma_{j+1}(\mu) = -1 + q\sigma_j(\mu + 1),$$

and therefore

$$\sigma_{j+i}(\mu) = -1 - q - \dots - q^{i-1} + q^i\sigma_j(\mu + i),$$

for any  $i = 0, 1, 2, \dots$ .

Since  $\Psi \in T\mathbb{Z}[[T]]$  and since the coefficient of  $T$  is 1, we have  $v(\Psi, \mu) = \mu$  for  $\mu \geq 0$ . For  $0 > \mu > -j$ ,  $j \geq 1$ , we have

$$\begin{aligned} (19) \quad v(p^{-j}\Psi(p^jT)^{q^j}, \mu) &= -j + v(\Psi(p^jT)^{q^j}, \mu) = -j + q^jv(\Psi(p^jT), \mu) \\ &= -j + q^jv(\Psi(S), j + \mu) = -j + (j + \mu)q^j \\ &> \mu = v(T, \mu), \end{aligned}$$

where we have used the variable  $S = p^jT$ .

**Remark 4.3.** For  $\mu = -j$ , we get equality in the previous formula.

Let us set, for  $j = 0, 1, 2, \dots$ ,

$$\ell_j(\mu) = -j + (j + \mu)q^j,$$

so that (19) becomes

$$(20) \quad v(p^{-j}\Psi(p^jT)^{q^j}, \mu) = \ell_j(\mu) > \ell_0(\mu) = \mu = v(T, \mu),$$

for  $0 > \mu > -j$ ,  $j \geq 1$ , with equality holding if  $\mu = -j$ . Notice that

$$\ell_0(\mu) = \mu = \sigma_1(\mu).$$

Because of (20) and (2), and by continuity of  $\mu \mapsto v(\Psi, \mu)$ , we have

$$v(\Psi, \mu) = v(T, \mu) = \mu = \sigma_1(\mu) \quad \text{for } \mu \geq -1.$$

We now reason by induction on  $n = 1, 2, \dots$ . We assume that, for any  $j = 1, 2, \dots, n$ , the side of projection  $[-j, 1 - j]$  on the  $\mu$ -axis of the valuation polygon of  $\Psi$  is the graph of  $\sigma_j(\mu)$ . This at least was proven for  $n = 1$ . We consider the various terms in the functional equation

$$\Psi = T - p^{-1}\Psi(pT)^q - p^{-2}\Psi(p^2T)^{q^2} - \sum_{j=3}^{\infty} p^{-j}\Psi(p^jT)^{q^j}.$$

We assume  $n > 1$ . For  $j = 1, 2, \dots, n$ , and  $-n - 1 < \mu < -n$ , we have

$$\begin{aligned} v(p^{-j}\Psi(p^jT)^{q^j}, \mu) &= -j + v(\Psi(p^jT)^{q^j}, \mu) = -j + q^jv(\Psi(p^jT), \mu) \\ &= -j + q^jv(\Psi(S), j + \mu) = -j + q^j\sigma_{n-j+1}(\mu + j), \end{aligned}$$

since  $j - n - 1 < j + \mu < j - n$ , and therefore the inductive assumption gives  $v(\Psi, j + \mu) = \sigma_{n-j+1}(\mu + j)$  in that interval. For  $j > n$ , and  $-n - 1 < \mu$ , we have instead, from (20),  $v(p^{-j}\Psi(p^jT)^{q^j}, \mu) = \ell_j(\mu)$ .




FIGURE 2. The Newton polygon  $Nw(f)$  of  $f$ .




FIGURE 3. The valuation polygon  $Val(f)$ .

**Lemma 4.4.** *Let  $n > 1$ . For  $j = 1, 2, \dots, n$  and for any  $\mu \in \mathbb{R}$ ,*

$$(21) \quad \sigma_{n+1}(\mu) < -j + q^j \sigma_{n-j+1}(\mu + j).$$

For  $j > n$  and  $\mu > -n - 1$ , we have

$$(22) \quad \sigma_{n+1}(\mu) < \ell_j(\mu).$$

For  $-n - 1 < \mu < -n$ ,

$$(23) \quad \sigma_{n+1}(\mu) < \mu.$$

*Proof.* Assertion (21) is clear, since the two affine functions  $\mu \mapsto \sigma_{n+1}(\mu)$  and  $\mu \mapsto -j + q^j \sigma_{n-j+1}(\mu + j)$  have the same slope  $q^n$ , while their values at  $\mu = -n$  are  $-q^{n-1} - q^{n-2} - \dots - q - 1$  and  $-j - q^{n-1} - q^{n-2} - \dots - q^{j+1} - q^j$ , respectively. Notice that

$$\begin{aligned} -j - q^{n-1} - q^{n-2} - \dots - q^{j+1} - q^j &= (q^j - j) - q^{n-1} - q^{n-2} - \dots - p - 1 \\ &> -q^{n-1} - q^{n-2} - \dots - q - 1, \end{aligned}$$

so that the conclusion follows.

We examine assertion (22), namely, that for  $j > n$  and  $\mu > -n - 1$ , we have

$$q^n(\mu + n) - q^{n-1} - q^{n-2} - \dots - q - 1 < -j + (j + \mu)q^j.$$

The previous inequality translates into

$$q^n(\mu + n) - q^{n-1} - q^{n-2} - \dots - q - 1 < -j + (j - n)q^j + (n + \mu)q^{n+(j-n)},$$

that is,

$$q^{n-1} + q^{n-2} + \dots + q + 1 - j + (j - n)q^j + (n + \mu)q^n(q^{j-n} - 1) > 0,$$

for  $\mu > -n - 1$ . Since the left-hand side is an increasing function of  $\mu$ , it suffices to show that the inequality holds for  $\mu = -n - 1$ , that is to prove that

$$(24) \quad q^{n-1} + q^{n-2} + \dots + q + 1 - j + (j - n)q^j - q^n(q^{j-n} - 1) > 0,$$

for any  $j > n > 1$ . We rewrite the left-hand side of (24) as

$$\begin{aligned} q^{n-1} + q^{n-2} + \dots + q + 1 - n + (n - j) + (j - n)q^j - q^j + q^n \\ = (q^{n-1} + q^{n-2} + \dots + q + 1 - n) + (q^j - 1)(j - n) + (q^n - q^j), \end{aligned}$$

where the four terms in round brackets on the right-hand side are each, obviously, positive numbers. The conclusion follows.

We finally show (23), namely that for  $-n - 1 < \mu < -n$ ,

$$q^n(\mu + n) - q^{n-1} - q^{n-2} - \dots - p - 1 < \mu.$$

It suffices to compare the values at  $\mu = -n - 1$  and at  $\mu = -n$ . We get

$$-q^n - q^{n-1} - q^{n-2} - \dots - q - 1 < -n - 1$$

and

$$-q^{n-1} - q^{n-2} - \dots - q - 1 < -n,$$

respectively, both obviously true. □

The previous calculation shows that the side of projection  $[-n - 1, -n]$  on the  $\mu$ -axis of the valuation polygon of  $\Psi$  is the graph of  $\sigma_{n+1}(\mu)$ . We have then crossed the inductive step Case  $n \Rightarrow$  Case  $n + 1$ , and Theorem 4.2 is proven. □

**Corollary 4.5.** *Proposition 1.3 holds true.*

*Proof.* We have seen that  $v_p(\Psi_p(x)) = v_p(x)$  if  $v_p(x) > 0$ . Then Proposition 1.3 follows from (7) and Lemma 1.5.  $\square$

**Corollary 4.6.** *For any  $i = 1, 2, \dots$ , and  $v_p(x) \geq -i$  (resp.  $v_p(x) > -i$ ), we have  $v_p(\Psi_q(x)) \geq -\frac{q^i-1}{q-1}$  (resp.  $v_p(\Psi_q(x)) > -\frac{q^i-1}{q-1}$ ). If  $v_p(x) > -1$ , we have  $v_p(\Psi_q(x)) = v_p(x)$ .*

*Proof.* The last part of the statement is a general fact for automorphisms of an open  $k$ -analytic disk  $D$  with one  $k$ -rational fixed point  $a \in D(k)$  (the disk  $v_p(x) > -1$  and  $x(a) = 0$ , in the present case) [8, Lemma 6.4.4].  $\square$

**4.7. Newton polygon of  $\Psi_q$ .** We now recall that to a Laurent series  $f = \sum_{i \in \mathbb{Z}} a_i T^i$  with coefficients  $a_i \in \mathbb{C}_p$ , converging in an annulus  $A := \alpha \leq v_p(T) \leq \beta$ , one associates two, dually related, polygons. The valuation polygon  $\mu \mapsto v(f, \mu)$  was recalled before. The *Newton polygon*  $\text{Nw}(f)$  of  $f$  is the convex closure in the standard affine plane  $\mathbb{R}^2$  of the points  $(-i, v(a_i))$  and  $(0, +\infty)$ . If  $a_i = 0$ , then  $v(a_i)$  is understood as  $+\infty$ . We define  $s \mapsto \text{Nw}(f, s)$  to be the function whose graph is the lower-boundary of  $\text{Nw}(f)$ . The main property of  $\text{Nw}(f)$  is that the length of the projection on the  $X$ -axis of the side of slope  $\sigma$  is the number of zeros of  $f$  of valuation  $= \sigma$ . The formula

$$v(f, \mu) = \inf_{i \in \mathbb{Z}} i \mu + v(a_i)$$

indicates (cf. [17]) that the relation between  $\text{Nw}(f)$  and  $\text{Val}(f)$  “almost” coincides with the duality formally described in the following lemma.

**Lemma 4.8** (Duality of polygons). *In the projective plane  $\mathbb{P}^2$ , with affine coordinates  $(X, Y)$ , we consider the polarity with respect to the parabola  $X^2 = -2Y$*

$$\mathbb{P}^2 \rightarrow (\mathbb{P}^2)^* \rightarrow \mathbb{P}^2, \quad \text{point}(\sigma, \tau) \mapsto \text{line}(Y = -\sigma X - \tau) \mapsto \text{point}(\sigma, \tau).$$

*Assume the graph  $\Gamma$  of a continuous convex piecewise affine function has consecutive vertices*

$$\dots, (-i_0, \varphi_0), (-i_1, \varphi_1), (-i_2, \varphi_2), (-i_3, \varphi_3), \dots,$$

*joined by the lines*

$$\dots, Y = \sigma_1 X + \tau_1, Y = \sigma_2 X + \tau_2, Y = \sigma_3 X + \tau_3, \dots$$

*Then the lines joining the points*

$$\dots, (-\sigma_1, -\tau_1), (-\sigma_2, -\tau_2), (-\sigma_3, -\tau_3), \dots$$

*are*

$$\dots, Y = i_1 X - \varphi_1, Y = i_2 X - \varphi_2, \dots,$$

*and the polarity transforms these back into*

$$\dots, (-i_1, \varphi_1), (-i_2, \varphi_2), \dots$$

We say that the graph  $\Gamma^*$  joining the vertices  $(\sigma_i, \tau_i), (\sigma_{i+1}, \tau_{i+1})$  by a straight segment is the dual graph of  $\Gamma$ . It is clear that the relation is reciprocal, that is,  $(\Gamma^*)^* = \Gamma$  and that  $\Gamma^*$  is a continuous concave piecewise affine function.

*Proof.* It is the magic of polarities. □

The precise relation between  $\text{Nw}(f)$  and  $\text{Val}(f)$  follows.

**Proposition 4.9.** *We have*

$$\text{Val}(f) = (-\text{Nw}(f))^*,$$

where  $-\text{Nw}(f)$  is the polygon obtained from  $\text{Nw}(f)$  by the transformation  $(X, Y) \mapsto (X, -Y)$ .

*Proof.* The most convincing proof follows from comparing Lemma 4.8 with Figures 2 and 3. □

We now apply the previous considerations to the two polygons associated to the function  $\Psi_q$ .

**Corollary 4.10.** *The Newton polygon  $\text{Nw}(\Psi_q)$  has vertices at the points*

$$V_i := \left( -q^i, iq^i - \frac{q^i - 1}{q - 1} \right) = (-q^i, iq^i - q^{i-1} - \dots - q - 1).$$

The equation of the side joining the vertices  $V_i$  and  $V_{i-1}$  is

$$Y = -iX - \frac{q^i - 1}{q - 1};$$

its projection on the  $X$ -axis is the segment  $[-q^i, -q^{i-1}]$ . So,  $\text{Nw}(\Psi)$  has the form described in Figure 4.

**Corollary 4.11.** *For any  $i = 0, 1, \dots$ , the map  $\Psi = \Psi_q$  induces coverings of degree  $q^i$ ,*

$$(25) \quad \Psi: \{x \in \mathbb{C}_p \mid v_p(x) > -i - 1\} \rightarrow \left\{ x \in \mathbb{C}_p \mid v_p(x) > -\frac{q^{i+1} - 1}{q - 1} \right\}$$

(in particular, an isomorphism

$$(26) \quad \Psi: \{x \in \mathbb{C}_p \mid v_p(x) > -1\} \xrightarrow{\sim} \{x \in \mathbb{C}_p \mid v_p(x) > -1\},$$

for  $i = 0$ ), finite maps of degree  $q^i$ ,

$$\begin{aligned} \Psi: \{x \in \mathbb{C}_p \mid -(i + 1) < v_p(x) < -i\} \\ \rightarrow \left\{ x \in \mathbb{C}_p \mid -\frac{q^{i+1} - 1}{q - 1} < v_p(x) < -\frac{q^i - 1}{q - 1} \right\}, \end{aligned}$$

and finite maps of degree  $q^{i+1} - q^i$ ,

$$\Psi: \{x \in \mathbb{C}_p \mid v_p(x) = -i - 1\} \rightarrow \left\{ x \in \mathbb{C}_p \mid -\frac{q^{i+1} - 1}{q - 1} \leq v_p(x) \right\}.$$




FIGURE 4. The Newton polygon  $Nw(\Psi_q)$  of  $\Psi_q$ .

*Proof.* The shape of the Newton polygon of  $\Psi$  indicates that, for any  $a \in \mathbb{C}_p$ , with  $v_p(a) > -1$ , the side of slope  $= v_p(a)$  of the Newton polygon of  $\Psi - a$  has projection of length 1 on the  $X$ -axis. So,  $\Psi: \{x \in \mathbb{C}_p \mid v_p(x) > -1\} \rightarrow \{x \in \mathbb{C}_p \mid v_p(x) > -1\}$  is bijective, hence biholomorphic. Now we recall from Corollary 4.6 that for any given  $i \geq 1$ ,

$$\Psi(\{x \in \mathbb{C}_p \mid v_p(x) > -i - 1\}) \subset \left\{x \in \mathbb{C}_p \mid v_p(x) > -\frac{q^{i+1} - 1}{q - 1}\right\}.$$

So, let  $a$  be such that  $-\frac{q^{i+1}-1}{q-1} < v_p(a) \leq -\frac{q^i-1}{q-1}$ , say  $v_p(a) = -\frac{q^i-1}{q-1} - \varepsilon$ , with  $\varepsilon \in [0, q^i)$ . Then, the Newton polygon of  $\Psi - a$  has a single side of slope  $> -i - 1$ , which has precisely slope  $= -\varepsilon q^{-i} - i$  and has projection of length  $q^i$  on the  $X$ -axis. So, the equation  $\Psi(x) = a$  has precisely  $q^i$  solutions  $x$  in the annulus  $-i - 1 < v_p(x) \leq -i$ . If, for the same  $i$ ,  $-\frac{q^i-1}{q-1} < v_p(a) \leq -\frac{q^{i-1}-1}{q-1}$ , the Newton polygon of  $\Psi - a$  has a side of slope  $-i$ , whose projection on the  $X$ -axis has length  $q^i - q^{i-1}$ , and a side of slope  $\sigma$ ,  $1 - i \geq \sigma > -i$ , whose projection on the  $X$ -axis has length  $q^{i-1}$ . So, again  $\Psi^{-1}(a)$  consists of  $q^i$  distinct points. We go on, for  $a$  in an annulus of the form  $-\frac{q^{i-j}-1}{q-1} < v_p(a) \leq -\frac{q^{i-j-1}-1}{q-1}$ , up to  $j = i - 2$ , i.e., to  $-\frac{q^2-1}{q-1} < v_p(a) \leq -1$ . In that case, the Newton polygon of  $\Psi - a$  has a side of slope  $-i$  of projection  $q^i - q^{i-1}$ , a side of slope  $1 - i$  of projection  $q^{i-1} - q^{i-2}$ , ..., a side of slope  $j - i$  of projection  $q^{i-j} - q^{i-j-1}$  on the  $X$ -axis, ..., up to a side of slope  $-1$  of projection  $q - 1$  on the  $X$ -axis. Finally, for  $v_p(a) > -1$ , there is still exactly one solution of  $\Psi(x) = a$ , with

$v_p(x) > -1$ . This means that  $\Psi$  induces a (ramified) covering of degree  $q^i$  in (25).  $\square$

**4.12. The addition law of  $\Psi$ .** We now extend the estimates of Corollary 4.11 to translates  $\Psi(a + x)$  of  $\Psi$ , for  $a \in \mathbb{Q}_p$ . Although we expect that the same discussion carries over to  $\Psi_q(a + x)$ , where  $a \in \mathbb{Q}_q$ , we assume for simplicity that  $q = p$  in the rest of this subsection.

**Proposition 4.13.** *Let  $m \in \mathbb{Z}_{>0}$  and let  $M(x_{-m} \dots, x_{-1}, x_0)$  be a monomial in  $\mathbb{Z}_p[x_{-m} \dots, x_{-1}, x_0]$  divisible by  $x_{-m}$  and of pure weight 1, where  $x_i$  weighs  $p^i$  for any  $i$ . Set*

$$M(x) := M(\Psi(p^m x), \dots, \Psi(x)).$$

Then, for any  $r = 1, 2, \dots$ ,

$$\begin{aligned} w_r(M(x)) &\geq m + 1 - (p - 1)r(m - r + 1) \\ &\quad + (p - 1) \left( \binom{m + 1}{2} - \binom{r}{2} \right) - \frac{p^{r+1} - 1}{p - 1} \\ &> m + 1 + (p - 1) \frac{(m - r)^2 + (m - r)}{2} - \frac{p^{r+1} - 1}{p - 1} \left( > -\frac{p^{r+1} - 1}{p - 1} \right), \end{aligned}$$

while, for  $r = 0, -1, -2, \dots$ , we get

$$w_r(M(x)) \geq m - r - (p - 1)mr + (p - 1) \binom{m + 1}{2} (\geq p(1 - r)).$$

*Proof.* This follows from the estimates of Corollary 4.11 via a totally self-contained, but lengthy, computation on isobaric polynomials of Witt-type. We refer to the upcoming paper [3] for the proof of a more general statement.  $\square$

We apply Proposition 4.13 to the study of the addition law of  $\Psi$ . From (7) and (8), we deduce, taking into account Proposition 3.5, that, for any  $c \in \mathbb{Q}_p$ ,

$$(27) \quad \Psi(x + c) = \lim_{i \rightarrow \infty} \varphi_i(\Psi(p^i x), \dots, \Psi(px), \Psi(x); \Psi(p^i c), \dots, \Psi(pc), \Psi(c)),$$

where

$$\begin{aligned} &\varphi_m(\Psi(p^m x), \dots, \Psi(px), \Psi(x); \Psi(p^m c), \dots, \Psi(pc), \Psi(c)) \\ &\quad - \varphi_{m-1}(\Psi(p^{m-1} x), \dots, \Psi(px), \Psi(x); \Psi(p^{m-1} c), \dots, \Psi(pc), \Psi(c)) \end{aligned}$$

is a sum of monomials  $M(x)$  as in Proposition 4.13.

**Theorem 4.14.**

- (1) *The function  $\Psi$  is bounded and uniformly continuous on any  $p$ -adic strip around  $\mathbb{Q}_p$ . In particular,*

$$\Psi(x) \in \mathcal{H}^{\text{bd}}.$$

- (2) *For any  $j = 0, 1, \dots$  and  $x \in \mathbb{Q}_p$ ,*

$$(28) \quad \Psi_p(x + p^j \mathbb{C}_p^\circ) \subset \Psi_p(x) + p^j \mathbb{C}_p^\circ.$$



*Proof.* For the first part of the statement, we observe that Proposition 4.13 shows that for any fixed  $r \geq 0$ , the sequence

$$i \mapsto \varphi_i(\Psi(p^i x), \dots, \Psi(px), \Psi(x); \Psi(p^i c), \dots, \Psi(pc), \Psi(c))$$

converges in the  $w_r$ -valuation. This means that for any  $\rho > 0$ , the previous sequence is a sequence of entire functions bounded on the  $p$ -adic strip  $\Sigma_\rho$  around  $\mathbb{Q}_p$ , which converges to  $\Psi(x + c)$  uniformly on  $\Sigma_\rho$ .

The second part of the statement was already proved in Corollary 4.5. It also follows from the estimates of Proposition 4.13 when  $r \leq 0$ .  $\square$

**4.15. The zeros of  $\Psi$ .** The following theorem is formulated in a way to make sense for  $q =$  any power of  $p$ . We expect that it is true in that generality. However, for the time being, we can only prove it for  $q = p$ .

**Theorem 4.16.** *In this statement, let  $q = p$ .*

- (1) *For any  $n = 1, 2, \dots$ , the map  $\Psi_q$  has  $q_n := q^n - q^{n-1}$  simple zeros of valuation  $-n$  in  $\mathbb{Q}_q$ . More precisely, for any system of representatives  $a_1, \dots, a_{q_n} \in \mathbb{Z}_q$  of  $(\mathbb{Z}_q/p^n\mathbb{Z}_q)^\times = W_{n-1}(\mathbb{F}_q)^\times$ , and any  $j = 1, \dots, q_n$ , the open disc  $D(a_j p^{-n}, p^-)$  contains a unique zero  $z_j^{(n)} \in \mathbb{Q}_q$  of  $\Psi_q$ . Then  $z_1^{(n)}, \dots, z_{q_n}^{(n)}$  are all the zeros of  $\Psi_q$  of valuation  $-n$ .*
- (2) *For  $n = 1, 2, \dots$  let  $z_1^{(n)}, \dots, z_{q_n}^{(n)}$  be the zeros of  $\Psi_q$  of valuation  $-n$ . We set*

$$\psi_n(x) = \prod_{j=1}^{q_n} \left( 1 - \frac{x}{z_j^{(n)}} \right) \in 1 + p^n x \mathbb{Z}_q[x].$$

*Then*

$$\Psi_q(x) = x \prod_{n=1}^{\infty} \psi_n(x)$$

*is the canonical convergent infinite Schnirelmann product expression [17, (4.13)] of  $\Psi_q(x)$  in the ring  $\mathbb{Q}_p\{x\}$ .*

- (3) *The inverse function  $\beta(T) = \beta_q(T)$  of  $\Psi_q(T)$  (i.e., the power series such that, in  $T\mathbb{Z}[[T]]$ ,  $\Psi_q(\beta_q(T)) = T = \beta_q(\Psi_q(T))$ ) belongs to  $T + T^2\mathbb{Z}[[T]]$ . Its disc of convergence is exactly  $v_p(T) > -1$ .*

*Proof.* We now prove the first statement in Theorem 4.16. We recall that here  $q = p$ , so that  $a_1, \dots, a_{p_n} \in \mathbb{Z}_p$ , with  $p_n = p^n - p^{n-1}$ , are a system of representatives of  $(\mathbb{Z}_p/p^n\mathbb{Z}_p)^\times$ .

**Lemma 4.17.** *For any  $m, n \in \mathbb{Z}_{>0}$ , with  $m \leq n$ , and any  $j = 1, \dots, p_n$ , the value of  $\Psi$  at the maximal point  $\xi_{a_j p^{-n}, p^m}$  (of Berkovich type 2) of the rigid disc  $D(a_j p^{-n}, (p^m)^+)$ , that is,  $-\log|\Psi(\xi_{a_j p^{-n}, p^m})| = w_m(\Psi(a_j p^{-n} + x))$ , is  $-\frac{p^m - 1}{p - 1} < 0$ .*

*Proof.* The proof follows from the addition law (27) in which

$$\Psi(a_j p^{-n}), \dots, \Psi(a_j p^{i-n}) \in \mathbb{Z}_p,$$

so that, for any  $i = 0, 1, 2, \dots$ ,

$$\varphi_i(\Psi(p^i x), \dots, \Psi(px), \Psi(x); \Psi(a_j p^{i-n}), \dots, \Psi(a_j p^{1-n}), \Psi(a_j p^{-n}))$$

is a sum of a dominant (at  $\xi_{a_j p^{-n}, p^m}$ ) term

$$\Psi(x) + \Psi(a_j p^{-n})$$

and of terms  $M(x)$  described, for  $m = n - i$ , in Proposition 4.13. □

From the harmonicity of the function  $|\Psi(x)|$  at the point  $\xi_{a_j p^{-n}, p^m}$ , the estimate of Lemma 4.17, and the fact that  $\Psi(a_j p^{-n}) \in \mathbb{Z}_p$ , we deduce that each of the  $p_n p^{m-n}$  open discs of radius  $p^m$ , centered at points of  $p^{-n} \mathbb{Z}_p \setminus p^{1-n} \mathbb{Z}_p$  contains at least one zero of  $\Psi_q$  in  $\mathbb{Q}_q$ . For  $m = 1$ , this proves the first part of the statement.

For the second part of the statement, we refer to [17, §4]. The fact that every  $\psi_n(x) \in \mathbb{Q}_p[x]$  is  $-n$ -extremal follows from the fact that its zeros are all of exact valuation  $-n$  [17, (2.7')].

The fact that  $\beta_p$  belongs to  $T + T^2 \mathbb{Z}[[T]]$  is obvious. The convergence of  $\beta_q$  for  $v_p(T) > -1$  follows from (26). The fact that it cannot converge in a bigger disk is a consequence of the fact that  $\Psi_q$  has  $q - 1$  zeros of valuation  $-1$ . □

**Corollary 4.18.** *All zeros of  $\Psi_q$  are simple and are contained in  $\mathbb{Q}_q$ . Each ball  $a + \mathbb{Z}_q \in \mathbb{Q}_q/\mathbb{Z}_q$  contains a single zero of  $\Psi_q$ .*

**Remark 4.19.** We believe that Theorem 4.16 holds, with essentially the same proof, for any power  $q$  of  $p$ . See Remark 3.9.

### 5. RINGS OF CONTINUOUS FUNCTIONS ON $\mathbb{Q}_p$

The point of this section is that of establishing the categorical limit/colimit formulas for the linear topologies of rings of  $p$ -adic functions on  $\mathbb{Q}_p$ . For topological algebra notions, we take the viewpoint and use the definitions explained in Appendix A (see also [4]).

We consider here a linearly topologized separated and complete ring  $k$ , whose family of open ideals we denote by  $\mathcal{P}(k)$ . In practice,  $k = \mathbb{Z}_p$  or  $= \mathbb{F}_p$ , or  $= \mathbb{Z}_p/p^r \mathbb{Z}_p$ , for any  $r \in \mathbb{Z}_{\geq 1}$ . More generally,  $A$  will be a complete and separated topological ring equipped with a  $\mathbb{Z}$ -linear topology, defined by a family of open additive subgroups of  $A$ . In particular, we have in mind  $A =$  a fixed finite extension  $K$  of  $\mathbb{Q}_p$ , whose topology is  $K^\circ$ -linear but not  $K$ -linear. Again, a possible  $k$  would be  $K^\circ$  or any  $K^\circ/(\pi_K)^r$ , for a parameter  $\pi = \pi_K$  of  $K$ , and  $r$  as before.

We will express our statements for an abelian topological group  $G$ , which is separated and complete in the  $\mathbb{Z}$ -linear topology defined by a countable family of profinite subgroups  $G_r$ , with  $G_r \supset G_{r+1}$ , for any  $r \in \mathbb{Z}$ . So,

$$G = \varprojlim_{r \rightarrow +\infty} G/G_r = \varinjlim_{r \rightarrow -\infty} G_r,$$

where  $G/G_r$  is discrete,  $G_r$  is compact, and limits and colimits are taken in the category of topological abelian groups separated and complete in a  $\mathbb{Z}$ -linear topology. We denote by  $\pi_r: G \rightarrow G/G_r$  the canonical projection. Then  $G$  is canonically a uniform space in which a function  $f: G \rightarrow A$  is uniformly continuous if and only if for any open subgroup  $J \subset A$ , the induced function  $G \rightarrow A/J$  factors via a  $\pi_r$  for some  $r = r(J)$ . A subset of  $G$  of the form  $\pi_r^{-1}(\{h\}) = g + G_r$ , for  $g \in G$  and  $h = \pi_r(g)$ , is sometimes called the *ball of radius  $G_r$  and center  $g$* . In particular,  $G$  is a locally compact, paracompact, 0-dimensional topological space. A general discussion of the duality between  $k$ -valued functions and measures on such a space, will appear in [5]. In practice, here  $G = \mathbb{Q}_p$  or  $\mathbb{Q}_p/p^r\mathbb{Z}_p$  or  $p^r\mathbb{Z}_p$ , with the obvious uniform and topological structure.

**Definition 5.1.** Let  $G$  and  $A$  be as before. Then we define  $\mathcal{C}(G, A)$  (resp.  $\mathcal{C}_{\text{unif}}^{\text{bd}}(G, A)$ ) as the  $A$ -algebra of continuous (resp. bounded and uniformly continuous) functions  $f: G \rightarrow A$ . We equip  $\mathcal{C}(G, A)$  (resp.  $\mathcal{C}_{\text{unif}}^{\text{bd}}(G, A)$ ) with the topology of uniform convergence on compact subsets of  $X$  (resp. on  $X$ ). For any  $r \in \mathbb{Z}$  and  $g \in G$ , we denote by  $\chi_{g+G_r}$  the characteristic function of  $g + G_r \in G/G_r$ . If  $G$  is discrete and  $h \in G$ , by  $e_h: G \rightarrow k$  we denote the function such that  $e_h(h) = 1$ , while  $e_h(x) = 0$  for any  $x \neq h$  in  $G$ .

**Remark 5.2.** It is clear that if  $A = k$  is a linearly topologized ring, any subset of  $k$  and therefore any function  $f: G \rightarrow k$  is bounded. So, we write  $\mathcal{C}_{\text{unif}}(G, k)$  instead of  $\mathcal{C}_{\text{unif}}^{\text{bd}}(G, k)$  in this case. If  $G$  is discrete, any function  $G \rightarrow k$  is (uniformly) continuous; still, the bijective map  $\mathcal{C}_{\text{unif}}(G, k) \rightarrow \mathcal{C}(G, k)$  is not an isomorphism in general, so we do keep the difference in notation. If  $G$  is compact, any continuous function  $G \rightarrow k$  is uniformly continuous, and  $\mathcal{C}_{\text{unif}}(G, k) \rightarrow \mathcal{C}(G, k)$  is an isomorphism, so there is no need to make any distinction.

**Lemma 5.3.** *With the notation as above, assume that  $G$  is discrete (so that the  $G_r$ 's are finite). Then  $\mathcal{C}(G, k)$  (resp.  $\mathcal{C}_{\text{unif}}(G, k)$ ) is the  $k$ -module of functions  $f: G \rightarrow k$  endowed with the topology of simple (resp. of uniform) convergence on  $G$ . So,*

$$\mathcal{C}(G, k) = \varprojlim_{r \rightarrow -\infty} \mathcal{C}(G_r, k) = \prod_h k e_h, \quad h \in G.$$

Similarly,

$$\mathcal{C}_{\text{unif}}(G, k) = \varprojlim_{I \in \mathcal{P}(k)} \prod_{h \in G} \square, u(k/I) e_h = \prod_{h \in G} \square, u k e_h,$$

where  $\prod_{h \in G} \square, u(k/I) e_h$  carries the discrete topology.

*Proof.* It is clear from the definitions. □

The next lemma is a simplified abstract form, in the framework of linearly topologized rings and modules, of the classical decomposition of a continuous function as a sum of characteristic functions of balls (see, for example, [13, §1.3.1]).

**Lemma 5.4.** *With the notation as above, assume that  $G$  is compact (so that the  $G/G_r$ 's are finite). Then*

$$\mathcal{C}(G, k) = \mathcal{C}_{\text{unif}}(G, k) = \varinjlim_{r \rightarrow +\infty}^u \mathcal{C}(G/G_r, k) = \varinjlim_{r \rightarrow +\infty}^u \bigoplus_{g+G_r \in G/G_r} k\chi_{g+G_r}.$$

For any  $r$ , the canonical morphism  $\mathcal{C}(G/G_r, k) \rightarrow \mathcal{C}(G, k)$  is injective.

*Proof.* This is also clear from the definitions. □

**Remark 5.5.** We observe that the inductive limit appearing in the formula hides the complication of formulas of the type

$$\chi_{g+G_r} = \sum_i \chi_{g_i+G_{r+1}} \quad \text{if } g + G_r = \bigcup_i g_i + G_{r+1},$$

which we do not need to make explicit for the present use (see [5] for a detailed discussion).

**Proposition 5.6.** *With the notation as above and  $G$  general, in the category  $\mathcal{CLM}_k^u$ , the following hold:*

(1) *We have*

$$\mathcal{C}(G, k) = \varprojlim_{r \rightarrow -\infty} \mathcal{C}(G_r, k)$$

for the restrictions  $\mathcal{C}(G_r, k) \rightarrow \mathcal{C}(G_{r+1}, k)$ . In particular, for any fixed  $r \in \mathbb{Z}$ ,

$$\mathcal{C}(G, k) = \prod_{g+G_r \in G/G_r} \mathcal{C}(g + G_r, k).$$

(2) *We have*

$$\mathcal{C}_{\text{unif}}(G, k) = \varinjlim_{r \rightarrow +\infty}^u \mathcal{C}_{\text{unif}}(G/G_r, k)$$

for the embeddings

$$\mathcal{C}_{\text{unif}}(G/G_r, k) \hookrightarrow \mathcal{C}_{\text{unif}}(G/G_{r+1}, k).$$

(3) *The natural morphism*

$$\mathcal{C}_{\text{unif}}(G, k) \rightarrow \mathcal{C}(G, k)$$

*is injective and has dense image.*

*Proof.* The first two parts follow from the universal properties of limits and colimits. The morphism in part (3) comes from the injective morphisms, for  $r \in \mathbb{Z}$ ,

$$\mathcal{C}_{\text{unif}}(G/G_r, k) \rightarrow \mathcal{C}(G, k)$$

and the universal property of colimits. The inductive limit of these morphisms in the category  $\mathcal{CLM}_k^u$  is a completion of the inductive limit taken in the category  $\text{Mod}_k$  of  $k$ -modules equipped with the  $k$ -linear inductive limit topology. Since the latter is separated and since the axiom AB5 holds for the abelian category  $\text{Mod}_k$ , we deduce that the morphism in part (3) is injective. The

morphism has dense image because, for any  $r \in \mathbb{Z}$  and for any  $s \in \mathbb{Z}_{\geq 0}$ , the composed morphism

$$\mathcal{C}_{\text{unif}}(G_r/G_{r+s}, k) \rightarrow \mathcal{C}_{\text{unif}}(G/G_{r+s}, k) \rightarrow \mathcal{C}(G, k) \rightarrow \mathcal{C}(G_r, k)$$

is the canonical map of Lemma 5.4

$$\mathcal{C}_{\text{unif}}(G_r/G_{r+s}, k) \rightarrow \mathcal{C}(G_r, k)$$

for the compact group  $G_r$  and its subgroup  $G_{r+s}$ . The fact that the set theoretic union  $\bigcup_{s \geq 0} \mathcal{C}_{\text{unif}}(G_r/G_{r+s}, k)$  is dense in  $\mathcal{C}(G_r, k)$  is built-in in the definition of  $\varinjlim^u$ .  $\square$

**Proposition 5.7.** *Let  $(H, \{H_r\}_r)$  be a locally compact group with the same properties as  $(G, \{G_r\}_r)$  above, so that  $(G \times H, \{G_r \times H_r\}_r)$  also has the same properties. Then we have a natural identification in  $\mathcal{CLM}_k^u$*

$$(29) \quad \mathcal{C}(G, k) \widehat{\otimes}_k^u \mathcal{C}(H, k) \xrightarrow{\sim} \mathcal{C}(G \times H, k)$$

and a continuous strictly closed embedding

$$(30) \quad \mathcal{C}_{\text{unif}}(G, k) \widehat{\otimes}_k^u \mathcal{C}_{\text{unif}}(H, k) \rightarrow \mathcal{C}_{\text{unif}}(G \times H, k).$$

*Proof.* We prove (29) first. By the first part of Proposition 5.6(1) and the fact that  $\widehat{\otimes}_k^u$  commutes with projective limits, we are reduced to the case of  $G$  and  $H$  being compact. We are then in the situation of Lemma 5.4 for both  $G$  and  $H$  (in particular, the  $G/G_r$ 's and the  $H/H_r$ 's are finite). We need to prove

$$\begin{aligned} \varinjlim_{r \rightarrow +\infty}^u \bigoplus_{(g,h) \in (G_r \times H_r)} k\chi_{(g,h)+G_r \times H_r} \\ = \varinjlim_{r \rightarrow +\infty}^u \bigoplus_{g \in G_r} k\chi_{g+G_r} \widehat{\otimes}_k^u \varinjlim_{r \rightarrow +\infty}^u \bigoplus_{h \in H_r} k\chi_{h+H_r}. \end{aligned}$$

Let  $M$  (resp.  $N$ ) be the left-hand side (resp. the right-hand side) in the previous equation. Then

$$M = \varprojlim_{I \in \mathcal{P}(k)} M/\overline{IM}, \quad N = \varprojlim_{I \in \mathcal{P}(k)} N/\overline{IN}.$$

We show that  $M/\overline{IM} \xrightarrow{\sim} N/\overline{IN}$  for any  $I \in \mathcal{P}(k)$ . We have

$$M/\overline{IM} = \varinjlim_r \bigoplus_{(g,h) \in (G_r \times H_r)} (k/I)\chi_{(g,h)+G_r \times H_r}.$$

Let

$$P := \varinjlim_r \bigoplus_{g \in G_r} k\chi_{g+G_r}, \quad Q := \varinjlim_r \bigoplus_{h \in H_r} k\chi_{h+H_r}.$$

Then

$$\begin{aligned} N/\overline{IN} &= P/\overline{IP} \otimes_{k/I} Q/\overline{IQ} \\ &= \varinjlim_r \bigoplus_{g \in G_r} (k/I)\chi_{g+G_r} \otimes_{k/I} \varinjlim_r \bigoplus_{h \in H_r} (k/I)\chi_{h+H_r} = M/\overline{IM}. \end{aligned}$$

This concludes the proof of (29).

We now pass to (30). We use Proposition 5.6 (2) to replace the map in the statement by

$$\begin{aligned} & \varinjlim_r^u \mathcal{C}_{\text{unif}}(G/G_r, k) \widehat{\otimes}_k^u \varinjlim_{r \rightarrow +\infty}^u \mathcal{C}_{\text{unif}}(H/H_r, k) \\ & \rightarrow \varinjlim_r^u \mathcal{C}_{\text{unif}}((G \times H)/(G_r \times H_r), k). \end{aligned}$$

By Lemma 5.3 this reduces to considering

$$\begin{aligned} & \varinjlim_r^u \varprojlim_{I \in \mathcal{P}(k)} \prod_{g \in G}^{\square, u} (k/I)e_{g+G_r} \widehat{\otimes}_k^u \varinjlim_{r \rightarrow +\infty}^u \varprojlim_{I \in \mathcal{P}(k)} \prod_{h \in H}^{\square, u} (k/I)e_{h+H_r} \\ & \rightarrow \varinjlim_r^u \varprojlim_{I \in \mathcal{P}(k)} \prod_{(g,h)}^{\square, u} (k/I)e_{(g+G_r, h+H_r)}. \end{aligned}$$

As before, let  $M$  (resp.  $N$ ) be the left-hand side (resp. the right-hand side) in the previous equation. Then

$$M = \varprojlim_{I \in \mathcal{P}(k)} M/\overline{IM}, \quad N = \varprojlim_{I \in \mathcal{P}(k)} N/\overline{IN}.$$

We show that  $M/\overline{IM} \hookrightarrow N/\overline{IN}$  in an embedding with the relative topology, for any  $I \in \mathcal{P}(k)$ . Now,

$$\begin{aligned} M/\overline{IM} &= \varinjlim_r^u \prod_g (k/I)e_{g+G_r} \widehat{\otimes}_{k/I}^u \varinjlim_r^u \prod_h (k/I)e_{h+H_r} \\ &= \varinjlim_r^u \left( \prod_g (k/I)e_{g+G_r} \widehat{\otimes}_{k/I}^u \prod_h (k/I)e_{h+H_r} \right) \end{aligned}$$

and

$$N/\overline{IN} = \varinjlim_r^u \prod_{(g,h)} (k/I)e_{(g+G_r, h+H_r)},$$

where  $\varinjlim_r^u$  and  $\widehat{\otimes}_{k/I}^u$  are taken in the category  $\mathcal{CLM}_{k/I}^u$ . So, our statement is reduced to the fact that, for  $k, G$ , and  $H$  discrete, if  $k^G$  (resp.  $k^H$ , resp.  $k^{G \times H}$ ) indicates the  $k$ -algebra of functions  $G \rightarrow k$  (resp.  $H \rightarrow k$ , resp.  $G \times H \rightarrow k$ ) with the discrete topology, then we have an inclusion

$$k^G \otimes_k k^H \hookrightarrow k^{G \times H}. \quad \square$$

We are especially interested in the following corollary.

**Corollary 5.8.**

(1) For any  $r \in \mathbb{Z}$ ,

$$\mathcal{C}_{\text{unif}}(\mathbb{Q}_p/p^{r+1}\mathbb{Z}_p, \mathbb{Z}_p) = \varprojlim_{s \rightarrow +\infty} \mathcal{C}(\mathbb{Q}_p/p^{r+1}\mathbb{Z}_p, \mathbb{Z}_p/p^s\mathbb{Z}_p),$$

where  $\mathcal{C}(\mathbb{Q}_p/p^{r+1}\mathbb{Z}_p, \mathbb{Z}_p/p^s\mathbb{Z}_p)$  is equipped with the discrete topology. It is the  $\mathbb{Z}_p$ -algebra of all maps  $\mathbb{Q}_p/p^{r+1}\mathbb{Z}_p \rightarrow \mathbb{Z}_p/p^s\mathbb{Z}_p$  equipped with the  $p$ -adic topology.

(2) For any  $r \in \mathbb{Z}$ ,

$$\mathcal{C}(\mathbb{Q}_p/p^{r+1}\mathbb{Z}_p, \mathbb{Z}_p) = \varprojlim_{s, t \rightarrow +\infty} \mathcal{C}(p^{-t}\mathbb{Z}_p/p^{r+1}\mathbb{Z}_p, \mathbb{Z}_p/p^s\mathbb{Z}_p),$$

where  $\mathcal{C}(p^{-t}\mathbb{Z}_p/p^{r+1}\mathbb{Z}_p, \mathbb{Z}_p/p^s\mathbb{Z}_p)$  is equipped with the discrete topology. It is the  $\mathbb{Z}_p$ -Hopf algebra of all maps  $\mathbb{Q}_p/p^{r+1}\mathbb{Z}_p \rightarrow \mathbb{Z}_p$  equipped with the topology of simple convergence on  $\mathbb{Q}_p/p^{r+1}\mathbb{Z}_p$  for the  $p$ -adic topology of  $\mathbb{Z}_p$ .

(3) We have

$$\mathcal{C}_{\text{unif}}(\mathbb{Q}_p, \mathbb{Z}_p) = \varinjlim_{r \rightarrow +\infty}^u \mathcal{C}_{\text{unif}}(\mathbb{Q}_p/p^{r+1}\mathbb{Z}_p, \mathbb{Z}_p);$$

(4) We have

$$\mathcal{C}(\mathbb{Q}_p, \mathbb{Z}_p) = \varprojlim_{s \rightarrow +\infty} \mathcal{C}(\mathbb{Q}_p, \mathbb{Z}_p/p^s\mathbb{Z}_p).$$

**Remark 5.9.** Formula (29) shows that  $\mathcal{C}(\mathbb{Q}_p, \mathbb{Z}_p)$  is a Hopf algebra object in  $\mathcal{CLM}_{\mathbb{Z}_p}^u$ .

**Remark 5.10.** We point out a tautological, but useful, formula which holds in  $\mathcal{C}(\mathbb{Q}_p/p^{r+1}\mathbb{Z}_p, \mathbb{F}_p)$ . For any  $h \in \mathbb{Q}_p/p^{r+1}\mathbb{Z}_p$ , let  $e_h$  denote as before the function  $\mathbb{Q}_p/p^{r+1}\mathbb{Z}_p \rightarrow \mathbb{F}_p$  such that  $e_h(h) = 1$  while  $e_h(x) = 0$  if  $x \in \mathbb{Q}_p/p^{r+1}$ ,  $x \neq h$ . For any  $i \leq r$ , the function

$$x_i : \mathbb{Q}_p/p^{r+1}\mathbb{Z}_p \rightarrow \mathbb{F}_p$$

was introduced in (14). We then have

$$x_i = \sum_{h \in \mathbb{Q}_p/p^{r+1}\mathbb{Z}_p} h_i e_h,$$

where  $h_i = x_i(h)$ .

**Lemma 5.11.** Let  $G$  and  $K$  be as above but assume  $G$  is discrete. Then in the category  $\mathcal{CLC}_K$ ,

$$\mathcal{C}(G, K) = \prod_{g \in G} K e_g$$

is a Fréchet  $K$ -algebra and

$$\mathcal{C}_{\text{unif}}^{\text{bd}}(G, K) = \ell_\infty(G, K)$$

is the Banach  $K$ -algebra of bounded sequences  $(a_g)_{g \in G}$  of elements of  $K$ , equipped with the componentwise sum and product and with the supnorm.

*Proof.* Obvious from the definitions. □

**Lemma 5.12.** Let  $G$  and  $K$  be as above, but assume  $G$  is compact. Then in the category  $\mathcal{CLC}_K$ ,

$$\mathcal{C}(G, K) = \mathcal{C}_{\text{unif}}^{\text{bd}}(G, K) = \ell_\infty^0(G, K)$$

is the Banach  $K$ -algebra of sequences  $(a_g)_{g \in G}$ , with  $a_g \in K$ , such that  $a_g \rightarrow 0$  along the filter of cofinite subsets of  $G$ , equipped with componentwise sum and product and with the supnorm.

*Proof.* This is a straight-forward generalization of the classical wavelet decomposition. See [13, Prop. 1.16].  $\square$

**Proposition 5.13.** *Let  $G$  and  $K$  be as in all this section. Then in the category  $\mathcal{C}\mathcal{L}\mathcal{C}_K$ , the following hold:*

(1) *We have*

$$\mathcal{C}(G, K) = \varprojlim_{r \rightarrow -\infty} \mathcal{C}(G_r, K)$$

*for the restrictions  $\mathcal{C}(G_r, K) \rightarrow \mathcal{C}(G_{r+1}, K)$ . In particular,  $\mathcal{C}(G, K)$  is a Fréchet  $K$ -algebra.*

(2) *We have*

$$\mathcal{C}_{\text{unif}}^{\text{bd}}(G, K) = \varinjlim_{r \rightarrow +\infty} \mathcal{C}_{\text{unif}}^{\text{bd}}(G/G_r, K)$$

*for the embeddings*

$$\mathcal{C}_{\text{unif}}^{\text{bd}}(G/G_r, K) \hookrightarrow \mathcal{C}_{\text{unif}}^{\text{bd}}(G/G_{r+1}, K),$$

*where the inductive limit of Banach  $K$ -algebras is strict. In particular,  $\mathcal{C}_{\text{unif}}^{\text{bd}}(G, K)$  is a complete bornological  $K$ -algebra.*

(3) *The natural morphism*

$$\mathcal{C}_{\text{unif}}^{\text{bd}}(G, K) \rightarrow \mathcal{C}(G, K)$$

*is injective and has dense image.*

*Proof.* It is clear. For the notion of a bornological topological  $K$ -vector space, we refer to [18, §6]. The fact that the notion is stable by strict inductive limits is Example 3 on page 39 of *loc. cit.* The statement on completeness is proved in [18, Lemma 7.9].  $\square$

**Proposition 5.14.** *Let  $G$  and  $H$  be locally compact groups as in Proposition 5.7. Then*

$$(31) \quad \mathcal{C}(G, K) \widehat{\otimes}_{\pi, K} \mathcal{C}(H, K) \xrightarrow{\sim} \mathcal{C}(G \times H, K),$$

*while the canonical map*

$$\mathcal{C}_{\text{unif}}^{\text{bd}}(G, K) \widehat{\otimes}_{\pi, K} \mathcal{C}_{\text{unif}}^{\text{bd}}(H, K) \rightarrow \mathcal{C}_{\text{unif}}^{\text{bd}}(G \times H, K)$$

*is a strictly closed embedding of complete bornological algebras.*

*Proof.* In the case of  $G$  and  $H$  being compact, this is detailed in the example after [18, Prop. 17.10]. In the general case, (31) follows by taking projective limits. The statement for  $\mathcal{C}_{\text{unif}}^{\text{bd}}(G, K)$  reduces instead to (30).  $\square$

We point out that  $(\mathcal{C}\mathcal{L}\mathcal{C}_K, \widehat{\otimes}_{\pi, K})$  is a  $K$ -linear symmetric monoidal category. From Remarks 5.7 and 5.14, we conclude the following proposition.



**Proposition 5.15.** *Let  $G$  be as in Definition 5.1, and let  $A$  be either  $k$  or  $K$ , as before, and  $\mathcal{C}(G, A)$  be as in loc. cit. We regard  $(\mathcal{CLM}_k^u, \widehat{\otimes}_k)$  and  $(\mathcal{CLC}_K, \widehat{\otimes}_{\pi, K})$  as symmetric monoidal categories. The coproduct, counit, and inversion*

$$\mathbb{F}(f)(x, y) = f(x + y), \quad \varepsilon(f) = f(0_G), \quad \rho(f)(x) = f(-x),$$

for any  $f \in \mathcal{C}(G, A)$  and any  $x, y \in G$ , define a structure of topological  $A$ -Hopf algebra on  $\mathcal{C}(G, A)$ , in the sense of the previous monoidal categories.

The following result describes the structure of the Hopf algebras of functions

$$\mathbb{Q}_p/p^{r+1}\mathbb{Z}_p \rightarrow \mathbb{Z}_p/p^{a+1}\mathbb{Z}_p,$$

for any  $r, a \in \mathbb{Z}$  and  $a \geq 0$ , in terms of the functions

$$x_i : \mathbb{Q}_p/p^{i+1}\mathbb{Z}_p \rightarrow \mathbb{F}_p$$

introduced in (14). See also Remark 5.10.

**Proposition 5.16.** *For any  $i \in \mathbb{Z}$ , let  $x_i$  be as in (14) and let  $X_i$  be indeterminates. For  $r \in \mathbb{Z}$  and  $i \in \mathbb{Z}_{\geq 0}$ , let  $\mathbb{F}_p(r, i)$  denote the  $\mathbb{F}_p$ -algebra*

$$\mathbb{F}_p[X_r, X_{r-1}, X_{r-2}, \dots, X_{r-i}] / (1 - X_r^{p-1}, 1 - X_{r-1}^{p-1}, \dots, 1 - X_{r-i}^{p-1}).$$

The dimension of  $\mathbb{F}_p(r, i)$  as a  $\mathbb{F}_p$ -vector space is  $(p - 1)^{i+1}$ . Let  $X_{r,i} := (X_{r-i}, X_{r-i+1}, \dots, X_{r-1}, X_r)$  be viewed as a Witt vector of length  $i + 1$  with coefficients in  $\mathbb{F}_p(r, i)$ . We make  $\mathbb{F}_p(r, i)$  into an  $\mathbb{F}_p$ -Hopf algebra by setting

$$\mathbb{P}X_{r,i} = X_{r,i} \otimes_{\mathbb{F}_p} 1 + 1 \otimes_{\mathbb{F}_p} X_{r,i}.$$

For any  $i = 0, 1, \dots$ , the  $\mathbb{F}_p$ -algebra map  $\mathbb{F}_p(r, i + 1) \rightarrow \mathbb{F}_p(r, i)$ , sending  $X_{r-j}$  to  $X_{r-j}$  if  $0 \leq j \leq i$  and  $X_{r-i-1}$  to 0, is a homomorphism of  $\mathbb{F}_p$ -Hopf algebras. Then in the category  $\mathcal{CLM}_{\mathbb{F}_p}^u$ ,

(1) the map

$$\mathbb{F}_p(r, i) \rightarrow \mathcal{C}(p^{r-i}\mathbb{Z}_p/p^{r+1}\mathbb{Z}_p, \mathbb{F}_p), \quad X_j \mapsto x_j,$$

for  $r - i \leq j \leq r$ , is an isomorphism of  $\mathbb{F}_p$ -Hopf algebras,

(2) the  $\mathbb{F}_p$ -Hopf algebra  $\mathcal{C}(\mathbb{Q}_p/p^{r+1}\mathbb{Z}_p, \mathbb{F}_p)$  equals

$$\mathbb{F}_p(r, \infty) := \varprojlim_{i \rightarrow +\infty} \mathbb{F}_p(r, i)$$

with the prodiscrete topology,

(3) the topological  $\mathbb{F}_p$ -algebra  $\mathcal{C}_{\text{unif}}(\mathbb{Q}_p/p^{r+1}\mathbb{Z}_p, \mathbb{F}_p)$  equals  $\mathbb{F}_p(r, \infty)$  equipped with the discrete topology.

*Proof.* Parts (1) and (2) are [7, Teorema 3.31]. Part (3) follows by forgetting the topology. □

**Remark 5.17.** Notice that the  $\mathbb{F}_p$ -algebras  $\mathbb{F}_p(r, i)$  are perfect.

**Corollary 5.18.** *For  $r \in \mathbb{Z}$  and  $i, a \in \mathbb{Z}_{\geq 0}$ , the following hold:*

- (1) *The topological  $\mathbb{Z}_p/p^{a+1}\mathbb{Z}_p$ -algebra  $\mathcal{C}_{\text{unif}}(\mathbb{Q}_p/p^{r+1}\mathbb{Z}_p, \mathbb{Z}_p/p^{a+1}\mathbb{Z}_p)$  equals*

$$W_a(\mathcal{C}_{\text{unif}}(\mathbb{Q}_p/p^{r+1}\mathbb{Z}_p, \mathbb{F}_p)) = W_a(\mathbb{F}_p(r, \infty))$$

*equipped with the discrete topology. Therefore,*

$$\mathcal{C}_{\text{unif}}(\mathbb{Q}_p/p^{r+1}\mathbb{Z}_p, \mathbb{Z}_p) = W(\mathcal{C}_{\text{unif}}(\mathbb{Q}_p/p^{r+1}\mathbb{Z}_p, \mathbb{F}_p)) = W(\mathbb{F}_p(r, \infty))$$

*equipped with the  $p$ -adic topology.*

- (2) *The  $\mathbb{Z}_p/p^{a+1}\mathbb{Z}_p$ -Hopf algebra  $\mathcal{C}(\mathbb{Q}_p/p^{r+1}\mathbb{Z}_p, \mathbb{Z}_p/p^{a+1}\mathbb{Z}_p)$  equals*

$$(32) \quad W_a(\mathcal{C}(\mathbb{Q}_p/p^{r+1}\mathbb{Z}_p, \mathbb{F}_p)) = W_a(\mathbb{F}_p(r, \infty))$$

*with the prodiscrete topology. Therefore,*

$$(33) \quad \mathcal{C}(\mathbb{Q}_p/p^{r+1}\mathbb{Z}_p, \mathbb{Z}_p) = W(\mathcal{C}(\mathbb{Q}_p/p^{r+1}\mathbb{Z}_p, \mathbb{F}_p)) = W(\mathbb{F}_p(r, \infty))$$

*equipped with the product topology of the prodiscrete topology of  $\mathbb{F}_p(r, \infty)$  on the components.*

**Definition 5.19.** We set

$$\mathcal{C} = \mathcal{C}(\mathbb{Q}_p, \mathbb{Z}_p).$$

For any  $r, a \in \mathbb{Z}$  with  $a \geq 0$ , we define a Fréchet  $\mathbb{Z}_p$ -subalgebra of  $\mathcal{C}$  by

$$\mathcal{C}_{r,a} := \{f \in \mathcal{C} \mid f(x + p^{r+1}\mathbb{Z}_p) \subset f(x) + p^{a+1}\mathbb{Z}_p \text{ for all } x \in \mathbb{Q}_p\}.$$

Let  $F$  be the set-theoretic map

$$F: \mathcal{C} \rightarrow \mathcal{C}, \quad f \mapsto f^p.$$

Then  $\mathcal{C}_{r,a+1} \subset \mathcal{C}_{r,a}$  and  $\mathcal{C}_{r,a} \subset \mathcal{C}_{r+1,a}$ ,

$$(34) \quad p^{a+1}\mathcal{C} \subset \mathcal{C}_{r,a}$$

is an ideal of  $\mathcal{C}_{r,a}$ , and  $F$  induces a map

$$F: \mathcal{C}_{r,a} \rightarrow \mathcal{C}_{r,a+1}.$$

There exists a canonical map

$$R_{r,a}: \mathcal{C}_{r,a} \rightarrow \mathcal{C}(\mathbb{Q}_p/p^{r+1}\mathbb{Z}_p, \mathbb{Z}_p/p^{a+1}\mathbb{Z}_p), \quad f \mapsto R_{r,a}(f),$$

such that

$$\pi_{a+1} \circ f = R_{r,a}(f) \circ \pi_{r+1},$$

which sits in the exact sequence

$$(35) \quad 0 \rightarrow p^{a+1}\mathcal{C} \rightarrow \mathcal{C}_{r,a} \xrightarrow{R_{r,a}} \mathcal{C}(\mathbb{Q}_p/p^{r+1}\mathbb{Z}_p, \mathbb{Z}_p/p^{a+1}\mathbb{Z}_p) = W_a(\mathbb{F}_p(r, \infty)) \rightarrow 0.$$

We conclude with the following proposition.

**Proposition 5.20.** *For any  $r \in \mathbb{Z}$  and any  $a \in \mathbb{Z}_{\geq 1}$ , the map  $f \mapsto \pi_1 \circ f$  induces an isomorphism*

$$\mathcal{C}_{r,a}/p\mathcal{C}_{r,a-1} \xrightarrow{\sim} \mathcal{C}_{\text{unif}}(\mathbb{Q}_p/p^{r+1}\mathbb{Z}_p, \mathbb{F}_p).$$

Similarly, for  $a = 0$ , we have

$$\mathcal{C}_{r,0}/p\mathcal{C} \xrightarrow{\sim} \mathcal{C}_{\text{unif}}(\mathbb{Q}_p/p^{r+1}\mathbb{Z}_p, \mathbb{F}_p).$$

The inverse of the isomorphism of discrete  $\mathbb{F}_p$ -algebras

$$\mathcal{C}_{r,a}/p\mathcal{C}_{r,a-1} \xrightarrow{\sim} \mathcal{C}_{r,0}/p\mathcal{C}$$

is provided by the map

$$F^a : \mathcal{C}_{r,0}/p\mathcal{C} \xrightarrow{\sim} \mathcal{C}_{r,a}/p\mathcal{C}_{r,a-1}, \quad f \mapsto f^{p^a}.$$

*Proof.* The first formula follows from (34) and (35). In fact,

$$\begin{aligned} \mathcal{C}_{r,a}/p\mathcal{C}_{r,a-1} &= (\mathcal{C}_{r,a}/p^a\mathcal{C})/p(\mathcal{C}_{r,a-1}/p^{a-1}\mathcal{C}) \\ &= W_a(\mathbb{F}_p(r, \infty))/pW_{a-1}(\mathbb{F}_p(r, \infty)) = \mathbb{F}_p(r, \infty). \end{aligned}$$

Similarly for the other formulas. □

By iteration, we get the following corollary.

**Corollary 5.21.** *We have*

$$\mathcal{C}_{r,a}/p^{a+1}\mathcal{C} \xrightarrow{\sim} \mathcal{C}(\mathbb{Q}_p/p^{r+1}\mathbb{Z}_p, \mathbb{Z}_p/p^{a+1}\mathbb{Z}_p) = W_a(\mathbb{F}_p(r, \infty)).$$

For any  $f \in \mathcal{C}_{r,a}$ , there exist  $f_0, f_1, \dots, f_a \in \mathcal{C}_{r,0}$ , well determined modulo  $p\mathcal{C}$ , such that

$$f \equiv f_0^{p^a} + pf_1^{p^{a-1}} + p^2f_2^{p^{a-2}} + \dots + p^af_a \pmod{p^{a+1}\mathcal{C}}.$$

### 6. $p$ -ADICALLY ENTIRE FUNCTIONS BOUNDED ON $\mathbb{Q}_p$

We prove here the statements announced in the introduction, namely, Theorem 2.17, Theorem 2.19, Proposition 2.20, Proposition 2.21, Proposition 2.23, Proposition 2.24, and Theorem 2.27. We assume  $q = p$  from now on, so, in particular,  $\Psi$  stands for  $\Psi_p$ .

We start with the proof of Theorem 2.17.

*Proof of Theorem 2.17.* It suffices to prove the statement over  $\mathbb{Z}_p$ . Notice that

$$\mathbb{Z}_p[\Psi(\lambda x) \mid \lambda \in \mathbb{Q}_p^\times] = \mathbb{Z}_p[\Psi(\lambda p^{-i}x) \mid i \in \mathbb{Z}, \lambda \in \mathbb{Z}_p^\times].$$

Both rings  $\mathbb{Z}_p[[\lambda x]_i \mid i \in \mathbb{Z}, \lambda \in \mathbb{Z}_p^\times]$  and  $\mathbb{Z}_p[\Psi(\lambda p^{-i}x) \mid i \in \mathbb{Z}, \lambda \in \mathbb{Z}_p^\times]$  are contained in the  $\mathbb{Z}_p$ -Banach ring  $\mathcal{C}_{\text{unif}}(\mathbb{Q}_p, \mathbb{Z}_p)$ , which may be identified with  $W(\mathcal{C}_{\text{unif}}(\mathbb{Q}_p, \mathbb{F}_p))$  equipped with the  $p$ -adic topology. Then  $AP_{\mathbb{Z}_p}$  consists of  $W(\mathbb{F}_p[[\lambda x]_i \mid i \in \mathbb{Z}, \lambda \in \mathbb{Z}_p^\times])$ . Notice that  $\mathbb{F}_p[[\lambda x]_i \mid i \in \mathbb{Z}, \lambda \in \mathbb{Z}_p^\times]$  is a perfect subring of the perfect ring  $\mathcal{C}_{\text{unif}}(\mathbb{Q}_p, \mathbb{F}_p)$ , since  $(\lambda x)_i^p = (\lambda x)_i$  for any  $i, \lambda$ . It suffices to prove the following.

**Lemma 6.1.** *For any fixed  $\lambda \in \mathbb{Z}_p^\times$ , the closure of  $\mathbb{Z}_p[\Psi(\lambda p^i x) \mid i = 0, 1, 2, \dots]$  in  $W(\mathcal{C}_{\text{unif}}(\mathbb{Q}_p, \mathbb{F}_p))$  coincides with  $W(\mathbb{F}_p[[\lambda x]_i \mid i = 0, -1, -2, \dots])$ .*

We may as well assume  $\lambda = 1$  and prove the following.

**Sublemma 6.2.** *The closure of  $\mathbb{Z}_p[\Psi(p^i x) \mid i = 0, 1, 2, \dots]$  in  $W(\mathcal{C}_{\text{unif}}(\mathbb{Q}_p, \mathbb{F}_p))$  coincides with  $W(\mathbb{F}_p[x_i \mid i = 0, -1, -2, \dots])$ .*

Let  $C$  be the closure of  $\mathbb{Z}_p[\Psi(p^i x) \mid i = 0, 1, 2, \dots]$  in  $W(\mathcal{C}_{\text{unif}}(\mathbb{Q}_p, \mathbb{F}_p))$ . The formula

$$[x_{-i}] = \lim_{N \rightarrow \infty} \Psi(p^i x)^N$$

shows that  $W(\mathcal{C}_{\text{unif}}(\mathbb{Q}_p, \mathbb{F}_p)) \subset C$ . It will suffice to show that, as functions  $\mathbb{Q}_p \rightarrow \mathbb{Z}_p$ ,

$$\Psi(x) \in W(\mathbb{F}_p[x_i \mid i = 0, -1, -2, \dots]).$$

We write the restriction of  $\Psi(x)$  to a function  $\mathbb{Q}_p \rightarrow \mathbb{Z}_p$  as

$$\Psi(x) = (\Psi_0(x), \Psi_1(x), \Psi_2(x), \dots),$$

with  $\Psi_i \in \mathcal{C}_{\text{unif}}(\mathbb{Q}_p, \mathbb{F}_p)$  and  $\Psi_0(x) = x_0$ . We have, from (2), the formula in  $\mathcal{C}_{\text{unif}}(\mathbb{Q}_p, \mathbb{Q}_p)$

$$(36) \quad \begin{aligned} \Psi(x) + p^{-1}\Psi(px)^p + \dots + p^{-i}\Psi(p^i x)^{p^i} + \dots \\ = x = (\dots, x_{-i}, \dots, x_{-2}, x_{-1}; x_0, *, *, \dots). \end{aligned}$$

From (36) we deduce that, as functions in  $\mathcal{C}_{\text{unif}}(\mathbb{Q}_p, p^{-i}\mathbb{Z}_p)$ ,

$$\begin{aligned} \Psi(x) + p^{-1}\Psi(px)^p + p^{-2}\Psi(p^2 x)^{p^2} + \dots + p^{-i}\Psi(p^i x)^{p^i} \\ = (x_{-i}, \dots, x_{-2}, x_{-1}; x_0, *, *, \dots). \end{aligned}$$

This shows, inductively on  $i$ , that

$$\Psi_i \in \mathbb{F}_p[x_j \mid j = 0, -1, -2, \dots, -i]. \quad \square$$

**Definition 6.3.** Let  $r \in \mathbb{Z}$  and  $a \in \mathbb{Z}_{\geq 1}$ . We define  $\mathcal{E}_{r,a}^\circ$  (resp.  $\mathcal{T}_{r,a}^\circ$ ) to be the  $\mathbb{Z}_p$ -subalgebra of  $\mathcal{E}_{p^r}^\circ$  (resp. of  $\mathcal{T}_{p^r}^\circ$ ) (cf. Definition 2.18) consisting of those functions  $f$  such that

$$f(x + p^{r+j}\mathbb{C}_p^\circ) \subset f(x) + p^{a+j}\mathbb{C}_p^\circ \quad \text{for all } x \in \mathbb{Q}_p \text{ and } j \in \mathbb{Z}_{\geq 1}.$$

**Remark 6.4.** For the rest of this section, the statements valid for the rings  $\mathcal{E}_{r,a}^\circ \subset \mathbb{Q}_p\{x\}$  hold equally well, and with the same proof, for the rings  $\mathcal{T}_{r,a}^\circ \subset \mathcal{O}(\Sigma_{p^{-r}})^\circ$ . For short, we deal with the former only.

Notice that

$$\mathcal{E}_{r,a+1}^\circ \subset \mathcal{E}_{r,a}^\circ \subset \mathcal{E}_{r+1,a+1}^\circ \quad \text{and} \quad p\mathcal{E}_{r,a}^\circ \subset \mathcal{E}_{r,a+1}^\circ,$$

and that we have a map  $F$  as in Definition 5.19 such that

$$F(\mathcal{E}_{r,a}^\circ) \subset \mathcal{E}_{r,a+1}^\circ.$$

**Remark 6.5.** We have

$$\mathcal{E}_{p^r}^\circ := \mathcal{E}_{r,0}^\circ.$$

We already proved (Proposition 1.3 and Corollary 4.5) that  $\Psi(x) \in \mathcal{E}_{0,0}^\circ$ . Therefore, for any  $i \in \mathbb{Z}_{\geq 0}$  and  $\ell = 0, 1, \dots, p - 1$ , the function  $\Psi(p^{i-r}x)^{\ell p^a}$  belongs to  $\mathcal{E}_{r-i,a}^\circ \subset \mathcal{E}_{r,a}^\circ$ .

**Lemma 6.6.** *If a sequence of functions  $n \mapsto f_n \in \mathcal{E}_{r,a}^\circ$  (resp.  $\in \mathcal{T}_{r,a}^\circ$ ) converges to  $f \in \mathbb{C}_p\{x\}$  (resp. to  $f \in \mathcal{O}(\Sigma_{p^{-r}})^\circ$ ) uniformly on bounded subsets of  $\mathbb{C}_p$  (resp. of  $\Sigma_{p^{-r}}$ ), then  $f \in \mathcal{E}_{r,a}^\circ$  (resp.  $\in \mathcal{T}_{r,a}^\circ$ ). Therefore,  $\mathcal{E}_{r,a}^\circ$  (resp.  $\mathcal{T}_{r,a}^\circ$ ) is a closed  $\mathbb{Z}_p$ -subalgebra of  $\mathbb{C}_p\{x\}$  (resp. of  $(\mathcal{O}(\Sigma_{p^{-r}})^\circ, \text{standard})$ ). The induced Fréchet algebra structure on  $\mathcal{E}_{r,a}^\circ$  (resp. on  $\mathcal{T}_{r,a}^\circ$ ) will be called standard.*

*Proof.* We deal, to fix ideas, with the case of  $\mathcal{E}_{r,a}^\circ$ . We show that for any  $c \in \mathbb{Q}_p$  and  $j = 0, 1, \dots$ ,

$$f(c + p^{r+j+1}\mathbb{C}_p^\circ) \subset f(c) + p^{a+j+1}\mathbb{C}_p^\circ.$$

By assumption, for any  $s, t \in \mathbb{N}$ , there exists  $N = N_{s,t}$  such that if  $n \geq N$ , then

$$(f_n - f)(p^{-s}\mathbb{C}_p^\circ) \subset p^t\mathbb{C}_p^\circ.$$

So, for  $c$  and  $j$  as before, let  $s$  be such that  $c + p^{r+j+1}\mathbb{C}_p^\circ \subset p^{-s}\mathbb{C}_p^\circ$ , and let  $t$  be  $\geq j + a + 1$ . Then, for any  $n \geq N_{s,t}$ ,

$$(f_n - f)(c + p^{r+j+1}\mathbb{C}_p^\circ) \subset (f_n - f)(p^{-s}\mathbb{C}_p^\circ) \subset p^t\mathbb{C}_p^\circ \subset p^{j+a+1}\mathbb{C}_p^\circ.$$

Therefore,  $f \in \mathcal{E}_{r,a}^\circ$ . □

Notice that Proposition 2.20 follows from Lemma 6.6, by taking  $a = 0$ .

Let  $r, a$  be as in Definition 6.3. Any function  $f \in \mathcal{E}_{r,a}^\circ$  induces a continuous function  $f|_{\mathbb{Q}_p} : \mathbb{Q}_p \rightarrow \mathbb{Z}_p$ . The  $\mathbb{Z}_p$ -linear map

$$(37) \quad \text{Res}^\circ : (\mathcal{E}_{r,a}^\circ, \text{standard}) \rightarrow \mathcal{C}_{r,a} \subset \mathcal{C}(\mathbb{Q}_p, \mathbb{Z}_p), \quad f \mapsto f|_{\mathbb{Q}_p},$$

is continuous and injective. By composition, we obtain, for any  $r \in \mathbb{Z}$  and any  $a, h = 0, 1, \dots, a$  morphism

$$(38) \quad R_{r,a} \circ \text{Res}^\circ : (\mathcal{E}_{r,a}^\circ, \text{standard}) \rightarrow \mathcal{C}(p^{r-h}\mathbb{Z}_p/p^{r+1}\mathbb{Z}_p, \mathbb{Z}_p/p^{a+1}\mathbb{Z}_p) = W_a(\mathbb{F}_p(r, h)),$$

where the right-hand side is equipped with the topology of (32). The kernel of that map is the set of  $g \in \mathcal{E}_{r,a}^\circ$  such that  $-\log\|g\|_{p^{h-r}} \geq a + 1$ . From (37) we also get maps of Fréchet  $\mathbb{Z}_p$ -algebras:

$$(39) \quad \text{Res}^\circ : (\mathcal{E}_\lambda^\circ, \{\| \cdot \|_{p^r\mathbb{Z}_p}\}_{r \in \mathbb{Z}})^\wedge \rightarrow \mathcal{C}(\mathbb{Q}_p, \mathbb{Z}_p), \quad f \mapsto f|_{\mathbb{Q}_p},$$

$$(40) \quad \text{Res}^\circ : (\mathcal{T}_\lambda^\circ, \{\| \cdot \|_{p^r\mathbb{Z}_p}\}_{r \in \mathbb{Z}})^\wedge \rightarrow \mathcal{C}(\mathbb{Q}_p, \mathbb{Z}_p), \quad f \mapsto f|_{\mathbb{Q}_p}.$$

**Lemma 6.7.** *Let  $r \in \mathbb{Z}$  and  $a \in \mathbb{Z}_{\geq 0}$  be as before.*

(1) *Any series of functions of the form*

$$\sum_{\ell=0}^{p-1} \sum_{i=0}^{\infty} c_{\ell,a,i} \Psi(p^{i-r}x)^{\ell p^a}, \quad c_{\ell,a,i} \in \mathbb{Z}_p,$$

*converges in the standard Fréchet topology of  $\mathbb{Q}_p\{x\}$  to an element of  $\mathcal{E}_{r,a}^\circ$  along the filter of cofinite subsets of  $\{0, 1, \dots, p-1\} \times \mathbb{Z}_{\geq 0}$ .*

- (2) For any element  $f \in \mathcal{C}_{r,a}$  and for any  $s = 0, 1, 2, \dots$ , there exist uniquely determined elements  $c_{\ell,b,i} = c_{\ell,b,i}^p \in \mathbb{Z}_p$  such that for

$$f_{r,a} := \sum_{b=0}^a \sum_{\ell=0}^{p-1} \sum_{i=0}^{\infty} c_{\ell,b,i} p^b \Psi(p^{i-r}x)^{\ell p^{a-b}} \in \mathcal{E}_{r,a}^{\circ},$$

where the infinite sum converges in the standard Fréchet topology of  $\mathcal{E}_{r,a}^{\circ}$ , we have

$$-\log\|(f - f_{r,a})\|_{p^{-r}\mathbb{Z}_p} \geq a + 1.$$

Same statement for  $\mathcal{E}_{r,a}^{\circ}$  replaced by  $\mathcal{T}_{r,a}^{\circ}$ .

- (3) For any element  $f \in \mathcal{C}_{r,a}$  and any  $h = 0, 1, \dots$ , there exist uniquely determined elements  $c_{\ell,b,i} = c_{\ell,b,i}^p \in \mathbb{Z}_p$ , such that for

$$f_{r,a,h} := \sum_{b=0}^a \sum_{\ell=0}^{p-1} \sum_{i=0}^h c_{\ell,b,i} p^b \Psi(p^{i-r}x)^{\ell p^{a-b}},$$

we have

$$-\log\|(f - f_{r,a,h})\|_{p^{h-r}\mathbb{Z}_p} \geq a + 1.$$

- (4) The map (38) is surjective.

- (5) The maps (39) and (40) are the isomorphisms of Theorem 2.19.

*Proof.* The first part is clear. As for the second, we observe that, for any  $b = 0, 1, \dots, a$ , the map  $R_{r,a} \circ \text{Res}^{\circ}$  transforms the function  $p^b \Psi(p^{i-r}x)^{\ell p^{a-b}}$ , for  $\ell = 0, 1, \dots, p - 1$ , into the Witt vector

$$(0, \dots, 0, w_b = x_{r-i}^{\ell}, 0, \dots, 0) \in W_a(\mathbb{F}_p(r, \infty)),$$

where  $x_{r-i}^{\ell}$  is placed at the  $b$ -th level. Since any  $y \in \mathbb{F}_p(r, \infty)$  admits a unique expression as a sum, convergent in the prodiscrete topology of  $\mathbb{F}_p(r, \infty)$ ,

$$y = \sum_{\ell=0}^{p-1} \sum_{i=0}^{\infty} \gamma_{\ell,i} x_{r-i}^{\ell}, \quad \gamma_{\ell,i} \in \mathbb{F}_p,$$

it is clear that any  $w = (w_0, w_1, \dots, w_a) \in W_a(\mathbb{F}_p(r, \infty))$  admits a unique expression as a sum

$$\sum_{b=0}^a \sum_{\ell=0}^{p-1} \sum_{i=0}^{\infty} [\gamma_{\ell,b,i}] (0, \dots, 0, w_b = x_{r-i}^{\ell}, 0, \dots, 0),$$

which in turn converges in the prodiscrete topology of  $W_a(\mathbb{F}_p(r, \infty))$ . More precisely, for any  $a, h = 0, 1, \dots$ , we can determine coefficients  $c_{\ell,b,i} \in \mathbb{Z}_p$  such that

$$w - \sum_{b=0}^a \sum_{\ell=0}^{p-1} \sum_{i=0}^h c_{\ell,b,i} x_{r-i}^{\ell}$$

has zero image in  $W_a(\mathbb{F}_p(r, h))$ . So, the function

$$f_{r,a,h} := \sum_{b=0}^a \sum_{\ell=0}^{p-1} \sum_{i=0}^h c_{\ell,b,i} p^b \Psi(p^{i-r} x) \ell p^{a-b}$$

is such that

$$\min\{v_p(f_{r,a,h}(x) - f(x)) \mid x \in p^{r-h}\mathbb{Z}_p + p^r\mathbb{C}_p^\circ\} \geq a + 1.$$

Finally, we already observed that the kernel of the map (38) consists of the elements  $g \in \mathcal{E}_{r,a}^\circ$  such that  $-\log\|g\|_{p^{h-r}\mathbb{Z}_p} \geq a + 1$ . This proves (2), (3) and (4).

As for the last part of the statement, we pick any  $f \in \mathcal{C}_{\text{unif}}(\mathbb{Q}_p, \mathbb{Z}_p)$  and a natural number  $N = 0, 1, \dots$ . Then there exists an  $M = 0, 1, \dots$  and  $f_M \in \mathcal{C}(\mathbb{Q}_p/p^M\mathbb{Z}_p, \mathbb{Z}_p)$  such that  $w_\infty(f - f_M) \geq N$ . It will suffice to determine an element  $g \in \mathbb{Z}_p[\Psi(\lambda x) \mid \lambda \in \mathbb{Q}_p^\times]$  such that  $w_\infty(g - f_M) \geq N$ . We then pick  $r \in \mathbb{Z}$  and  $a \in \mathbb{Z}_{\geq 0}$  so that  $r + 1 \geq M$  and  $a + 1 \geq N$ . The statement follows from the surjectivity of (38). This concludes the proof.  $\square$

As a corollary, we obtain the proof of Propositions 2.23 and 2.24. We now give the proof of Proposition 2.21.

*Proof of Proposition 2.21.* We discuss  $(\mathcal{E}_\lambda^\circ, \text{standard})$  in order to fix ideas. The case of  $(\mathcal{T}_\lambda^\circ, \text{standard})$  is analogous. The coproduct of  $\mathcal{E}_\lambda^\circ$  originates from (7), which we spell out as

$$\begin{aligned} x \mapsto & \Psi(x \widehat{\otimes}_{\mathbb{Z}_p} 1 + 1 \widehat{\otimes}_{\mathbb{Z}_p} x) \\ & = \Phi(\Psi(x \widehat{\otimes}_{\mathbb{Z}_p} 1), \Psi(px \widehat{\otimes}_{\mathbb{Z}_p} 1), \dots; \Psi(1 \widehat{\otimes}_{\mathbb{Z}_p} x), \Psi(1 \widehat{\otimes}_{\mathbb{Z}_p} px), \dots) \\ & = \Phi(\Psi(x) \widehat{\otimes}_{\mathbb{Z}_p} 1, \Psi(px) \widehat{\otimes}_{\mathbb{Z}_p} 1, \dots; 1 \widehat{\otimes}_{\mathbb{Z}_p} \Psi(x), \widehat{\otimes}_{\mathbb{Z}_p} \Psi(1px), \dots), \end{aligned}$$

and the identification (10). The fact that  $\mathcal{E}_\lambda^\circ$  only depends upon  $|\lambda|$  follows from the fact that, for any  $f \in \mathbb{C}\{x\}$ , the map  $\mathbb{Q}_p \rightarrow \mathbb{C}\{x\}$ ,  $a \mapsto f(ax)$ , is continuous. For any  $n \in \mathbb{Z}$ , the map  $n\iota: \Psi(\lambda^{-1}p^j x) \mapsto \Psi(\lambda^{-1}p^j nx)$ , for any  $j = 0, 1, \dots$ , is an endomorphism of  $\mathcal{E}_\lambda^\circ$ . By continuity, we obtain a map  $a\iota: \mathcal{E}_\lambda^\circ \rightarrow \mathcal{E}_\lambda^\circ$ , for any  $a \in \mathbb{Z}_p$ . If  $m, n \in \mathbb{Z}$  are such that  $mn = 1 + ap^N$ , for  $a \in \mathbb{Z}$  and  $N \in \mathbb{Z}$ ,  $N \gg 0$ , then  $\Psi(\lambda^{-1}p^j mnx)$  is close to  $\Psi(\lambda^{-1}p^j x)$ . Again by continuity we find that if  $a \in \mathbb{Z}_p^\times$ ,  $a\iota$  is an automorphism of  $\mathcal{E}_\lambda^\circ$ .  $\square$

We finally prove our uniform approximation Theorem 2.27.

*Proof.* We discuss the integral case only; the bounded case follows directly. We first observe that a  $\mathcal{CLM}_{\mathbb{Z}_p}^u$ -morphism

$$(\mathcal{APH}_{0, \mathbb{Z}_p}, \text{strip}) = \varinjlim_{\rho \rightarrow 0}^u (\mathcal{APH}_{\mathbb{Z}_p}(\Sigma_\rho), \text{strip}) \rightarrow (AP_{\mathbb{Z}_p}, w_\infty)$$

exists because so does, for any  $\rho > 0$ , the morphism  $(\mathcal{APH}_{\mathbb{Z}_p}(\Sigma_\rho), \text{strip}) \rightarrow (AP_{\mathbb{Z}_p}, w_\infty)$ . Moreover, that morphism is injective. An element of  $\mathcal{APH}_{0, \mathbb{Z}_p}$  is

represented by a sequence  $P_{\rho_n} \in \mathbb{Z}_p[\Psi(x/\lambda) \mid \lambda \in \mathbb{Q}_p^\times]$  with  $\rho_n$  decreasing to 0, such that for any  $\varepsilon > 0$ , there exists  $N_\varepsilon$  such that for any  $m \geq n \geq N_\varepsilon$ ,

$$\|P_{\rho_n} - P_{\rho_m}\|_{\mathbb{Q}_p, \rho_m} < \varepsilon.$$

Let  $f \in AP_{\mathbb{Z}_p}$  and let  $N \in \mathbb{Z}_{>0}$ . By the definition of u.a.p. functions, there exists a polynomial

$$P_N := \sum_{\lambda \in \mathbb{Q}_p^\times} a_\lambda \Psi(x/\lambda),$$

where  $a_\lambda \in \mathbb{Z}_p = 0$  for almost all  $\lambda$ , such that

$$w_\infty(f - P_N) > N.$$

By (28) of Theorem 4.14, for any  $N > 0$ , there exists  $\rho_N > 0$  such that  $v_p(P_N(a+x) - P_N(a)) > N$  for any  $a \in \mathbb{Q}_p$  and  $x \in \mathbb{C}_p$ ,  $|x| \leq \rho_N$ . We may assume that the sequence  $N \rightarrow \rho_N$  decreases to 0. We deduce that for  $M \geq N$ ,

$$-\log\|P_N - P_M\|_{\mathbb{Q}_p, \rho_M} > N.$$

So, the sequence  $N \mapsto P_N$  represents a germ  $P \in \mathcal{APH}_{0, \mathbb{Z}_p}$  whose restriction to  $\mathbb{Q}_p$  is  $f$ . □

**Remark 6.8.** We are not asserting here that there should be a  $p$ -adic strip  $\Sigma_\rho$  around  $\mathbb{Q}_p$  on which  $f$  extends analytically. In fact, an inductive limit in the category  $\mathcal{CLM}_{\mathbb{Z}_p}^u$  is not necessarily supported by a set-theoretic inductive limit (see Appendix A.1 below) and similarly for a locally convex inductive limit of Banach spaces.

#### APPENDIX A. NON ARCHIMEDEAN TOPOLOGICAL ALGEBRA

A prime number  $p$  is fixed throughout this paper and  $q = p^f$  is a power of  $p$ . So,  $\mathbb{Q}_q$  will denote the unramified extension of  $\mathbb{Q}_p$  of degree  $f$ , and  $\mathbb{Z}_q$  will be its ring of integers. Unless otherwise specified, a *ring* is meant to be commutative with 1.

**A.1. Linear topologies.** Let  $k$  be a separated and complete linearly topologized ring; we will denote by  $\mathcal{P}(k)$  the family of open ideals of  $k$ . We will consider the category  $\mathcal{CLM}_k^u$  of separated and complete linearly topologized  $k$ -modules  $M$  such that the map multiplication by scalars

$$k \times M \rightarrow M, \quad (r, m) \mapsto rm,$$

is *uniformly continuous* for the product uniformity of  $k \times M$ . Morphisms of  $\mathcal{CLM}_k^u$  are continuous  $k$ -linear maps. This is the classical category of [12, Ch. III, §2]. See [4] for more details.

**Remark A.2.** All over this paper we will assume that in a topological ring  $R$  (resp. topological  $R$ -module  $M$ ), the product (resp. the scalar product) map  $R \times R \rightarrow R$  (resp.  $R \times M \rightarrow M$ ) is at least continuous for the product topology of  $R \times R$  (resp. of  $R \times M$ ); morphisms will be continuous morphisms of rings (resp. of  $R$ -modules).



By a *nonarchimedean* (*n.a.*) ring  $R$  (resp.  $R$ -module  $M$ ) we mean a topological ring  $R$  (resp.  $R$ -module  $M$ ) equipped with a topology for which a basis of neighborhoods of 0 consists of additive subgroups and additive translations are homeomorphisms. So, any valued nonarchimedean field  $K$  is a n.a. ring in the previous sense and, if  $K$  is non-trivially valued, the category  $\mathcal{LC}_K$  of locally convex  $K$ -vector spaces [18] is a full subcategory of the category of n.a.  $K$ -modules. But, such a field  $K$  is never a linearly topologized ring. The ring of integers  $K^\circ$  is indeed linearly topologized, but no nonzero object of  $\mathcal{LC}_K$  is an object of  $\mathcal{CLM}_{K^\circ}^u$ .

**Definition A.3.** Let  $R$  be a topological ring and  $M$  be a topological  $R$ -module. A closed topological  $R$ -submodule  $N$  of  $M$  is said to be *strictly* closed if it is endowed with the subspace topology of  $M$ .

For any object  $M$  of  $\mathcal{CLM}_k^u$ ,  $\mathcal{P}(M)$  will denote the family of open  $k$ -submodules of  $M$ . The category  $\mathcal{CLM}_k^u$  admits all limits and colimits. The former are calculated in the category of  $k$ -modules but not the latter. So, a limit will be denoted by  $\varprojlim$  while a colimit will carry an apex  $(-)^u$  as in  $\varinjlim^u$ . In particular, for any family  $M_\alpha, \alpha \in A$ , of objects of  $\mathcal{CLM}_k^u$ , the direct sum and direct product will be denoted by

$$\bigoplus_{\alpha \in A}^u M_\alpha, \quad \prod_{\alpha \in A} M_\alpha,$$

respectively. We explicitly notice that  $\bigoplus_{\alpha \in A}^u M_\alpha$  is the completion of the algebraic direct sum  $\bigoplus_{\alpha \in A} M_\alpha$  of the algebraic  $k$ -modules  $M_\alpha$ 's, equipped with the  $k$ -linear topology for which a fundamental system of open  $k$ -modules consists of the  $k$ -submodules

$$\bigoplus_{\alpha \in A} (U_\alpha + IM_\alpha)$$

such that  $U_\alpha \in \mathcal{P}(M_\alpha)$  for all  $\alpha$ , and  $I \in \mathcal{P}(k)$  is independent of  $\alpha$ . Then the  $k$ -module underlying  $\bigoplus_{\alpha \in A}^u M_\alpha$  in general properly contains the algebraic direct sum  $\bigoplus_{\alpha \in A} M_\alpha$ . It will also be useful to introduce the *uniform box product* of the same family

$$\prod_{\alpha \in A}^{\square, u} M_\alpha,$$

which, set-theoretically, coincides with  $\prod_{\alpha \in A} M_\alpha$ , but whose family of open submodules consists of all  $U := \prod_{\alpha \in A} U_\alpha$ , with  $U_\alpha \in \mathcal{P}(M_\alpha)$ , such that there exists  $I_U \in \mathcal{P}(k)$  such that  $I_U M_\alpha \subset U_\alpha$  for any  $\alpha \in A$ . The category  $\mathcal{CLM}_k^u$ , equipped with the tensor product  $\widehat{\otimes}_k^u$  of [16, 0.7.7] (see also [12, Ch. III, §2, Exer. 28]) is a symmetric monoidal category. The category of monoids of  $\mathcal{CLM}_k^u$  is denoted by  $\mathcal{ACL M}_k^u$ .

For two objects  $M$  and  $N$  of  $\mathcal{CLM}_k^u$ , we have

$$M \widehat{\otimes}_k^u N = \varprojlim_{P \in \mathcal{P}(M), Q \in \mathcal{P}(N)} M/P \otimes_k N/Q,$$

so that  $\widehat{\otimes}_k^u$  commutes with filtered projective limits in  $\mathcal{CLM}_k^u$ .

**A.4. Semivaluations.** We describe here full subcategories of  $\mathcal{CLM}_k^u$ , and special base rings  $k$ , of most common use. We denote by  $\mathbb{Z}_{(p)} = \mathbb{Q} \cap \mathbb{Z}_p$ , the localization of  $\mathbb{Z}$  at  $(p)$ . Then  $\mathbb{C}_p$  will be the completion of a fixed algebraic closure of  $\mathbb{Q}_p$ . On  $\mathbb{C}_p$ , we use the absolute value  $|x| = |x|_p = p^{-v_p(x)}$  for the  $p$ -adic valuation  $v = v_p$ , with  $v_p(p) = 1$ , and  $x \in \mathbb{C}_p$ .

**Definition A.5.** A *semivaluation* on a ring  $R$  is a map  $w: R \rightarrow \mathbb{R} \cup \{+\infty\}$  such that  $w(0) = +\infty$ ,  $w(x+y) \geq \min(w(x), w(y))$  and  $w(xy) \geq w(x) + w(y)$  for any  $x, y \in R$ . We will say that a semivaluation is *positive* if it takes its values in  $\mathbb{R}_{\geq 0} \cup \{+\infty\}$ .

**Remark A.6.**

- (1) If  $w_1, w_2, \dots, w_n$  are a finite set of semivaluations on the ring  $R$ , so is their infimum

$$w := \inf_{i=1, \dots, n} w_i.$$

- (2) The *trivial valuation*  $v_0: R \rightarrow \{0, +\infty\}$ , which exists on any ring  $R$ , is (in our sense!) a positive semivaluation.
- (3) We will indifferently use the multiplicative notation  $|x|_w = \exp(-w(x))$ .

For any semivaluation  $w$  of a ring  $R$ , the family

$$R_{w,c} := \{x \in R \mid w(x) \geq c\}$$

for  $c \in \mathbb{R}$  is a fundamental set of open subgroups for a group topology of  $R$ . Moreover,  $R_{w,0}$  is a subring of  $R$  and all  $R_{w,c}$  are  $R_{w,0}$ -submodules of  $R$ . A (*multi-*) *semivalued ring*  $(R, \{w_\alpha\}_{\alpha \in A})$  is a ring  $R$  equipped with a family  $\{w_\alpha\}_{\alpha \in A}$  of semivaluations. A semivalued ring is endowed with the topology in which any  $x \in R$  has a fundamental system of neighborhoods consisting of the subsets

$$x + \bigcap_{\alpha \in F} R_{\alpha, c_\alpha},$$

where  $F$  varies among finite subsets of  $A$  and, for any  $\alpha \in F$ ,  $c_\alpha \in \mathbb{R}$ . A *Fréchet ring* (resp. *Banach ring*) is a ring  $R$  which is separated and complete in the topology induced by a countable family of semivaluations (resp. by a single semivaluation). If the semivaluations  $w_\alpha$  are all positive, the Fréchet (resp. Banach) ring  $(R, \{w_\alpha\}_{\alpha \in A})$  is linearly topologized. We will call it a *linearly topologized Fréchet* (resp. *Banach*) *ring*. When  $R$  is an algebra over a Banach ring  $(S, v)$ , and the semivaluations  $w_\alpha$  satisfy

$$w_\alpha(xy) = v(x) + w_\alpha(y) \quad \text{for all } x \in S, y \in R,$$

we also say that  $R = (R, \{w_\alpha\}_{\alpha \in A})$  is a *Fréchet* (resp. *Banach*) *S-algebra*. In the particular case when  $(S, v)$  is a complete non-trivially valued real-valued field  $(K, v)$ , a Fréchet or Banach *S-algebra* is a Fréchet or Banach algebra over  $K$  in the classical sense. Notice however that we allow  $v$  to be the trivial valuation of  $S$  or  $K$ . We denote by  $\mathcal{CLC}_K$  the category of locally convex topological  $K$ -vector spaces of [18], where morphisms are continuous  $K$ -linear maps, which are moreover separated and complete.

**Lemma A.7.** *Let  $(S, v)$  be a Banach ring and let  $(R, \{w_n\}_{n=1,2,\dots})$  be a Fréchet  $S$ -algebra. Let  $(R_n, w_n)$  be the separated completion of  $R$  in the locally convex topology induced by the semivaluation  $w_n$ . Assume  $w_n(r) \geq w_m(r)$  for any  $r \in R$  if  $n \leq m$ . Then the identity of  $R$  extends to a morphism  $R_m \rightarrow R_n$  of Banach  $S$ -algebras and  $R$  is the limit, in the category of n.a.  $S$ -algebras, of the filtered projective system  $(R_n)_n$ .*

*In particular, a  $S$ -subalgebra  $T$  of  $R$  is dense in  $R$  if and only if it is dense in  $R_n$  for any  $n$ .*

**A.8. Tensor products.** Let  $(S, v)$  be a complete real-valued ring and let  $R = (R, \{w_\alpha\}_{\alpha \in A})$  and  $R' = (R', \{w'_\beta\}_{\beta \in B})$  be two Fréchet  $S$ -algebras. Then we define a Fréchet  $S$ -algebra  $R \widehat{\otimes}_{\pi, S} R'$  as the completion of the  $S$ -algebra  $R \otimes_S R'$  in the topology induced by the following semivaluations [11, §2.1.7], for any  $(\alpha, \beta) \in A \times B$ ,

$$w_{\alpha, \beta}(g) = \sup \left( \min_{1 \leq i \leq n} w_\alpha(x_i) + w'_\beta(y_i) \right),$$

where the supremum runs over all possible representations

$$g = \sum_{i=1}^n x_i \otimes y_i, \quad x_i \in R, \quad y_i \in R'.$$

The following proposition follows immediately from Lemma A.7.

**Proposition A.9.** *Let  $(S, v)$  be a Banach ring and let  $(R, \{w_n\}_{n=1,2,\dots})$ ,  $(R', \{w'_n\}_{n=1,2,\dots})$  be two Fréchet  $S$ -algebra satisfying the assumption of Lemma A.7. Then, with the same notation,  $R \widehat{\otimes}_{\pi, S} R'$  is the limit, in the category of n.a.  $S$ -algebras, of the filtered projective system of Banach  $S$ -algebras  $(R_n \widehat{\otimes}_{\pi, S} R'_n)_n$ .*

Notice that

- (1) if  $R$  and  $R'$  are Fréchet algebras over a complete real-valued field  $(K, v)$ , with nontrivial valuation  $v$ ,  $R \widehat{\otimes}_{\pi, K} R'$  coincides with both the completed projective and the inductive tensor product of [18] (cf. Lemma 17.2 and Lemma 17.6 of *loc. cit.*);
- (2) if  $R$  and  $R'$  are linearly topologized Fréchet algebras over a linearly topologized Banach ring  $(S, v)$ ,  $R \widehat{\otimes}_{\pi, S} R'$  coincides with  $R \widehat{\otimes}_S^u R'$ .

#### APPENDIX B. CLASSICAL THEORY OF ALMOST PERIODIC FUNCTIONS

The main character of this paper, our function  $\Psi$ , shows many analogies with the classical holomorphic almost periodic functions of Bohr, Bochner, and Besicovitch [9]. In fact, many of the subtle function theoretic difficulties which appear in the  $p$ -adic setting are also encountered in classical Harmonic Analysis. We feel that a short presentation of the basics of the classical theory might be useful. See also the survey article [14].

**B.1. Fejér's Theorem.** Let  $(\mathcal{C}_{\text{unif}}^{\text{bd}}(\mathbb{R}, \mathbb{R}), \|\cdot\|_{\mathbb{R}})$  be the Banach algebra of bounded uniformly continuous functions  $\mathbb{R} \rightarrow \mathbb{R}$ , equipped with the supnorm on  $\mathbb{R}$ . For  $\lambda \in \mathbb{R}_{>0}$ , let  $\mathcal{P}_{\mathbb{R}, \lambda} \subset \mathcal{C}_{\text{unif}}^{\text{bd}}(\mathbb{R}, \mathbb{R})$  be the strictly closed Banach subalgebra of continuous periodic functions of period  $\lambda$ .

Let us recall the classical *Fejér's theorem* [20, §13.31]. Let  $f \in \mathcal{P}_{\mathbb{R}, \lambda}$ . The *Fourier expansion* of  $f$  is the formal trigonometric series

$$\frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos\left(\frac{2\pi n}{\lambda} z\right) + b_n \sin\left(\frac{2\pi n}{\lambda} z\right),$$

with

$$a_n = \frac{2}{\lambda} \int_0^{\lambda} f(t) \cos\left(\frac{2\pi n}{\lambda} z\right) dt, \quad b_n = \frac{2}{\lambda} \int_0^{\lambda} f(t) \sin\left(\frac{2\pi n}{\lambda} z\right) dt.$$

The sequence of the partial sums

$$S_N(f) = \frac{a_0}{2} + \sum_{n=1}^N a_n \cos\left(\frac{2\pi n}{\lambda} z\right) + b_n \sin\left(\frac{2\pi n}{\lambda} z\right),$$

does not necessarily converge to  $f$  uniformly on  $\mathbb{R}$ . However, the Cesaro means

$$\sigma_n = \frac{S_0 + \cdots + S_{n-1}}{n}$$

converge to  $f$  uniformly on  $\mathbb{R}$ . In particular, we have the following result.

**Theorem B.2.**  $\mathbb{R}[\cos(\frac{2\pi}{\lambda}x), \sin(\frac{2\pi}{\lambda}x)]$  is dense in the  $\mathbb{R}$ -Banach algebra  $(\mathcal{P}_{\mathbb{R}, \lambda}, \|\cdot\|_{\mathbb{R}})$ .

We will show below that a suitably reformulated  $p$ -adic analog of Theorem B.2 holds true  $p$ -adically.

**Definition B.3** (Bohr's definition of u.a.p. functions). A continuous function  $f: \mathbb{R} \rightarrow \mathbb{R}$  is *uniformly almost periodic* (u.a.p. for short) if, for any  $\varepsilon > 0$ , there exists  $\ell_{\varepsilon} > 0$  such that for any interval  $I \subset \mathbb{R}$  of length  $\ell_{\varepsilon}$ , there exists  $\tau \in I$  such that

$$|f(x + \tau) - f(x)| < \varepsilon \quad \text{for all } x \in \mathbb{R}.$$

It is easy to check that the set of uniformly almost periodic functions  $\mathbb{R} \rightarrow \mathbb{R}$  is a closed subalgebra  $AP_{\mathbb{R}}$  of  $(\mathcal{C}_{\text{unif}}^{\text{bd}}(\mathbb{R}, \mathbb{R}), \|\cdot\|_{\mathbb{R}})$  [9, Ch. I, §1, Thm. 4°, 5°]. We define  $AP_{\mathbb{C}} \subset (\mathcal{C}_{\text{unif}}^{\text{bd}}(\mathbb{R}, \mathbb{C}), \|\cdot\|_{\mathbb{R}})$  similarly.

The following result is Bohr's "approximation theorem". We refer to [9, §I.5] for its proof and for a detailed description of the contributions of S. Bochner and H. Weyl.

**Theorem B.4.**  $(AP_{\mathbb{R}}, \|\cdot\|_{\mathbb{R}})$  identifies with the completion of the normed ring

$$\left(\mathbb{R}\left[\cos\left(\frac{2\pi}{\lambda}x\right), \sin\left(\frac{2\pi}{\lambda}x\right) \mid \lambda \in \mathbb{R}^{\times}\right], \|\cdot\|_{\mathbb{R}}\right).$$

Similarly for  $(AP_{\mathbb{C}}, \|\cdot\|_{\mathbb{R}})$ .

We propose  $p$ -adic analogs of those Banach algebras and of the latter theorem.

**B.5. Dirichlet series.** Let  $\mathbb{C}\{x\}$  be the Fréchet  $\mathbb{C}$ -algebra of entire functions  $\mathbb{C} \rightarrow \mathbb{C}$ , equipped with the topology of uniform convergence on compact subsets of  $\mathbb{C}$ . The rotation  $z \mapsto iz$  transforms trigonometric series into series of exponentials and Bohr's definition naturally propagates into the following.

**Definition B.6** ([9, §III.2, Thm. 1°]). For any interval  $(a, b) \subset \mathbb{R}$ , an analytic function  $f$  on the strip  $(a, b) \times i\mathbb{R} \subset \mathbb{C}$  is *almost periodic holomorphic* on  $(a, b)$  if, for any  $\varepsilon > 0$ , there exists  $\ell_\varepsilon > 0$  such that for any interval  $I \subset \mathbb{R}$  of length  $\ell_\varepsilon$ , there exists  $\tau \in I$  such that

$$|f(x + i\tau) - f(x)| < \varepsilon \quad \text{for all } x \in (a, b) \times i\mathbb{R}.$$

We let  $\mathcal{APH}_{\mathbb{C}}((a, b))$  denote the  $\mathbb{C}$ -algebra of almost periodic holomorphic functions on  $(a, b)$ .

Notice that  $\mathcal{APH}_{\mathbb{C}}((a, b))$  is a closed subalgebra of the Fréchet algebra  $\mathcal{O}((a, b) \times i\mathbb{R})$ ; the induced Fréchet algebra structure is called *standard*. We may equip  $\mathcal{APH}_{\mathbb{C}}((a, b))$  with the finer Fréchet algebra structure of uniform convergence on substrips  $(a', b') \times i\mathbb{R}$ , for  $a < a' < b' < b$ . We informally call this topology *the strip topology*.

The following polynomial approximation theorem [9, §III.3, Thm. 3°] holds.

**Theorem B.7.**  $\mathcal{APH}_{\mathbb{C}}((a, b))$  is the Fréchet completion of the  $\mathbb{C}$ -polynomial algebra generated by the restrictions to  $(a, b) \times i\mathbb{R}$  of all continuous characters of  $\mathbb{R}$ , namely, by the maps

$$e_\lambda: (a, b) \times i\mathbb{R} \rightarrow \mathbb{C}, \quad z \mapsto e^{\lambda z},$$

for  $\lambda \in \mathbb{R}^\times$ , equipped with the strip topology.

The assignment  $(a, b) \mapsto \mathcal{APH}_{\mathbb{C}}((a, b))$  uniquely extends to a sheaf of Fréchet  $\mathbb{C}$ -algebras on  $\mathbb{R}$ .

**Definition B.8.**

- (1) We denote by  $\mathcal{APH}_{0, \mathbb{C}}$  the stalk of the sheaf  $\mathcal{APH}_{\mathbb{C}}$  at 0 equipped with the locally convex inductive limit topology of the system of Fréchet algebras  $\mathcal{APH}_{\mathbb{C}}((-\varepsilon, \varepsilon))$  as  $\varepsilon \rightarrow 0^+$ .
- (2) We denote by  $\mathcal{APH}_{\mathbb{C}} \subset \mathbb{C}\{x\}$  the Fréchet algebra of global sections of  $\mathcal{APH}_{\mathbb{C}}$  equipped with the strip topology.

Notice that we have a natural injective morphism, induced by restriction of functions and the properties of the inductive limit

$$\mathcal{APH}_{\mathbb{C}} \rightarrow \mathcal{APH}_{0, \mathbb{C}}.$$

The next corollary follows from the combined theorems of approximation Theorem B.4 and Theorem B.7.

**Corollary B.9.**  $(\mathcal{AP}_{\mathbb{C}}, \| \cdot \|_{\mathbb{R}})$  identifies with the completion of the normed ring  $(\mathcal{APH}_{0, \mathbb{C}}, \| \cdot \|_{\mathbb{R}})$ .

**Remark B.10.** Sections of the sheaf  $\mathcal{AP}\mathcal{H}_{\mathbb{C}}$  on open subsets of  $\mathbb{R}$  may be viewed as generalized Dirichlet series [9, §III.3]. A  $p$ -adic analog on  $\mathbb{Q}_p$  of the sheaf  $\mathcal{AP}\mathcal{H}_{\mathbb{C}}$  of Dirichlet series on  $\mathbb{R}$ , might be useful in the theory of  $p$ -adic L-functions.

### APPENDIX C. NUMERICAL CALCULATIONS BY M. CANDILERA

The following calculations were performed with *Mathematica*<sup>©</sup>. We computed the first coefficients of the series  $\Psi_p(T) = \sum_{n=1}^{\infty} b_n T^n$ , for  $p = 2$ , up to the term of degree  $2^5$ , and for  $p = 3$ , up to degree  $3^4$ . We also evaluated the a few coefficients of  $\Psi_5(T)$  and  $\Psi_7(T)$ . We give here tables of the  $p$ -adic orders of the coefficients  $b_n$  for  $p = 2, 3$ . For those values of  $p$ , we also draw the graph of the function  $n \mapsto v_p(b_n)$  and compare it with the Newton polygon of  $\Psi_p$  (flipped around the  $y$ -axis). We confirm experimentally the calculation of the corresponding valuation polygons.

**C.1. Very first coefficients.** For  $p = 2$ , we have

$$\begin{aligned} \Psi_2(T) = & T - 2 \cdot T^2 + 2^4 \cdot T^3 - 11 \cdot 2^5 \cdot T^4 + 7 \cdot 2^{11} \cdot T^5 \\ & - 7 \cdot 37 \cdot 2^{12} \cdot T^6 + 3 \cdot 751 \cdot 2^{16} \cdot T^7 - 301627 \cdot 2^{17} \cdot T^8 \\ & + 308621 \cdot 2^{26} \cdot T^9 + 2^{27} \cdot T^{10} \cdot u(T), \end{aligned}$$

for a unit  $u(T) \in \mathbb{Z}_{(2)}[[T]]^{\times}$ .

For  $p = 3$ ,

$$\begin{aligned} \Psi_3(T) = & T - 3^2 \cdot T^3 + 3^7 \cdot T^5 - 2^2 \cdot 7 \cdot 3^{11} \cdot T^7 + 2 \cdot 7 \cdot 13 \cdot 113 \cdot 3^{14} \cdot T^9 \\ & - 5 \cdot 89 \cdot 1249 \cdot 3^{22} \cdot T^{11} + 5 \cdot 117 \cdot 217667 \cdot 3^{28} \cdot T^{13} + \dots \end{aligned}$$

For  $p = 5$ ,

$$\begin{aligned} \Psi_5(T) = & T - 5^4 \cdot T^5 + 5^{13} \cdot T^9 - 53 \cdot 59 \cdot 5^{21} \cdot T^{13} \\ & + 3 \cdot 11 \cdot 97 \cdot 1123 \cdot 1699 \cdot 5^{29} \cdot T^{17} + 5^{37} \cdot T^{21} \cdot u(T), \end{aligned}$$

for a unit  $u(T) \in \mathbb{Z}_{(5)}[[T]]^{\times}$ .

For  $p = 7$ ,

$$\Psi_7(T) = T - 7^6 \cdot T^7 + 7^{19} \cdot T^{13} - 2 \cdot 31 \cdot 37 \cdot 359 \cdot 7^{31} \cdot T^{19} + 7^{43} \cdot T^{25} \cdot u(T),$$

for a unit  $u(T) \in \mathbb{Z}_{(7)}[[T]]^{\times}$ .

C.2. First 24 coefficients of  $\Psi_2(t)$  and 2-adic order of the 32 first.




FIGURE 5. Newton and valuation polygons of  $\Psi_2$ .

$\Psi_2(t) = \sum_{n \geq 1} b_n t^n$							
$b_1$	0	$b_9$	26	$b_{17}$	61	$b_{25}$	101
$b_2$	1	$b_{10}$	27	$b_{18}$	62	$b_{26}$	102
$b_3$	4	$b_{11}$	33	$b_{19}$	70	$b_{27}$	110
$b_4$	5	$b_{12}$	34	$b_{20}$	71	$b_{28}$	111
$b_5$	11	$b_{13}$	42	$b_{21}$	81	$b_{29}$	121
$b_6$	12	$b_{14}$	43	$b_{22}$	82	$b_{30}$	122
$b_7$	16	$b_{15}$	48	$b_{23}$	89	$b_{31}$	128
$b_8$	17	$b_{16}$	49	$b_{24}$	90	$b_{32}$	129

$$\begin{aligned}
 b_1 &= 1, & b_2 &= -2, & b_3 &= 16 = 2^4, & b_4 &= -352 = -2^5 \cdot 11, \\
 b_5 &= 14336 = 2^{11} \cdot 7, & b_6 &= -1060864 = -2^{12} \cdot 7 \cdot 37, \\
 b_7 &= 147652608 = 2^{16} \cdot 3 \cdot 751, \\
 b_8 &= -39534854144 = -2^{17} \cdot 301627, \\
 b_9 &= 20711204716544 = 2^{26} \cdot 308621, \\
 b_{10} &= -21454855889485824 = -2^{27} \cdot 3^2 \cdot 13 \cdot 701 \cdot 1949, \\
 b_{11} &= 44195700516541431808 = 2^{33} \cdot 5145056699, \\
 b_{12} &= -181554407879323198423040 = -2^{34} \cdot 5 \cdot 41 \cdot 2273 \cdot 22679509, \\
 b_{13} &= 1489469015852141109009448960 = 2^{42} \cdot 5 \cdot 67733208918623, \\
 b_{14} &= -24421319844213105128638664146944 = -2^{43} \cdot 3^2 \cdot 8179 \cdot 37716952983613, \\
 b_{15} &= 800530746908074643997623203521363968 = 2^{48} \cdot 31 \cdot 71 \cdot 1619 \cdot 826201 \cdot 966018887,
 \end{aligned}$$

$$\begin{aligned}
 b_{16} &= -52473187457503996327647036404796036743168 \\
 &= -2^{49} \cdot 31 \cdot 397 \cdot 13687 \cdot 2882489 \cdot 191972726039, \\
 b_{17} &= 6878395240848057051122842718175351390427152384 \\
 &= 2^{61} \cdot 3 \cdot 47 \cdot 59 \cdot 919 \cdot 24709 \cdot 15791216459521333, \\
 b_{18} &= -1803212578568825704559863338710346864852507172012032 \\
 &= 2^{62} \cdot 3^2 \cdot 19 \cdot 97 \cdot 173 \cdot 1665967 \cdot 581220517 \cdot 140723269997, \\
 b_{19} &= 945424354393817092018179744741353462710753588534117924864 \\
 &= 2^{70} \cdot 7^2 \cdot 23 \cdot 15973 \cdot 44485316159805664956515547941, \\
 b_{20} &= -991360632780906301560343330625129510790528483073480047449866240 \\
 &= 2^{71} \cdot 5 \cdot 167 \cdot 14503 \cdot 15445577653440901 \cdot 2244675152281633901, \\
 b_{21} &= 2079045830009718214618472297232655379089817022368004517660824096997376 \\
 &= 2^{81} \cdot 109 \cdot 23549 \cdot 167442376921 \cdot 2000645152343730624200879183, \\
 b_{22} &= -8720175189463740580963423057535032711261236371520206719551905031269050744832 \\
 &= 2^{82} \cdot 47 \cdot 1867 \cdot 105323 \cdot 2119591 \cdot 80618233393589 \cdot 1141865166972250409671, \\
 b_{23} &= 73150235997673008411264495083486904164758556563477195586370441676376428384144588800 \\
 &= 2^{89} \cdot 3^3 \cdot 5^2 \cdot 175082340917111384848376265817809832605816887352831773, \\
 b_{24} &= -1227258187586069935509530355473988020883482157853428276444146736521211077001846045664083968 \\
 &= 2^{90} \cdot 54617 \cdot 76121647308197 \cdot 238451637287968840726339672350427699951944293.
 \end{aligned}$$

C.3. 3-adic values of the first 81 coefficients of  $\Psi_3(T)$ .




FIGURE 6. The Newton and valuation polygons of  $\Psi_3$ .



$b_3$	2	$b_{23}$	58	$b_{43}$	135	$b_{63}$	213
$b_5$	7	$b_{25}$	64	$b_{45}$	141	$b_{65}$	223
$b_7$	11	$b_{27}$	68	$b_{47}$	151	$b_{67}$	231
$b_9$	14	$b_{29}$	79	$b_{49}$	159	$b_{69}$	238
$b_{11}$	22	$b_{31}$	87	$b_{51}$	166	$b_{71}$	247
$b_{13}$	28	$b_{33}$	94	$b_{53}$	175	$b_{73}$	255
$b_{15}$	33	$b_{35}$	103	$b_{55}$	183	$b_{75}$	262
$b_{17}$	40	$b_{37}$	111	$b_{57}$	190	$b_{77}$	271
$b_{19}$	46	$b_{39}$	118	$b_{59}$	199	$b_{79}$	279
$b_{21}$	51	$b_{41}$	127	$b_{61}$	207	$b_{81}$	284

**Acknowledgements.** It is a pleasure to acknowledge that the proofs in Sections 3 and 4 of this text are based on a discussion with Philippe Robba which took place in April 1980. I am strongly indebted to him for this and for his friendship. I thank my colleague Giuseppe De Marco for his patient explanations on classical Fourier theory. I thank the MPIM of Bonn for hospitality during March 2018, when this article was completed.

#### REFERENCES

- [1] F. Baldassarri, *Interpretazione funzionale di certe iperalgebre e dei loro anelli di bivettori di Witt*, Tesi di laurea, Padova, 1974.
- [2] F. Baldassarri, Una funzione intera  $p$ -adica a valori interi, *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)* **2** (1975), no. 2, 321–331. MR0412157
- [3] F. Baldassarri, The Artin–Hasse isomorphism of perfectoid open unit disks and a Fourier-type theory for continuous functions on  $\mathbb{Q}_p$ , preprint.
- [4] F. Baldassarri and M. Cailotto, Closed exact categories of modules over generalized adic rings, preprint.
- [5] F. Baldassarri and M. Cailotto, Duality for functions and measures on locally compact 0-dimensional spaces with values in generalized adic rings, preprint.
- [6] I. Barsotti, Metodi analitici per varietà abeliane in caratteristica positiva. Capitoli I, II, *Ann. Scuola Norm. Sup. Pisa (3)* **18** (1964), 1–25. MR0168566
- [7] I. Barsotti, Metodi analitici per varietà abeliane in caratteristica positiva. Capitolo III, IV, *Ann. Scuola Norm. Sup. Pisa (3)* **19** (1965), 277–330. MR0209292
- [8] V. G. Berkovich, *Spectral theory and analytic geometry over non-Archimedean fields*, *Mathematical Surveys and Monographs*, 33, American Mathematical Society, Providence, RI, 1990. MR1070709
- [9] A. S. Besicovitch, *Almost periodic functions*, Dover Publications, Inc., New York, 1955. MR0068029
- [10] H. Bohr, Zur Theorie der Fastperiodischen Funktionen, *Acta Math.* **46** (1925), no. 1–2, 101–214. MR1555201
- [11] S. Bosch, U. Güntzer, and R. Remmert, *Non-Archimedean analysis*, *Grundlehren der Mathematischen Wissenschaften*, 261, Springer-Verlag, Berlin, 1984. MR0746961
- [12] N. Bourbaki, *Éléments de mathématique*, reprint, Masson, Paris, 1985. MR0782297
- [13] P. Colmez, Fonctions d’une variable  $p$ -adique, *Astérisque* No. 330 (2010), 13–59. MR2642404
- [14] R. L. Cooke, Almost-periodic functions, *Amer. Math. Monthly* **88** (1981), no. 7, 515–526. MR0628019
- [15] J.-M. Fontaine, *Groupes  $p$ -divisibles sur les corps locaux*, Société Mathématique de France, Paris, 1977. MR0498610

- [16] A. Grothendieck and J. A. Dieudonné, *Eléments de géométrie algébrique. I*, Grundlehren der Mathematischen Wissenschaften, 166, Springer-Verlag, Berlin, 1971. MR3075000
- [17] M. Lazard, Les zéros des fonctions analytiques d'une variable sur un corps valué complet, Inst. Hautes Études Sci. Publ. Math. No. 14 (1962), 47–75. MR0152519
- [18] P. Schneider, *Nonarchimedean functional analysis*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2002. MR1869547
- [19] P. Schneider, *Galois representations and  $(\varphi, \Gamma)$ -modules*, Cambridge Studies in Advanced Mathematics, 164, Cambridge University Press, Cambridge, 2017. MR3702016
- [20] E. C. Titchmarsh, *The theory of functions*, reprint of the second (1939) edition, Oxford University Press, Oxford, 1958. MR3155290

Received May 12, 2019; accepted May 24, 2019

Francesco Baldassarri  
Dipartimento di Matematica  
Università di Padova  
Via Trieste, 63, 35121 Padova, Italy

Maurizio Candilera  
Dipartimento di Matematica  
Università di Padova  
Via Trieste, 63, 35121 Padova, Italy