

^{52g}Mn production routes for multi-modal imaging applications

F. Barbaro^{1,2}, L. De Nardo^{1,3}, L. Meléndez-Alafort^{4,5}, L. Canton¹, M.P. Carante⁶, A. Colombi^{2,6}, A. Fontana⁶

¹ INFN, Sezione di Padova, Padova, Italy.

² Dipartimento di Fisica dell'Università di Pavia, Pavia, Italy.

³ Dipartimento di Fisica dell'Università di Padova, Padova, Italy.

⁴ Istituto Oncologico Veneto IOV IRCCS, Padova, Italy.

⁵ INFN, Laboratori Nazionali di Legnaro, Legnaro (Padova), Italy.

⁶ INFN, Sezione di Pavia, Pavia, Italy

The radionuclide ^{52g}Mn is of significant medical interest for the innovative PET-MRI multi-modal imaging technique. In this study we compare its standard cyclotron production route $^{nat}\text{Cr}(p,x)^{52g}\text{Mn}$ with the alternative reaction $^{nat}\text{V}(\alpha,x)^{52g}\text{Mn}$. The theoretical calculations are performed by a suitable tuning of the nuclear level density parameters of the TALYS reaction code, with the aim to obtain a good agreement with the experimental cross sections. The production route with ^{nat}V results in a more favorable radionuclidic purity than with ^{nat}Cr . Dosimetric studies are performed to establish the time frame in which ^{52g}Mn can be used with an acceptable dose to the patient.

Sezione 5

francesca.barbaro@pd.infn.it