

UNIVERSITÀ DEGLI STUDI DI PADOVA

DIPARTIMENTO DI FISICA “G. GALILEI”

Scuola di dottorato di ricerca in fisica, ciclo XXII

**Cosmological correlation functions in scalar and
vector inflationary models**

Direttore della Scuola:
Ch.mo Prof. Attilio Stella

Supervisori:
Ch.mo Prof. Sabino Matarrese
Dr. Nicola Bartolo

Dottoranda:
Emanuela Dimastrogiovanni

Contents

1	Introduction	4
2	Schwinger-Keldysh formalism	7
3	Scalar loop corrections to P_ζ	9
4	Perturbative expansion of the Lagrangian in $P(X, \phi)$ theories	12
4.1	Arnowitt-Deser-Misner (ADM) decomposition for $P(X, \phi)$ theories	13
4.1.1	Solving Hamiltonian and momentum constraint equations	15
4.1.2	Reduction to the canonical case	17
4.2	Fourth-order expansion of the Lagrangian in the canonical case	18
5	Tensor loop corrections to P_ζ	19
6	Complete expression for P_ζ at one loop	25
7	Study of perturbations in anisotropic cosmologies	27
8	Inflation and primordial vector fields	29
8.1	Self-coupled vector field models	30
8.2	Vector-field coupled to gravity	31
8.3	Ackerman-Carroll-Wise (ACW) model	31
8.4	Models with varying gauge coupling	32
8.5	$SU(2)$ vector model	33
9	Correlation functions of ζ in the $SU(2)$ model	35
9.1	The power spectrum	36
9.2	Bispectrum and trispectrum: Abelian contributions	38
9.3	Bispectrum and trispectrum: Non-Abelian contributions	40
9.3.1	Trispectrum from vector bosons: exchange diagram	43
9.3.2	Trispectrum from vector bosons: point-interaction diagram	46
10	Amplitude of non-Gaussianity: f_{NL} and τ_{NL}	49
11	Shape of non-Gaussianity and statistical anisotropy features	53
11.1	Momentum dependence of the bispectrum and trispectrum	53
11.2	Features and level of anisotropy	55
12	Trispectrum for $f(\tau)$ models of gauge interactions	60
13	Conclusions	64
14	Acknowledgments	67

15 Appendices	68
15.1 Computation of the interaction Hamiltonian	68
15.2 Study of leading slow roll order vertices in the fourth order action	70
15.3 Complete expressions of one-loop two-vertex diagrams to leading order .	72
15.4 Background and first order perturbation equations for the gauge fields . .	73
15.5 Calculation of the number of e-foldings of single-(scalar)field driven inflation in the presence of a vector multiplet	76
15.6 Complete expressions for the functions appearing in the bispectrum from quartic interactions	78
15.7 More details on computing the analytic expressions of vector-exchange diagrams	79
15.8 Complete expressions for functions appearing in point-interaction diagrams	80
16 References	82

to Matteo

1. Introduction

In the standard cosmological model, at very early times the Universe undergoes a quasi de Sitter exponential expansion driven by a scalar field, the inflaton, with an almost flat potential. The quantum fluctuations of this field are thought to be at the origin of both the Large Scale Structures and the Cosmic Microwave Background (CMB) fluctuations that we are able to observe at the present epoch [1]. CMB measurements indicate that the primordial density fluctuations are of order 10^{-5} , have an almost scale-invariant power spectrum and are fairly consistent with Gaussianity and statistical isotropy [2, 3, 4, 5, 6, 7, 8, 9]. All of these features find a convincing explanation within the inflationary paradigm. Nevertheless, deviations from the basic single-(scalar)field slow-roll model of inflation are allowed by experimental data. On one hand, it is then important to search for observational signatures that can help discriminate among all the possible scenarios; on the other hand, it is important to understand what the theoretical predictions are in this respect for the different models.

Non-Gaussianity and statistical anisotropy are two powerful signatures. A random field is defined “Gaussian” if it is entirely described by its two-point function, higher order connected correlators being equal to zero. Primordial non-Gaussianity [10, 11] is theoretically predicted by inflation: it arises from the interactions of the inflaton with gravity and from self-interactions. However, it is observably too small in the single-field slow-roll scenario [12, 13, 152]. Alternatives to the latter have been proposed that predict higher levels of non-Gaussianity such as multifield scenarios [15, 16, 17, 18, 19, 20, 21], curvaton models [22, 23, 24, 25, 26, 27] and models with non-canonical Lagrangians [28, 29, 30, 31, 32]. Many efforts have been directed to the study of higher order (three and four-point) cosmological correlators in these models [33, 34, 35, 36, 37, 31, 152, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48] and towards improving the prediction for the two-point function, through quantum loop calculations [49, 50, 51, 13, 52, 53, 54, 55, 56]. From WMAP, the bounds on the bispectrum amplitude are given by $-4 < f_{NL}^{loc} < 80$ [8] and by $-125 < f_{NL}^{equil} < 435$ [9] at 95% CL, respectively in the local and in the equilateral configurations. For the trispectrum, WMAP provides $-5.6 \times 10^5 < g_{NL} < 6.4 \times 10^5$ [57] (g_{NL} is the “local” trispectrum amplitude from cubic contributions), whereas from Large-Scale-Structures data $-3.5 \times 10^5 < g_{NL} < 8.2 \times 10^5$ [58], at 95% CL. Planck [59] is expected to set further bounds on primordial non-Gaussianity.

Statistical isotropy has always been considered one of the key features of the CMB fluctuations. The appearance of some “anomalies” [60, 61, 62] in the observations though, after numerous and careful data analysis, suggests a possible a breaking of this symmetry that might have occurred at some point of the Universe history, possibly at very early times. This encouraged a series of attempts to model this event, preferably by incorporating it in theories of inflation. Let us shortly describe the above mentioned “anomalies”. First of all, the large scale CMB quadrupole appears to be “too low” and

the octupole “too planar”; in addition to that, there seems to exist a preferred direction along which quadrupole and octupole are aligned [63, 64, 60, 65, 66]. Also, a “cold spot”, i.e. a region of suppressed power, has been observed in the southern Galactic sky [61, 67]. Finally, an indication of asymmetry in the large-scale power spectrum and in higher-order correlation functions between the northern and the southern ecliptic hemispheres was found [68, 62, 69]. Possible explanations for these anomalies have been suggested such as improper foreground subtraction, WMAP systematics, statistical flukes; the possibilities of topological or cosmological origins for them have been proposed as well. Moreover, considering a power spectrum anisotropy due to the existence of a preferred spatial direction \hat{n} and parametrized by a function $g(k)$ as

$$P(\vec{k}) = P(k) \left(1 + g(k)(\hat{k} \cdot \hat{n})^2 \right), \quad (1.1)$$

the five-year WMAP temperature data have been analyzed in order to find out what the magnitude and orientation of such an anisotropy could be. The magnitude has been found to be $g = 0.29 \pm 0.031$ and the orientation aligned nearly along the ecliptic poles [70]. Similar results have been found in [71], where it is pointed out that the origin of such a signal is compatible with beam asymmetries (uncorrected in the maps) which should therefore be investigated before we can find out what the actual limits on the primordial g are.

Several fairly recent works have taken the direction of analysing the consequences, in terms of dynamics of the Universe and of cosmological fluctuations, of an anisotropic pre-inflationary or inflationary era. A cosmic no-hair conjecture exists according to which the presence of a cosmological constant at early times is expected to dilute any form of initial anisotropy [72]. This conjecture has been proven to be true for many (all Bianchi type cosmologies except for the the Bianchi type-IX, for which some restrictions are needed to ensure the applicability of the theorem), but not all kinds of metrics and counterexamples exist in the literature [73, 74, 75]. Moreover, even in the event isotropization should occur, there is a chance that signatures from anisotropic inflation or from an anisotropic pre-inflationary era might still be visible today [76, 77, 78, 79]. In the same context of searching for models of the early Universe that might produce some anisotropy signatures at late time, new theories have been proposed such as spinor models [80, 81, 82, 83], higher p-forms [84, 85, 86, 87, 88, 89] and primordial vector field models.

Within vector field models, higher order correlators had been computed in [90, 91, 92, 93, 94] and, more recently, in [95, 96] for $U(1)$ vector fields. We considered $SU(2)$ vector field models in [97, 98]. Non-Abelian theories offer a richer amount of predictions compared to the Abelian case. Indeed, self interactions provide extra contributions to the bispectrum and trispectrum of curvature fluctuations that are naturally absent in the Abelian case. We verified that these extra contributions can be equally important in a large subset of the parameter space of the theory and, in some case, can even become the dominant ones.

The promising perspective of achieving more and more precise measurements for the cosmological observables thanks to Planck and future experiments and the search for signatures that may help identify the correct inflationary model, have also motivated studies of higher order corrections to cosmological correlation functions and to the power spectrum in particular. Indeed, loop corrections to the correlators arise from the interactions involving the fields during inflation and therefore carry some important information about the physics of the very early Universe.

Loop corrections may lead to interesting effects which scale like the power of the number of e-folds between horizon exit of a given mode k and the end of inflation [99, 100, 101, 102, 103]. The interest in loop corrections to the correlators of cosmological perturbations generated during an early epoch of inflation has been recently stimulated by two papers of Weinberg [104, 105]. The reason is that one-loop corrections to the power spectrum of the curvature perturbation ζ seem to show some infra-red divergences which scale like $\ln(kL)$, where L^{-1} is some infra-red comoving momentum cut-off [106, 107, 108, 109]. However, it has been discussed in [110, 111] (see also [112]) that such potentially large corrections do not appear in quantities that are directly observable.

As to the power spectrum of curvature perturbations, one-loop corrections have been computed in single-field slow-roll inflation by D. Seery [54, 55] and by N. Bartolo and myself [56], in single-field slow-roll inflation. In [56] we completed the analysis carried out in [54, 55], where the metric tensor fluctuations had been neglected for simplicity, by including them in the calculations and proving that their contribution is as important as the one from the scalar perturbations. In the context of loop-calculations, we have also been working on corrections to the power spectrum in theories with non-canonical Lagrangians, which allow for higher and possibly observable corrections [113].

It can be safely stated that in standard single-field slow-roll inflation, the perturbative expansion is well-behaved, in the sense that the agreement with observations found at tree-level for the power-spectrum is not spoiled by the radiative corrections and, on a more general basis, higher order loop corrections introduce smaller and smaller corrections as the perturbation series expansion progresses. This is not generically true in more general theories, such as for instance models with non-canonical Lagrangians, for which bounds need to be requested on the parameters of the theory in order to preserve the validity of the perturbative approach [114, 115].

This thesis collects the main results of our work on loop corrections to the power spectrum in theories of scalar inflation [56, 113] and on primordial non-Gaussianity and anisotropy predictions from theories of inflation where vector fields can play a role in the production of the late time cosmological fluctuations [97, 98]. The δN and the Schwinger-Keldysh formalisms are the main tools of our computation and will be briefly reviewed.

2. Schwinger-Keldysh formalism

The temperature fluctuations in the CMB are rather small, of order 10^{-5} . Theoretical predictions for the power-spectrum of curvature perturbations during inflation provide a very good match at tree level: this suggests that it is correct to use perturbation theory to evaluate cosmological correlators. A formalism conveniently employed to implement the perturbative approach is the Schwinger-Keldysh, also dubbed as “in-in”, formalism. It was first formulated in [49, 50, 51], later applied by J. Maldacena in [13] to the calculation of the bispectrum of curvature fluctuations and revived by S. Weinberg in [52, 53]. In this formalism the expectation value of a field operator $\Theta(t)$ is given by

$$\langle \Omega | \Theta(t) | \Omega \rangle = \left\langle 0 \left| \left[\bar{T} \left(e^{i \int_0^t H_I(t') dt'} \right) \right] \Theta_I(t) \left[T \left(e^{-i \int_0^t H_I(t') dt'} \right) \right] \right| 0 \right\rangle, \quad (2.1)$$

where $|\Omega\rangle$ represents the vacuum of the interacting theory, T and \bar{T} are time-ordering and anti-ordering operators, the subscript I indicates the fields in interaction picture and H_I is the interaction Hamiltonian. The interaction picture has the advantage of allowing to deal with free fields only; the fields can be thus Fourier expanded in terms of quantum creation and annihilation operators

$$\delta\phi(\vec{x}, t) = \int d^3k e^{i\vec{k}\vec{x}} \left[a_{\vec{k}} \delta\phi_{\vec{k}}(t) + a_{-\vec{k}}^+ \delta\phi_{\vec{k}}^*(t) \right],$$

with commutation rules

$$\left[a_{\vec{k}}, a_{\vec{k}'}^+ \right] = (2\pi)^2 \delta^{(3)}(\vec{k} - \vec{k}').$$

The in-in formula has many similarities with the S-matrix in quantum field theory in terms of mathematical structure and perturbative approach, but they also have fundamental differences: the S-matrix corresponds to a transition amplitude between an initial and a final state; a cosmological correlation function is instead the expectation value of a given observable at a given time; moreover, asymptotic states in cosmology are only defined at very early times, when the same initial conditions as in Minkowsky spacetime apply for the free fields.

Using the positive and negative path technique of the in-in formalism [52, 53], the expectation value above can be recast in the form

$$\langle \Omega | \Theta(t) | \Omega \rangle = \left\langle 0 \left| T \left(\Theta_I(t) e^{-i \int_0^t dt' (H_I^+(t') - H_I^-(t'))} \right) \right| 0 \right\rangle, \quad (2.2)$$

where the plus and minus signs indicate modified Feynman propagators, i.e. modified rules of contraction between interacting fields; schematically we have

$$\langle T(\phi_1 \phi_2 \dots \phi_n) \rangle = \sum_{ij, lm, \dots} [\widehat{\phi_i \phi_j}, \widehat{\phi_l \phi_m}, \dots], \quad (2.3)$$

where the sum is taken over all of the possible sets of field contractions and

$$\begin{aligned} \phi^+(\eta') \widehat{\phi^+}(\eta'') &= G^>(\eta', \eta'') \Theta(\eta' - \eta'') + G^<(\eta', \eta'') \Theta(\eta'' - \eta'), \\ \phi^+(\eta') \widehat{\phi^-}(\eta'') &= G^<(\eta', \eta''), \\ \phi^-(\eta') \widehat{\phi^+}(\eta'') &= G^>(\eta', \eta''), \\ \phi^-(\eta') \widehat{\phi^-}(\eta'') &= G^<(\eta', \eta'') \Theta(\eta' - \eta'') + G^>(\eta', \eta'') \Theta(\eta'' - \eta'). \end{aligned}$$

In momentum space we have

$$G_k^>(\eta', \eta'') \equiv \delta\phi_k(\eta')\delta\phi_k^*(\eta''),$$
$$G_k^<(\eta', \eta'') \equiv \delta\phi_k^*(\eta')\delta\phi_k(\eta'').$$

It is important to remember that, when we apply this formalism, the external fields are always supposed to be treated like +fields.

3. Scalar loop corrections to P_ζ

The power spectrum for the comoving curvature perturbation ζ is defined by

$$\langle \zeta_{\vec{k}_1}(t) \zeta_{\vec{k}_2}(t) \rangle = (2\pi)^3 P_\zeta(k) \delta^{(3)}(\vec{k}_1 + \vec{k}_2), \quad (3.1)$$

This and all other correlation functions presented in this thesis are computed using the δN formula. $\zeta(\vec{x})$ at a given time t can be interpreted as a geometrical quantity indicating the fluctuations in the local expansion of the universe; in fact, if $N(\vec{x}, t^*, t)$ is the number of e-foldings of expansion evaluated between times t^* and t , where the initial hypersurface is chosen to be flat and the final one is uniform density, we have

$$\zeta(\vec{x}, t) = N(\vec{x}, t^*, t) - N(t^*, t) \equiv \delta N(\vec{x}, t). \quad (3.2)$$

The number of e-foldings $N(\vec{x}, t^*, t)$ depends on all the fields and their perturbations on the initial slice. In principle, since the fields are governed by second order differential equations, it should also depend on their first time derivatives, but if we assume that slow-roll conditions apply, then the time derivatives will not count as independent quantities.

Let us apply Eq. (3.2) to the computation of P_ζ in single-field slow-roll inflation (the Lagrangian for the scalar field is given by $L_\phi = (1/2)g^{\mu\nu}\partial_\mu\phi\partial_\nu\phi - V(\phi)$)

$$\begin{aligned} \langle \zeta_{\vec{k}_1}(t) \zeta_{\vec{k}_2}(t) \rangle &= \int \frac{d^3x_1}{(2\pi)^3} \frac{d^3x_2}{(2\pi)^3} e^{-i(\vec{k}_1\vec{x}_1 + \vec{k}_2\vec{x}_2)} \\ &\left\langle \left(\sum_n \frac{N^{(n)}(t^*, t)}{n!} (\delta\phi(\vec{x}_1, t^*))^n \right), \left(\sum_m \frac{N^{(m)}(t^*, t)}{m!} (\delta\phi(\vec{x}_2, t^*))^m \right) \right\rangle. \end{aligned} \quad (3.3)$$

The sums can be expanded to the desired order. Up to one loop we have

$$\begin{aligned} \langle \zeta_{\vec{k}_1}(t) \zeta_{\vec{k}_2}(t) \rangle &= N^{(1)2} \langle \delta\phi_{\vec{k}_1} \delta\phi_{\vec{k}_2} \rangle_* \\ &+ \frac{1}{2!} N^{(1)} N^{(2)} \int d^3q \langle \delta\phi_{\vec{k}_1} \delta\phi_{\vec{q}} \delta\phi_{\vec{k}_2 - \vec{q}} \rangle_* + (\vec{k}_1 \leftrightarrow \vec{k}_2) \\ &+ \frac{1}{3!} N^{(1)} N^{(3)} \int d^3q d^3p \langle \delta\phi_{\vec{k}_1} \delta\phi_{\vec{q}} \delta\phi_{\vec{p}} \delta\phi_{\vec{q} + \vec{p} - \vec{k}_2} \rangle_* + (\vec{k}_1 \leftrightarrow \vec{k}_2) \\ &+ \frac{1}{(2!)^2} (N^{(2)})^2 \int d^3q d^3p \langle \delta\phi_{\vec{q}} \delta\phi_{\vec{k}_1 - \vec{q}} \delta\phi_{\vec{p}} \delta\phi_{\vec{k}_2 - \vec{p}} \rangle_*. \end{aligned} \quad (3.4)$$

where a star indicates evaluation around the time of horizon crossing. Eq. (3.4) can finally be rewritten as [116, 109]

$$\begin{aligned} \langle \zeta_{\vec{k}_1}(t) \zeta_{\vec{k}_2}(t) \rangle &= (2\pi)^3 \delta^{(3)}(\vec{k}_1 + \vec{k}_2) \left[(N^{(1)})^2 (P_{\text{tree}}(k_1) + P_{\text{one-loop}}(k_1)) \right. \\ &+ N^{(1)} N^{(2)} \int \frac{d^3q}{(2\pi)^3} B_\phi(k_1, q, |\vec{k}_1 - \vec{q}|) \\ &+ \frac{1}{2} (N^{(2)})^2 \int \frac{d^3q}{(2\pi)^3} P_{\text{tree}}(q) P_{\text{tree}}(|\vec{k}_1 - \vec{q}|) \\ &\left. + N^{(1)} N^{(3)} P_{\text{tree}}(k) \int \frac{d^3q}{(2\pi)^3} P_{\text{tree}}(q) \right], \end{aligned} \quad (3.5)$$




Figure 1. Diagrammatic representation of the one loop corrections to the power spectrum of $\delta\phi$ from scalar modes to leading ($\sim \epsilon^0$) order in slow-roll.




Figure 2. Next-to-leading ($\sim \sqrt{\epsilon}$) order one loop corrections from scalar modes to the power spectrum of $\delta\phi$.

$P_{\text{tree}}(k)$ is the tree level power spectrum (3.6)

$$\langle \delta\phi_{\vec{k}_1}^- \delta\phi_{\vec{k}_2}^- \rangle_* = (2\pi)^3 P(k) \delta^{(3)}(\vec{k}_1 + \vec{k}_2) = (2\pi)^3 \frac{H_*^2}{2k^3} \delta^{(3)}(\vec{k}_1 + \vec{k}_2), \quad (3.6)$$

where H_* is the Hubble parameter evaluated at horizon exit (when $k = aH$). The variance per logarithmic interval in k is given by $\mathcal{P}(k) = (k^3/2\pi^2)P(k)$. The one loop contribution to the power spectrum is given by

$$P_{\text{one-loop}}(k) = P_{\text{scalar}}(k) + P_{\text{tensor}}(k), \quad (3.7)$$

where the first term on the right-hand side, P_{scalar} , accounts for the contributions coming from the inflaton self-interactions and were computed by D. Seery in [108, 109]

$$P_{\text{scalar}} = \frac{H_*^4}{k^3} [g_1 \ln(k) + g_2], \quad (3.8)$$

where g_1 and g_2 are numerical factors. Their diagrammatic representation is given in Fig.1 for the leading order and in Fig.2 for the next-to-leading order corrections. The loop corrections P_{tensor} , arising from interactions between the tensor (graviton) modes and the scalar field, were ignored for simplicity in [108, 106], however they should be included since they are not slow-roll suppressed compared to loops of scalar modes. Their computation was presented for the first time in our paper [56] and will be reviewed in the first part of this thesis.

Both P_{scalar} and P_{tensor} are evaluated at around the time of horizon crossing and as such they are due to genuine quantum effects.

The contributions in the third and fourth lines of Eq. (3.5), also dubbed as “classical one-loop”, can be considered as classical loop contributions arising after the perturbation modes leave the horizon. The distinction between classical and quantum loops is

understood as for example in [109]: quantum loops find their origin in the Lagrangian interaction terms between the inflaton perturbations and the gravitational modes or from self-interaction of $\delta\phi$; classical loops are corrections merely coming from the expansion of ζ using the δN formula and originate from zeroth order terms in the Schwinger-Keldysh formula.

Finally, the second line of (3.5) includes the integral of $B_\phi(k_1, k_2, k_3)$, the bispectrum of the scalar field defined by

$$\langle \delta\phi_{\vec{k}_1}^- \delta\phi_{\vec{k}_2}^- \delta\phi_{\vec{k}_3}^- \rangle \equiv (2\pi)^3 \delta^{(3)}(\vec{k}_1 + \vec{k}_2 + \vec{k}_3) B_\phi(k_1, k_2, k_3) \quad (3.9)$$

and from [13] we have

$$B_\phi \simeq \frac{\sqrt{\epsilon} H_*^4 F(k_i)}{m_P}, \quad (3.10)$$

where m_P is the Planck mass, ϵ is the slow-roll parameter ($\epsilon \equiv -\dot{H}/H^2$) and F is a function of the momenta moduli k_i of dimension $(\text{mass})^{-6}$.

4. Perturbative expansion of the Lagrangian in $P(X, \phi)$ theories

We will now review the computation of the tensor loop corrections to P_ζ . For our purposes, the exponentials in Eq. (2.2) need to be expanded up to second order in the interaction Hamiltonian H_I

$$\begin{aligned} \langle \Omega | \Theta(\eta) | \Omega \rangle_{1L} &= i \langle 0 | T \left[\Theta \int_{-\infty}^{\eta} d\eta' \left(H_I^+(\eta') - H_I^-(\eta') \right) \right] | 0 \rangle \\ &+ \frac{(-i)^2}{2} \langle 0 | T \left[\Theta \int_{-\infty}^{\eta} d\eta' \left(H_I^+(\eta') - H_I^-(\eta') \right) \int_{-\infty}^{\eta} d\eta'' \left(H_I^+(\eta'') - H_I^-(\eta'') \right) \right] | 0 \rangle, \end{aligned} \quad (4.1)$$

where $\Theta(t) \equiv \delta\phi_{k_1}^-(\eta) \delta\phi_{k_2}^-(\eta)$. One-loop power-spectrum diagrams require an expansion of the interaction Hamiltonian to third and fourth order in the field fluctuations, i.e. $H_I \equiv H_I^{(3)} + H_I^{(4)}$. We provide in Figs. (3) and (4) the diagrammatic representation of the leading order corrections that we will find for the diagrams with tensor loops in single-field slow-roll inflation. The continuous lines represent scalar propagators, whereas the dotted lines indicate tensor propagators. In order to derive this result and the analytic expressions for these diagrams, we need to first calculate and expand H_I up to fourth order in the field perturbations $\delta\phi$ and $\delta\gamma$. The starting point is the Lagrangian of the system.

We will begin with a more general Lagrangian for the scalar field than the usual $L_\phi = (1/2)g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - V(\phi)$, by introducing a non-conventional kinetic term, i.e.

$$S = \frac{1}{2} \int d^3x dt \sqrt{-g} \left[M_P^2 R + 2P(X, \phi) \right], \quad (4.2)$$

where $P = P(X, \phi)$ is a generic function of the scalar field and of $X \equiv \frac{1}{2}g_{\mu\nu} \partial^\mu \phi \partial^\nu \phi$ and R is the Ricci scalar in four dimensions. Notice that the action (4.2) reduces to the standard case if $P = X - V$, where V is the potential for the scalar field.

Theories of inflation where the Lagrangian kinetic term is a generic function of the scalar field and its first derivatives, like in Eq. (4.2), are string theory-inspired. They represent interesting alternatives to the basic inflationary scenario because of their non-Gaussianity predictions. The crucial quantity in this sense is represented by the speed of sound $c_s^2 \equiv (\partial_X P) / (\partial_X P + 2X \partial_{XX} P)$, which is allowed to vary between 0 and 1. The perturbative expansion of the interaction Hamiltonian in this kind of models has coefficients proportional to inverse powers of the sound speed and therefore, for small values of c_s , allows both for non-negligible loop corrections to the power spectrum of the curvature fluctuations [113] and for large values for the amplitudes of three [31] and four [41, 42, 43, 44, 45, 46] point functions. In this thesis, we will carry out the calculations of the interaction Hamiltonian for these general theories up to a certain point and then, for simplicity in the presentation, focus on the canonical case (the remaining computations for more general Lagrangians will be found in [113]).

Let us list the background equations for the system

$$2\dot{H} + 3H^2 = -P, \quad (4.3)$$

$$3H^2 = 2XP_X - P, \quad (4.4)$$

$$\begin{aligned} \dot{X}(P_X + 2XP_{XX}) + 2\sqrt{3}(2XP_X - P)^{1/2}XP_X \\ = \sqrt{2X}(P_\phi - 2XP_{X\phi}), \end{aligned} \quad (4.5)$$

where a dot indicates a derivative w.r.t. cosmic time and, to zeroth order, we have $X \equiv \frac{\dot{\phi}^2}{2}$.

The so called *flow-parameters* are defined as

$$\varepsilon \equiv -\frac{\dot{H}}{H^2}, \quad (4.6)$$

$$\eta \equiv \frac{\dot{\varepsilon}}{\varepsilon H}. \quad (4.7)$$

These quantities reduce to the slow-roll parameters in the standard case, so it is natural to assume $|\varepsilon| \ll 1$ and $|\eta| \ll 1$. It is not correct to talk about *slow-roll* if P is left as a generic function of X and ϕ , since the smallness of ε and η does not necessarily indicate that $\dot{\phi}^2 \ll H^2$ and $|\dot{\phi}| \ll |3H\dot{\phi}|$. It can be convenient to decompose ε as the sum $\varepsilon = \varepsilon_\phi + \varepsilon_X$, where

$$\varepsilon_\phi \equiv -\frac{\dot{\phi}}{H^2} \frac{\partial H}{\partial \phi}, \quad (4.8)$$

$$\varepsilon_X \equiv -\frac{\dot{X}}{H^2} \frac{\partial H}{\partial X}. \quad (4.9)$$

The parameters that are expected to appear in the perturbative expansion of the Lagrangian are

$$c_s^2 = \frac{P_X}{P_X + 2XP_{XX}}, \quad (4.10)$$

$$s \equiv \frac{\dot{c}_s}{c_s H}, \quad (4.11)$$

$$u \equiv 1 - \frac{1}{c_s^2}, \quad (4.12)$$

$$\Sigma \equiv XP_X + 2X^2P_{XX}, \quad (4.13)$$

$$\lambda \equiv X^2P_{XX} + \frac{2}{3}X^3P_{XXX}, \quad (4.14)$$

$$\Pi \equiv X^3P_{XXX} + \frac{2}{5}X^4P_{XXXX}, \quad (4.15)$$

where c_s is the sound speed. c_s is allowed to vary between 0 and 1, so the quantity $|u|$ can freely range between 0 and ∞ . The only assumption we make is $s \ll 1$, from c_s being constant in the standard case.

4.1. Arnowitt-Deser-Misner (ADM) decomposition for $P(X, \phi)$ theories

The Lagrangian in Eq. (4.2) will now undergo a perturbative expansion in terms of the field fluctuations $\delta\phi(\vec{x}, t) \equiv \phi(\vec{x}, t) - \phi_0(t)$ (ϕ_0 is the homogeneous background value for the field) and of the metric fluctuations.




Figure 3. Diagrammatic representation of the leading order (tensor mode) corrections from $H_I^{(4)}$ to the power spectrum of $\delta\phi$.




Figure 4. Diagrammatic representation of the leading order (tensor mode) corrections from $H_I^{(3)}$ to the power spectrum of $\delta\phi$. Notice that this diagram is not slow-roll suppressed compared to the one in Fig. 3, whereas this is not the case for the scalar modes (see Figs. 1 and 2).

It is convenient to adopt the 3+1 Arnowitt-Deser-Misner (ADM) splitting for the metric. In the spatially flat gauge the perturbed metric is

$$ds^2 = -N^2 dt^2 + h_{ij}(dx^i + N^i dt)(dx^j + N^j dt), \quad (4.16)$$

$$h_{ij} = a^2(t)(e^\gamma)_{ij}, \quad (4.17)$$

where $a(t)$ is the scale factor, γ_{ij} is a tensor perturbation with $\partial_i \gamma_{ij} = \gamma_{ii} = 0$ (traceless and divergenceless) and $\det(e^\gamma)_{ij} = 1$. Notice that repeated lower indices are summed up with a Kronecker delta, so $\partial_i \gamma_{ij}$ stands for $\delta^{ik} \partial_i \gamma_{kj}$ and $\gamma_{ij} = \delta^{ij} \gamma_{ij}$.

In the ADM formalism, the action (4.2) becomes [13]

$$S = \frac{1}{2} \int dt d^3x \sqrt{h} \left[NR^{(3)} + 2NP + N^{-1} (E_{ij} E^{ij} - E^2) \right], \quad (4.18)$$

where $R^{(3)}$ is the curvature scalar associated with the three dimensional metric h_{ij} and

$$E_{ij} = \frac{1}{2} (\dot{h}_{ij} - \nabla_i N_j - \nabla_j N_i),$$

$$E = h^{ij} E_{ij}.$$

A dot indicates derivatives w.r.t. time t , all the spatial indices are raised and lowered with h_{ij} and units of $M_{Pl}^{-2} \equiv 8\pi G = 1$ will be from now on employed. To 4th order we have

$$R^{(3)} = -\frac{1}{4} \partial_i \gamma_{al} \partial_i \gamma_{al}. \quad (4.19)$$

The lapse and shift functions, N and N^i can be written as

$$N = 1 + \alpha,$$

$$N_j = \partial_j \theta + \beta_j,$$

where α , θ and β are functions of time and space (β_j is divergenceless). We have exploited the gauge freedom to set two scalar and two vector modes to zero, thus leaving one scalar mode from N , one scalar and two vector modes from N_j and two tensor modes (the two independent polarizations of the graviton) from h_{ij} , together with the inflaton field perturbation $\delta\phi$. N and N_i are non-dynamical degrees of freedom and can be expressed in terms of the other modes ($\delta\phi$ and γ_{ij}), once the Hamiltonian and the momentum constraints (we derive them in the next section) are solved.

4.1.1. Solving Hamiltonian and momentum constraint equations

Momentum and Hamiltonian constraints are derived from varying the action w.r.t. the shift and lapse functions respectively. It turns out that, in order to expand the action to a given order n , it is only necessary to perturb N and N_i up to order $n - 2$ [13, 31]. Therefore we will solve the constraints to second order in the metric and scalar field fluctuations.

Let us begin with the expansions

$$\begin{aligned}\alpha &= \alpha_1 + \alpha_2, \\ \beta_i &= \beta_{1i} + \beta_{2i}, \\ \theta &= \theta_1 + \theta_2.\end{aligned}$$

where α_1 and α_2 are respectively first and second order in the fields fluctuations (similarly for β_{1i} and β_{2i} , and for θ_1 and θ_2). Let us then expand P to second order. P is a generic function of X and ϕ . We first need the expansion of X

$$\begin{aligned}X &= -g^{\mu\nu}\partial_\mu\phi\partial_\nu\phi = -\frac{1}{2}\left[g^{00}\dot{\phi}^2 + 2g^{0i}\partial_i\phi\dot{\phi} + g^{ij}\partial_i\phi\partial_j\phi\right] \\ &= -\frac{1}{2}\left[-N^{-2}(\dot{\phi} + \delta\dot{\phi})^2 + 2N^{-2}N^i\partial_i\delta\phi(\dot{\phi} + \delta\dot{\phi}) + \left(h^{ij} - \frac{N^iN^j}{N^2}\right)\partial_i\delta\phi\partial_j\delta\phi\right] \\ &= X_0 + \Delta X\end{aligned}\tag{4.20}$$

where $N^i \equiv h^{ij}N_j$, X_0 is the zeroth order part, i.e. $X_0 = \frac{\dot{\phi}^2}{2}$ and ΔX is the perturbation to the desired order ($\Delta X = \Delta X_1 + \Delta X_2 + \Delta X_3 + \dots$). Notice that $\phi(t, \vec{x}) = \phi_0(t) + \delta\phi(t, \vec{x})$, but for simplicity we will suppress the subscript '0' in the background value of the field.

The expressions for the perturbations ΔX_i become

$$\Delta X_1 = 2X_0\left[\frac{\delta\dot{\phi}}{\dot{\phi}} - \alpha_1\right]\tag{4.21}$$

$$\Delta X_2 = X_0\left[\left(\frac{\delta\dot{\phi}}{\dot{\phi}}\right)^2 - 4\alpha_1\frac{\delta\dot{\phi}}{\dot{\phi}} - 2\alpha_2 + 3\alpha_1^2 - 2N_1^i\partial_j\frac{\delta\dot{\phi}}{\dot{\phi}}\right] - \frac{1}{a^2\dot{\phi}^2}\partial_i\delta\phi\partial_i\delta\phi\tag{4.22}$$

and so on for ΔX_3 and higher order terms. The expansion of $P(X, \phi)$ up to second order becomes

$$P(X, \phi) = P_0 + P_X|_0\Delta X + P_\phi|_0\Delta\phi + \frac{1}{2!}P_{XX}|_0(\Delta X)^2 + \frac{1}{2!}P_{\phi\phi}|_0\Delta\phi^2 + P_{X\phi}|_0\Delta X\Delta\phi\tag{4.23}$$

where as usual the subscript '0' indicated the zeroth order, $P_X = \partial_X P$, $P_\phi = \partial_\phi P$ and similarly for the second order derivatives, $\Delta\phi = \delta\phi$ and ΔX needs to be expanded up to the needed order.

We are now ready to write the momentum and Hamiltonian constraints

$$\nabla_i \left[N^{-1} \left(E_j^i - \delta_j^i E \right) \right] = N^{-1} P_X \left[\dot{\phi} - N^l \partial_l \phi \right] \partial_j \phi \quad (4.24)$$

$$R^{(3)} + 2P - 4P_X X - N^{-2} \left(E_{ij} E^{ij} - E^2 \right) - 2P_X h^{ij} \partial_i \phi \partial_j \phi = 0 \quad (4.25)$$

The momentum constraint to first order reads

$$2H \partial_j \alpha_1 - \frac{1}{2a^2} \partial^2 \beta_{1j} = P_X \dot{\phi} \partial_j \delta \phi, \quad (4.26)$$

where $H = \dot{a}/a$ is the Hubble parameter. Eq. (4.26) can be solved to derive α_1 . Taking the derivative ∂^j of both sides of (4.26) and using the divergenceless condition for β , we have

$$\alpha_1 = \frac{P_X \dot{\phi} \delta \phi}{2H}. \quad (4.27)$$

Using the solution found for α_1 , we find $\partial^2 \beta_{1j} = 0$, from which we can set $\beta_{1j} = 0$. Here $\partial^2 \equiv \delta^{ij} \partial_i \partial_j$, which we will indicate in the rest of the thesis also as $\partial_i \partial_i$, and from now on we define $\beta_i \equiv \beta_{2i}$ for simplicity.

The momentum constraint to second order is

$$\begin{aligned} & 2H \partial_j \alpha_2 - 4H \alpha_1 \partial_j \alpha_1 - \frac{1}{a^2} \partial_j \alpha_1 \partial^2 \theta_1 + \frac{1}{a^2} \partial_i \alpha_1 \partial_i \partial_j \theta_1 - \frac{1}{2} \partial_i \alpha_1 \dot{\gamma}_{ij} \\ & - \frac{1}{2a^2} \partial^2 \beta_j + \frac{1}{4} \dot{\gamma}_{ik} \partial_i \gamma_{kj} - \frac{1}{4} \dot{\gamma}_{ik} \partial_i \dot{\gamma}_{kj} - \frac{1}{4} \dot{\gamma}_{ik} \partial_j \gamma_{ik} + \frac{1}{2a^2} \partial_i \theta_1 \partial^2 \gamma_{ij} \\ & = P_X \partial_j \delta \phi \delta \dot{\phi} + 2X P_{XX} \partial_j \delta \phi \delta \dot{\phi} - 2X P_{XX} \dot{\phi} \alpha_1 \partial_j \delta \phi - P_X \dot{\phi} \alpha_1 \partial_j \delta \phi \\ & + P_{X\phi} \dot{\phi} \delta \phi \partial_j \delta \phi. \end{aligned} \quad (4.28)$$

The solutions are

$$\begin{aligned} \alpha_2 = & \frac{\alpha_1^2}{2} + \frac{1}{2Ha^2} \partial^{-2} \left[\partial^2 \alpha_1 \partial^2 \theta_1 - \partial_i \partial_j \alpha_1 \partial_i \partial_j \theta_1 \right] + \frac{P_X}{2H} \partial^{-2} \Sigma \\ & + \frac{1}{4H} \partial^{-2} \left[\dot{\gamma}_{ij} \partial_i \partial_j \alpha_1 \right] - \frac{1}{4a^2 H} \partial^{-2} \left[\partial_i \partial_j \theta_1 \partial^2 \gamma_{ij} \right] \\ & + \frac{1}{8H} \partial^{-2} \left[\partial_j \dot{\gamma}_{ik} \partial_j \gamma_{ik} \right] + \frac{P_{X\phi} \dot{\phi}}{2H} \partial^{-2} \left[(\partial_j \delta \phi)^2 + \delta \phi \partial^2 \delta \phi \right] \\ & + \frac{X P_{XX}}{H} \partial^{-2} \left[\partial^2 \delta \phi \delta \dot{\phi} + \partial_j \delta \phi \partial_j \delta \dot{\phi} - \dot{\phi} \left(\partial_j \alpha_1 \partial_j \delta \phi + \alpha_1 \partial^2 \delta \phi \right) \right], \end{aligned} \quad (4.29)$$

where $\Sigma \equiv \partial^2 \delta \phi \delta \dot{\phi} + \partial_j \delta \phi \partial_j \delta \dot{\phi}$, and

$$\begin{aligned} \frac{1}{2a^2} \partial^2 \beta_j = & 2H \partial_j \alpha_2 - 4H \alpha_1 \partial_j \alpha_1 - \frac{1}{a^2} \partial_j \alpha_1 \partial^2 \theta_1 + \frac{1}{a^2} \partial_i \alpha_1 \partial_i \partial_j \theta_1 \\ & - \frac{1}{2} \partial_i \alpha_1 \dot{\gamma}_{ij} + \frac{1}{4} \dot{\gamma}_{ik} \partial_i \gamma_{kj} - \frac{1}{4} \dot{\gamma}_{ik} \partial_i \dot{\gamma}_{kj} - \frac{1}{4} \dot{\gamma}_{ik} \partial_j \gamma_{ik} \\ & + \frac{1}{2a^2} \partial_i \theta_1 \partial^2 \gamma_{ij} - P_X \partial_j \delta \phi \delta \dot{\phi} - 2X P_{XX} \partial_j \delta \phi \delta \dot{\phi} \\ & + 2X P_{XX} \dot{\phi} \alpha_1 \partial_j \delta \phi + P_X \dot{\phi} \alpha_1 \partial_j \delta \phi - P_{X\phi} \dot{\phi} \delta \phi \partial_j \delta \phi. \end{aligned} \quad (4.30)$$

Let us now move to the Hamiltonian constraint which provides

$$\frac{4H}{a^2}\partial^2\theta_1 = -4XP_X \left(\frac{\delta\dot{\phi}}{\dot{\phi}} - \alpha_1 \right) + 2P_\phi\delta\phi - 8P_{XX}X^2 \left(\frac{\delta\dot{\phi}}{\dot{\phi}} - \alpha_1 \right) - 4XP_{X\phi}\delta\phi - 12H^2\alpha_1, \quad (4.31)$$

to first order and

$$\begin{aligned} -\frac{4H}{a^2}\partial^2\theta_2 = & (-2\alpha_1) \left[4XP_X \frac{\delta\dot{\phi}}{\dot{\phi}} + 20P_{XX}X^2 \frac{\delta\dot{\phi}}{\dot{\phi}} + 2XP_{X\phi}\delta\phi + 8P_{XXX}X^3 \frac{\delta\dot{\phi}}{\dot{\phi}} \right. \\ & \left. + 4P_{XX\phi}X^2\delta\phi + \frac{4H}{a^2}\partial^2\theta_1 \right] - \frac{4X(P_X + 2XP_{XX})}{a^2\dot{\phi}} \partial_i\theta_1\partial_i\delta\phi - \frac{1}{a^4}(\partial^2\theta_1)^2 \\ & + \frac{1}{a^2} \left[-\gamma_{iq}\partial_q\partial_i\theta_1 + \frac{1}{a^2}(\partial_i\partial_j\theta_1)^2 \right] + (-6H^2 + 2XP_X + 4X^2P_{XX}) [3\alpha_1^2 - 2\alpha_2] \\ & + 4\alpha_1^2 (3X^2P_{XX} + 2X^3P_{XXX}) + \frac{\delta\dot{\phi}^2}{\dot{\phi}^2} [2XP_X + 16X^2P_{XX} + 8X^3P_{XXX}] \\ & + \frac{\delta\dot{\phi}\delta\phi}{\dot{\phi}} [4XP_{X\phi} + 8X^2P_{XX\phi}] - \frac{(\partial_i\delta\phi)^2}{a^2\dot{\phi}^2} [4X^2P_{XX} - 2XP_X] + \delta\phi^2 [-P_{\phi\phi} \\ & + 2XP_{X\phi\phi}] + \frac{1}{4} [\dot{\gamma}_{ij}\dot{\gamma}_{ij} + \frac{1}{a^2}\partial_a\gamma_{iq}\partial_a\gamma_{iq}] - \frac{4H}{a^2}\gamma_{ij}\partial_i\partial_j\theta_1, \end{aligned} \quad (4.32)$$

to second order.

4.1.2. Reduction to the canonical case

In the canonical case, to zeroth order in perturbation theory $P = (\dot{\phi}^2/2) - V(\phi)$, so $P_X = 1$ and $P_{\phi^n} = -\partial^n V/\partial\phi^n$ with all other derivatives of P being zero. The solutions above therefore reduce to [56]

$$\alpha_1 = \frac{1}{2H}\dot{\phi}\delta\phi, \quad (4.33)$$

$$\frac{4H}{a^2}\partial^2\theta_1 = -2V_\phi\delta\phi - 2\dot{\phi}\delta\dot{\phi} + 2\alpha_1(-6H^2 + (\dot{\phi})^2), \quad (4.34)$$

$$\begin{aligned} \alpha_2 = & \frac{\alpha_1^2}{2} + \frac{1}{2H}\partial^{-2}\Sigma + \frac{1}{2Ha^2}\partial^{-2} [\partial^2\alpha_1\partial^2\theta_1 - \partial_i\partial_j\alpha_1\partial_i\partial_j\theta_1] \\ & + \frac{1}{4H}\partial^{-2} [\dot{\gamma}_{ij}\partial_i\partial_j\alpha_1] - \frac{1}{4Ha^2}\partial^{-2} [\partial_i\partial_j\theta_1\partial^2\gamma_{ij}] + \frac{1}{8H}\partial^{-2} [\partial_j\dot{\gamma}_{ik}\partial_j\gamma_{ik}], \end{aligned} \quad (4.35)$$

$$\begin{aligned} \frac{4H}{a^2}\partial^2\theta_2 = & 2\alpha_1 \left[2\dot{\phi}\delta\dot{\phi} + \frac{4H}{a^2}\partial^2\theta_1 \right] + \frac{2}{a^2}\dot{\phi}\partial_i\theta_1\partial_i\delta\phi - \frac{1}{a^4}\partial_i\partial_j\theta_1\partial_i\partial_j\theta_1 \\ & + \frac{1}{a^4}(\partial^2\theta_1)^2 - (3\alpha_1^2 - 2\alpha_2)(\dot{\phi}^2 - 6H^2) - \delta\dot{\phi}^2 - \frac{1}{a^2}\partial_i\delta\phi\partial_i\delta\phi \\ & - V_{\phi\phi}\delta\phi^2 - \frac{1}{4a^2}\partial_a\gamma_{iq}\partial_a\gamma_{iq} - \frac{1}{4}\dot{\gamma}_{ij}\dot{\gamma}_{ij} + \frac{1}{a^2}\dot{\gamma}_{iq}\partial_i\partial_q\theta_1, \end{aligned} \quad (4.36)$$

$$\frac{1}{2a^2}\partial^4\beta_j = \frac{1}{a^2}\partial^2\alpha_1\partial_j\partial^2\theta_1 - \frac{1}{a^2}\partial_m\partial_j\alpha_1\partial_m\partial^2\theta_1 + \frac{1}{a^2}\partial_m\alpha_1\partial_m\partial_j\partial^2\theta_1$$

$$\begin{aligned}
& -\frac{1}{a^2}\partial_j\alpha_1\partial^4\theta_1 - \frac{1}{a^2}\partial_m\partial_j\partial_i\alpha_1\partial_i\partial_m\theta_1 + \frac{1}{a^2}\partial^2\partial_i\alpha_1\partial_i\partial_j\theta_1 \\
& - \frac{1}{a^2}\partial_i\partial_j\alpha_1\partial_i\partial^2\theta_1 + \frac{1}{a^2}\partial_m\partial_i\alpha_1\partial_m\partial_i\partial_j\theta_1 + \partial_m\partial_j\delta\dot{\phi}\partial_m\delta\phi \\
& - \partial^2\delta\dot{\phi}\partial_j\delta\phi + \partial_j\delta\dot{\phi}\partial^2\delta\phi - \partial_m\delta\dot{\phi}\partial_m\partial_j\delta\phi - \frac{1}{2}\partial^2(\gamma_{ij}\partial_i\alpha_1) \\
& - \frac{1}{2a^2}\partial^2(\partial_j\gamma_{bq}\partial_q\partial_b\theta_1) + \frac{1}{2a^2}\partial^2(\partial^2\gamma_{jk}\partial_k\theta_1) - \frac{1}{4}\partial^2(\gamma_{il}\partial_i\gamma_{jl}) \\
& + \frac{1}{4}\partial^2(\gamma_{ik}\partial_i\gamma_{kj}). \tag{4.37}
\end{aligned}$$

where $V_{\phi\phi} \equiv \partial^2 V / \partial \phi^2$ and ∂^{-2} is the inverse of the laplacian operator. Notice that the equations (4.33) through (4.37) agree with the results obtained in [152], if we set γ_{ij} to zero.

4.2. Fourth-order expansion of the Lagrangian in the canonical case

The expansion of the action up to 4th order can be now derived by plugging the solutions (4.27) through (4.32) in (4.18). The final expression is quite lengthy and can be found in [113]. We will here only report the 4th order expansion of the action in the canonical case

$$\begin{aligned}
S_4 = a^3 \int dt d^3x & \left[-\frac{1}{24}V_{\phi\phi\phi\phi}\delta\phi^4 + \frac{1}{2a^2}\partial_{(i}\beta_{j)}\partial_i\beta_j + \frac{1}{2a^4}\partial_j\theta_1\partial_j\delta\phi\partial_m\theta_1\partial_m\delta\phi \right. \\
& - \frac{1}{a^2}\delta\dot{\phi}(\partial_j\theta_2 + \beta_j)\partial_j\delta\phi + \left(\alpha_1^2\alpha_2 - \frac{1}{2}\alpha_2^2\right)(-6H^2 + \dot{\varphi}^2) \\
& + \frac{\alpha_1}{2}\left[-\frac{1}{3}V_{\phi\phi\phi}\delta\phi^3 - 2V_\phi\alpha_1^2\delta\phi + \alpha_1\left(-\frac{1}{a^2}\partial_i\delta\phi\partial_i\delta\phi - V_{\phi\phi}\delta\phi^2\right) \right. \\
& - \frac{2}{a^4}(\partial_i\partial_j\theta_2\partial_i\partial_j\theta_1 - \partial^2\theta_1\partial^2\theta_2 + \partial_i\beta_j\partial_i\partial_j\theta_1) + \frac{2}{a^2}(\dot{\phi}(\partial_j\theta_2 + \beta_j)\partial_j\delta\phi \\
& + \delta\dot{\phi}\partial_j\theta_1\partial_j\delta\phi) \left. \right] + \alpha_1^2\left[\frac{1}{2a^2}(\gamma_{qi}\partial_a\partial_i\gamma_{aq} - \frac{1}{2}\partial_a\gamma_{iq}\partial_a\gamma_{iq}) - \frac{1}{4}\dot{\gamma}_{ij}\dot{\gamma}_{ij} + \dot{\gamma}_{iq}\partial_i\partial_q\theta_1\right] \\
& - \frac{1}{a^2}\left[\frac{1}{2}\gamma_{ik}\gamma_{kj}\partial_j\delta\phi\partial_i\delta\phi - \alpha_1\dot{\gamma}_{ij}\partial_j\delta\phi\partial_i\delta\phi + \alpha_2\partial_i\delta\phi\partial_i\delta\phi \right. \\
& - \dot{\phi}\partial_j\delta\phi(\gamma_{ij}\partial_i\theta_2 + \gamma_{ij}\beta_i + \gamma_{ij}\partial_i\theta_1) - \partial_k\theta_2\dot{\gamma}_{ab}\partial_b\gamma_{ak} - \beta_k\dot{\gamma}_{ab}\partial_b\gamma_{ak} \\
& - \frac{1}{2}\dot{\gamma}_{ab}\beta_k\partial_k\gamma_{ab} - \alpha_1(H\dot{\gamma}_{ab}\partial_a\partial_b\theta_2 + \dot{\gamma}_{ab}\partial_a\partial_b\theta_2 + \dot{\gamma}_{ab}\partial_a\beta_b) + \frac{1}{2}\dot{\gamma}_{ab}\partial_k\gamma_{ab}\partial_k\theta_2 \left. \right] \\
& + \frac{1}{2a^4}\left(-8\dot{\gamma}_{ip}\partial_i\partial_j\theta_1\partial_p\partial_j\theta_2 - 4\dot{\gamma}_{ip}\partial_i\partial_j\theta_1\partial_p\beta_j - 4\dot{\gamma}_{ip}\partial_p\partial_j\theta_1\partial_j\beta_i \right. \\
& - \partial_q\theta_1\partial_i\dot{\gamma}_{jq}\partial_i\partial_j\theta_2 - \partial_q\theta_2\partial_i\dot{\gamma}_{jq}\partial_i\partial_j\theta_1 - \beta_q\partial_i\dot{\gamma}_{jq}\partial_i\partial_j\theta_1 \\
& - \partial_q\theta_1\partial_i\dot{\gamma}_{jq}\partial_i\beta_j - \partial_q\theta_1\partial_i\dot{\gamma}_{jq}\partial_j\beta_i + \partial_q\theta_1\partial_q\dot{\gamma}_{ij}\partial_i\partial_j\theta_2 \\
& \left. + 2\partial_q\theta_2\partial_q\dot{\gamma}_{ij}\partial_i\partial_j\theta_1 + 2\partial_q\theta_1\partial_q\dot{\gamma}_{ij}\partial_i\beta_j + 2\beta_q\partial_q\dot{\gamma}_{ij}\partial_i\partial_j\theta_1\right). \tag{4.38}
\end{aligned}$$

Similarly, the loop computation will be from now on performed considering this simpler case.

5. Tensor loop corrections to P_ζ

Let us then consider the terms in the interaction Hamiltonian H_I that involve the tensor modes. The third order action in single-field slow-roll inflation with non-zero graviton fluctuations was calculated in [13]; we will focus on the leading order term in the slow-roll parameters, so we have

$$H_I^{(3)}(\eta) \equiv \frac{a^2(\eta)}{2} \int d^3x \gamma_{ij} \partial_i \delta\phi \partial_j \delta\phi. \quad (5.1)$$

The fourth order action is given by Eq. (15.2.1). Notice that some of the interaction terms involving the tensor modes in (15.2.1) appear with time derivatives, therefore the construction of the path integral formula requires additional care compared to the case where time derivatives only appear in the kinetic term of the Lagrangian. This issue will be discussed in Appendix 15.1. Also, it is possible to show that in Eq. (15.2.1), of all the leading terms in the slow-roll expansion, only one will provide a non-zero contribution to the loop correction (see Appendix 15.2 for a detailed analysis) and contribute to the interaction Hamiltonian to fourth order which becomes

$$H_I^{(4)}(\eta) \equiv \frac{a^2(\eta)}{4} \int d^3x \gamma_{ik} \gamma_{kj} \partial_i \delta\phi \partial_j \delta\phi \quad (5.2)$$

where the tensor fluctuations are

$$\gamma_{ij}(\vec{x}, t) = \int d^3k e^{i\vec{k}\vec{x}} \sum_{\lambda} \left[\varepsilon_{ij}(\hat{k}, \lambda) b_{\vec{k}, \lambda} \gamma_k(t) + \varepsilon_{ij}^*(-\hat{k}, \lambda) b_{-\vec{k}, \lambda}^+ \gamma_k^*(t) \right], \quad (5.3)$$

with

$$\left[b_{\vec{k}, \lambda}, b_{\vec{k}', \lambda'}^+ \right] = (2\pi)^2 \delta^{(3)}(\vec{k} - \vec{k}') \delta_{\lambda, \lambda'}. \quad (5.4)$$

The equation of motion for the eigenfunctions $\delta\phi_k(t)$ can be derived in the approximation of de-Sitter space from the second-order action

$$S_2 = \int d\eta' \frac{1}{(H\eta)^2} \left[(\delta\phi')^2 - (\partial_i \delta\phi)^2 \right], \quad (5.5)$$

(where $d\eta = dt/a(t)$ is the conformal time) and they are given by the well-known expression

$$u_k(\eta) = \frac{H}{\sqrt{2k^3}} (1 + ik\eta) e^{-ik\eta}. \quad (5.6)$$

In the same approximation, the eigenfunctions for the tensor modes $\gamma_k(\eta)$ are given by $u_k^T \equiv 2u_k$.

Let us now begin with the one-loop one-vertex part of the diagram (given in Fig. 3) which we label with the subscript $(1L, 1v)$; this can be written as [104], [105]

$$\langle \delta\phi_{\vec{k}_1}(\eta^*) \delta\phi_{\vec{k}_2}(\eta^*) \rangle_{(1L, 1v)} = i \int d\eta' \left\langle \left[H_I^{(4)}(\eta'), \delta\phi_{\vec{k}_1}(\eta^*) \delta\phi_{\vec{k}_2}(\eta^*) \right] \right\rangle. \quad (5.7)$$

We will study this in detail

$$\begin{aligned}
\langle \delta\phi_{\vec{k}_1}^-(\eta^*)\delta\phi_{\vec{k}_2}^-(\eta^*) \rangle_{(1L,1v)} &= 2i \int_{-\infty}^{\eta^*} d\eta' a^2(\eta') \int \frac{d^3x}{(2\pi)^3} \int d^3q_1 d^3q_2 d^3q_3 d^3q_4 \\
&\times e^{-i\sum_n \vec{q}_n \cdot \vec{x}} P_{ij}(iq_3^i)(iq_4^j) u_{k_1}(\eta^*) u_{q_3}^*(\eta') u_{k_2}(\eta^*) u_{q_4}^*(\eta') u_{q_1}(\eta') u_{q_2}^*(\eta') \\
&\times \delta^{(3)}(\vec{k}_1 + \vec{q}_3) \delta^{(3)}(\vec{k}_2 + \vec{q}_4) \delta^{(3)}(\vec{q}_1 + \vec{q}_2) + c.c., \tag{5.8}
\end{aligned}$$

where the extra factor of 2 accounts for the number of equivalent diagrams obtained by permuting the field contractions, $u_k(\eta)$ is given by Eq. (5.6) and

$$P_{ij} = \sum_{\lambda, \lambda'} \epsilon_{ik}^\lambda(\hat{q}) \epsilon_{kj}^{*\lambda'}(\hat{q}) = 2 \sin^2 \theta. \tag{5.9}$$

Integration and the use of the delta function lead to a simpler form

$$\begin{aligned}
\langle \delta\phi_{\vec{k}_1}^-(\eta^*)\delta\phi_{\vec{k}_2}^-(\eta^*) \rangle_{(1L,1v)} &= -i \delta^{(3)}(\vec{k}_1 + \vec{k}_2) \frac{H_*^4}{2k^4} \int \frac{d^3q}{q^3} \sin^2 \theta \int_{-\infty}^{\eta^*} \frac{d\eta'}{\eta'^2} e^{2ik(\eta' - \eta^*)} (1 + ik\eta^*)^2 \\
&\times (1 - ik\eta')^2 (1 + iq\eta') (1 - iq\eta') + c.c., \tag{5.10}
\end{aligned}$$

This equation is exact except for the approximation of using the de Sitter space formula for the scale factor, $a(\eta) = -(H\eta)^{-1}$, and evaluating the Hubble radius $H(\eta')$ at the time η^* . The reason why this is allowed is the following: the contribution to the integral w.r.t. time from regions well before horizon crossing is negligible compared to the contribution due to the region around horizon crossing [13, 104, 105]; in addition to that, we are choosing η to be just a few e-folds after horizon crossing, so we can assume that the Hubble radius (as well as any of the slow-roll parameters of the theory) will not undergo a big variation during this interval of time. The same approximation will be applied to the two-vertex diagrams.

We first solve the time integral. It is convenient to perform a change of variable like in [106], i.e. we set $x' = -k\eta'$ and $x^* = -k\eta^*$ so that

$$\begin{aligned}
\langle \delta\phi_{\vec{k}_1}^-(\eta^*)\delta\phi_{\vec{k}_2}^-(\eta^*) \rangle_{(1L,1v)} &= \delta^{(3)}(\vec{k}_1 + \vec{k}_2) \frac{H_*^4}{2k^4} \int \frac{d^3q}{q^3} \sin^2 \theta \operatorname{Im} \left[\int_{x^*}^{\infty} \frac{dx'}{k} \frac{k^2}{x'^2} e^{2i(x' - x^*)} \right. \\
&\times \left. (1 + ix^*)^2 (1 - ix')^2 (1 + i\frac{q}{k}x') (1 - i\frac{q}{k}x') \right]. \tag{5.11}
\end{aligned}$$

After integrating the imaginary part, we end up with the following result

$$\begin{aligned}
\langle \delta\phi_{\vec{k}_1}^-(\eta^*)\delta\phi_{\vec{k}_2}^-(\eta^*) \rangle_{(1L,1v)} &= \delta^{(3)}(\vec{k}_1 + \vec{k}_2) \frac{H_*^4}{2k^4} \int \frac{d^3q}{q^3} \sin^2 \theta \frac{2k^2(3 + x^{*2}) + q^2(5 + 5x^{*2} + 2x^{*4})}{4k} \\
&= \delta^{(3)}(\vec{k}_1 + \vec{k}_2) \frac{H_*^4}{2k^4} 2\pi \frac{4}{3} \left[\frac{k}{2} (3 + x^{*2}) \int \frac{dq}{q} + \frac{1}{4k} (5 + 5x^{*2} + 2x^{*4}) \int dq \right],
\end{aligned}$$

where the factor $4/3$ comes from integrating with respect to the azimuthal angle θ (notice that the reference frame in momentum space has been chosen in such a way that the external wave vector \vec{k} lies along the positive z axis). We now solve the momentum integrals. Both the logarithmic and the quadratic one exhibit ultraviolet divergences and the logarithmic part diverges also at very low momenta. Ultraviolet divergences

can be treated as in flat space; the infrared logarithmic divergence is fixed introducing a momentum lower cutoff ℓ^{-1} to be interpreted as a ‘box size’ [110, 111, 117, 118, 119] which can be fixed to be not much larger than the present horizon [110, 111]. As an example, consider the first integral of Eq. (5.12) which is convenient to split as follows

$$\int_{\ell^{-1}}^{\Lambda} \frac{dq}{q} = \int_{\ell^{-1}}^k \frac{dq}{q} + \int_k^{\Lambda} \frac{dq}{q}, \quad (5.12)$$

where we have introduced an upper cutoff Λ . The first integral gives $\ln(k\ell)$; the second integral can be renormalized introducing a counterterm $-\ln(\Lambda/k_0)$, where k_0 is a renormalization constant. The final result for Eq.(5.12) reads

$$\langle \delta\phi_{\vec{k}_1}(\eta^*) \delta\phi_{\vec{k}_2}(\eta^*) \rangle_{(1L,1v)} = \pi \delta^{(3)}(\vec{k}_1 + \vec{k}_2) \frac{2H^4}{3k^3} (3 + x^{*2}) [\ln(k\ell) - \ln(k) + \alpha],$$

where α is a left over constant from renormalization.

Let us now focus on the one-loop contribution from the 3rd order action with the gravitons (see Fig. 4 for its diagrammatic representation)

$$\begin{aligned} \langle \delta\phi_{\vec{k}_1}(\eta^*) \delta\phi_{\vec{k}_2}(\eta^*) \rangle_{(1L,2v)} &= \frac{(-i)^2}{2} \langle T [\delta\phi_{\vec{k}_1}(\eta^*) \delta\phi_{\vec{k}_2}(\eta^*) \int_{-\infty}^{\eta^*} d\eta' (H_I^+(\eta') - H_I^-(\eta')) \\ &\quad \times \int_{-\infty}^{\eta^*} d\eta'' (H_I^+(\eta'') - H_I^-(\eta''))] \rangle \\ &= \frac{(-i)^2}{2} \langle T [\delta\phi_{\vec{k}_1}(\eta^*) \delta\phi_{\vec{k}_2}(\eta^*) (A + B + C + D)] \rangle, \end{aligned} \quad (5.13)$$

where

$$A \equiv \int_{-\infty}^{\eta^*} d\eta' H_I^+ \int_{-\infty}^{\eta^*} d\eta'' H_I^+, \quad (5.14)$$

$$B \equiv \int_{-\infty}^{\eta^*} d\eta' H_I^- \int_{-\infty}^{\eta^*} d\eta'' H_I^-, \quad (5.15)$$

$$C \equiv - \int_{-\infty}^{\eta^*} d\eta' H_I^+ \int_{-\infty}^{\eta^*} d\eta'' H_I^-, \quad (5.16)$$

$$D \equiv - \int_{-\infty}^{\eta^*} d\eta' H_I^- \int_{-\infty}^{\eta^*} d\eta'' H_I^+. \quad (5.17)$$

It is easy to check that $B = A^*$ and $C = C^* = D$. We can write Eq. (5.13) as

$$\begin{aligned} \langle \delta\phi_{\vec{k}_1}(\eta^*) \delta\phi_{\vec{k}_2}(\eta^*) \rangle_{(1L,2v)} &= 4(-i)^2 \delta^{(3)}(\vec{k}_1 + \vec{k}_2) k^4 \int d^3q \sin^4\theta \\ &\quad \times \int_{-\infty}^{\eta^*} \frac{d\eta'}{(H\eta')^2} \int_{-\infty}^{\eta^*} \frac{d\eta''}{(H\eta'')^2} (w_f^A + w_f^B + w_f^C + w_f^D), \end{aligned} \quad (5.18)$$

where the factor $\sin^4\theta$ comes from contractions of the polarization tensors with external momenta [120]

$$\epsilon_{ij}(\vec{q}) k^i k^j = \frac{k^2}{\sqrt{2}} \left[1 - \left(\frac{\vec{q} \cdot \vec{k}}{qk} \right)^2 \right] = \frac{k^2}{\sqrt{2}} \sin^2\theta, \quad (5.19)$$

and the wave functions w_f are

$$\begin{aligned}
w_f^A(\eta', \eta'') &= u_k(\eta^*) u_k^*(\eta') u_k(\eta^*) u_k^*(\eta'') [u_{|\vec{k}-\vec{q}|}(\eta') u_{|\vec{k}-\vec{q}|}^*(\eta'') u_q(\eta') u_q^*(\eta'') \\
&\quad \times \Theta(\eta' - \eta'') + u_{|\vec{k}-\vec{q}|}^*(\eta') u_{|\vec{k}-\vec{q}|}(\eta'') u_q^*(\eta') u_q(\eta'') \Theta(\eta'' - \eta')], \\
w_f^B(\eta', \eta'') &= u_k^*(\eta^*) u_k(\eta') u_k^*(\eta^*) u_k(\eta'') [u_{|\vec{k}-\vec{q}|}(\eta') u_{|\vec{k}-\vec{q}|}^*(\eta'') u_q(\eta') u_q^*(\eta'') \\
&\quad \times \Theta(\eta'' - \eta') + u_{|\vec{k}-\vec{q}|}^*(\eta') u_{|\vec{k}-\vec{q}|}(\eta'') u_q^*(\eta') u_q(\eta'') \Theta(\eta' - \eta'')], \\
w_f^C(\eta', \eta'') &= -u_k(\eta^*) u_k^*(\eta') u_k^*(\eta^*) u_k(\eta'') u_{|\vec{k}-\vec{q}|}^*(\eta') u_{|\vec{k}-\vec{q}|}(\eta'') u_q^*(\eta') u_q(\eta''), \\
w_f^D(\eta', \eta'') &= -u_k^*(\eta^*) u_k(\eta') u_k(\eta^*) u_k^*(\eta'') u_{|\vec{k}-\vec{q}|}(\eta') u_{|\vec{k}-\vec{q}|}^*(\eta'') u_q(\eta') u_q^*(\eta''),
\end{aligned}$$

so $w_f^C(\eta', \eta'') = w_f^D(\eta'', \eta')$ and is a real number and $w_f^B(\eta', \eta'') = w_f^{A*}(\eta'', \eta')$. We will label the two contributions by A and C , so that the one-loop contribution with two vertices to the two point function will be broken into two parts

$$\langle \delta\phi_{\vec{k}_1}^-(\eta^*) \delta\phi_{\vec{k}_2}^-(\eta^*) \rangle_{(1L,2v)} = \langle \delta\phi_{\vec{k}_1}^-(\eta^*) \delta\phi_{\vec{k}_2}^-(\eta^*) \rangle_{(1L,2v)}^A + \langle \delta\phi_{\vec{k}_1}^-(\eta^*) \delta\phi_{\vec{k}_2}^-(\eta^*) \rangle_{(1L,2v)}^C. \quad (5.20)$$

Let's look in details at the two parts.

$$\begin{aligned}
\langle \delta\phi_{\vec{k}_1}^-(\eta^*) \delta\phi_{\vec{k}_2}^-(\eta^*) \rangle_{(1L,2v)}^A &= -\delta^{(3)}(\vec{k}_1 + \vec{k}_2) \frac{H_*^4}{2k^2} \int \frac{d^3q}{q^3} \frac{\sin^4\theta}{|\vec{k}-\vec{q}|^3} e^{-2ik\eta^*} (1 + ik\eta^*)^2 \\
&\quad \times \int_{-\infty}^{\eta^*} \frac{d\eta'}{\eta'^2} e^{i\eta'(k-q-|\vec{k}-\vec{q}|)} (1 - ik\eta') (1 + iq\eta') (1 + i|\vec{k}-\vec{q}|\eta') \\
&\quad \times \int_{-\infty}^{\eta'} \frac{d\eta''}{\eta''^2} e^{i(k+q+|\vec{k}-\vec{q}|)\eta''} (1 - ik\eta'') (1 - iq\eta'') (1 + i|\vec{k}-\vec{q}|\eta'') + c.c.
\end{aligned} \quad (5.21)$$

The second time integral has $e^{ig\eta''} \left[-\frac{1}{\eta''} + \frac{c}{g}\eta'' - i\left(\frac{gb-c}{g^2}\right) \right]$ as its primitive function, where $g \equiv k + q + |\vec{k} - \vec{q}|$, $b \equiv -qk - (q+k)|\vec{k} - \vec{q}|$ and $c \equiv qk|\vec{k} - \vec{q}|$. This should be evaluated between $-\infty$ and η' . It is soon evident that the lower bound represents a problem for this evaluation. We need to remind ourself, though, that the choice of the integration time contour needs to be deformed and to cross the complex plane to account for the right choice of the vacuum [13]. This is done by integrating in a slightly imaginary direction, i.e. taking $\eta'' \rightarrow \eta'' + i\epsilon|\eta''|$, where ϵ is a fixed small real number; so for example

$$\int_{-\infty}^{\eta} d\eta' e^{ik\eta'} = \frac{e^{ik\eta}}{ik}. \quad (5.22)$$

With this contour prescription, our integral in η'' vanishes at $-\infty$. Performing the same change of variables as in (8.14)

$$\begin{aligned}
\langle \delta\phi_{\vec{k}_1}^-(\eta^*) \delta\phi_{\vec{k}_2}^-(\eta^*) \rangle_{(1L,2v)}^A &= -\delta^{(3)}(\vec{k}_1 + \vec{k}_2) \frac{H_*^4}{k^2} \int \frac{d^3q}{q^3} \frac{\sin^4\theta}{|\vec{k}-\vec{q}|^3} \text{Re} \left[\int_{x^*}^{\infty} \frac{dx'}{k} \frac{k}{x'} 2e^{2i(x'-x^*)} (1 + ix^*)^2 \right. \\
&\quad \times \left. \left(\frac{k}{x'} - \frac{c}{kg}x' - i\left(\frac{gb-c}{g^2}\right) \right) \left(1 - i\frac{d}{k}x' + \frac{s}{k^2}x'^2 - i\frac{c}{k^3}x'^3 \right) \right],
\end{aligned} \quad (5.23)$$

where $d \equiv q - k + |\vec{k} - \vec{q}|$ and $s \equiv kq + (k - q)|\vec{k} - \vec{q}|$. The result of the integration w.r.t. time is a polynomial function of $\sin 2x^*$, $\cos 2x^*$, $\text{Si}(2x^*)$, $\text{Ci}(2x^*)$ and their products with coefficients which depend on g, b, c, d, s, k . Notice that in the large scale limit $x^* \rightarrow 0$ a singularity similar to the one found in [106] shows up in our result (see also [121] for a recent discussion on these kind of singularities). However, by evaluating the power spectrum of $\delta\phi$ just a few e-folds after horizon crossing, we are safe from these kind of singular behaviour [109].

The next step consist in performing the momentum integral. The integrals we need to evaluate are of the following kind

$$\int \frac{d^3q}{q^3} \frac{\sin^4\theta}{|\vec{q} - \vec{k}|^3} f(\vec{q}), \quad (5.24)$$

where $f(\vec{q})$ is a sum of functions of momentum. Let us begin for simplicity by considering the constant term of the sum, i.e. let us study

$$\int \frac{d^3q}{q^3} \frac{\sin^4\theta}{|\vec{q} - \vec{k}|^3}. \quad (5.25)$$

For the specific case of equation (5.25) the integrand function has singularities at $\vec{q} = 0$ and at $\vec{q} = \vec{k}$ and shows no ultraviolet singularities. Based on an approximate evaluation performed considering a sphere of radius ℓ^{-1} around $\vec{q} = 0$, where $\ell^{-1} \ll k$, the integral is proportional to a function $\ln(k\ell)$. The same result can be obtained working in a small sphere around $\vec{q} = \vec{k}$ after a change of variables $\vec{q}_0 = \vec{q} - \vec{k}$. The contribution from large values of q is negligible w.r.t. the ones from the singular points, so the integral over the whole momentum space is expected to be proportional to $\ln(k\ell)$. The exact value of the integral can be found after a change of variable from the (q, θ) to the (q, p) space, where $p \equiv |\vec{q} - \vec{k}|$ and is equal to $(16\pi/225k^3)(1 + 30 \ln(k\ell)) \sim k^{-3}(10^{-1} + 10 \ln(k\ell))$.

Integrating Eq. (5.23), we find ultraviolet power law and logarithmic singularities in addition to infrared logarithmic contributions. The final result of the integration is a function of $x^* = e^{-N_*}$, where $N_* = \ln(a_*/a_k)$ is the number of e-foldings from horizon crossing

$$\langle \delta\phi_{\vec{k}_1}(\eta^*) \delta\phi_{\vec{k}_2}(\eta^*) \rangle_{(1L, 2v)}^A = \pi \delta^{(3)}(\vec{k}_1 + \vec{k}_2) \frac{H_*^4}{k^3} (a_1 \ln(k) + a_2 \ln(k\ell) + a_3), \quad (5.26)$$

where a_1, a_2 and a_3 are functions of x^* (see Appendix 15.3). We are calculating the two point function for the scalar field a few e-foldings after horizon crossing, so x^* may be chosen to range between 10^{-1} and 10^{-2} . In this range $a_1 \sim \mathcal{O}(1)$ and negative, $a_2 = -16/(15x_*^2) + (8/15)(5 - 8\text{Ci}(2x_*))$ and $a_3 = -8/(225x_*^2) + \mathcal{O}(1) + \rho$, where ρ is a left-over scheme-dependent renormalization constant of the kind present in equation (5.13).

Let us now move to part C of Eq. (5.13) which we give below

$$\begin{aligned}
\langle \delta\phi_{\vec{k}_1}(\eta^*) \delta\phi_{\vec{k}_2}(\eta^*) \rangle_{(1L,2v)}^C &= \delta^{(3)}(\vec{k}_1 + \vec{k}_2) \frac{H_*^4}{2k^2} \int \frac{d^3q}{q^3} \frac{\sin^4\theta}{|\vec{k} - \vec{q}|^3} (1 + (k\eta^*)^2) \\
&\times \int_{-\infty}^{\eta^*} \frac{d\eta'}{\eta'^2} e^{ig\eta'} Q(\eta') \int_{-\infty}^{\eta^*} \frac{d\eta''}{\eta''^2} e^{-ig\eta''} Q^*(\eta'') \\
&= \delta^{(3)}(\vec{k}_1 + \vec{k}_2) \frac{H_*^4}{2k^2} \int \frac{d^3q}{q^3} \frac{\sin^4\theta}{|\vec{k} - \vec{q}|^3} (1 + (k\eta^*)^2) \\
&\times \left[\left(\text{Re} \int d\eta' e^{ig\eta'} \frac{Q(\eta')}{\eta'^2} \right)^2 + \left(\text{Im} \int d\eta' e^{ig\eta'} \frac{Q(\eta')}{\eta'^2} \right)^2 \right], \quad (5.27)
\end{aligned}$$

where $Q(\eta') \equiv 1 + ig\eta' + b\eta'^2 - ic\eta'^3$. Let us integrate over conformal time

$$\int_{x^*}^{\infty} \frac{dx'}{k} \frac{k^2}{x'^2} e^{-i\frac{g}{k}x'} \left[1 + i\frac{g}{k}x' + \frac{b}{k^2}x'^2 - i\frac{c}{k^3}x'^3 \right] = \frac{e^{-i\frac{g}{k}x^*}}{k^2} \left(-\frac{k^3}{x} + \frac{ck}{g}x^* + i\frac{(gb-c)k^2}{g^2} \right), \quad (5.28)$$

where again the integration has been performed by continuing η' to the complex plane, i.e. ($\eta' \rightarrow \eta' + i\epsilon|\eta'|$), and then taking the limit $\epsilon \rightarrow 0$.

We are now ready to integrate over momentum

$$\begin{aligned}
\langle \delta\phi_{\vec{k}_1}(\eta^*) \delta\phi_{\vec{k}_2}(\eta^*) \rangle_{(1L,2v)}^C &= \delta^{(3)}(\vec{k}_1 + \vec{k}_2) \frac{H_*^4}{2k^2} \int \frac{d^3q}{q^3} \frac{\sin^4\theta}{|\vec{k} - \vec{q}|^3} (1 + x^{*2}) \\
&\times \left(\frac{k^4}{x^{*2}} - \frac{2k^2c}{g} - \frac{2k^2bc}{g^3} + \frac{k^2c^2}{g^4} + \frac{k^2b^2}{g^2} + \frac{x^{*2}c^2}{g^2} \right). \quad (5.29)
\end{aligned}$$

Similarly to what we have done in part A, one can check that there are no ultraviolet singularities in the remaining five integrals although some infrared logarithmic contributions are still present and the final result is

$$\langle \delta\phi_{\vec{k}_1}(\eta^*) \delta\phi_{\vec{k}_2}(\eta^*) \rangle_{(1L,2v)}^C = \pi \delta^{(3)}(\vec{k}_1 + \vec{k}_2) \frac{H_*^4}{k^3} (c_1 + c_2 \ln(k\ell)), \quad (5.30)$$

where $c_1 = (1/225)(8/x^{*2} + 107 + 50x^{*2})$ and $c_2 = (16/15x^{*2}) + (4/15)$. Notice that the $(x^*)^{-2}$ coefficients in c_1 and c_2 exactly cancels the $(x^*)^{-2}$ coefficients in a_2 and a_3 . This is not surprising: based on [106, 109], we expect we might observe a logarithmic singularity if we push $x^* \rightarrow 0$ in our results (which is indeed present in the $\text{Ci}(2x^*)$ term of a_2), but no power-law singularities are actually expected.

6. Complete expression for P_ζ at one loop

Let us now collect our results in the final formula for the power spectrum of the curvature perturbation ζ computed up to one-loop level. This can be derived from Eq. (3.4), which follows from the δN formula. Summing the main results of the previous section, Eqs. (5.13), (5.26) and (5.30), we obtain the one-loop graviton correction to the inflaton power spectrum

$$\langle \delta\phi_{\vec{k}_1}^-(\eta^*) \delta\phi_{\vec{k}_2}^-(\eta^*) \rangle_{1L} = \pi \delta^{(3)}(\vec{k}_1 + \vec{k}_2) \frac{H^4}{k^3} [f_1 \ln(k) + f_2 \ln(k\ell) + f_3], \quad (6.1)$$

where

$$f_1 = -\frac{2}{15} (25 + 15x^{*2} + 4x^{*4}), \quad (6.2)$$

$$f_2 = 2 + (2/3)x^{*2} + a_2 + c_2, \quad (6.3)$$

and f_3 is given by a left-over scheme-dependent renormalization constant plus contributions of order $\mathcal{O}(1)$ (see Appendix 15.3 for the complete expressions of a_2 , c_2 and f_3). If we calculate the two point function of $\delta\phi$ a few e-foldings after horizon crossing, i.e. x^* ranges for example between 10^{-1} and 10^{-2} , f_1 reduces to a negative constant of order $\mathcal{O}(1)$ and $f_2 \sim 4(1 - \text{Ci}(2x^*)) \sim \mathcal{O}(10)$. In the limit where $x^* \rightarrow 1$ both f_1 and f_2 turn out to be of order unity.

In order to understand which is the dominant contribution in Eq. (3.5) and how big it is, one needs to (i) know the slow-roll order of the coefficients $N^{(i)}$: $N^{(1)} \sim \epsilon^{-1/2}$, $N^{(2)} \sim \epsilon^0$, $N^{(3)} \sim \epsilon^{1/2}$; (ii) compute the integrals involving the power spectrum $P(q)$. This is discussed in details in Ref. [109] (for the case of scalar perturbations only), see in particular Sec IV of [109]. It turns out that the crucial quantity is represented by the number of e-foldings of inflation between the times of horizon exit of the mode ℓ^{-1} , which corresponds to the infrared cutoff, and the time of horizon exit of the mode k we want to observe. However, to deal with observable quantities one has to choose ℓ not much bigger than the present cosmological horizon H_0^{-1} [110, 111].

The relevant point about Eq. (6.1) is that it gives in Eq. (3.5) a contribution which is of the same order of magnitude as those coming from loops which accounts for scalar perturbations only. Since in terms of the slow-roll parameters $(N^{(1)})^2 \sim \epsilon_*^{-1}$ the magnitude of the one-loop graviton correction turns out to be

$$\Delta P_\zeta^{\text{1loop}}(k) \sim \frac{2\pi^2}{k^3} \alpha(k) \frac{1}{\epsilon_*} \mathcal{P}_*^2(k), \quad (6.4)$$

where we have used Eq. (3.6) for the power spectrum of the inflaton field. In Eq. (6.4) $\alpha(k)$ includes the various coefficients of Eq. (6.1), and it is $\mathcal{O}(1)$. Eq. (6.4) allows a more direct comparison with the results of Ref. [109], showing that the graviton contributions to the one-loop corrections are comparable to the ones computed only from scalar interactions. Notice that also for the tensor contributions we find terms of the form $\ln(k)$.

Summarizing our work: beyond linear order, the tensor perturbation modes produced during inflation unavoidably mix with scalar modes; this fact alone would require to include the tensor modes for a self-consistent computation. Most importantly, despite a naive expectation suggested by the fact that the power spectrum of the tensor modes is suppressed on large scales with respect to that of the curvature (scalar) perturbations, our results show explicitly that their inclusion is necessary since their contribution is not at all negligible with respect to the loop corrections arising from interactions involving the inflaton field only.

7. Study of perturbations in anisotropic cosmologies

The Bianchi models represent a classification of all homogeneous and anisotropic cosmologies. In the cosmic no-hair conjecture, any initial background, in the presence of a positive cosmological constant, eventually evolves in a de Sitter universe, where all the initial existing anisotropies are rapidly washed out. This conjecture was proven to be true for all of the Bianchi models except for the Bianchi-IX by Wald in [72]. For the Bianchi-IX this result is also true under the assumption that the cosmological constant overcomes the spatial curvature terms. More recently, there has been a revived interest in homogeneous but anisotropic models of the early Universe and several attempts to understand what kind of observational signatures these models might produce. A close look has in fact been given to the fluctuations in the metric and in the energy tensor developing during a hypothetical anisotropic stage either in a pre-inflationary epoch [78, 79] or during inflation itself [76, 77, 144]. The Bianchi-I model is the simplest of all anisotropic and homogeneous models and has often been the choice for such kind of studies. In the Bianchi-I model, the metric has a form

$$ds^2 = dt^2 - e^{2\alpha} \left(e^{2\beta} \right)_{ij} dx^i dx^j \quad (7.1)$$

where α is a function of time as well as β_{ij} . The latter is a 3×3 diagonal traceless matrix that anisotropizes the volume expansion

$$\left(e^{2\beta} \right)_{ij} = \delta_{ij} e^{2\beta_i}, \quad \sum_{i=1}^3 \beta_i = 0. \quad (7.2)$$

A metric described by Eq. (7.1) but with constant functions α and β is often presented in the form

$$ds^2 = dt^2 - t^{2p_1} dx_1^2 + t^{2p_2} dx_2^2 + t^{2p_3} dx_3^2, \quad (7.3)$$

where the coefficients p_1, p_2, p_3 are numerical constants satisfying the relation

$$p_1 + p_2 + p_3 = p_1^2 + p_2^2 + p_3^2 = 1. \quad (7.4)$$

The three parameters can be equal in pairs in the cases $(-\frac{1}{3}, \frac{2}{3}, \frac{2}{3})$ and $(0, 0, 1)$. In all other cases they are distinct, one being negative and the other two being positive. The universe described by (7.3) is spatially flat and, for any possible value of the coefficients p_i , its volume element is equal to $\sqrt{-g^3} d^3x = t d^3x$, where g^3 is the determinant of the three-metric.

In [78], we consider a pre-inflationary epoch characterized by an initially expanding type-I Bianchi universe that evolves with an energy density not yet dominated by a cosmological constant. If by the time the cosmological constant has become the dominant form of energy the metric is still homogeneous, the Universe will eventually enter a de Sitter epoch; the fluctuations of this metric and their growth, however, can play an important role in determining the evolution of the Universe at this early stage. The question we ask concerns the kind of initial conditions are needed for the Universe to remain homogeneous by the time the cosmological constant eventually dominates the total energy density. We employ a metric as in Eq. (7.1) for the background and solve

the first order Einstein equations for the perturbations considering a pressureless fluid. As expected, the anisotropy in the background is responsible for a coupling at the linear order of one of the tensor modes with the density contrast, so there is a correlation between the scalar and the gravitational perturbations. Moreover, the evolutions of the two tensor modes, the “free” one and the one that is coupled to the scalar, are very different from each other and very much depend on the scale of interest. It turns out that, for a reasonably large set of initial conditions, the growth of these perturbations could be fast enough so as to lead the Universe into an inhomogeneous state before the cosmological constant becomes the dominant form of energy.

A similar scenario was investigated in [79], where both expanding and contracting Bianchi-I Universes are studied during a pre-inflationary era. A scalar field dominates the matter content. In particular, the evolution of the gravitational perturbations of the metric (not to be confused with the gravitational waves generated from the vacuum fluctuation during inflation) is analysed, to point out that they may grow to significantly alter the geometry of spacetime before inflation begins and, in any case, they may leave an imprint on the observed CMB power spectrum.

Also in [76, 77] a perturbative analysis was performed for Bianchi-I cosmologies leading to anisotropic inflation; in addition to that, predictions for the power spectrum of curvature fluctuations and gravity waves produced during inflation are provided for this models. Finally, a “vector hair” model and its observational consequences in terms of anisotropic signatures in the cosmological correlation functions and primordial gravitational waves, were recently discussed in [144].

8. Inflation and primordial vector fields

The attempt to explain some of the CMB “anomalous” features as the indication of a break of statistical isotropy is the main reason behind ours and many of the existing inflationary models populated by vector fields, but not the only one. The first one of these models [122] was formulated with the goal of producing inflation by the action of vector fields, without having to invoke the existence of a scalar field. The same motivations inspired the works that followed [123, 124, 125]. Lately, models where primordial vector fields can leave an imprint on the CMB have been formulated as an alternative to the basic inflationary scenario, in the search for interesting non-Gaussianity predictions [90, 91, 92, 93, 94, 95, 96, 97, 98]. Finally, vector fields models of dark energy have been proposed [126, 127, 128, 129, 130, 131]. All this appears to us as a rich bag of motivations for investigating these scenarios.

Before we quickly sketch some of them and list the results so far achieved in this direction, it is important to briefly indicate and explain the main issues and difficulties that these models have been facing. We will also shortly discuss the mechanisms of production of the curvature fluctuations in these models.

Building a model where primordial vector fields can drive inflation and/or produce the observed spectrum of large scale fluctuations requires a more complex Lagrangian than the basic gauge invariant $\mathcal{L}_{vector} = -(\sqrt{-g}/4)F^{\mu\nu}F_{\mu\nu}$. In fact, for a conformally invariant theory as the one described by \mathcal{L}_{vector} , vector fields fluctuations are not excited on superhorizon scales. It is then necessary to modify the Lagrangian. For some of the existing models, these modifications have been done to the expense of destabilizing the theory, by “switching on” unphysical degrees of freedom. This was pointed out in [132, 133, 134], where a large variety of vector field models was analyzed in which longitudinal polarization modes exist that are endowed with negative squared masses (the “wrong” signs of the masses are imposed for the theory to satisfy the constraints that allow a suitable background evolution). It turns out that, in a range of interest of the theory, these fields acquire negative total energy, i.e. behave like “ghosts”, the presence of which is known to be responsible for an unstable vacuum. A related problem for some of these theories is represented by the existence of instabilities affecting the equations of motion of the ghost fields [132, 133, 134].

In the remaining part of this section, we are going to present some of these models together with some recent attempts to overcome their limits.

In all of the models we will consider, primordial vector fields fluctuations end up either being entirely responsible for or only partially contributing to the curvature fluctuations at late times. This can happen through different mechanisms. If the vector fields affects the universe expansion during inflation, its contribution ζ_A to the total ζ can be derived from combining the definition of the number of e-foldings ($N = \int H dt$) with the Einstein equation ($H^2 = (8\pi G/3)(\rho_\phi + \rho_A)$), ρ_A being the energy density of the vector field and

ρ_ϕ the inflaton energy density) and using the δN expansion of the curvature fluctuation in terms of both the inflaton and the vector fields fluctuations (see Sec. 9). To lowest order we have [92]

$$\zeta_A = \frac{A_i}{2m_P^2} \delta A_i, \quad (8.1)$$

where a single vector field has been taken into account for simplicity (m_P is the reduced Planck mass, A is the background value of the field and δA its perturbation). When calculating the amplitude of non-Gaussianity in Sec. 10, we will refer to this case as “vector inflation” for simplicity.

A different fluctuation production process is the curvaton mechanism which was initially formulated for scalar theories but it is also applicable to vectors [135, 136]. Specifically, inflation is driven by a scalar field, whereas the curvaton field(s) (now played by the vectors), has a very small (compared to the Hubble rate) mass during inflation. Towards the end of the inflationary epoch, the Hubble rate value starts decreasing until it equates the vector mass; when this eventually happens, the curvaton begins to oscillate and it will then dissipate its energy into radiation. The curvaton becomes responsible for a fraction of the total curvature fluctuation that is proportional to a parameter, r , related to the ratio between the curvaton energy density and the total energy density of the universe at the epoch of the curvaton decay [92]

$$\zeta_A = \frac{r}{3} \frac{\delta \rho_A}{\rho_A}, \quad (8.2)$$

where $r \equiv 3\rho_A/(3\rho_A + 4\rho_\phi)$. Anisotropy bounds on the power spectrum favour small values of r .

From Eqs. (8.1) and (8.2) we can see that, depending on which one of these two mechanisms of production of the curvature fluctuations is considered, different coefficients will result in the δN expansion (see Eq. (9.4)).

In this section we will describe both models where inflation is intended to be vector-field driven and those models in which, instead, the role of the inflaton is played by a scalar field, whereas the energy of the vector is a subdominant contribution to the total energy density of the universe during the entire inflationary phase.

8.1. Self-coupled vector field models

A pioneer work on vector field driven inflation was formulated by L. H. Ford [122], who considered a single self-coupled field A_μ with a Lagrangian

$$L_{vector} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + V(\psi) \quad (8.3)$$

where $F_{\mu\nu} \equiv \partial_\mu B_\nu - \partial_\nu B_\mu$ and the potential V is a function of $\psi \equiv B_\alpha B^\alpha$. Different scenarios of expansion are analyzed by the author for different functions V . The universe expands anisotropically at the end of the inflationary era and this anisotropy either survives until late times or is damped out depending on the shape and the location of

the minima of the potential.

The study of perturbations in a similar model was proposed by Dimopoulos in [135] where he showed that for a Lagrangian

$$L_{vector} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} + \frac{1}{2}m^2 B_\mu B^\mu \quad (8.4)$$

and for $m^2 \simeq -2H^2$, the transverse mode of the vector field is governed by the same equation of motion as a light scalar field in a de Sitter stage. A suitable superhorizon power spectrum of fluctuations could therefore arise. In order to prevent production of large scale anisotropy, in this model the vector field plays the role of the curvaton while inflation is driven by a scalar field.

8.2. Vector-field coupled to gravity

The Lagrangian in Eq. (8.4) may be also intended, at least during inflation, as including a non-minimal coupling of the vector field to gravity; indeed the mass term can be rewritten as

$$L_{vector} \supset \frac{1}{2} \left(m_0^2 + \xi R \right) B_\mu B^\mu \quad (8.5)$$

where, for the whole duration of the inflationary era, the bare mass m_0 is assumed to be much smaller than the Hubble rate and the Ricci scalar $R = -6 \left[\frac{\ddot{a}}{a} + \left(\frac{\dot{a}}{a} \right)^2 \right]$ can be approximated as $R \simeq -12H^2$. For the specific value $\xi = 1/6$, Eq. (8.4) is retrieved.

For the Lagrangian just presented, Golovnev et al [123] proved that the problem of excessive anisotropy production in the case where inflation is driven by vector fields can be avoided if either a triplet of mutually orthogonal or a large number N of randomly oriented vector fields is considered.

The Lagrangian (8.5) with $\xi = 1/6$ was also employed in [136], where inflation is scalar-field-driven and a primordial vector field affects large-scale curvature fluctuations and, similarly, in [137], which includes a study of the backreaction of the vector field on the dynamics of expansion, by introducing a Bianchi type-I metric.

8.3. Ackerman-Carroll-Wise (ACW) model

A model was proposed in [138] where Lagrange multipliers (λ) are employed to determine a fixed norm primordial vector field $B_\mu B^\mu = m^2$

$$L_{vector} \supset \lambda \left(B^\mu B_\mu - m^2 \right) - \rho_\Lambda \quad (8.6)$$

where ρ_Λ is a vacuum energy. The expansion rate in this scenario is anisotropic: if we orient the x -axis of the spatial frame along the direction determined by the vector field, we find two different Hubble rates: along the x -direction it is equal to

$$H_b^2 = \frac{\rho_\Lambda}{m_P^2} \frac{1}{P(\mu)}, \quad (8.7)$$

and it is given by $H_a = (1 + c\mu^2)H_b$ along the orthogonal directions; $\mu \equiv m/m_P$, P is a polynomial function of μ and c is a parameter appearing in the kinetic part of the Lagrangian that we omitted in (8.6) (see [138] for its complete expression). As expected, an isotropic expansion is recovered if the vev of the vector field is set to zero.

8.4. Models with varying gauge coupling

Most of the models mentioned so far successfully solve the problem of attaining a slow-roll regime for the vector-fields without imposing too many restrictions on the parameters of the theory and of avoiding excessive production of anisotropy at late times. None of them though escapes those instabilities related to the negative energy of the longitudinal modes (although a study of the instabilities for fixed-norm field models was done in [139] where some stable cases with non-canonical kinetic terms were found). As discussed in [132, 133, 134], in the self-coupled model a ghost appears at small (compared to the horizon) wavelengths; in the non-minimally coupled and in the fixed-norm cases instead the instability concerns the region around horizon crossing.

Models with varying gauge coupling can overcome the problem of instabilities and have recently attracted quite some attention. In [90], the authors consider a model of hybrid inflation [140, 141, 142, 143] with the introduction of a massless vector field

$$L \supset \frac{1}{2} (\partial_\mu \phi \partial^\mu \phi + \partial_\mu \chi \partial^\mu \chi) - \frac{1}{4} f^2(\phi) F_{\mu\nu} F^{\mu\nu} + V(\phi, \chi, B_\mu) \quad (8.8)$$

where ϕ is the inflaton and χ is the so-called “waterfall” field. The potential V is chosen in such a way as to preserve gauge invariance; this way the longitudinal mode disappears and instabilities are avoided.

Similarly, Kanno et al [144] consider a vector field Lagrangian of the type

$$L_{vector} = -\frac{1}{4} f^2(\phi) F^{\mu\nu} F_{\mu\nu}, \quad (8.9)$$

but in a basic scalar field driven inflation model. Very recently, in [145, 146] the linear perturbations in these kind of models have been investigated.

Finally, in [92, 96] varying mass vector field models have been introduced

$$L_{vector} = -\frac{1}{4} f^2(\phi) F^{\mu\nu} F_{\mu\nu} + \frac{1}{2} m^2 B_\mu B^\mu, \quad (8.10)$$

where $f \simeq a^\alpha$ and $m \simeq a$ (a is the scale factor and α is a numerical coefficient). The special cases $\alpha = 1$ and $\alpha = -2$ are of special interest. In fact, introducing the fields \tilde{A}_μ and A_μ , related to one another by $\tilde{A}_\mu \equiv f B_\mu = a A_\mu$ (\tilde{A}_μ and A_μ are respectively the comoving and the physical vectors), it is possible to verify that the physical gauge fields are governed by the same equations of motion as a light scalar field in a de Sitter background. Vector fields in this theory can then generate the observed (almost) scale invariant primordial power spectrum.

8.5. $SU(2)$ vector model

Let us consider some models where inflation is driven by a scalar field in the presence of an $SU(2)$ vector multiplet [97, 98]. A fairly general Lagrangian can be the following

$$S = \int d^4x \sqrt{-g} \left[\frac{m_P^2 R}{2} - \frac{f^2(\phi)}{4} g^{\mu\alpha} g^{\nu\beta} \sum_{a=1,2,3} F_{\mu\nu}^a F_{\alpha\beta}^a - \frac{M^2}{2} g^{\mu\nu} \sum_{a=1,2,3} B_\mu^a B_\nu^a + L_\phi \right], \quad (8.11)$$

where L_ϕ is the Lagrangian of the scalar field and $F_{\mu\nu}^a \equiv \partial_\mu B_\nu^a - \partial_\nu B_\mu^a + g_c \varepsilon^{abc} B_\mu^b B_\nu^c$ (g_c is the $SU(2)$ gauge coupling). Both f and the effective mass M can be viewed as generic functions of time. The fields B_μ^a are comoving and related to the physical fields by $A_\mu^a = (B_0^a, B_i^a/a)$. The free field operators can be Fourier expanded in their creation and annihilation operators

$$\delta A_i^a(\vec{x}, \eta) = \int \frac{d^3q}{(2\pi)^3} e^{i\vec{q}\cdot\vec{x}} \sum_{\lambda=L,R,long} \left[e_i^\lambda(\hat{q}) a_{\vec{q}}^{a,\lambda} \delta A_\lambda^a(q, \eta) + e_i^{*\lambda}(-\hat{q}) (a_{-\vec{q}}^{a,\lambda})^\dagger \delta A_\lambda^{*a}(q, \eta) \right], \quad (8.12)$$

where the polarization index λ runs over left (L), right (R) and longitudinal ($long$) modes and

$$\left[a_{\vec{k}}^{a,\lambda}, (a_{\vec{k}'}^{a',\lambda'})^\dagger \right] = (2\pi)^3 \delta_{a,a'} \delta_{\lambda,\lambda'} \delta^{(3)}(\vec{k} - \vec{k}'). \quad (8.13)$$

Here η the conformal time ($d\eta = dt/a(t)$). Once the functional forms of f and M have been specified, the equations of motion for the vector bosons can be written. We compute cosmological correlation functions up to fourth order considering an action as in (8.11). The expression of the correlators that we derive, prior to explicitating the wavefunction for the gauge bosons, apply to any $SU(2)$ theory with an action as in (8.11), both for what we will call the ‘‘Abelian’’ and for the ‘‘non-Abelian’’ contributions. In particular, the structure of the interaction Hamiltonian is independent of the functional dependence of f and M and determines the general form of and the anisotropy coefficients appearing in the final ‘‘non-Abelian’’ expressions (see Sec. 9). When it comes to explicitate the wavefunctions, a choice that can help keeping the result as easy to generalize as possible is the following

$$\delta B^T = -\frac{\sqrt{\pi x}}{2\sqrt{k}} \left[J_{3/2}(x) + iJ_{-3/2}(x) \right], \quad (8.14)$$

for the transverse mode and

$$\delta B^\parallel = n(x) \delta B^T, \quad (8.15)$$

for the longitudinal mode (n is a unknown function of $x \equiv -k\eta$) [97, 98]. Let us see why. As previously stated, for $f \simeq a^\alpha$ and with $\alpha = 0, 1, -2$, it is possible to verify that the (physical) transverse mode behaves exactly like a light scalar field in a de Sitter background (see also Appendix 15.4). Considering the solution (8.14) then takes into account at least these special cases. As to the longitudinal mode, a parametrization was adopted as in (8.15) in order to keep the analysis more general and given that, because of the instability issues, introducing this degree of freedom into the theory requires special attention. We are going to keep the longitudinal mode ‘‘alive’’ in the calculations we

present, by considering a nonzero function $n(x)$, and focus on the simplest case of $f = 1$. This case is known to be affected by quantum instabilities in the longitudinal mode, anyway we choose $f = 1$ for the sake of simplicity in our presentation. The results can be easily generalized to gauge invariant models (please refer to Sec. 12 for a sample generalization of some of the calculations to massless $f \simeq a^{(1,-2)}$ models).

9. Correlation functions of ζ in the $SU(2)$ model

We are now ready to compute the power spectrum, bispectrum and trispectrum for the curvature fluctuations ζ generated during inflation

$$\langle \zeta_{\vec{k}_1} \zeta_{\vec{k}_2} \rangle = (2\pi)^3 \delta^{(3)}(\vec{k}_1 + \vec{k}_2) P_\zeta(\vec{k}), \quad (9.1)$$

$$\langle \zeta_{\vec{k}_1} \zeta_{\vec{k}_2} \zeta_{\vec{k}_3} \rangle = (2\pi)^3 \delta^{(3)}(\vec{k}_1 + \vec{k}_2 + \vec{k}_3) B_\zeta(\vec{k}_1, \vec{k}_2, \vec{k}_3) \quad (9.2)$$

$$\langle \zeta_{\vec{k}_1} \zeta_{\vec{k}_2} \zeta_{\vec{k}_3} \zeta_{\vec{k}_4} \rangle = (2\pi)^3 \delta^{(3)}(\vec{k}_1 + \vec{k}_2 + \vec{k}_3 + \vec{k}_4) T_\zeta(\vec{k}_1, \vec{k}_2, \vec{k}_3, \vec{k}_4). \quad (9.3)$$

Notice that, on the right-hand side of (9.1) through (9.3), we indicated a dependence from the direction of the wavevectors; in models of inflation where isotropy is preserved, the power spectrum and the bispectrum only depend on the moduli of the wave vectors. This will not be the case for the $SU(2)$ model.

The δN formula (3.2) will be applied to our inflaton+ $SU(2)$ vector model

$$\begin{aligned} \zeta(\vec{x}, t) = & N_\phi \delta\phi + N_a^\mu \delta A_\mu^a + \frac{1}{2} N_{\phi\phi} (\delta\phi)^2 + \frac{1}{2} N_{ab}^{\mu\nu} \delta A_\mu^a \delta A_\nu^b + N_{\phi a}^\mu \delta\phi \delta A_\mu^a \\ & + \frac{1}{3!} N_{\phi\phi\phi} (\delta\phi)^3 + \frac{1}{3!} N_{abc}^{\mu\nu\lambda} \delta A_\mu^a \delta A_\nu^b \delta A_\lambda^c + \frac{1}{2} N_{\phi\phi a}^\mu (\delta\phi)^2 \delta A_\mu^a + \frac{1}{2} N_{\phi ab}^{\mu\nu} \delta\phi \delta A_\mu^a \delta A_\nu^b \\ & + \frac{1}{3!} N_{\phi\phi\phi\phi} (\delta\phi)^4 + \frac{1}{3!} N_{abcd}^{\mu\nu\lambda\eta} \delta A_\mu^a \delta A_\nu^b \delta A_\lambda^c \delta A_\eta^d + \dots, \end{aligned} \quad (9.4)$$

where now

$$N_\phi \equiv \left(\frac{\partial N}{\partial \phi} \right)_{t^*}, \quad N_a^\mu \equiv \left(\frac{\partial N}{\partial A_\mu^a} \right)_{t^*}, \quad N_{\phi a}^\mu \equiv \left(\frac{\partial^2 N}{\partial \phi \partial A_\mu^a} \right)_{t^*} \quad (9.5)$$

and so on for higher order derivatives.

Our plan is to show the derivation the correlation functions of ζ from the ones of $\delta\phi$ and δA_i^a , after a replacement of the δN expansion (9.4) in Eqs. (9.1) through (9.3).

The correlation functions can be evaluated using the Schwinger-Keldysh formula (2.1), that we recall here for convenience

$$\langle \Omega | \Theta(t) | \Omega \rangle = \left\langle 0 \left[\bar{T} \left(e^{i \int_0^t H_I(t') dt'} \right) \right] \Theta_I(t) \left[T \left(e^{-i \int_0^t H_I(t') dt'} \right) \right] \right| 0 \rangle. \quad (9.6)$$

When calculating the spectra of ζ , the perturbative expansions in Eq. (9.4) and (9.6) will be carried out to only include tree-level contributions, neglecting higher order ‘‘loop’’ terms, either classical, i.e. from the δN series, or of quantum origin, i.e. from the Schwinger-Keldysh series. Assuming that the $SU(2)$ coupling g_c is ‘‘small’’ and that we are dealing with ‘‘small’’ fluctuations in the fields and given the fact that a slow-roll regime is being assumed, it turns out that it is indeed safe for the two expansions to be truncated at tree-level.

The correlation functions of ζ will then result as the sum of scalar, vector and (scalar and vector) mixed contributions. As to the vector part, this will be made up of terms that are merely generated by the δN expansion, i.e. they only include the zeroth order of the in-in formula (we call these terms ‘‘Abelian’’, being them retrievable in the $U(1)$

case), and by (“non-Abelian”) terms arising from the Schwinger-Keldysh operator expansion beyond zeroth order, i.e. from the gauge fields self-interactions.

Let us now discuss the level of generality of the results we will present in the next sections, w.r.t. the choice of a specific Lagrangian.

The expression for the Abelian contributions provided in Secs. 9.1 and 9.2 apply to any $SU(2)$ model of gauge interactions with no direct coupling between scalar and vector fields (extra terms would be otherwise needed in Eqs. (9.18) and (9.19)). The next stage in the Abelian contributions computation would be to explicitate the derivatives of the e-foldings number and the wavefunctions of the fields: they both depend on the equations of motion of the system, therefore the fixing of a specific model is required at this point.

As to the non-Abelian contributions, the results in Eqs. (9.33) and (9.34) are completely general except for assuming, again, that no direct vector-scalar field coupling exists. The structure of Eqs. (9.49) and (9.52) is instead due to the choice of a non-Abelian gauge group. The expressions of the anisotropy coefficients I_n and L_n in Eqs. (9.49) and (9.52) depend on the specific non-Abelian gauge group (for $SU(2)$ one of the I_n is given in Eq. (9.51)). Finally, the specific expressions of the isotropic functions F_n (a sample of which is shown in Eq. (9.50)) and G_n were derived considering the Lagrangian (8.11) with $f = 1$ and the eigenfunctions for the vector bosons provided in Eqs. (8.14) and (8.15).

9.1. The power spectrum

The power spectrum of ζ can be straightforwardly derived at tree-level, using the δN expansion (9.4), from the inflaton and the vector fields power spectra

$$P_\zeta(\vec{k}) = P^{iso}(k) \left[1 + g^{ab} (\hat{k} \cdot \hat{N}_a) (\hat{k} \cdot \hat{N}_b) + is^{ab} \hat{k} \cdot (\hat{N}_a \times \hat{N}_b) \right]. \quad (9.7)$$

The isotropic part of the previous expression has been factorized in

$$P^{iso}(k) \equiv N_\phi^2 P_\phi(k) + (\vec{N}_c \cdot \vec{N}_d) P_+^{cd}, \quad (9.8)$$

where we have defined the following combinations

$$P_\pm^{ab} \equiv (1/2)(P_R^{ab} \pm P_L^{ab}), \quad (9.9)$$

from the power spectra for the right, left and longitudinal polarization modes

$$P_R^{ab} \equiv \delta_{ab} \delta A_R^a(k, t^*) \delta A_R^{b*}(k, t^*), \quad (9.10)$$

$$P_L^{ab} \equiv \delta_{ab} \delta A_L^a(k, t^*) \delta A_L^{b*}(k, t^*), \quad (9.11)$$

$$P_{long}^{ab} \equiv \delta_{ab} \delta A_{long}^a(k, t^*) \delta A_{long}^{b*}(k, t^*). \quad (9.12)$$

The anisotropic parts are weighted by the coefficients

$$g^{ab} \equiv \frac{N^a N^b (P_{long}^{ab} - P_+^{ab})}{N_\phi^2 P_\phi + (\vec{N}_c \cdot \vec{N}_d) P_+^{cd}}, \quad (9.13)$$

$$s^{ab} \equiv \frac{N^a N^b P_-^{ab}}{N_\phi^2 P_\phi + (\vec{N}_c \cdot \vec{N}_d) P_+^{cd}}, \quad (9.14)$$

(where a sum is intended over indices c and d but not over a and b). Eq. (9.8) can also be written as

$$P^{iso}(k) = N_\phi^2 P_\phi \left[1 + \beta_{cd} \frac{P_+^{cd}}{P_\phi} \right], \quad (9.15)$$

after introducing the parameter

$$\beta_{cd} \equiv \frac{\vec{N}_c \cdot \vec{N}_d}{N_\phi^2}. \quad (9.16)$$

Notice that what when we say ‘‘isotropic’’, as far as the expression for the power spectrum is concerned, we simply mean ‘‘independent’’ of the direction of the wave vector. In this case instead, the vector bosons introduce three preferred spatial directions: the r.h.s. of Eq. (9.7) depends on their orientation w.r.t. the wave vector.

As expected, the coefficients g^{ab} and s^{ab} that weight the anisotropic part of the power spectrum are related to β_{cd} , i.e. to the parameters that quantify how much the expansion of the universe is affected by the vector bosons compared to the scalar field.

Assuming no parity violation in the model, we have $s^{ab} = 0$; the parameters g^{ab} and β_{ab} are instead unconstrained. In the $U(1)$ case and for parity conserving theories, Eq. (9.7) reduces to [92]

$$P_\zeta(\vec{k}) = P_\zeta^{iso}(k) \left[1 + g \left(\hat{k} \cdot \hat{n} \right) \right] \quad (9.17)$$

where \hat{n} indicates the preferred spatial direction; also one can check that in this simple case, if $P_+ \simeq P_\phi$ and $P_{long} = kP_+$ ($k \neq 1$), the relation $g = (k - 1)\beta/(1 + \beta)$ holds, where $\beta \equiv (N_A/N_\phi)^2$ (the anisotropy coefficient g is not to be confused with the $SU(2)$ coupling constant g_c). If it is safe to assume $|g| \ll 1$ (see discussion following Eq. (1.1) and references [70, 71]), a similar upper bound can also be placed on β .

In the case where more than one special directions exists, as in the $SU(2)$ model, no such analysis on the anisotropy data has been so far carried out, the g^a parameters cannot then be constrained, unless assuming that the three directions converge into a single one; in that case a constraint could be placed on the sum $|g| \equiv |\sum_a g^a|$, where $a = 1, 2, 3$ and $P_\zeta(\vec{k}) = P_\zeta^{iso}(k) \left[1 + g^a \left(\hat{k} \cdot \hat{n}_a \right) \right]$.

In the next sections we will present the results for the tree-level contributions to the bispectrum and to the trispectrum of ζ .

These can be classified in two categories, that we indicate as ‘‘Abelian’’ and ‘‘non-Abelian’’. The former are intended as terms that merely arise from the δN expansion and are thus retrievable in the Abelian case; the latter are derived from the linear and quadratic expansions (in terms of the gauge bosons interaction Hamiltonian) of the Schwinger-Keldysh formula and are therefore peculiar to the non-Abelian case.

9.2. Bispectrum and trispectrum: Abelian contributions

By plugging the δN expansion (9.4) in Eqs. (9.2) and (9.3), we have

$$\begin{aligned}
B_\zeta(\vec{k}_1, \vec{k}_2, \vec{k}_3) \supset & \frac{1}{2} N_\phi^2 N_{\phi\phi} [P_\phi(k_1)P_\phi(k_2) + perms.] \\
& + \frac{1}{2} N_a^\mu N_b^\nu N_{cd}^{\rho\sigma} [\Pi_{\mu\rho}^{ac}(\vec{k}_1)\Pi_{\nu\sigma}^{bd}(\vec{k}_2) + perms.] \\
& + \frac{1}{2} N_\phi N_a^\mu N_b^\nu [P_\phi(k_1)\Pi_{\mu\nu}^{ab}(\vec{k}_2) + perms.] \\
& + N_\phi N_\phi^2 B_\phi(k_1, k_2, k_3), \tag{9.18}
\end{aligned}$$

for the bispectrum and

$$\begin{aligned}
T_\zeta(\vec{k}_1, \vec{k}_2, \vec{k}_3, \vec{k}_4) \supset & N_\phi^4 T_\phi(\vec{k}_1, \vec{k}_2, \vec{k}_3, \vec{k}_4) \\
& + N_\phi^3 N_{\phi\phi} [P_\phi(k_1)B_\phi(|\vec{k}_1 + \vec{k}_2|, k_3, k_4) + perms.] \\
& + N_\phi^2 N_a^\mu N_b^\nu [P_{\mu\nu}^{ab}(\vec{k}_3)B_\phi(k_1, k_2, |\vec{k}_3 + \vec{k}_4|) + perms.] \\
& + N_\phi^2 N_{\phi\phi}^2 [P_\phi(k_1)P_\phi(k_2)P_\phi(|\vec{k}_1 + \vec{k}_3|) + perms.] \\
& + N_\phi^3 N_{\phi\phi\phi} [P_\phi(k_1)P_\phi(k_2)P_\phi(k_3) + perms.] \\
& + N_\phi^2 N_{\phi a}^\mu N_{\phi b}^\nu [P_{\mu\nu}^{ab}(\vec{k}_1 + \vec{k}_3)P_\phi(k_1)P_\phi(k_2) + perms.] \\
& + N_a^\mu N_b^\nu N_{\phi c}^\rho N_{\phi d}^\sigma [P_{\mu\rho}^{ac}(\vec{k}_1)P_{\nu\sigma}^{bd}(\vec{k}_2)P_\phi(|\vec{k}_1 + \vec{k}_3|) + perms.] \\
& + N_\phi^2 N_a^\mu N_b^\nu [P_\phi(k_1)P_\phi(k_2)P_{\mu\nu}^{ab}(\vec{k}_3) + perms.] \\
& + N_\phi N_a^\mu N_b^\nu N_{\phi cd}^{\rho\sigma} [P_{\mu\rho}^{ac}(\vec{k}_1)P_{\nu\sigma}^{bd}(\vec{k}_2)P_\phi(k_3) + perms.] \\
& + N_{\phi\phi} N_\phi N_{\phi a}^\mu N_b^\nu [P_\phi(k_2)P_\phi(|\vec{k}_1 + \vec{k}_2|)P_{\mu\nu}^{ab}(\vec{k}_4) + perms.] \\
& + N_{ab}^{\mu\nu} N_c^\rho N_{\phi d}^\sigma N_\phi [P_{ac}^{\mu\rho}(\vec{k}_2)P_{bd}^{\nu\sigma}(\vec{k}_1 + \vec{k}_2)P_\phi(k_4) + perms.] \\
& + N_a^\mu N_b^\nu N_{cd}^{\rho\sigma} N_{ef}^{\delta\eta} [P_{\mu\rho}^{ac}(\vec{k}_1)P_{\nu\delta}^{be}(\vec{k}_2)P_{\sigma\eta}^{df}(\vec{k}_1 + \vec{k}_3) + perms.] \\
& + N_a^\mu N_b^\nu N_c^\rho N_{def}^{\sigma\delta\eta} [P_{\mu\sigma}^{ad}(\vec{k}_1)P_{\nu\delta}^{be}(\vec{k}_2)P_{\rho\eta}^{cf}(\vec{k}_3) + perms.], \tag{9.19}
\end{aligned}$$

for the trispectrum.

Before we proceed with explicitating these quantities and for the rest of the thesis, the N_0^a coefficients will be set to zero. It is in fact possible to verify that the temporal mode $B_0^a = 0$ is a solution to the equations of motion for the vector bosons, after slightly restricting the parameter space of the theory (see Appendix 15.4). The adoption of this kind of solutions, which is related to the assumption of a slow-roll regime for the vector fields, implies that the derivatives of N w.r.t. the temporal mode can be set to zero.

Let us now provide some definition for the quantities introduced in (9.18)-(9.19): we are going to switch from the greek indices μ, ν, \dots to the latin ones, generally used for labelling the three spatial directions, in order to stress that all of the vector quantities will be from now on three-dimensional

$$\Pi_{ij}^{ab}(\vec{k}) \equiv T_{ij}^{even}(\vec{k})P_+^{ab} + iT_{ij}^{odd}(\vec{k})P_-^{ab} + T_{ij}^{long}(\vec{k})P_{long}^{ab}, \tag{9.20}$$

where

$$T_{ij}^{even}(\vec{k}) \equiv e_i^L(\hat{k})e_j^{*L}(\hat{k}) + e_i^R(\hat{k})e_j^{*R}(\hat{k}), \tag{9.21}$$

$$T_{ij}^{odd}(\vec{k}) \equiv i \left[e_i^L(\hat{k}) e_j^{*L}(\hat{k}) - e_i^R(\hat{k}) e_j^{*R}(\hat{k}) \right], \quad (9.22)$$

$$T_{ij}^{long}(\vec{k}) \equiv e_i^l(\hat{k}) e_j^{*l}(\hat{k}). \quad (9.23)$$

The polarization vectors are $e^L(\hat{k}) \equiv \frac{1}{\sqrt{2}}(\cos\theta \cos\phi - i \sin\phi, \cos\theta \sin\phi + i \cos\phi, -\sin\theta)$, $e^R(\hat{k}) = e^{*L}(\hat{k})$ and $e^l(\hat{k}) = \hat{k} = (\sin\theta \cos\phi, \sin\theta \sin\phi, \cos\theta)$, from which we have

$$T_{ij}^{even}(\vec{k}) = \delta_{ij} - \hat{k}_i \hat{k}_j, \quad (9.24)$$

$$T_{ij}^{odd}(\vec{k}) = \epsilon_{ijk} \hat{k}_k, \quad (9.25)$$

$$T_{ij}^{long}(\vec{k}) = \hat{k}_i \hat{k}_j. \quad (9.26)$$

The purely scalar terms in Eqs. (9.18)-(9.19) are already known from the literature ‡.

As to the mixed (scalar-vector) terms, they can be ignored if one considers a Lagrangian where there is no direct coupling between the inflaton and the gauge bosons but the latter condition is not sufficient for concluding that the mixed derivatives are null. As an example, it is useful to refer to [148] which, among other things, includes an analytic study for the case of a set of slowly rolling fields with a separable quadratic potential. The number of e-folding is written as a sum of integrals over the different fields, to be evaluated between their values at an initial (generally set at around horizon crossing) and a final times. For each field, the value at the final time depends on the total field configuration at the initial time, so the mixed derivative of N can in principle be non-zero. Anyway, if the final time approaches the end of inflation, it is reasonable to assume that, by then, the fields have stabilized to their equilibrium value and no longer carry the memory of their evolution. If this happens, the sum of integrals which defines N becomes independent of the final field configuration and its mixed derivatives can therefore be shown to be zero. It turns out that we are allowed the same kind of analytic study, if we work with the Lagrangian in Eq. (8.11) and introduce some slow-roll assumptions for the vector fields (see Appendix 15.4 for a discussion about these assumptions and Appendix 15.5 for the actual calculation of N and its derivatives).

Let us then look at the (purely) vector part. Its anisotropy features can be stressed by rewriting them as follows

$$B_\zeta(\vec{k}_1, \vec{k}_2, \vec{k}_3) \supset \frac{1}{2} N_a^i N_b^j N_{cd}^{kl} \Pi_{ik}^{ac}(\vec{k}_1) \Pi_{jl}^{bd}(\vec{k}_2) = M_k^c N_{cd}^{kl} M_l^d \quad (9.27)$$

$$\begin{aligned} T_\zeta(\vec{k}_1, \vec{k}_2, \vec{k}_3, \vec{k}_4) &\supset N_a^\mu N_b^\nu N_{cd}^{\rho\sigma} N_{ef}^{\delta\eta} P_{\mu\rho}^{ac}(\vec{k}_1) P_{\nu\delta}^{be}(\vec{k}_2) P_{\sigma\eta}^{df}(\vec{k}_1 + \vec{k}_3) \\ &+ N_a^\mu N_b^\nu N_c^\rho N_{def}^{\sigma\delta\eta} P_{\mu\sigma}^{ad}(\vec{k}_1) P_{\nu\delta}^{be}(\vec{k}_2) P_{\rho\eta}^{cf}(\vec{k}_3) \\ &= M_i^c L_{ce}^{ij} M_j^e + M_i^f M_j^e M_k^d N_{fed}^{ijk}, \end{aligned} \quad (9.28)$$

where

$$M_k^c(\vec{k}) \equiv N_a^i P_{ik}^{ac}(\vec{k}) = P_+^{ac}(k) \left[\delta_{ik} N_a^i + p^{ac}(k) \hat{k}_k (\hat{k} \cdot \vec{N}_a) + i q^{ac}(k) (\hat{k} \times \vec{N}_a)_k \right] \quad (9.29)$$

‡ In single-field slow-roll inflation $P_\phi = H_*^2/2k^3$, where H_* is the Hubble rate evaluated at horizon exit; the bispectrum and the trispectrum of the scalar field (B_ϕ and T_ϕ) can be found in [12, 13, 147, 152, 39]. For the bispectrum see also Eq. (3.10)

$$\begin{aligned}
L_{ce}^{jl}(\vec{k}) &\equiv N_{cd}^{ji} P_{ik}^{df}(\vec{k}) N_{fe}^{kl} \\
&= P_+^{df}(\vec{k}) [\vec{N}_{cd}^j \cdot \vec{N}_{ef}^l + p^{df}(k) (\hat{k} \cdot \vec{N}_{cd}^j) (\hat{k} \cdot \vec{N}_{ef}^l) + iq^{df}(k) \hat{k} \cdot \vec{N}_{cd}^j \times \vec{N}_{ef}^l]. \tag{9.30}
\end{aligned}$$

In the previous equations, we defined

$$p^{ac}(k) \equiv \frac{P_{long}^{ac} - P_+^{ac}}{P_+^{ac}}, \tag{9.31}$$

$$q^{ac}(k) \equiv \frac{P_-^{ac}}{P_+^{ac}}, \tag{9.32}$$

with $\vec{N}_a \equiv (N_a^1, N_a^2, N_a^3)$ and $\vec{N}_{cd}^j \equiv (N_{cd}^{j1}, N_{cd}^{j2}, N_{cd}^{j3})$.

Notice that, as for the power spectrum (9.7), also in Eqs. (9.27)-(9.28) the anisotropic parts of the expressions are weighted by coefficients that are proportional either to P_- or to $(P_{long} - P_+)$. When these two quantities are equal to zero, the (Abelian) bispectrum and trispectrum are therefore isotropized. $P_- = 0$ in parity conserving theories, like the ones we have been describing. According to the parametrization (8.15) of the longitudinal mode, we have $P_{long} - P_+ = (|n(x)|^2 - 1)P_+$.

9.3. Bispectrum and trispectrum: Non-Abelian contributions

We list the non-Abelian terms for the bispectrum

$$B_\zeta(\vec{k}_1, \vec{k}_2, \vec{k}_3) \supset N_a^i N_b^j N_c^k B_{ijk}^{abc}(\vec{k}_1, \vec{k}_2, \vec{k}_3) \tag{9.33}$$

and for the trispectrum

$$\begin{aligned}
T_\zeta(\vec{k}_1, \vec{k}_2, \vec{k}_3, \vec{k}_4) &\supset N_a^i N_b^j N_c^k N_d^l T_{ijkl}^{abcd}(\vec{k}_1, \vec{k}_2, \vec{k}_3, \vec{k}_4) \\
&\quad + N_a^i N_b^j N_\phi N_{\phi c}^k \left[P_\phi(k_3) B_{ijk}^{abc}(\vec{k}_1, \vec{k}_2, \vec{k}_3 + \vec{k}_4) + perms. \right] \\
&\quad + N_a^i N_b^j N_c^k N_{de}^{lm} \left[P_{il}^{ad}(\vec{k}_1) B_{jkm}^{bce}(\vec{k}_1 + \vec{k}_2, \vec{k}_3, \vec{k}_4) + perms. \right]. \tag{9.34}
\end{aligned}$$

The computation of the vector bosons spectra

$$\langle \delta A_i^a \delta A_j^b \delta A_k^c \rangle = \delta^{(3)}(\vec{k}_1 + \vec{k}_2 + \vec{k}_3) B_{ijk}^{abc}, \tag{9.35}$$

$$\langle \delta A_i^a \delta A_j^b \delta A_k^c \delta A_l^d \rangle = \delta^{(3)}(\vec{k}_1 + \vec{k}_2 + \vec{k}_3 + \vec{k}_4) T_{ijkl}^{abcd}, \tag{9.36}$$

will be reviewed in this section. This requires the expansion of the in-in formula up to second order in the interaction Hamiltonian

$$\begin{aligned}
\langle \Theta(\eta^*) \rangle &\supset i \langle T \left[\Theta \int_{-\infty}^{\eta^*} d\eta' \left(H_{int}^+(\eta') - H_{int}^-(\eta') \right) \right] \rangle \\
&\quad + \frac{(-i)^2}{2} \langle T \left[\Theta \int_{-\infty}^{\eta^*} d\eta' \left(H_{int}^+(\eta') - H_{int}^-(\eta') \right) \int_{-\infty}^{\eta^*} d\eta'' \left(H_{int}^+(\eta'') - H_{int}^-(\eta'') \right) \right] \rangle. \tag{9.37}
\end{aligned}$$

The interaction Hamiltonian needs to be expanded up to fourth order in the fields fluctuations, i.e. $H_{int} = H_{int}^{(3)} + H_{int}^{(4)}$, where

$$H_{int}^{(3)} = g_c \varepsilon^{abc} g^{ik} g^{jl} \left(\partial_i \delta B_j^a \right) \delta B_k^b \delta B_l^c + g_c^2 \varepsilon^{eab} \varepsilon^{ecd} g^{ik} g^{jl} B_i^a \delta B_j^b \delta B_k^c \delta B_l^d \tag{9.38}$$

$$H_{int}^{(4)} = g_c^2 \varepsilon^{eab} \varepsilon^{ecd} g^{ij} g^{kl} \delta B_i^a \delta B_k^b \delta B_j^c \delta B_l^d. \tag{9.39}$$

To tree-level, the relevant diagrams are pictured in Figs. 5 and 6. By looking at Eqs. (9.38) and (9.39), we can see that there is a bispectrum diagram that is lower



Figure 5. Diagrammatic representations of the tree-level contributions to the vector fields bispectrum.



Figure 6. Diagrammatic representations of the tree-level contributions to the vector fields trispectrum: vector-exchange (on the left) and contact-interaction (on the right) diagrams.

in terms of power of the $SU(2)$ coupling ($\sim g_c$) compared to the trispectrum ($\sim g_c^2$); as a matter of fact, for symmetry reasons that we are going to discuss later in this section, g_c^2 interaction terms are needed to provide a non-zero contributions to the bispectrum. The propagators for “plus” and “minus” fields are

$$\delta B_i^{a,+}(\eta') \widehat{\delta B_j^{b,+}}(\eta'') = \tilde{\Pi}_{ij}^{ab}(\eta', \eta'') \Theta(\eta' - \eta'') + \bar{\Pi}_{ij}^{ab}(\eta', \eta'') \Theta(\eta'' - \eta'), \quad (9.40)$$

$$\delta B_i^{a,+}(\eta') \widehat{\delta B_j^{b,-}}(\eta'') = \bar{\Pi}_{ij}^{ab}(\eta', \eta''), \quad (9.41)$$

$$\delta B_i^{a,-}(\eta') \widehat{\delta B_j^{b,+}}(\eta'') = \tilde{\Pi}_{ij}^{ab}(\eta', \eta''), \quad (9.42)$$

$$\delta B_i^{a,-}(\eta') \widehat{\delta B_j^{b,-}}(\eta'') = \bar{\Pi}_{ij}^{ab}(\eta', \eta'') \Theta(\eta' - \eta'') + \tilde{\Pi}_{ij}^{ab}(\eta', \eta'') \Theta(\eta'' - \eta'), \quad (9.43)$$

or

$$\tilde{\Pi}_{ij}^{ab}(\vec{k}) \equiv T_{ij}^{even}(\hat{k}) \tilde{P}_+^{ab} + iT_{ij}^{odd}(\hat{k}) \tilde{P}_{ij}^{ab} + T_{ij}^{long}(\hat{k}) \tilde{P}_{ij}^{ab} \quad (9.44)$$

$$\bar{\Pi}_{ij}^{ab}(\vec{k}) \equiv T_{ij}^{even}(\hat{k}) \bar{P}_+^{ab} + iT_{ij}^{odd}(\hat{k}) \bar{P}_{ij}^{ab} + T_{ij}^{long}(\hat{k}) \bar{P}_{ij}^{ab} \quad (9.45)$$

in Fourier space. In the previous equations we set $\tilde{P}_\pm^{ab} \equiv (1/2)(\tilde{P}_R^{ab} \pm \tilde{P}_L^{ab})$, $\tilde{P}_R^{ab} = \delta_{ab} \delta B_R^{ab}(k, \eta^*) \delta B_R^{*ab}(k, \eta)$ and $\bar{P}_\pm^{ab} = (\tilde{P}_\pm^{ab})^*$ (similar definitions apply for \tilde{P}_L^{ab} and \tilde{P}_{long}^{ab}).

We are now ready to show the computation of the following contributions to the bispectrum and trispectrum of ζ

$$\langle \zeta_{\vec{k}_1} \zeta_{\vec{k}_2} \zeta_{\vec{k}_3} \rangle \supset N_a^i N_b^j N_c^k \langle \delta A_i^a(\vec{k}_1) \delta A_j^b(\vec{k}_2) \delta A_k^c(\vec{k}_3) \rangle, \quad (9.46)$$

$$\langle \zeta_{\vec{k}_1} \zeta_{\vec{k}_2} \zeta_{\vec{k}_3} \zeta_{\vec{k}_4} \rangle \supset N_a^i N_b^j N_c^k N_d^l \langle \delta A_i^a(\vec{k}_1) \delta A_j^b(\vec{k}_2) \delta A_k^c(\vec{k}_3) \delta A_l^d(\vec{k}_4) \rangle. \quad (9.47)$$

Eq. (9.46) becomes

$$\begin{aligned} \langle \zeta_{\vec{k}_1}^- \zeta_{\vec{k}_2}^- \zeta_{\vec{k}_3}^- \rangle \supset & N_a^i N_b^j N_c^k \frac{\delta^{(3)}(\vec{k}_1 + \vec{k}_2 + \vec{k}_3)}{a^3(\eta^*)} \left[\int d\eta a^4(\eta) \tilde{\Pi}_{im}(\vec{k}_1) \tilde{\Pi}_j^l(\vec{k}_2) \tilde{\Pi}_k^m(\vec{k}_3) \right. \\ & \left. \times \left(g_c \varepsilon^{abc} k_{1l} + g_c^2 \varepsilon^{eda} \varepsilon^{ebc} B_l^d \right) \right] + \text{perms.} + c.c. \end{aligned} \quad (9.48)$$

Even before performing the time integration, one realizes that, because of the antisymmetric properties of the Levi-Civita tensor, the $\sim g_c$ contribution on the r.h.s. of Eq. (9.48) is equal to zero once the sum over all the possible permutations has been performed. The vector bosons bispectrum is therefore proportional to g_c^2 . The final result from (9.48) has the following form

$$\langle \zeta_{\vec{k}_1}^- \zeta_{\vec{k}_2}^- \zeta_{\vec{k}_3}^- \rangle \supset (2\pi)^3 \delta^{(3)}(\vec{k}_1 + \vec{k}_2 + \vec{k}_3) g_c^2 H_*^2 \sum_n F_n(k_i, \eta^*) I_n(\hat{k}_i \cdot \hat{k}_j, \vec{A}_i \cdot \vec{A}_j, \hat{k}_i \cdot \vec{A}_j) \quad (9.49)$$

where the F_n 's are isotropic functions of time and of the moduli of the wave vectors ($i = 1, 2, 3$) and the I_n 's are anisotropic coefficients. The sum in the previous equation is taken over all possible combinations of products of three polarization indices, i.e. $n \in (EEE, EEl, ELE, \dots, lll)$, where E stands for ‘‘even’’, l for ‘‘longitudinal’’. The complete expressions for the terms appearing in the sum are quite lengthy and we list them in Appendix 15.6. As an example, we report here one of these terms

$$F_{lll} = -n^6(x^*) \frac{1}{24k^6 k_1^2 k_2^2 k_3^2 x^{*2}} [A_{EEE} + (B_{EEE} \cos x^* + C_{EEE} \sin x^*) E_i x^*] \quad (9.50)$$

$$\begin{aligned} I_{lll} = & \varepsilon^{aa'b'} \varepsilon^{ac'e} \left[\left((\hat{k}_1 \cdot \vec{N}^{a'}) (\hat{k}_3 \cdot \vec{N}^{b'}) (\hat{k}_2 \cdot \vec{N}^{c'}) (\hat{k}_1 \cdot \hat{k}_2) (\hat{k}_3 \cdot \hat{A}^e) \right. \right. \\ & \left. \left. - (\hat{k}_3 \cdot \vec{N}^{a'}) (\hat{k}_2 \cdot \vec{N}^{b'}) (\hat{k}_1 \cdot \vec{N}^{c'}) (\hat{k}_1 \cdot \hat{k}_2) (\hat{k}_3 \cdot \hat{A}^e) \right) + (1 \leftrightarrow 3) + (2 \leftrightarrow 3) \right] \end{aligned} \quad (9.51)$$

where A_{EEE} , B_{EEE} and C_{EEE} are functions of x^* and of the momenta $k_i \equiv |\vec{k}_i|$ (they are all reported in Appendix 15.6), E_i is the exponential-integral function. As we will discuss in more details in Sec. 11.2, one of the more interesting features of these models is that the bispectrum and the trispectrum turn out to have an amplitude that is modulated by the preferred directions that break statistical isotropy.

Let us now move to the trispectrum. Again, we count two different kinds of contributions, the first from $\sim g_c$ and the second from $\sim g_c^2$ interaction terms, respectively in $H_{int}^{(3)}$ and $H_{int}^{(4)}$. The former produce vector-exchange diagrams, the latter are represented by contact-interaction diagrams (see Fig. 6). Their analytic expressions are different, but they both have a structure similar to (9.49)

$$\begin{aligned} \langle \zeta_{\vec{k}_1}^- \zeta_{\vec{k}_2}^- \zeta_{\vec{k}_3}^- \zeta_{\vec{k}_4}^- \rangle \supset & (2\pi)^3 \delta^{(3)}(\vec{k}_1 + \vec{k}_2 + \vec{k}_3 + \vec{k}_4) g_c^2 H_*^2 \\ & \times \sum_n G_n(k_i, k_{\hat{1}2}, k_{\hat{1}4}, \eta^*) L_n(\hat{k}_i \cdot \hat{k}_j, \vec{A}_i \cdot \vec{A}_j, \hat{k}_i \cdot \vec{A}_j) \end{aligned} \quad (9.52)$$

where we define $k_{\hat{1}2} \equiv |\vec{k}_1 + \vec{k}_2|$ and $k_{\hat{1}4} \equiv |\vec{k}_1 + \vec{k}_4|$. We will present the details of the computation of the gauge fields trispectrum and the explicit expression of the functions appearing in (9.52) in the next section.

9.3.1. Trispectrum from vector bosons: exchange diagram

Let us begin with the two vertex diagram. Using the language of Eq. (9.37), it can be put in the form

$$\langle \Theta(\eta^*) \rangle \supset \frac{(-i)^2}{2} \langle T \left[\Theta \int_{-\infty}^{\eta^*} d\eta' \left(H^+(\eta') - H^-(\eta') \right) \int_{-\infty}^{\eta^*} d\eta'' \left(H^+(\eta'') - H^-(\eta'') \right) \right] \rangle \quad (9.53)$$

where now $H \equiv H_{int}^{(3)}$, $\Theta \equiv \delta A_\mu^a \delta A_\nu^b \delta A_\rho^c \delta A_\sigma^d$ and the inclusion symbol as usual points out that what stands on the right-hand side is only one of the contributions to $\langle \Theta(\eta^*) \rangle$.

Eq. (9.53) can be rewritten as follows

$$\langle \Theta(\eta^*) \rangle \supset \frac{(-i)^2}{2} \langle T [\Theta (A + B + C + D)] \rangle \quad (9.54)$$

where

$$\begin{aligned} A &\equiv \int_{-\infty}^{\eta^*} d\eta' H^+(\eta') \int_{-\infty}^{\eta^*} d\eta'' H^+(\eta'') \\ B &\equiv \int_{-\infty}^{\eta^*} d\eta' H^-(\eta') \int_{-\infty}^{\eta^*} d\eta'' H^-(\eta'') \\ C &\equiv - \int_{-\infty}^{\eta^*} d\eta' H^+(\eta') \int_{-\infty}^{\eta^*} d\eta'' H^-(\eta'') \\ D &\equiv - \int_{-\infty}^{\eta^*} d\eta' H^-(\eta') \int_{-\infty}^{\eta^*} d\eta'' H^+(\eta'') \end{aligned}$$

For each one of the integrals listed above, due to the presence of both the fields and their spatial derivatives in $H_{int}^{(3)}$, there are three different sets of contractions of the external with the vertex field-operators: for the first set, the field-operators with derivatives in the vertices are both contracted with external fields; for the second one, only one of the two field-operators with derivatives contracts with an external field (the other contracts with another internal field); for the third set, the field-operators with derivatives contract with each other.

A sample set of contractions of the first type is provided in the following equation

$$\begin{aligned} T_{ijkl}^{abcd} &\supset \frac{g_c^2}{2a^4(\eta^*)} \varepsilon^{a'b'c'} \varepsilon^{a''b''c''} k_m k_{m'} \\ &\times \int d\eta' a^A(\eta') \int d\eta'' a^A(\eta'') g^{mp} g^{m'p'} g^{nq} g^{n'q'} \tilde{\Pi}_{in}^{aa'} \tilde{\Pi}_{jp}^{bb'} \tilde{\Pi}_{kn'}^{ca''} \tilde{\Pi}_{lp'}^{db''} \tilde{\Pi}_{qq'}^{c'c''} \end{aligned} \quad (9.55)$$

where the first four $\tilde{\Pi}$ s correspond to contractions between external and internal fields whereas the last one indicates the contraction between the two remaining internal fields. The a^{-4} factor comes from expressing the external (physical) fields in terms of the comoving ones. As a reminder, we define $(2\pi)^3 \delta^{(3)}(\vec{k}_1 + \vec{k}_2 + \vec{k}_3 + \vec{k}_4) T_{ijkl}^{abcd} \equiv \langle \delta A_i^a \delta A_j^b \delta A_k^c \delta A_l^d \rangle$. The expression in (9.55) can be rewritten as follows

$$\begin{aligned} T_{ijkl}^{abcd} &\supset \frac{g_c^2}{2a^4(\eta^*)} \varepsilon^{a'b'c'} \varepsilon^{a''b''c''} k_m k_{m'} \delta^{aa'} \delta^{bb'} \delta^{ca''} \delta^{db''} \delta^{c'c''} \\ &\times \sum_{\alpha, \beta, \gamma, \delta, \sigma} \left(\int d\eta' \int d\eta'' \right)_{\alpha\beta\gamma\delta\sigma} T_{in}^\alpha T_{jm}^\beta T_{kn'}^\gamma T_{lp'}^\delta T_{nn'}^\sigma \end{aligned} \quad (9.56)$$

where the greek indices of the sum indicate either the transverse (E) or longitudinal (L) modes, the $\left(\int d\eta' \int d\eta''\right)$ stand for the integrals over the wave functions, the chosen time variable being $x \equiv -k\eta$ ($k \equiv \sum_{i=1,\dots,4} k_i$, $k_i \equiv |\vec{k}_i|$).

Let us define the coefficients $T_{ijkl}^{\alpha\beta\gamma\delta\sigma} \equiv k_m k_{m'} T_{in}^\alpha T_{jm}^\beta T_{kn'}^\gamma T_{lm'}^\delta T_{nn'}^\sigma$. They should be calculated for each one of the three different sets of contractions and for each permutation within the specific set. This is a straightforward but rather lengthy and not particularly interesting calculation. A convenient way to proceed could be the following: we first compute the time integrals in order to find out which one among the combinations of longitudinal and transverse mode functions in the string $[\alpha, \beta, \gamma, \delta, \sigma]$ provides the highest amplitude for the trispectrum (in order to be able to perform this comparison we work, as it is usually done when trying to quantify the amplitude of a three or of a four-point function, in the so called ‘‘equilateral configuration’’, which for the trispectrum means taking $k_1 = k_2 = k_3 = k_4$); for the combination with the highest amplitude, we then calculate the coefficients $T_{ijkl}^{\alpha\beta\gamma\delta\sigma}$ for all the different sets of contractions and sum over all the permutations.

Let us now perform our calculations. The wavefunctions we are going to adopt were introduced in Sec. 8.5 (see Eqs. (8.14) and (8.15)). It is possible to verify that $B = A^*$ and $D = C^*$ and that integrals of type A are consistently smaller in amplitude than integrals of type C . We therefore report the combined contribution $C + D = 2\text{Re}[C]$ for one of the permutations

$$\begin{aligned} \left(\int d\eta' \int d\eta''\right)_{EEEE} &= \frac{1}{8k_1^3 k_2^3 k_3^3 k_4^3 k_{12}^3 (k_{12} + k_1 + k_2)(k_{12} + k_3 + k_4)x^{*8}} \\ &\times [(M - 2E)[(N - 2F)(AB + CD) + (2H + L)(CB - AD)] \\ &+ (2G + P)[(N - 2F)(AD - CB) + (2H + L)(AB + CD)] \end{aligned} \quad (9.57)$$

$$\left(\int d\eta' \int d\eta''\right)_{EEEEl} = n^2(x^*) \left(\int d\eta' \int d\eta''\right)_{EEEE} \quad (9.58)$$

$$\left(\int d\eta' \int d\eta''\right)_{EEEl} = n^4(x^*) \left(\int d\eta' \int d\eta''\right)_{EEEE} \quad (9.59)$$

$$\left(\int d\eta' \int d\eta''\right)_{EEll} = n^6(x^*) \left(\int d\eta' \int d\eta''\right)_{EEEE} \quad (9.60)$$

$$\left(\int d\eta' \int d\eta''\right)_{Eull} = n^8(x^*) \left(\int d\eta' \int d\eta''\right)_{EEEE} \quad (9.61)$$

$$\left(\int d\eta' \int d\eta''\right)_{ulll} = n^{10}(x^*) \left(\int d\eta' \int d\eta''\right)_{EEEE} \quad (9.62)$$

where $A, B, C, D, E, F, G, H, L, M, N$ and P are functions of x^* and of the momenta moduli to be provided in Appendix 15.7 (see Eqs. (15.7.1) through (15.7.9)). Obviously, the value of the integrals does not change when permuting its labels $\alpha\beta\gamma\delta\sigma$, apart from a different power of the coefficient $n(x^*)$. We need now to find out if there is one, among the integrals in Eqs. (9.57) through (9.62), that has the largest amplitude, i.e. understand if something can be said about the order of magnitude of $n(x^*)$. We could try to extrapolate some information about $n(x^*)$ from what happens at very late times.

In the models discussed in Ref. [135, 92] it turns out that the longitudinal mode is $\delta B^{\parallel} = \sqrt{2}\delta B^T$.[§] If this is the correct asymptotic behaviour and we find it reasonable to extrapolate back until the horizon crossing epoch, it is then correct to conclude that in this case the amplitude is the largest for the integral among the ones listed in Eqs. (9.57) through (9.62) containing the highest powers of n , i.e. for $(\int d\eta' \int d\eta'')$ _{uuu}. The coefficients we intend to calculate are then of the kind T_{ijkl}^{llll} only. We list them below for the three different sets of contractions we introduced above and for one particular permutation (see again Appendix 15.7 for more details)

$$T_{ijkl}^{llll(1)} = k_1 k_3 k_{1234} (\hat{k}_1 \cdot \hat{k}_{\hat{1}2}) (\hat{k}_1 \cdot \hat{k}_2) (\hat{k}_3 \cdot \hat{k}_4) (\hat{k}_3 \cdot \hat{k}_{\hat{1}2}), \quad (9.63)$$

$$T_{ijkl}^{llll(2)} = k_3 k_{\hat{1}2} k_{1234} (\hat{k}_1 \cdot \hat{k}_{\hat{1}2}) (\hat{k}_2 \cdot \hat{k}_{\hat{1}2}) (\hat{k}_3 \cdot \hat{k}_4) (\hat{k}_3 \cdot \hat{k}_{\hat{1}2}), \quad (9.64)$$

$$T_{ijkl}^{llll(3)} = k_{\hat{1}2} k_{\hat{1}2} k_{1234} (\hat{k}_1 \cdot \hat{k}_{\hat{1}2}) (\hat{k}_2 \cdot \hat{k}_{\hat{1}2}) (\hat{k}_3 \cdot \hat{k}_{\hat{1}2}) (\hat{k}_4 \cdot \hat{k}_{\hat{1}2}). \quad (9.65)$$

We adopted the following notation: $k_s \equiv |\vec{k}_s|$, $\hat{k}_s \equiv \vec{k}_s/k_s$, the index s running over the four external momenta; $k_{ss's''s'''} \equiv \hat{k}_{s_i} \hat{k}_{s'_j} \hat{k}_{s''_k} \hat{k}_{s'''_l}$, with $s, s', s'', s''' = 1, 2, 3, 4$ and with the indices i, j, k, l indicating the spatial components of the vectors; $\vec{k}_{\hat{s}s'} \equiv \vec{k}_s + \vec{k}_{s'}$, $k_{\hat{s}s'} \equiv |\vec{k}_s + \vec{k}_{s'}|$ and so $\hat{k}_{\hat{s}s'} \equiv \vec{k}_{\hat{s}s'}/k_{\hat{s}s'}$.

It is possible to prove that, once the Levi-Civita coefficients and the sum over the permutations are taken into account, only the first set of contractions is left. The final result after these cancellations can be written in the following form

$$\begin{aligned} \langle \delta A_i^a \delta A_j^b \delta A_k^c \delta A_l^d \rangle_* \supset (2\pi)^3 \delta^{(3)}(\vec{k}_1 + \vec{k}_2 + \vec{k}_3 + \vec{k}_4) g_c^2 \left(\frac{H_* x^*}{k} \right)^4 \varepsilon^{abc''} \varepsilon^{cdc''} \left[I \times k_{1234} \times \left(\sum_{i=1}^4 t_i \right) \right. \\ \left. + II \times k_{1324} \times \left(\sum_{i=5}^8 t_i \right) + III \times k_{1432} \times \left(\sum_{i=9}^{12} t_i \right) \right]. \quad (9.66) \end{aligned}$$

All the possible permutations have been included in the previous equation and, as a reminder, the indices i, j, k, l are hidden in $k_{ss's''s'''}$ on the right-hand side. We define

$$\begin{aligned} I \equiv n^{10} \times \left(\frac{1}{8k_1^3 k_2^3 k_3^3 k_4^3 k_{\hat{1}2}^3 (k_{\hat{1}2} + k_1 + k_2)(k_{\hat{1}2} + k_3 + k_4) x^{*8}} \right) \\ \times [(M - 2E)[(N - 2F)(AB + CD) + (2H + L)(CB - AD)] \\ + (2G + P)[(N - 2F)(AD - CB) + (2H + L)(AB + CD)] \quad (9.67) \end{aligned}$$

(from Eqs. (9.57)-(9.62)). The function II is defined from I by exchanging k_2 with k_3 and $k_{\hat{1}2}$ with $k_{\hat{1}3}$; similarly, III is defined from I by exchanging k_2 with k_4 and $k_{\hat{1}2}$ with $k_{\hat{1}4}$, so they are all functions of the horizon crossing time $x^* \equiv -k\eta^*$ and of the moduli of the external momenta and of their sums. This amounts to seven independent variables, x^* , k_1 , k_2 , k_3 , k_4 , $k_{\hat{1}2}$ and $k_{\hat{1}4}$. The coefficients t_i ($i = 1, \dots, 12$) come from $T_{ijkl}^{\alpha\beta\gamma\delta\sigma}$ and

[§] As another example, in models with varying kinetic function and mass [95, 96], we have verified that $n(x) \gg 1$ at late times and for a vector field that is light until the end of inflation.

so they are also functions of the momenta moduli (see Eqs. (15.7.10) through (15.7.21) for their expressions). Finally, the anisotropic part of Eq. (9.66) is represented by the $k_{ss's''s'''}$ terms, which, in the final expression for the curvature perturbation trispectrum, have their spatial indices contracted with the derivatives N_i^a of the number of e-foldings w.r.t. the vector fields as follows

$$\begin{aligned} \langle \zeta_{\vec{k}_1} \zeta_{\vec{k}_2} \zeta_{\vec{k}_3} \zeta_{\vec{k}_4} \rangle \supset N_i^a N_j^b N_k^c N_l^d \langle \delta A_i^a \delta A_j^b \delta A_k^c \delta A_l^d \rangle_* \supset (2\pi)^3 \delta^{(3)}(\vec{k}_1 + \vec{k}_2 + \vec{k}_3 + \vec{k}_4) g_c^2 \left(\frac{H_* x^*}{k} \right)^4 \\ \times I \times \left(\sum_{i=1}^4 t_i \right) \times \Delta_I + perms. \end{aligned} \quad (9.68)$$

where the anisotropic term in the first permutation is

$$\Delta_I \equiv \varepsilon^{abc''} \varepsilon^{cdc''} N_i^a N_j^b N_k^c N_l^d k_{1234}. \quad (9.69)$$

It can be interesting to rewrite Δ_I in terms of all its variables

$$\Delta_I = \sum_{(a<b)a,b=1}^3 \left[(N^a)^2 (N^b)^2 \times \prod_{[i,j]=[1,2],[3,4]} \det(M_I^{i,j,a,b}) \right] \quad (9.70)$$

The M_I 's are 2×2 matrices whose entries are represented by the cosines of the angles between the wavevectors and the \vec{N}^a

$$M_I^{i,j,a,b} \equiv \begin{vmatrix} \cos \theta_{ia} & \cos \theta_{ja} \\ \cos \theta_{ib} & \cos \theta_{jb} \end{vmatrix}.$$

i.e. $\cos \theta_{ia} \equiv \hat{k}_i \cdot \hat{N}^a$ and so on.

The two permutations in Eq. (9.68) can be written in a similar fashion with anisotropic coefficients

$$\begin{aligned} \Delta_{II} \equiv \varepsilon^{abc''} \varepsilon^{cdc''} N_i^a N_j^b N_k^c N_l^d k_{1324} &= \sum_{(a<b)a,b=1}^3 \left[(N^a)^2 (N^b)^2 \times \prod_{[i,j]=[1,3],[2,4]} \det(M_{II}^{i,j,a,b}) \right], \\ \Delta_{III} \equiv \varepsilon^{abc''} \varepsilon^{cdc''} N_i^a N_j^b N_k^c N_l^d k_{1432} &= \sum_{(a<b)a,b=1}^3 \left[(N^a)^2 (N^b)^2 \times \prod_{[i,j]=[1,4],[3,2]} \det(M_{III}^{i,j,a,b}) \right]. \end{aligned}$$

The number of angular variables is equal to 12. These are to be added to the six scalar variables from the isotropic part of (9.68) ($k_1, k_2, k_3, k_4, k_{\hat{1}2}$ and $k_{\hat{1}4}$) and to three parameters represented by the lengths of the vectors \vec{N}^a in Δ . The anisotropy coefficients here become equal to zero in the event of an alignment of the gauge vectors along a unique direction.

9.3.2. Trispectrum from vector bosons: point-interaction diagram

Let us now move to the one-vertex diagrams

$$\langle \Theta(\eta^*) \rangle \supset i \langle T \left[\Theta \int_{-\infty}^{\eta^*} d\eta' \left(H^+(\eta') - H^-(\eta') \right) \right] \rangle, \quad (9.71)$$

where now $H \equiv H_{int}^{(4)}$ and again $\Theta \equiv \delta A_\mu^a \delta A_\nu^b \delta A_\rho^c \delta A_\sigma^d$. After working out the Wick contractions, this becomes

$$N_a^\mu N_b^\nu N_c^\rho N_d^\sigma T_{\mu\nu\rho\sigma}^{abcd}(\vec{k}_1, \vec{k}_2, \vec{k}_3, \vec{k}_4) \supset g_c^2 \left(\frac{H_* x^*}{k} \right)^4 \epsilon^{a'bc} \epsilon^{a'da} N_a^m N_b^n N_c^o N_d^p \\ \times \sum_{\alpha\beta\gamma\delta} \left(\int dx \right)_{\alpha\beta\gamma\delta} T_{mi}^\alpha T_{nj}^\beta T_{oi}^\gamma T_{pj}^\delta + \text{permutations.} \quad (9.72)$$

Let us list the coefficients $T_{mnop}^{\alpha\beta\gamma\delta} \equiv T_{mi}^\alpha T_{nj}^\beta T_{oi}^\gamma T_{pj}^\delta$ for one of the permutations

$$T_{mnop}^{EEEE} = \delta_{mo}\delta_{np} - \delta_{mo}\hat{k}_{p4}\hat{k}_{n4} - \delta_{mo}\hat{k}_{n2}\hat{k}_{p2} + \delta_{mo}\hat{k}_{n2}\hat{k}_{p4}\hat{k}_2 \cdot \hat{k}_4 - \delta_{np}\hat{k}_{o3}\hat{k}_{m3} \\ + k_{3434} + k_{3232} - \hat{k}_2 \cdot \hat{k}_4 (k_{3234} + k_{1214}) - \delta_{np}\hat{k}_{m1}\hat{k}_{o1} + k_{1414} + k_{1212} \\ + \delta_{np}\hat{k}_{m1}\hat{k}_{o3}\hat{k}_1 \cdot \hat{k}_3 - \hat{k}_1 \cdot \hat{k}_3 (k_{1232} + k_{1434}) + \hat{k}_1 \cdot \hat{k}_3 \hat{k}_2 \cdot \hat{k}_4, \quad (9.73)$$

$$T_{mnop}^{EEEEI} = \delta_{mo}\hat{k}_{p4}\hat{k}_{n4} - \delta_{mo}\hat{k}_{n2}\hat{k}_{p4}\hat{k}_2 \cdot \hat{k}_4 - k_{3434} + \hat{k}_2 \cdot \hat{k}_4 (k_{3234} + k_{1214}) - k_{1414} \\ + k_{1434}\hat{k}_1 \cdot \hat{k}_3 - k_{1234}\hat{k}_1 \cdot \hat{k}_3 \hat{k}_2 \cdot \hat{k}_4, \quad (9.74)$$

$$T_{mnop}^{EEIE} = \delta_{np}\hat{k}_{o3}\hat{k}_{m3} - \delta_{np}\hat{k}_{m1}\hat{k}_{o3}\hat{k}_1 \cdot \hat{k}_3 - k_{3434} + \hat{k}_1 \cdot \hat{k}_3 (k_{1434} + k_{1232}) - k_{3232} \\ + k_{3234}\hat{k}_2 \cdot \hat{k}_4 - k_{1234}\hat{k}_1 \cdot \hat{k}_3 \hat{k}_2 \cdot \hat{k}_4, \quad (9.75)$$

$$T_{mnop}^{EIEE} = \delta_{mo}\hat{k}_{n2}\hat{k}_{p2} - \delta_{mo}\hat{k}_{n2}\hat{k}_{p4}\hat{k}_2 \cdot \hat{k}_4 - k_{3232} + \hat{k}_2 \cdot \hat{k}_4 (k_{3234} + k_{1214}) - k_{1212} \\ + k_{1232}\hat{k}_1 \cdot \hat{k}_3 - k_{1234}\hat{k}_1 \cdot \hat{k}_3 \hat{k}_2 \cdot \hat{k}_4, \quad (9.76)$$

$$T_{mnop}^{IEEE} = \hat{k}_{m1}\hat{k}_{o1}\delta_{np} - \hat{k}_{m1}\hat{k}_{o3}k_{13}\delta_{np} - k_{1414} + \hat{k}_1 \cdot \hat{k}_3 (k_{1434} + k_{1232}) - k_{1212} \\ + k_{1214}\hat{k}_2 \cdot \hat{k}_4 - k_{1234}\hat{k}_1 \cdot \hat{k}_3 \hat{k}_2 \cdot \hat{k}_4, \quad (9.77)$$

$$T_{mnop}^{EELL} = k_{3434} - k_{3234}\hat{k}_2 \cdot \hat{k}_4 - k_{1434}\hat{k}_1 \cdot \hat{k}_3 + k_{1234}\hat{k}_1 \cdot \hat{k}_3 \hat{k}_2 \cdot \hat{k}_4, \quad (9.78)$$

$$T_{mnop}^{EIEL} = \delta_{mo}\hat{k}_{n2}\hat{k}_{p4}\hat{k}_2 \cdot \hat{k}_4 - \hat{k}_2 \cdot \hat{k}_4 (k_{3234} + k_{1214}) + k_{1234}\hat{k}_1 \cdot \hat{k}_3 \hat{k}_2 \cdot \hat{k}_4, \quad (9.79)$$

$$T_{mnop}^{ELLE} = k_{3232} - k_{3234}\hat{k}_2 \cdot \hat{k}_4 - k_{1232}\hat{k}_1 \cdot \hat{k}_3 + k_{1234}\hat{k}_1 \cdot \hat{k}_3 \hat{k}_2 \cdot \hat{k}_4, \quad (9.80)$$

$$T_{mnop}^{LLEE} = k_{1212} - k_{1214}\hat{k}_2 \cdot \hat{k}_4 - k_{1232}\hat{k}_1 \cdot \hat{k}_3 + k_{1234}\hat{k}_1 \cdot \hat{k}_3 \hat{k}_2 \cdot \hat{k}_4, \quad (9.81)$$

$$T_{mnop}^{LEEL} = k_{1414} - k_{1214}\hat{k}_2 \cdot \hat{k}_4 - k_{1434}\hat{k}_1 \cdot \hat{k}_3 + k_{1234}\hat{k}_1 \cdot \hat{k}_3 \hat{k}_2 \cdot \hat{k}_4, \quad (9.82)$$

$$T_{mnop}^{LEIE} = \delta_{np}\hat{k}_{m1}\hat{k}_{o3}\hat{k}_1 \cdot \hat{k}_3 - \hat{k}_1 \cdot \hat{k}_3 (k_{1434} + k_{1232}) + k_{1234}\hat{k}_1 \cdot \hat{k}_3 \hat{k}_2 \cdot \hat{k}_4, \quad (9.83)$$

$$T_{mnop}^{LLEE} = k_{1232}\hat{k}_1 \cdot \hat{k}_3 - k_{1234}\hat{k}_1 \cdot \hat{k}_3 \hat{k}_2 \cdot \hat{k}_4, \quad (9.84)$$

$$T_{mnop}^{LEEL} = k_{1214}\hat{k}_2 \cdot \hat{k}_4 - k_{1234}\hat{k}_1 \cdot \hat{k}_3 \hat{k}_2 \cdot \hat{k}_4, \quad (9.85)$$

$$T_{mnop}^{LELL} = k_{1434}\hat{k}_1 \cdot \hat{k}_3 - k_{1234}\hat{k}_1 \cdot \hat{k}_3 \hat{k}_2 \cdot \hat{k}_4, \quad (9.86)$$

$$T_{mnop}^{LEll} = k_{3234}\hat{k}_2 \cdot \hat{k}_4 - k_{1234}\hat{k}_1 \cdot \hat{k}_3 \hat{k}_2 \cdot \hat{k}_4, \quad (9.87)$$

$$T_{mnop}^{Llll} = k_{1234}\hat{k}_1 \cdot \hat{k}_3 \hat{k}_2 \cdot \hat{k}_4. \quad (9.88)$$

When evaluating the integrals $(\int dx)_{\alpha\beta\gamma\delta}$, we use again the wavefunctions previously introduced in Eqs. (8.14) and (8.15). The final result is

$$\left(\int dx \right)_{EEEE} = \frac{1}{24k^5 k_1^2 k_2^2 k_3^2 k_4^2 x^{*7}} [Q_{EEEE} + A_{EEEE} \text{cix}^* (B_{EEEE} \cos x^* + C_{EEEE} \sin x^*) \\ + D_{EEEE} \text{six}^* (E_{EEEE} \cos x^* + F_{EEEE} \sin x^*)], \quad (9.89)$$

$$\left(\int dx\right)_{EEEE} = n^2(x^*) \left(\int dx\right)_{EEEE}, \quad (9.90)$$

$$\left(\int dx\right)_{EEU} = n^4(x^*) \left(\int dx\right)_{EEEE}, \quad (9.91)$$

$$\left(\int dx\right)_{UE} = n^6(x^*) \left(\int dx\right)_{EEEE}, \quad (9.92)$$

$$\left(\int dx\right)_{UU} = n^8(x^*) \left(\int dx\right)_{EEEE}, \quad (9.93)$$

where Q_{EEEE} , A_{EEEE} , B_{EEEE} , C_{EEEE} , D_{EEEE} , E_{EEEE} and F_{EEEE} are functions of x^* and of the momenta $k_i \equiv |\vec{k}_i|$, ci and si stand respectively for the CosIntegral and the SinIntegral functions. The expressions of these functions can be found in Appendix 15.8. It is again important noticing that the anisotropy coefficients become zero if the gauge fields are all aligned.

Finally, summing up the coefficients in Eqs. (9.73) through (9.88), one realizes that if the longitudinal and the transverse mode evolve in the same way, the total contribution from the point-interaction diagram is isotropic

$$\begin{aligned} \langle \zeta_{\vec{k}_1} \zeta_{\vec{k}_2} \zeta_{\vec{k}_3} \zeta_{\vec{k}_4} \rangle &\supset (2\pi)^3 \delta^3(\vec{k}_1 + \vec{k}_2 + \vec{k}_3 + \vec{k}_4) g_c^2 \left(\frac{H_* x^*}{k}\right)^4 \epsilon^{a'bc} \epsilon^{a'da} (\vec{N}^c \cdot \vec{N}^a) (\vec{N}^b \cdot \vec{N}^d) \\ &\times \frac{1}{24k^5 k_1^2 k_2^2 k_3^2 k_4^2 x^{*7}} [Q_{EEEE} + A_{EEEE} ci x^* (B_{EEEE} \cos x^* + C_{EEEE} \sin x^*) \\ &\quad + D_{EEEE} si x^* (E_{EEEE} \cos x^* + F_{EEEE} \sin x^*)] + \text{permutations}. \quad (9.94) \end{aligned}$$

10. Amplitude of non-Gaussianity: f_{NL} and τ_{NL}

Our definitions for the non-Gaussianity amplitudes are

$$\frac{6}{5}f_{NL} = \frac{B_\zeta(\vec{k}_1, \vec{k}_2, \vec{k}_3)}{P^{iso}(k_1)P^{iso}(k_2) + perms.} \quad (10.1)$$

$$\tau_{NL} = \frac{2T_\zeta(\vec{k}_1, \vec{k}_2, \vec{k}_3, \vec{k}_4)}{P^{iso}(k_1)P^{iso}(k_2)P^{iso}(k_{14}) + 23 \text{ perms.}} \quad (10.2)$$

The choice of normalizing the bispectrum and the trispectrum by the isotropic part of the power spectrum, instead of using its complete expression P_ζ , is motivated by the fact that the latter would only introduce a correction to the previous equations proportional to the anisotropy parameter g , which is a small quantity.

The parameters f_{NL} and τ_{NL} receive contributions both from scalar (“ s ”) and from vector (“ v ”) fields

$$f_{NL} = f_{NL}^{(s)} + f_{NL}^{(v)}, \quad (10.3)$$

$$\tau_{NL} = \tau_{NL}^{(s)} + \tau_{NL}^{(v)}. \quad (10.4)$$

The latter can again be distinguished into Abelian (A) and non-Abelian (NA)

$$f_{NL}^{(v)} = f_{NL}^{(A)} + f_{NL}^{(NA)}, \quad (10.5)$$

$$\tau_{NL}^{(v)} = \tau_{NL}^{(A_1)} + \tau_{NL}^{(A_2)} + \tau_{NL}^{(NA_1)} + \tau_{NL}^{(NA_2)}. \quad (10.6)$$

The contribution $f_{NL}^{(A)}$ comes from Eq. (9.27), $f_{NL}^{(NA)}$ from (9.49), $\tau_{NL}^{(A_1)}$ and $\tau_{NL}^{(A_2)}$ from (9.28), finally $\tau_{NL}^{(NA_1)}$ from (9.52) and $\tau_{NL}^{(NA_2)}$ from the last line of (9.34).

In order to keep the vector contributions manageable and simple in their structure, all gauge and vector indices will be purposely neglected in this section and so the angular functions appearing in the anisotropy coefficients will be left out of the final amplitude results. This is acceptable considering that these functions will in general introduce numerical corrections of order one. Nevertheless, it is important to keep in mind that the amplitudes also depend on the angular parameters of the theory.

We will now focus on the dependence of f_{NL} and τ_{NL} from the non-angular parameters of the theory and quickly draw a comparison among the different contributions listed in Eqs. (10.3) through (10.6).

The expression of the number of e-foldings depends on the specific model and, in particular, on the mechanism of production of the fluctuations. Two possibilities have been described in Sec. 8. For “vector inflation” we have

$$N_a^i = \frac{A_i^a}{2m_P^2}, \quad N_{ab}^{ij} = \frac{\delta_{ab}\delta^{ij}}{2m_P^2} \quad (10.7)$$

Table 1. Order of magnitude of f_{NL} in different scenarios.

	f_{NL}^s	f_{NL}^A	f_{NL}^{NA}
general case	$\frac{1}{(1+\beta)^2} \frac{N_{\phi\phi}}{N_\phi^2}$	$\frac{\beta}{(1+\beta)^2} \frac{N_{AA}}{N_\phi^2}$	$\frac{\beta^2}{(1+\beta)^2} g_c^2 \left(\frac{m}{H}\right)^2$
v.inflation	$\frac{\epsilon}{\left(1+\left(\frac{A}{m_P} \sqrt{\epsilon}\right)^2\right)^2}$	$\frac{\epsilon^2}{\left(1+\left(\frac{A}{m_P} \sqrt{\epsilon}\right)^2\right)^2} \left(\frac{A}{m_P}\right)^2$	$\frac{\epsilon^2 g_c^2}{\left(1+\left(\frac{A}{m_P} \sqrt{\epsilon}\right)^2\right)^2} \left(\frac{A^2}{m_P H}\right)^2$
v.curvaton	$\frac{\epsilon}{\left(1+\left(\frac{Am_P}{A_{tot}^2}\right)^2 \epsilon r^2\right)^2}$	$\frac{\epsilon^2 r^3}{\left(1+\left(\frac{Am_P}{A_{tot}^2}\right)^2 \epsilon r^2\right)^2} \left(\frac{Am_P^2}{A_{tot}^3}\right)^2$	$\frac{\epsilon^2 r^3 g_c^2}{\left(1+\left(\frac{Am_P}{A_{tot}^2}\right)^2 \epsilon r^2\right)^2} \left(\frac{A^2 m_P^2}{A_{tot}^3 H}\right)^2$

Table 2. Order of magnitude of the vector contributions to τ_{NL} in different scenarios.

	$\tau_{NL}^{NA_1}$	$\tau_{NL}^{NA_2}$	$\tau_{NL}^{A_1}$	$\tau_{NL}^{A_2}$
general case	$10^3 \frac{\beta^2 \epsilon g_c^2}{(1+\beta)^3} \left(\frac{m_P}{H}\right)^2$	$10^{-5} \frac{\beta^{3/2} \epsilon^{3/2} g_c^2}{(1+\beta)^3} \left(\frac{A}{H}\right) \left(\frac{m_P}{H}\right) m_P^2 N_{AA}$	$\frac{\beta \epsilon^2}{(1+\beta)^3} m_P^4 N_{AA}^2$	$\frac{\beta^{3/2} \epsilon^{3/2}}{(1+\beta)^3} m_P^3 N_{AAA}$
v.inflation	same as above	$10^{-5} \frac{\beta^{3/2} \epsilon^{3/2} g_c^2}{(1+\beta)^3} \left(\frac{A}{H}\right) \left(\frac{m_P}{H}\right)$	$\frac{\beta \epsilon^2}{(1+\beta)^3}$	0
v.curvaton	same as above	$10^{-5} \frac{r \beta^{3/2} \epsilon^{3/2} g_c^2}{(1+\beta)^3} \left(\frac{A}{H}\right) \left(\frac{m_P}{H}\right) \left(\frac{m_P}{A}\right)^2$	$\frac{r^2 \beta \epsilon^2}{(1+\beta)^3} \left(\frac{m_P}{A}\right)^4$	$\frac{r \beta^{3/2} \epsilon^{3/2}}{(1+\beta)^3} \left(\frac{m_P}{A}\right)^3$

Table 3. Order of magnitude of the ratios f_{NL}^v/f_{NL}^s in different scenarios.

	f_{NL}^A/f_{NL}^s	f_{NL}^{NA}/f_{NL}^s
general case	$\beta \frac{N_{AA}}{N_{\phi\phi}}$	$\beta^2 g_c^2 \left(\frac{m}{H}\right)^2 \frac{N_\phi^2}{N_{\phi\phi}}$
v.inflation	β	$\frac{\beta^2 g_c^2}{\epsilon} \left(\frac{m_P}{H}\right)^2$
v.curvaton	$\beta r \left(\frac{m_P}{A}\right)^2$	$\frac{\beta^2 g_c^2}{\epsilon r} \left(\frac{A}{H}\right)^2$

(see Appendix 15.5 for their derivation). In the vector curvaton model the same quantities become [92, 97]

$$N_a^i = \frac{2}{3} r \frac{A_i^a}{\sum_b |\vec{A}^b|^2}, \quad N_{ab}^{ij} = \frac{1}{3} r \frac{\delta_{ab} \delta^{ij}}{\sum_c |\vec{A}^c|^2}. \quad (10.8)$$

Neglecting tensor and gauge indices, the expressions above can be simplified as $N_A \simeq A/m_P^2$ and $N_{AA} \simeq 1/m_P^2$ in vector inflation, $N_A \simeq r/A$ and $N_{AA} \simeq r/A^2$ in the vector curvaton model. Also we have $N_{AAA} = 0$ in vector inflation and $N_{AAA} \simeq r/A^3$ in vector curvaton.

We are now ready to provide the final expressions for the amplitudes: in Table 1 we list all the contributions to f_{NL} , Table 2 includes the vector contributions to τ_{NL} , the scalar contributions being given by

$$\tau_{NL}^{(s)} = \frac{\epsilon}{(1+\beta)^3} + \frac{\epsilon^2}{(1+\beta)^3}. \quad (10.9)$$

In the expressions appearing in the tables, numerical coefficients of order one have not been reported. Also, m is by definition equal to m_P in vector inflation and to A/\sqrt{r}

Table 4. Order of magnitude of the ratios τ_{NL}^v/τ_{NL}^s in different scenarios.

	$\tau_{NL}^{NA1}/\tau_{NL}^s$	$\tau_{NL}^{NA2}/\tau_{NL}^s$	$\tau_{NL}^{A1}/\tau_{NL}^s$	$\tau_{NL}^{A2}/\tau_{NL}^s$
general case	$10^3 \beta^2 g_c^2 \left(\frac{m_P}{H}\right)^2$	$10^{-5} \beta^{3/2} \epsilon^{1/2} g_c^2 \left(\frac{A}{H}\right) \left(\frac{m_P}{H}\right) m_P^2 N_{AA}$	$\beta \epsilon m_P^4 N_{AA}^2$	$\beta^{3/2} \epsilon^{1/2} m_P^3 N_{AAA}$
v.inflation	same as above	$10^{-5} \beta^{3/2} \epsilon^{1/2} g_c^2 \left(\frac{A}{H}\right) \left(\frac{m_P}{H}\right)$	$\beta \epsilon$	0
v.curvaton	same as above	$10^{-5} r \beta^{3/2} \epsilon^{1/2} g_c^2 \left(\frac{A}{H}\right) \left(\frac{m_P}{H}\right) \left(\frac{m_P}{A}\right)^2$	$r^2 \beta \epsilon \left(\frac{m_P}{A}\right)^4$	$r \beta^{3/2} \epsilon^{1/2} \left(\frac{m_P}{A}\right)^3$

in the vector curvaton model; $N_\phi \simeq (m_P \sqrt{\epsilon})^{-1}$ and $N_{\phi\phi} \simeq m_P^{-2}$, with $\epsilon \equiv (\dot{\phi}^2)/(2m_P^2 H^2)$.

The quantities involved in the amplitude expressions are g , β , r , ϵ , g_c , m_P/H , A/m_P and A/H . We already know that g and β are to be considered smaller than one (see discussion after Eq. (9.17)). Similarly, as mentioned after Eq. (8.2), r has to remain small at least until inflation ends so as to attain an ‘‘almost isotropic’’ expansion. The slow-roll parameter ϵ and the $SU(2)$ coupling g_c are small respectively to allow the inflaton to slowly roll down its potential and for perturbation theory to be valid. The ratio m_P/H is of order 10^5 (assuming $\epsilon \sim 10^{-1}$). Finally, A/m_P and A/H have no stringent bounds. A reasonable choice could be to assume that the expectation value of the gauge fields is no larger than the Planck mass, i.e. $A/m_P \leq 1$. As to the A/H ratio, different possibilities are allowed, including the one where it is of order one (see Sec. 6 of [97] for a discussion on this).

Let us now compare the different amplitude contributions. The ratios between scalar and vector contributions are shown in Table 3 for the bispectrum and Table 4 for the trispectrum. We can observe that the dominance of a given contribution w.r.t. another one very much depends on the selected region of parameter space. It turns out that it is allowed for the vector contributions to be larger than the scalar ones and also for the non-Abelian contributions to be larger than the Abelian ones. This is discussed more in details in Sec. 6 of [97]. An interesting point is, for instance, the following: ignoring tensor and gauge indices, the ratio $g_c A/H$, that appears in many of the Tables entries, is a quantity smaller than one; if we consider the different configurations identified by gauge and vector indices, we realize that this is not always true, in fact the value of this ratio can be $\gg 1$ in some configurations (see also Appendix 15.4 and Eqs. (15.4.9) through (15.4.11) in particular).

Finally, it is interesting to compare bispectrum and trispectrum amplitudes (see Table 5). Again, it is allowed for the ratios appearing in Table 5 to be either large or small, depending on the specific location within the parameter space of the theory. For instance, the combination of a small bispectrum with a large trispectrum is permitted. The latter is an interesting possibility: if the bispectrum was observably small, we could still hope the information about non-Gaussianity to be accessible thanks to the trispectrum.

Another interesting feature of this model is that the bispectrum and the trispectrum

Table 5. Order of magnitude of the ratios $\tau_{NL}^v / (f_{NL}^{NA})^2$ in different scenarios.

	$\tau_{NL}^{NA1} / (f_{NL}^{NA})^2$	$\tau_{NL}^{NA2} / (f_{NL}^{NA})^2$	$\tau_{NL}^{A1} / (f_{NL}^{NA})^2$	$\tau_{NL}^{A2} / (f_{NL}^{NA})^2$
v.i.	$10^9 \frac{\epsilon(1+\beta)}{g_c^2 \beta^2} \left(\frac{H}{m_P}\right)^2$	$10 \frac{\epsilon^{3/2}(1+\beta)}{\beta^{5/2} g_c^2} \left(\frac{A}{H}\right) \left(\frac{H}{m_P}\right)^3$	$10^6 \frac{\epsilon^2(1+\beta)}{\beta^3 g_c^4} \left(\frac{H}{m_P}\right)^4$	0
v.c.	$10^9 \frac{r^2 \epsilon(1+\beta)}{g_c^2 \beta^2} \frac{m_P^2}{A^2} \frac{H^2}{A^2}$	$10 \frac{r^5 \epsilon^{3/2}(1+\beta)}{\beta^{5/2} g_c^2} \frac{H^3}{A^3} \frac{m_P}{H} \frac{m_P^2}{A^2}$	$10^6 \frac{r^6 \epsilon^2(1+\beta)}{\beta^3 g_c^4} \left(\frac{m_P}{A}\right)^4 \left(\frac{H}{A}\right)^4$	$10^6 \frac{r^3 \epsilon^{3/2}(1+\beta)}{g_c^2 \beta^{5/2}} \frac{m_P^3}{A^3} \frac{H^4}{A^4}$

depend on the same set of quantities. If these correlation functions were independently known, that information could then be used to test the theory and place some bounds on its parameters.

11. Shape of non-Gaussianity and statistical anisotropy features

Studying the shape of non-Gaussianity means understanding the features of momentum dependence of the bispectrum and higher order correlators (see e.g. [150]). If they also depend on variables other than momenta, it is important to determine how these other variables affect the profiles for any given momentum set-up. This is the case as far as the bispectrum and the trispectrum of the gauge fields are concerned, given the fact that they are functions, besides of momenta, also of a large set of angular variables (see Eqs. (9.49) and (9.52)).

11.1. Momentum dependence of the bispectrum and trispectrum

We show the study of the momentum dependence of the F_n and G_n functions in Eqs. (9.49) and (9.52) first and then analyze the angular variables dependence of the spectra, once the momenta have been fixed in a given configuration. A natural choice would be to consider the configuration where the correlators are maximized.

The maxima can be easily determined for the bispectrum by plotting the isotropic functions F_n and G_n in terms of two of their momenta. These plots are provided in Fig. 7, where the variables are $x_2 \equiv k_2/k_1$ and $x_3 \equiv k_3/k_1$. Each one of the plots corresponds to a single isotropic functions of the sum in Eq. (9.49). It is apparent that the maxima are mostly located in the in the so-called local region, i.e. for $k_1 \sim k_2 \gg k_3$; three out of the eight graphs do not have their peaks in this configuration but, at the same time, they show negligible amplitudes compared to the “local” peaked graphs.

The situation is much more complex for the trispectrum, being the number of momentum variables larger than three ($k_1, k_2, k_3, k_4, k_{\hat{1}2}$ and $k_{\hat{1}4}$). The momentum dependence of the isotropic functions can be studied by selecting different configurations for the tetrahedron made up by the four momentum vectors, in such a way as to narrow the number of independent momentum variables down to two. A list of possible configurations was presented in [44]. We consider two of them, the “equilateral” and the “specialized planar”.

In the equilateral configuration the four sides of the tetrahedron have the same length ($k_1 = k_2 = k_3 = k_4$), therefore $x \equiv k_{\hat{1}2}/k_1$ and $y \equiv k_{\hat{1}4}/k_1$ can be chosen as variables for the plots. The plots of the isotropic functions of contact interaction and vector exchange contributions are provided in Fig. 8. The former (*c.i.*) shows a constant behaviour in this configuration, being independent of $k_{\hat{1}2}$ and $k_{\hat{1}4}$. The latter (*v.e.(I)*, *v.e.(II)* and *v.e.(III)*) diverge as $k_{\hat{1}i}^{-3}$ ($i = 1, 2, 3$ respectively for the three plots) in the limit of a flat tetrahedron, i.e. $(k_{\hat{1}i}/k_1) \rightarrow 0$.

In the specialized planar configuration, the tetrahedron is flattened and, in addition to that, three of the six momentum variables are set equal to one another ($k_1 = k_3 = k_{\hat{1}4}$); this leaves two independent variables, which can be $x \equiv k_2/k_1$ and $y \equiv k_3/k_1$. There is a double degeneracy in this configuration, due to the fact that the quadrangle can have internal angles larger than or smaller/equal to π , as we can see from the plus and minus




Figure 7. Plot of $r_n \equiv \Theta(x_2 - x_3)\Theta(x_3 - 1 + x_2)x_2^2x_3^2R_n(x_2, x_3)$, where we define $R_n = k_1^6 F_n$. The Heaviside step functions Θ help restricting the plot domain to the region (x_2, x_3) that is allowed for the triangle $\vec{k}_1 + \vec{k}_2 + \vec{k}_3 = 0$ (in particular, we set $x_3 < x_2$). We also set $x^* = 1$.




Figure 8. Plots of the isotropic functions appearing in the vector fields trispectrum (from Eq. (9.52)): *c.i.* is the contribution from contact-interaction diagrams, *v.e.(I)*, *v.e.(II)* and *v.e.(III)* are the contributions from the vector-exchange diagrams. The equilateral configuration has been considered in this figure.

signs in the expressions for k_{12} and k_{13} [44]

$$\frac{k_{12}}{k_1} = \sqrt{1 + \frac{x^2 y^2}{2} \pm \frac{xy}{2} \sqrt{(4-x^2)(4-y^2)}}, \quad (11.1)$$

$$\frac{k_{13}}{k_1} = \sqrt{x^2 + y^2 - \frac{x^2 y^2}{2} \mp \frac{xy}{2} \sqrt{(4-x^2)(4-y^2)}}. \quad (11.2)$$

The two cases are plotted in Figs. 9 and 10. Notice that divergences generally occur as $x, y \rightarrow 0$, as $x \rightarrow y$ and $(x, y) \rightarrow (2, 2)$.

11.2. Features and level of anisotropy

Statistical homogeneity and isotropy are considered characterizing features of the CMB fluctuations distribution, if one ignores the issues raised by the “anomalous” detections we presented in the introduction.

Homogeneity of the correlation functions equates translational invariance and hence total momentum conservation, as enforced by the delta functions appearing on the left-hand sides of Eqs. (9.1) through (9.3). This invariance property can then be pictured as the three momentum vectors forming a closed triangle for the bispectrum and the four momenta arranged in a tetrahedron for the trispectrum (see Fig. 11).




Figure 9. Plots of the contact interaction and of the vector-exchange contributions in the specialized planar configuration (plus sign).

Statistical isotropy corresponds to invariance w.r.t. rotations in space of the momentum (for the power spectrum) and of the triangle or tetrahedron made up by the momenta, respectively for the bispectrum and the trispectrum. This symmetry can be broken, as it for example happens in the $SU(2)$ case, by assuming the existence of preferred spatial directions in the early universe that might be revealed in the CMB observations. When this happens, the correlation functions are expected to be sensitive to the spatial orientation of the wave number or of the momenta triangles and tetrahedrons w.r.t. these special directions. Analytically, the bispectrum and the trispectrum will depend on the angles among the vector bosons and the wave vectors (besides the angles among the gauge bosons themselves), as shown in the coefficients I_n and L_n appearing in Eqs. (9.49) and (9.52). This implies that both the amplitude and the shape of bispectrum and trispectrum will be affected by these mutual spatial orientations. The modulation of the shapes by the directions that break statistical anisotropy was discussed with some examples both for the bispectrum and the trispectrum in our papers [97, 98]. These examples are here reported in Figs. 12 and 13.

In Fig. 12 we show the plot of the vector contribution to the bispectrum of ζ , properly normalized in the configuration

$$\vec{N}_3 = N_A(0, 0, 1) \tag{11.3}$$

$$\vec{N}_1 = \vec{N}_2 = N_A(\sin \theta \cos \phi, \sin \theta \sin \phi, \cos \theta), \tag{11.4}$$




Figure 10. Plots of the contact interaction and of the vector-exchange contributions in the specialized planar configuration (minus sign).

where, the (x, y, z) coordinate frame is chosen to be $\hat{k}_3 = \hat{x}$ and $\hat{k}_1 = \hat{k}_2 = \hat{z}$ and δ is the angle between $\vec{N}_{1,2}$ and \hat{k}_3 . The coefficients I_n in this configuration become

$$I_{EEE} = m^2 N_A^4 \left[-20 - 24 \cos \delta + 2 \cos \theta - 12 \cos^2 \delta + 12 \cos^2 \theta - 2 \cos^3 \theta + 6 \cos \theta \cos^2 \delta - 2 \cos^2 \theta \cos \delta + 2 \cos \delta \cos^3 \theta \right], \quad (11.5)$$

$$I_{III} = m^2 N_A^4 \left[4 \cos^2 \delta \right], \quad (11.6)$$

$$I_{IIE} = m^2 N_A^4 \left[4 - 2 \cos \theta - 6 \cos^2 \theta - 4 \cos^2 \delta \right], \quad (11.7)$$

$$I_{IEI} = m^2 N_A^4 \left[-4 \cos^2 \delta \right], \quad (11.8)$$

$$I_{IEU} = m^2 N_A^4 \left[-4 \cos^2 \delta \right], \quad (11.9)$$

$$I_{EEI} = m^2 N_A^4 \left[-2 \cos^2 \delta \right], \quad (11.10)$$

$$I_{EIE} = m^2 N_A^4 \left[4 - 4 \cos^2 \theta - 8 \cos^2 \delta \right], \quad (11.11)$$

$$I_{IEE} = m^2 N_A^4 \left[2 + \cos \theta - 3 \cos^2 \theta - 4 \cos^2 \delta \right]. \quad (11.12)$$

where $m^2 \equiv (A)/(N_A)$, A being the background value of the \vec{A}_a 's evaluated at horizon crossing. The analytic expression of the 'non-Abelian' bispectrum normalized to the ratio $(g_c^2 H^2 m^2 N_A^4)/(k_1^6 x_2^2 x_3^2)$, as a function of the angles θ and δ and for fixed values of



Figure 11. Representation of momentum conservation for the bispectrum (the three momenta form a closed triangle) and for the trispectrum (the momenta form a tetrahedron).

x^* , x_2 and x_3 is

$$B_\zeta(\theta, \delta) \simeq g_c^2 \frac{H^2}{k_1^6} \frac{m^2 N_A^4}{10^{-1}} \left[\cos^2 \delta (8 \cos \theta - 1.4 \times 10^3) + 3 \cos \delta (\cos^3 \theta - \cos^2 \theta - 11) \right. \\ \left. - 11 \cos 2\delta - 40 - 6 \cos 2\theta - \cos \theta (3 \cos^2 \theta - 30 \cos \theta - 10) \right], \quad (11.13)$$

where we set $x^* = 1$, while x_2 and x_3 were chosen in the 'squeezed' region, $x_2 = 0.9$ and $x_3 = 0.1$.

In Fig. 13 we provide a similar plot, but for the trispectrum from vector-exchange contributions and in a different configuration

$$\begin{aligned} \hat{N}_2 \cdot \hat{k}_i &= 0 \quad (i = 1, \dots, 4) \\ \hat{N}_1 \cdot \hat{k}_1 &= \cos \delta, & \hat{N}_1 \cdot \hat{k}_2 &= 0 \\ \hat{N}_3 \cdot \hat{k}_2 &= \cos \theta, & \hat{N}_3 \cdot \hat{k}_1 &= 0. \end{aligned} \quad (11.14)$$

In addition to that, let us assume that all the \vec{N}_a have the same magnitude N_A . In this configuration, we have

$$\Delta_I = \Delta_{III} = N_A^4 \cos^2 \theta \cos^2 \delta, \quad \Delta_{II} = 0, \quad (11.15)$$

therefore the expression in Eq. (9.68) becomes

$$T_\zeta \supset g_c^2 H_*^4 N_A^4 [ISO] \cos^2 \theta \cos^2 \delta \quad (11.16)$$

where the expression in brackets includes an isotropic term (which is rotationally invariant)

$$ISO \equiv \left(\frac{x^*}{k} \right)^4 \left(I \sum_{i=1}^4 t_i + III \sum_{i=9}^{12} t_i \right). \quad (11.17)$$

Fig. 13 plots the trispectrum contribution in Eq. (11.16) normalized to its isotropic part.

Another comment should be added concerning statistical anisotropy in the model. Notice that both the bispectrum and the trispectrum can be written as the sum of a purely isotropic and an anisotropic parts. The orders of magnitude of these two parts can, for instance, be read from Table 2 for the trispectrum: each one among $\tau_{NL}^{NA_2}$, $\tau_{NL}^{A_1}$ and $\tau_{NL}^{A_2}$ provide the order of magnitude of the level of both their isotropic and




Figure 12. Plot of $f(\theta, \delta) \equiv [(B_\zeta(\theta, \delta, x^*, x_2, x_3)x_2^2x_3^2k_1^6)/(g_c^2H^2m^2N_A^4)]$ evaluated at $(x^* = 1, x_2 = 0.9, x_3 = 0.1)$ in a sample angular configuration (see Eqs. (11.3) through (11.13)).




Figure 13. Plot of the anisotropic part of the trispectrum from the contribution due to vector-exchange diagrams in a sample angular configuration (see Eqs. (11.14) through (11.17)).

anisotropic contributions, which are therefore comparable; $\tau_{NL}^{NA_1}$ instead quantifies a purely anisotropic contribution which, as discussed in Sec. 10, can be comparable to the other three parts, if not the dominant one. A similar discussion applies to the bispectrum (see f_{NL}^A and f_{NL}^{NA} in Table 1). We can then conclude that, for the three and for the four point function, there is room in the parameter space of the theory for the anisotropic contributions to be as large as, or even larger than, the isotropic ones.

12. Trispectrum for $f(\tau)$ models of gauge interactions

We will now show that it is quite straightforward to extend the calculations we performed for $f = 1$ to cases where f is not a constant. One interesting model is the one studied in [149] and also recently discussed in [90], where the field is effectively massless ($m_0 = \xi = 0$) so the action (8.11) for the gauge field becomes

$$S = \int d^4x \sqrt{-g} \left[-\frac{1}{4} f^2(\phi) g^{\mu\nu} g^{\rho\sigma} F_{\mu\nu}^a F_{\rho\sigma}^a + \dots \right], \quad (12.1)$$

where again $F_{\mu\nu}^a \equiv \partial_\mu B_\nu^a - \partial_\nu B_\mu^a + g_c \varepsilon^{abc} B_\mu^b B_\nu^c$.

Let us introduce the fields \tilde{A}_i^a and A_i^a , related by the equations $\tilde{A}_i^a \equiv f B_i^a = a A_i^a$. The A_i^a are the physical fields.

We can expand the perturbations of \tilde{A}_i^a in terms of creation and annihilation operators in the usual way

$$\delta \tilde{A}_i^a(\eta, \vec{x}) = \int \frac{d^3q}{(2\pi)^3} \sum_{\lambda=R,L} \left[e_i^\lambda(\hat{q}) a_{\vec{q}}^{a\lambda} \delta \tilde{A}_\lambda^a(\eta, q) + h.c. \right]. \quad (12.2)$$

If $f = f_0 a^\alpha$, with α equal either to 1 or -2 (f_0 is a constant), it is possible to prove [90] that the equation of motion for $\delta \tilde{A}_\lambda^a$ is the same as the one for δB^T , where by δB^T we mean the transverse mode function in Eq. (8.14). This is equivalent to saying that, under the assumption $\alpha = 1, -2$, the physical gauge fields are governed by the same equation of motion as a light scalar field in a de Sitter space and so they generate a scale invariant power spectrum. Let us sketch the calculation of the trispectrum in this theory.

The general expressions of the Abelian terms still hold, except that the power spectrum in Eq. (9.1) reduces to

$$P_{ij}^{ab} = T_{ij}^{ab} P_+, \quad (12.3)$$

having gauged the longitudinal modes away.

Let us now have a look at the non-Abelian part. First of all, we need to set $n(x^*) = 0$. The interaction Hamiltonian to third and fourth order are the same as in Eqs. (9.38) and (9.39), but with extra $f^2(\tau)$ factors. The anisotropy coefficients such as I_n and L_n that survive after setting the longitudinal mode to zero do not change. On the other hand, the wavefunctions for the gauge fields are now given by $\delta B = \delta \tilde{A}/f = \delta B^T/f_0 a^\alpha$. The new trispectrum therefore differs because of extra scale factors inside and outside the time integrals, which in general imply a different power of H_* in the final results and a different momentum dependence in the isotropic part of the expressions.

Let us take into account the trispectrum. The three non-Abelian vector contributions from Eq. (9.34) in the $f = 1$ case can be schematically written as follows

$$\langle \delta B^4 \rangle_{line3} \simeq (\delta B^3) \int d\eta (\delta B)^3 = (\delta B^T)^3 \int d\eta (\delta B^T)^3, \quad (12.4)$$

$$\begin{aligned} \langle \delta B^4 \rangle_{v.e.} &\simeq (\delta B)^4 \int d\eta' (\delta B)^3 \int d\eta'' (\delta B)^3 \\ &= (\delta B^T)^4 \int d\eta' (\delta B^T)^3 \int d\eta'' (\delta B^T)^3, \end{aligned} \quad (12.5)$$




Figure 14. Plots of the isotropic functions of some of the vector-exchange contributions in the equilateral configuration, for the $f \simeq a^{-2}$ model. In this and in the next figures, “v.e.(I,II,III)” represent the isotropic functions associated with the very last term in square brackets in Eq. (12.13); “v.e.(new)” represents the isotropic function associated with the k_{1144} , k_{2244} , k_{1133} and k_{2233} terms in the second line of Eq. (12.13).

$$\langle \delta B^4 \rangle_{c.i.} \simeq (\delta B)^4 \int d\eta (\delta B)^4 = (\delta B^T)^4 \int d\eta (\delta B^T)^4, \quad (12.6)$$

where we indicate with the subscript *line3* the contribution from the third line of Eq. (9.34) and with *v.e.* and *c.e.* respectively the vector exchange and the contact interaction contributions to the first line of (9.34). $\langle \delta B^4 \rangle \equiv \langle \delta B_i^a(\vec{k}_1) \delta B_j^b(\vec{k}_2) \delta B_k^c(\vec{k}_3) \delta B_l^d(\vec{k}_4) \rangle$ and we have omitted all the gauge and vector indices, as well as complex conjugate symbols, for simplicity. Let us see now how Eqs. (12.4)-(12.6) change if $f = f_0 a^\alpha$ ($\alpha = 1, -2$)

$$\langle \delta B^4 \rangle_{line3} \simeq (\delta B)^3 \int d\eta f^2 (\delta B)^3 \simeq \left(\frac{\delta B^T}{a^\alpha} \right)^3 \int d\eta a^{2\alpha} \left(\frac{\delta B^T}{a^\alpha} \right)^3, \quad (12.7)$$

$$\begin{aligned} \langle \delta B^4 \rangle_{v.e.} &\simeq (\delta B)^4 \int d\eta' f^2 (\delta B)^3 \int d\eta'' f^2 (\delta B)^3 \\ &\simeq \left(\frac{\delta B^T}{a^\alpha} \right)^4 \int d\eta' a^{2\alpha} \left(\frac{\delta B^T}{a^\alpha} \right)^3 \int d\eta'' a^{2\alpha} \left(\frac{\delta B^T}{a^\alpha} \right)^3, \end{aligned} \quad (12.8)$$

$$\langle \delta B^4 \rangle_{c.i.} \simeq (\delta B)^4 \int d\eta f^2 (\delta B)^4 \simeq \left(\frac{\delta B^T}{a^\alpha} \right)^4 \int d\eta a^{2\alpha} \left(\frac{\delta B^T}{a^\alpha} \right)^4. \quad (12.9)$$

Using $a = (-H\eta)^{-1}$ in the previous equations, we get




Figure 15. Plots of the isotropic functions of some of the vector-exchange contributions in the specialized planar configuration (plus sign), for the $f \simeq a^{-2}$ model.

$$\langle \delta B^4 \rangle_{line3} \simeq H_*^{4\alpha} \left(\delta B^T(-\eta^*)^\alpha \right)^3 \int d\eta \left(\delta B^T \right)^3 (-\eta)^\alpha, \quad (12.10)$$

$$\begin{aligned} \langle \delta B^4 \rangle_{v.e.} &\simeq H_*^{6\alpha} \left(\delta B^T(-\eta^*)^\alpha \right)^4 \int d\eta' \left(\delta B^T \right)^3 (-\eta')^\alpha \\ &\times \int d\eta'' \left(\delta B^T \right)^3 (-\eta'')^\alpha, \end{aligned} \quad (12.11)$$

$$\langle \delta B^4 \rangle_{c.i.} \simeq H_*^{6\alpha} \left(\delta B^T(-\eta^*)^\alpha \right)^4 \int d\eta \left(\delta B^T \right)^4 (-\eta)^{2\alpha}. \quad (12.12)$$

Let us now consider more in details the $\alpha = -2$ case for contact-interaction and vector-exchange contributions. The expressions for the anisotropy coefficients are respectively given by

$$\begin{aligned} T_{ijkl}^{EEEE} &= k_1 k_3 (\hat{k}_1 \cdot \hat{k}_3 - \hat{k}_1 \cdot \hat{k}_{12} \hat{k}_3 \cdot \hat{k}_{12}) [\delta_{ij} \delta_{kl} - \delta_{ij} \hat{k}_{k4} \hat{k}_{l4} - \delta_{ij} \hat{k}_{k3} \hat{k}_{l3} - \delta_{kl} \hat{k}_{i2} \hat{k}_{j2} - \delta_{kl} \hat{k}_{i1} \hat{k}_{j1} \\ &+ \delta_{ij} \hat{k}_{k3} \hat{k}_{l4} \hat{k}_3 \cdot \hat{k}_4 + \delta_{kl} \hat{k}_{i1} \hat{k}_{j2} \hat{k}_1 \cdot \hat{k}_2 + k_{1144} + k_{2244} + k_{1133} + k_{2233} - k_{2234} \hat{k}_3 \cdot \hat{k}_4 \\ &- k_{1134} \hat{k}_3 \cdot \hat{k}_4 - k_{1244} \hat{k}_1 \cdot \hat{k}_2 + k_{1233} \hat{k}_1 \cdot \hat{k}_2 + k_{1234} \hat{k}_1 \cdot \hat{k}_2 \hat{k}_3 \cdot \hat{k}_4] \end{aligned} \quad (12.13)$$

and by Eq. (9.73), for one of the possible permutations. These expressions are more complicated w.r.t T_{ijkl}^{llll} in Eq. (9.63) and T_{mnop}^{llll} in Eq. (9.88) for the longitudinal modes. As a result, when studying the shape of the trispectrum, for the isotropic functions appearing in it, several diagrams need to be taken into account, one for each term in




Figure 16. Plots of the isotropic functions of some of the vector-exchange contributions in the specialized planar configuration (minus sign), for the $f \simeq a^{-2}$ model.

T_{ijkl}^{EEEE} and T_{ijkl}^{EEEEEE} . For comparison with the $f = 1$ case, we plotted the isotropic functions associated with the very last term in square brackets in Eq. (12.13) (see “ $-v.e(I)$ ”, “ $-v.e(II)$ ” and “ $-v.e(III)$ ” in Figs. 14, 15 and 16). By comparing these plots with the ones in Figs. 8, 9 and 10, it is evident that they have very similar shapes. On the other hand, when we consider the isotropic functions associated with terms that are not present in the $f = 1$ case, several differences arise in the plots; we provide a sample in Figs. 14, 15 and 16 with the “ $v.e.(new)$ ” plots, which represent isotropic functions associated with the k_{1144} , k_{2244} , k_{1133} and k_{2233} terms in the second line of Eq. (12.13). We verified that similar observations can be made concerning the shapes of the contact-interaction contributions.

13. Conclusions

Cosmology has entered what it is called its “precision era”: over the past few years, observations of the CMB and of other cosmological probes have been performed which have greatly improved the previous bounds on some of the fundamental parameters characterizing the early Universe cosmology. These bounds are expected to become a lot stricter with the advent of new experiments, such as the ongoing Planck satellite mission. The theoretical search for models of the early Universe has consequently been concentrated on, among other things, trying to produce more and more accurate predictions.

In this optics, alternative models to the basic single-field slow-roll inflationary scenario have been and are being investigated, with a strong focus on the computation of higher order (three and four-point) correlation functions and on higher order (beyond tree-level) corrections. All of these “higher order” predictions are expected to be particularly revealing of the early Universe physics, given that they are an indication of the type of fields and interactions populating the specific inflationary scenarios.

In this thesis I have presented an overview of the existing results concerning some of these “higher order” predictions; in particular, I focused on our works on loop corrections in inflationary scalar field models and on the primordial non-Gaussianity and statistical anisotropy predictions in some inflationary models populated with non-Abelian vector fields.

Loop corrections to the power spectrum of the comoving curvature fluctuation ζ in single-field inflation arise both from the inflaton self-interactions and from the coupling of the scalar field with gravitons. We have calculated the corrections from tensor loops, previously neglected in the literature for simplicity reasons. It turns out that one-loop corrections from tensor-scalar interactions are of the same order of magnitude as those arising from scalar self-interactions, therefore they cannot be neglected in a self-consistent calculation.

One loop corrections have been found to be suppressed by an $(H/m_P)^2$ factor compared to the tree-level result; they exhibit a slightly different dependence from the external momentum, because of the presence of a logarithmic factor (which does not spoil scale invariance) and they turn out vary with time at most as fast as the logarithm of the scale factor. One-loop diagrams generally present both infrared logarithmic and ultraviolet power-law (in addition to logarithmic) divergences. The ultraviolet divergences can be treated using ordinary regularization and renormalization techniques as in flat-space quantum field theory. The infrared divergences are “cured” by introducing an infrared cutoff represented by the smallest observable physical mode, i.e. by considering a finite space, a sort of “box” of observation, with a size equal to the current horizon length. As expected, tensor modes provide, as well as the scalar modes, observably small corrections at one loop level, as long as the basic inflationary scenario is concerned. Their

computation is nevertheless instructive and provides a path that can be followed in the investigation of “non-standard” models of inflation. More appealing results are for instance expected in inflationary scenarios described by non-canonical Lagrangians, some of which, $P(X, \phi)$ models, have been described and analysed in this thesis.

Motivated by an interest in models that combine non-Gaussianity and statistical anisotropy predictions for the CMB fluctuations, we have worked on theories of inflation where primordial vector fields effectively participate in the production of the ζ perturbations, eventually focusing on some $SU(2)$ vector fields models. The two, three and four point correlation functions in these models result as the sum of scalar and vector contributions. The latter are of two kinds, “Abelian” (i.e. arising from the zeroth order terms in the Schwinger-Keldysh expansion) and “non-Abelian” (i.e. originating from the self-interactions of the vector fields). The bispectrum and the trispectrum final results are presented as a sum of products of isotropic functions of the momenta (F_n and G_n in the text) multiplied by anisotropy coefficients (which we indicated by I_n and L_n) that depend on the angles among all gauge and wave vectors. The amplitude of non-Gaussianity has been evaluated through the parameters f_{NL} and τ_{NL} ; in particular we have discussed the dependence of these functions from the non-angular parameters of the theory. We have provided the comparisons among the different (scalar versus vector, Abelian versus non-Abelian) contributions to f_{NL} and τ_{NL} , noticing that any one of them can be the dominant contribution depending on the selected region of parameter space. In particular, we have stressed how the anisotropic contributions to the bispectrum and the trispectrum can overcome the isotropic parts. An interesting feature of these models is that the bispectrum and the trispectrum depend on the same set of parameters and their amplitudes are therefore strictly related to one another. We have presented the shapes of both the bispectrum and the trispectrum. The isotropic functions appearing in their final expressions have been analyzed separately from their anisotropy coefficients. The bispectrum isotropic functions have been found to preferably show a local shape. The trispectrum ones have been plotted selecting equilateral and specialized planar configurations. Finally, the full expressions (i.e. completed by the anisotropy coefficients) of bispectrum and trispectrum have been presented in specific momenta configuration, in order to provide a hint of the modulation of shapes and amplitudes operated by anisotropy.

In our view, the most promising features of these models consists in the possibility of providing non-Gaussianity and statistical anisotropy predictions that are related to one another because of the fact that they share the same underlying theory. Models that combine both types of predictions could be more easily testable and, from non-Gaussianity measurement, more stringent statistical anisotropy predictions could be produced or viceversa.

Inflationary models that do not spoil the current agreement with experimental data constitute a huge variety but many of them have very distinctive features that might

be confirmed or ruled out by observations. Both the the non-Gaussianity and statistical anisotropy predictions and the nature and the amount of higher corrections to the cosmological correlation functions can be ranked among these distinctive features and certainly deserve further investigations, looking forward for a confront with new and promising experimental data.

While completing this thesis, the Wilkinson Microwave Anisotropy Probe (WMAP) team published its seven-year data analyses [153, 154]. The new bounds on non-Gaussianity from this study are given by $-10 < f_{NL}^{equil} < 74$ and $-214 < f_{NL}^{loc} < 266$, at 95% CL.

14. Acknowledgments

A sincere and warm thank to my supervisors Nicola Bartolo and Sabino Matarrese for letting me in the most enjoyable and fruitful working environment someone could ever hope for, for being there for me anytime I needed, for being two unique models of commitment and love for physics and for being such incredibly nice people.

A special thank to Toni Riotto: I have always admired him as a scientist and, in addition to this, I came to truly appreciate him as a person and as a teacher, especially during the last year of my Ph.D, when I had the great chance to know him better and to work for him. Thanks also for always finding a spot to talk with me.

To Nicola, Sabino and Toni: I know how lucky I have been to be part of your team during these three years and I hope, if not now, at least in the future, to convince you that all the time and energies you dedicated to help me during my Ph.D and the trust you gave me have been well spent.

15. Appendices

15.1. Computation of the interaction Hamiltonian

The propagator of two fields ϕ_1 and ϕ_2 is defined by (see for example [151])

$$\langle \phi_1 \phi_2 \rangle = \int D\phi D\Pi e^{i \int d^4x (\Pi \dot{\phi} - H)}, \quad (15.1.1)$$

where Π is the momentum conjugate to ϕ and H is the Hamiltonian density. If H is quadratic in Π , as it happens for example in flat space-time for a field governed by a Lagrangian $L = \int d^4x \left(\frac{1}{2} \partial_\mu \partial^\mu \phi - V(\phi) \right)$ the square in the exponent can be completed and the integral in Π evaluated and all is left is

$$\langle \phi_1 \phi_2 \rangle = \int D\phi e^{iL}. \quad (15.1.2)$$

So, if an interaction term with time derivatives appears in the Lagrangian, Π and ϕ are independent fields in the path integral. This will provide some extra vertices that need to be accounted for in the Feynman diagrams. We sketch a derivation of these extra vertices. It will turn out to be similar to what is done in [108], although complicated by the presence of gravitons. To keep the calculations easier we will at first ignore all spatial derivative and tensor indices, this will also make the notation simpler. Also, we will momentarily ignore all numerical real coefficients; it is instead very important to keep track of imaginary coefficients, time derivatives and powers of the scale factor a , so we will make sure they are all accounted for in our analysis.

The total action is $S = \int d\eta (L_\gamma + L_\phi)$, where

$$\begin{aligned} L_\gamma &= a^2 \gamma'^2 + \Gamma_\gamma \gamma' \delta \phi'^2 + \Gamma_\gamma \gamma'^3 + \Gamma_\phi \delta \phi' \gamma'^2 + \Gamma_{\gamma\phi} \delta \phi' \gamma' + \Gamma_{\gamma\gamma} \gamma'^2 \\ &\quad + \lambda_{\phi\gamma} \delta \phi' + \lambda_{\phi\phi\gamma} \gamma' + \lambda_{\gamma\gamma} \gamma', \\ L_\phi &= a^2 \delta \phi'^2 + \Gamma_1 \delta \phi'^2 + \Gamma_2 \delta \phi'^2 + \omega \delta \phi'^3 + \lambda \delta \phi', \end{aligned} \quad (15.1.3)$$

where $f' \equiv df/d\eta$ and where we define

$$\Gamma_\gamma \sim a\gamma, \quad (15.1.4)$$

$$\Gamma_\phi \sim a\delta\phi, \quad (15.1.5)$$

$$\Gamma_1 \sim \dot{\phi} a^2 \delta\phi, \quad (15.1.6)$$

$$\Gamma_2 \sim a^2 \delta\phi^2, \quad (15.1.7)$$

$$\Gamma_{\gamma\phi} \sim a^2 \delta\phi\gamma, \quad (15.1.8)$$

$$\Gamma_{\gamma\gamma} \sim a^2 \gamma^2, \quad (15.1.9)$$

$$\omega \sim a\delta\phi, \quad (15.1.10)$$

$$\lambda_{\phi\phi\phi} \sim a\delta\phi^3, \quad (15.1.11)$$

$$\lambda_{\phi\gamma\gamma} \sim a\delta\phi\gamma^2, \quad (15.1.12)$$

$$\lambda_{\phi\phi\gamma} \sim a\gamma\delta\phi^2, \quad (15.1.13)$$

$$\lambda_{\gamma\gamma\gamma} \sim a\gamma^3. \quad (15.1.14)$$

Notice that in equations (15.1.4) through (15.1.14) we use the equivalence symbol meaning that we skip details about integrations in momenta and real coefficients.

The conjugate momenta are

$$\Pi^\gamma \equiv \frac{\delta L}{\delta \gamma'} = a^2 \gamma' + \lambda_{\gamma\gamma} + \lambda_{\phi\phi\gamma} + \Gamma_{\gamma\phi} \delta\phi' + \Gamma_\gamma \delta\phi'^2 + \Gamma_{\phi\gamma'} \delta\phi' + \Gamma_{\gamma\gamma} \gamma'^3 + \Gamma_\gamma \gamma'^2, \quad (15.1.15)$$

$$\Pi^\phi \equiv \frac{\delta L}{\delta(\delta\phi')} = \Gamma_\gamma \gamma' \delta\phi' + \Gamma_{\gamma\phi} \gamma' + \lambda_{\phi\phi\gamma} + a^2 \delta\phi' + \Gamma_1 \delta\phi' + \Gamma_2 \delta\phi' + \lambda_{\phi\phi\phi} + \omega \delta\phi'^2 \quad (15.1.16)$$

We solve perturbatively the equations (15.1.15) and (15.1.16) in order to derive γ' and $\delta\phi'$ to fourth order

$$\begin{aligned} \gamma' &= a^{-2} \left[\Pi^\gamma + \lambda_{\gamma\gamma} + \lambda_{\phi\phi\gamma} + a^{-2} \Gamma_{\gamma\phi} \Pi^\phi + a^{-4} \Gamma_\gamma \Pi^\phi \Pi^\phi + a^{-4} \Gamma_\phi \Pi^\gamma \Pi^\phi \right. \\ &\quad + a^{-2} \Gamma_{\gamma\gamma} \Pi^\gamma + a^{-4} \Gamma_\gamma \Pi^\gamma \Pi^\gamma + a^{-4} \Gamma_{\gamma\phi} \Gamma_1 \Pi^\phi + a^{-4} \Gamma_\phi \Pi^\gamma \Pi^\phi \\ &\quad \left. + a^{-6} \Gamma_\phi \Gamma_1 \Pi^\gamma \Pi^\phi + a^{-2} \Gamma_{\gamma\gamma} \Pi^\phi \right], \end{aligned} \quad (15.1.17)$$

$$\begin{aligned} \delta\phi' &= a^{-2} \left[\Pi^\phi + a^{-4} \Gamma_\gamma \Pi^\gamma \Pi^\phi + a^{-6} \Gamma_\gamma \Gamma_1 \Pi^\gamma \Pi^\phi + a^{-2} \Gamma_{\gamma\phi} \Pi^\gamma + \lambda_{\phi\phi\gamma} \right. \\ &\quad + a^{-2} \Gamma_1 \Pi^\phi + a^{-4} \Gamma_1 \Gamma_1 \Pi^\phi + a^{-2} \Gamma_2 \Pi^\phi + \lambda_{\phi\phi\phi} + a^{-4} \omega \Pi^\phi \Pi^\phi \\ &\quad + a^{-4} \Gamma_{\gamma\phi} \Gamma_1 \Pi^\gamma + a^{-2} \Gamma_1 \lambda_{\phi\phi\gamma} + a^{-6} \Gamma_1 \Gamma_1 \Gamma_1 \Pi^\phi + a^{-4} \Gamma_1 \Gamma_2 \Pi^\phi \\ &\quad \left. + a^{-2} \Gamma_1 \lambda_{\phi\phi\phi} + a^{-6} \Gamma_1 \omega \Pi^\phi \Pi^\phi \right]. \end{aligned} \quad (15.1.18)$$

The next steps are: derive the hamiltonian $H = \Pi^\gamma \gamma' + \Pi^\phi \delta\phi' - L(\gamma, \delta\phi, \gamma', \delta\phi')$, where we need to plug in the solution (15.1.17) and (15.1.18) for γ' and $\delta\phi'$; construct the action as $S = S_0 + S_\Pi$, where $S_0 = \int d\eta (L_\gamma + L_\phi)$ and S_Π includes the terms that depend on the conjugate momenta of the fields (a change of variables similar to the one that Seery performs in [108] over the conjugate momenta will also be necessary).

Let's consider the vertices in S_Π that are involved in the corrections to the one loop point function for the scalar field

$$\begin{aligned} S_\Pi \supset \int d\eta' &\left[a^{-4} \Gamma_1 \delta\phi' \Pi^\phi + a^{-4} \Gamma_1 \Pi^\phi \Pi^\phi + a^{-2} \Gamma_1 \gamma' \Pi^\phi + a^{-4} \Gamma_2 \Pi^\phi \Pi^\phi \right. \\ &\left. + a^{-4} \Gamma_\phi \delta\phi' \Pi^\gamma \Pi^\gamma + a^{-4} \omega \delta\phi' \Pi^\phi \Pi^\phi \right]. \end{aligned} \quad (15.1.19)$$

The first three vertices belong to the third order part of the action; $a^{-4} \Gamma_1 \delta\phi' \Pi^\phi$ and $a^{-2} \Gamma_1 \gamma' \Pi^\phi$, provide a correction to the two point function at one loop with two vertices. Because of the presence of Γ_1 which involves a factor of $\dot{\phi}$, it is subleading in slow roll order w.r.t. the corrections coming from fourth order vertices. We will therefore neglect these diagrams. The same applies to the second vertex, $a^{-4} \Gamma_1 \Pi^\phi \Pi^\phi$, although this may contribute to correcting the one point function

$$\langle \delta\phi_{\vec{k}}(\eta^*) \rangle \supset C_1 \frac{H_*^2}{k^3} \sqrt{\epsilon} \int d^3 q f_1(\vec{q}) \int_{-\infty}^{\eta^*} d\eta' \delta(0) (1 - ik\eta') e^{ik\eta'} + c.c., \quad (15.1.20)$$

where C_1 is a numerical real coefficient and $\delta(0)$ is the Dirac delta function deriving from the propagator of the Π^ϕ 's and f_1 is a scalar function of the internal momentum. The main contribution to the integral is due to times around horizon crossing since at early times the rotation to imaginary plane of the contour integral makes the exponent

decrease rapidly to zero and moreover η^* was chosen to be just a few e-folding after horizon crossing. Also, since the integrand function goes to zero as η' approaches zero, we get a good approximation of this integral taking the upper limit $\eta^* \rightarrow 0$. The result is purely imaginary and it cancels out with its complex conjugate.

Let us now move to the fourth order vertices. From $a^{-4}\Gamma_2\Pi^\phi\Pi^\phi$ we have

$$\langle \delta\phi_{\vec{k}_1}^-(\eta^*)\delta\phi_{\vec{k}_2}^-(\eta^*) \rangle \supset \frac{H^4}{k^6} C_2 \int d^3q f_2(\vec{q}) \int_{-\infty}^{\eta^*} d\eta' \delta(0) (1 - ik\eta')^2 e^{2ik\eta'} + c.c. \quad (15.1.21)$$

The same consideration as in (15.1.20) apply to the integral above, which gives a zero contribution, as well as the following diagrams (corresponding to the last two vertices in (15.1.19))

$$\begin{aligned} \langle \delta\phi_{\vec{k}_1}^-(\eta^*)\delta\phi_{\vec{k}_2}^-(\eta^*) \rangle \supset & \frac{H^4}{k^6} C_3 \int d^3q f_3(\vec{q}) \int_{-\infty}^{\eta^*} d\eta' \delta(0) \eta'^2 (1 - ik\eta') e^{2ik\eta'} \\ & + \frac{H^4}{k^6} C_4 \int d^3q f_4(\vec{q}) \int_{-\infty}^{\eta^*} d\eta' \delta(0) \eta'^2 (1 - ik\eta') e^{2ik\eta'} + c.c. \end{aligned} \quad (15.1.22)$$

15.2. Study of leading slow roll order vertices in the fourth order action

We are interested in computing the correlators just a few e-foldings after the scales we consider cross the horizon, so we can assume that the slow roll parameters remain small and can be treated as constants during this length of time. It is then correct to limit our interest to the leading order slow-roll contribution to the action .

Let us start from the study of the slow-roll order of the fluctuations derived as solution to the constraint equations:

$$\begin{aligned} \alpha_1 &= \sqrt{\varepsilon} Q_{\alpha_1}[\delta\phi], \\ \theta_1 &= \sqrt{\varepsilon} Q_{\theta_1}[\delta\phi], \\ \alpha_2 &= \varepsilon R_{\alpha_2}[\delta\phi^2] + \sqrt{\varepsilon} S_{\alpha_2}[\delta\phi, \gamma] + T_{\alpha_2}[\delta\phi^2], \\ \theta_2 &= \varepsilon R_{\theta_2}[\delta\phi^2] + \sqrt{\varepsilon} S_{\theta_2}[\delta\phi, \gamma] + T_{\theta_2}[\delta\phi^2] + \varepsilon^2 U_{\theta_2}[\delta\phi^2] + V_{\theta_2}[\gamma^2] \\ &\quad + \varepsilon^{3/2} W_{\theta_2}[\delta\phi, \gamma], \\ \beta_j &= \varepsilon R_j[\delta\phi^2] + \sqrt{\varepsilon} S_j[\delta\phi, \gamma] + T_j[\delta\phi^2] + V_j[\gamma^2], \end{aligned}$$

where $S[\delta\phi, \gamma]$ is a linear function of $\delta\phi$ and/or its derivatives and a linear function of γ and/or its derivatives, $R[\delta\phi^2]$ is a quadratic function of $\delta\phi$ and/or its derivatives and so on. Notice that the first order fluctuations are subleading ($\sim \sqrt{\varepsilon}$) w.r.t. the second order ones ($\sim \varepsilon^0$). This criterium allows a suppression of a large number of terms in the 4th order action based on keeping the leading order (i.e. $\sim \varepsilon^0$) terms only

$$\begin{aligned} S_4 &= a^3 \int dt d^3x \left[\frac{1}{4a^2} (\partial_i \beta_j + \partial_j \beta_i) \partial_i \beta_j - \frac{1}{a^2} \delta\dot{\phi} (\partial_j \theta_2 + \beta_j) \partial_j \delta\phi + 3H^2 \alpha_2^2 \right. \\ &\quad - \frac{1}{a^2} \left(\frac{1}{4} \gamma_{ik} \gamma_{kj} \partial_j \delta\phi \partial_i \delta\phi + \alpha_2 \partial_i \delta\phi \partial_i \delta\phi - \partial_k \theta_2 \gamma_{ab} \partial_b \gamma_{ak} + \frac{1}{2} \gamma_{ab} \partial_k \gamma_{ab} \partial_k \theta_2 \right) \\ &\quad \left. + \frac{1}{a^2} \left(\beta_k \gamma_{ab} \partial_b \gamma_{ak} - \frac{1}{2} \gamma_{ab} \beta_k \partial_k \gamma_{ab} \right) \right]. \end{aligned}$$

It can be easily shown that the terms in the action that do not contain the gravitons reproduce the ones in equation (37) of [152]. The contribution to the power spectrum due to these vertices has been calculated by these authors, but only for the scalar part. We will then focus on all the tensor contributions from these and from the remaining terms. Interaction vertices with both two and four tensor fluctuations will be obtained once the expressions for α_2 , θ_2 and β_j are plugged in the action. The terms in the action that we need for constructing Feynman diagrams with one loop of gravitons are

$$S_{\gamma^2} = a^3 \int d^3x dt \left[-\frac{1}{4a^2} \beta_j \partial^2 \beta_j - \frac{1}{a^2} \delta \dot{\phi} \partial_j \delta \phi \partial_j \theta_2 - \frac{1}{4a^2} \partial_j \delta \phi \partial_i \delta \phi \gamma_{ik} \gamma_{kj} \right. \\ \left. - \frac{1}{a^2} \delta \dot{\phi} \partial_j \delta \phi \beta_j + \frac{1}{2a^2} \left(2\gamma_{ab} \partial_a \gamma_{ak} (\partial_k \theta_2 + \beta_k) - \gamma_{ab} \partial_k \gamma_{ab} (\beta_k + \partial_k \theta_2) \right) \right]. \quad (15.2.1)$$

Let's plug the expressions for β_j and θ_2 into (15.2.1) considering the terms with two gravitons. The result is an ensemble of vertices which can in principle contribute to the one loop corrections to the power spectrum of the scalar field. Apart from $\partial_j \delta \phi \partial_i \delta \phi \gamma_{ik} \gamma_{kj}$, all of the other terms contain time derivatives of one, two or three of the four fields

$$\beta_j \partial^2 \beta_j \supset a^4 \left[\partial^{-4} \left(\partial_m \partial_j \delta \dot{\phi} \partial_m \delta \phi - \partial^2 \delta \dot{\phi} \partial_j \delta \phi + \partial_j \delta \dot{\phi} \partial^2 \delta \phi \right. \right. \\ \left. \left. - \partial_m \delta \dot{\phi} \partial_m \partial_j \delta \phi \right) (\gamma_{ik} \partial_i \gamma_{kj} - \gamma_{il} \partial_i \gamma_{lj}) \right. \\ \left. + \partial^{-2} \left(\partial_m \partial_j \delta \dot{\phi} \partial_m \delta \phi - \partial^2 \delta \dot{\phi} \partial_j \delta \phi + \partial_j \delta \dot{\phi} \partial^2 \delta \phi \right. \right. \\ \left. \left. - \partial_m \delta \dot{\phi} \partial_m \partial_j \delta \phi \right) \partial^{-2} (\gamma_{ik} \partial_i \gamma_{kj} - \gamma_{il} \partial_i \gamma_{lj}) \right], \quad (15.2.2)$$

$$\delta \dot{\phi} \partial_j \delta \phi \partial_j \theta_2 \supset \delta \dot{\phi} \partial_j \delta \phi \frac{1}{16H} \partial^{-2} \partial_j \left[\frac{1}{2a^2} \partial_a \gamma_{iq} \partial_a \gamma_{iq} + \gamma_{ij} \gamma_{ij} \right] \quad (15.2.3)$$

$$\delta \dot{\phi} \partial_j \delta \phi \beta_j \supset \delta \dot{\phi} \partial_j \delta \phi \frac{a^2}{2} \partial^{-2} \partial_j \left[\gamma_{ik} \partial_i \gamma_{kj} - \gamma_{ik} \partial_i \gamma_{kj} \right], \quad (15.2.4)$$

$$\gamma_{ab} \partial_a \gamma_{bk} \partial_k \theta_2 \supset \gamma_{ab} \partial_a \gamma_{bk} \frac{a^2}{4H} \partial^{-2} \partial_k \left[-6H \partial^{-2} \Sigma - \delta \dot{\phi}^2 - \frac{1}{a^2} \partial_i \delta \phi \partial_i \delta \phi \right] \quad (15.2.5)$$

$$\gamma_{ab} \partial_a \gamma_{bk} \beta_k \supset \gamma_{ab} \partial_a \gamma_{bk} 2a^2 \partial^{-4} \left[\partial_m \partial_k \delta \dot{\phi} \partial_m \delta \phi - \partial^2 \delta \dot{\phi} \partial_k \delta \phi \right. \\ \left. + \partial_k \delta \dot{\phi} \partial^2 \delta \phi - \partial_m \delta \dot{\phi} \partial_m \partial_k \delta \phi \right], \quad (15.2.6)$$

$$\gamma_{ab} \partial_k \gamma_{ab} \beta_k \supset \gamma_{ab} \partial_k \gamma_{ab} \partial^{-4} \left[\partial_m \partial_k \delta \dot{\phi} \partial_m \delta \phi - \partial^2 \delta \dot{\phi} \partial_k \delta \phi + \partial_k \delta \dot{\phi} \partial^2 \delta \phi \right. \\ \left. - \partial_m \delta \dot{\phi} \partial_m \partial_k \delta \phi \right] \quad (15.2.7)$$

$$\gamma_{ab} \partial_k \gamma_{ab} \partial_k \theta_2 \supset \gamma_{ab} \partial_k \gamma_{ab} \left[-6H \partial^{-2} \left(\partial^2 \delta \phi \delta \dot{\phi} + \partial_j \delta \phi \partial_j \delta \dot{\phi} \right) \right. \\ \left. - \delta \dot{\phi}^2 - \frac{1}{a^2} \partial_i \delta \phi \partial_i \delta \phi \right]. \quad (15.2.8)$$

We will now prove that the vertices that include time derivatives do not actually contribute to the two point function. First of all notice that the tensor fields carry polarization tensors ϵ_{ij} with the property $q^i \epsilon_{ij} = 0$ and are always contracted with

other tensor fields in the calculations; this implies that, if a partial derivative index is contracted with a tensor index, that diagram will be zero. Based on this observation, we can ignore several of the vertices with time derivatives. We are eventually left with only two of them, that we will call V_1 , V_2 and V_3

$$V_1 \sim \partial_j \left(\delta \dot{\phi} \partial_j \delta \phi \right) \partial^{-2} \left(\partial_a \gamma_{bc} \partial_a \gamma_{bc} \right), \quad (15.2.9)$$

$$V_2 \sim \partial_j \left(\delta \dot{\phi} \partial_j \delta \phi \right) \partial^{-2} \left(\dot{\gamma}_{ab} \gamma_{ab} \right), \quad (15.2.10)$$

$$V_3 \sim \dot{\gamma}_{ab} \partial_k \gamma_{ab} \left(\beta_k + \partial_k \theta_2 \right). \quad (15.2.11)$$

where (15.2.11) is given by the sum of (15.2.7) and (15.2.8). Notice that the γ_{ij} fields need to be contracted between each other and that $\sum_{\lambda, \lambda'} \epsilon_{iq}^{\lambda*} \epsilon_{iq}^{\lambda'} = \text{constant}$ [120]; the derivatives of $\delta \phi$ contract with derivatives of γ , so this produces $\vec{k} \cdot \vec{q}$ factors. Therefore we have

$$\langle \delta \phi_{\vec{k}_1}(\eta^*) \delta \phi_{\vec{k}_2}(\eta^*) \rangle_{V_1+V_2} \sim i \delta^{(3)}(\vec{k}_1 + \vec{k}_2) H_*^4 \int \frac{d^3 q}{q^3} f_1(q^2) \vec{k} \cdot \vec{q} \int_{-\infty}^{\eta^*} d\eta' f_2(\eta') + c.c. \quad (15.2.12)$$

where $f_1(q^2)$ and $f_2(\eta')$ are some functions of q^2 and η' . This contribution is evidently zero for symmetry reasons.

15.3. Complete expressions of one-loop two-vertex diagrams to leading order

In the following we provide the explicit expression for Eqs. (5.26) and (5.30). Eq. (5.26) reads as

$$\langle \delta \phi_{\vec{k}_1}(\eta^*) \delta \phi_{\vec{k}_2}(\eta^*) \rangle_{(1L,2v)}^A = \pi \delta^{(3)}(\vec{k}_1 + \vec{k}_2) \frac{H_*^4}{k^3} \left(a_1 \ln(k) + a_2 \ln(k\ell) + a_3 \right), \quad (15.3.1)$$

where

$$a_1 = -\frac{4}{15} \left(5 + 5x^{*2} + 2x^{*4} \right), \quad (15.3.2)$$

$$a_2 = \frac{8}{15x^{*2}} \left[-2 + \left(5 - 8\sigma_c \tilde{\sigma}_c + 4\pi\sigma_s - 8\sigma_s \tilde{\sigma}_s \right) x^{*2} - 8(\pi\sigma_c + 2\tilde{\sigma}_c \sigma_s - 2\sigma_c \tilde{\sigma}_c) x^{*3} + (1 + 8\sigma_c \tilde{\sigma}_c - 4\pi\sigma_s + 8\sigma_s \tilde{\sigma}_s) x^{*4} \right], \quad (15.3.3)$$

$$a_3 = \frac{1}{1800x^{*2}} \left[-64 - \left(-3120 + 15136\sigma_c \tilde{\sigma}_c - 450\pi^2 \sigma_c \tilde{\sigma}_c - 7568\pi\sigma_s + 225\pi^3 \sigma_s + 15136\sigma_s \tilde{\sigma}_s \right) x^{*2} - \left(15136\pi\sigma_c - 450\pi^3 \sigma_c + 30272\tilde{\sigma}_c \sigma_s - 900\pi^2 \tilde{\sigma}_c \sigma_s - 30272\sigma_c \tilde{\sigma}_s + 900\pi^2 \sigma_c \tilde{\sigma}_s \right) x^{*3} - \left(-672 - 15136\sigma_c \tilde{\sigma}_c + 450\pi^2 \sigma_c \tilde{\sigma}_c + 7568\pi\sigma_s - 225\pi^3 \sigma_s - 15136\sigma_s \tilde{\sigma}_c + 450\pi^2 \sigma_s \tilde{\sigma}_s \right) x^{*4} - 208x^{*6} \right] + \rho, \quad (15.3.4)$$

and ρ is a constant left over from renormalization of ultraviolet divergences. We have defined

$$\sigma_s \equiv \sin 2x^*,$$

$$\begin{aligned}\sigma_c &\equiv \cos 2x^*, \\ \tilde{\sigma}_s &\equiv \text{Si}(2x^*), \\ \tilde{\sigma}_c &\equiv \text{Ci}(2x^*),\end{aligned}$$

where Si and Ci stand for the sine-integral and the cosine-integral functions, i.e.

$$\begin{aligned}\text{Si}(x) &= \int_0^x \frac{\sin(t)}{t} dt, \\ \text{Ci}(x) &= \int_0^x \frac{\cos(t) - 1}{t} dt + \ln(x) + \gamma,\end{aligned}$$

with γ indicating the Euler Gamma function.

The expression for Eq. (5.30) is

$$\langle \delta\phi_{k_1}^-(\eta^*) \delta\phi_{k_2}^-(\eta^*) \rangle_{(1L,2\nu)}^C = \pi \delta^{(3)}(\vec{k}_1 + \vec{k}_2) \frac{H^4}{k^3} (c_1 + c_2 \ln(k\ell)), \quad (15.3.5)$$

where

$$c_1 = \frac{1}{225} \left(\frac{8}{x^{*2}} + 107 + 50x^{*2} \right), \quad (15.3.6)$$

$$c_2 = \frac{4}{15} \left(\frac{4}{x^{*2}} + 1 \right). \quad (15.3.7)$$

Finally the quantity f_3 appearing in Eq. (6.1) is given by

$$f_3 = a_3 + c_1 + \alpha', \quad (15.3.8)$$

where $\alpha' \equiv 2(1 + x^{*2}/3)\alpha$ from Eq. (5.13).

15.4. Background and first order perturbation equations for the gauge fields

The equations of motion for the gauge fields and for a Lagrangian as in Eq. (8.11) with $f = 1$ have been completely derived for the $U(1)$ case in [135]. We are going to carry out a similar calculation for the $SU(2)$ case

$$\begin{aligned}\frac{1}{\sqrt{-g}} \partial_\mu \left[\sqrt{-g} g^{\mu\alpha} g^{\nu\beta} \left(F_{\alpha\beta}^{(AB)a} + g_c \varepsilon^{abc} B_\alpha^b B_\beta^c \right) \right] + M^2 g^{\mu\nu} B_\mu^a \\ + g_c \varepsilon^{abc} g^{\gamma\nu} g^{\delta\beta} F_{\gamma\delta}^{(AB)b} B_\beta^c + g_c^2 \varepsilon^{abc} \varepsilon^{bb'c'} g^{\nu\alpha} g^{\delta\beta} B_\delta^c B_\alpha^{b'} B_\beta^{c'} = 0\end{aligned} \quad (15.4.1)$$

where $F_{\mu\nu}^{(AB)a} \equiv \partial_\mu B_\nu^a - \partial_\nu B_\mu^a$.

The $\nu = 0$ component of the equations of motion is

$$\begin{aligned}\partial_j \dot{B}_j^a - \partial_j \partial_j B_0^a + a^2 M^2 B_0^a + g_c \varepsilon^{abc} \left[- \left(\partial_j B_j^b \right) B_0^c - 2B_j^b \partial_j B_0^c - \dot{B}_j^b B_j^c \right. \\ \left. + g_c \varepsilon^{cb'c'} B_j^b B_0^{b'} B_j^{c'} \right] = 0\end{aligned} \quad (15.4.2)$$

where $B_\mu^a = B_\mu^a(\vec{x}, t)$.

Let us now move to the spatial ($\nu = i$) part of (15.4.1)

$$\begin{aligned}\ddot{B}_i^a + H \dot{B}_i^a - \frac{1}{a^2} \partial_j \partial_j B_i^a + M^2 B_i^a - \partial_i \dot{B}_0^a - H \partial_i B_0^a + \frac{1}{a^2} \partial_i \partial_j B_j^a \\ + g_c \varepsilon^{abc} \left[H B_0^b B_i^c + \dot{B}_0^b B_i^c + B_0^b \dot{B}_i^c \right] - g_c \frac{\varepsilon^{abc}}{a^2} \left[\left(\partial_j B_j^b \right) B_i^c + B_j^b \partial_j B_i^c \right]\end{aligned}$$

$$\begin{aligned}
& +g_c \varepsilon^{abc} [(\partial_i B_0^b) B_0^c - \dot{B}_i^b B_0^c] - g_c \frac{\varepsilon^{abc}}{a^2} [(\partial_i B_j^b) B_j^c - (\partial_j B_i^b) B_j^c] \\
& + g_c^2 \varepsilon^{abc} \varepsilon^{bb'c'} [B_0^c B_i^{b'} B_0^{c'}] - \frac{g_c^2}{a^2} \varepsilon^{abc} \varepsilon^{bb'c'} [B_j^c B_i^{b'} B_j^{c'}] = 0
\end{aligned} \tag{15.4.3}$$

If we contract Eq.(15.4.1) with ∂_ν , we get the integrability condition

$$\begin{aligned}
& (aM)^2 \dot{B}_0^a - M^2 \partial_i B_i^a + 3H (\partial_i \partial_i B_0^a - \partial_i \dot{B}_i^a) + g_c \varepsilon^{abc} [2H (\partial_i B_i^b B_0^c + B_i^b \partial_i B_0^c + \dot{B}_j^b B_j^c) - \partial_j B_0^b \dot{B}_j^c \\
& - \ddot{B}_j^b B_j^c + \partial_j \dot{B}_0^b B_j^c - \partial^2 B_0^b B_0^c - \partial_i \dot{B}_i^b B_0^c + \frac{1}{a^2} (B_i^b \partial^2 B_i^c + \partial_i B_j^b \partial_j B_i^c + B_j^b \partial_j \partial_i B_i^c + \partial^2 B_j^b B_j^c \\
& - \partial_i \partial_j B_i^b B_j^c)] + g_c^2 \varepsilon^{abc} \varepsilon^{bb'c'} [a^2 (\dot{B}_0^c B_0^{b'} B_0^{c'} + B_0^c \dot{B}_0^{b'} B_0^{c'} + B_0^c B_0^{b'} \dot{B}_0^{c'}) + 2H B_i^c B_0^{b'} B_i^{c'} \\
& - \dot{B}_i^c B_0^{b'} B_i^{c'} - B_i^c \dot{B}_0^{b'} B_i^{c'} - B_i^c B_0^{b'} \dot{B}_i^{c'} - \partial_i B_0^c B_i^{b'} B_0^{c'} - B_0^c \partial_i B_i^{b'} B_0^{c'} - B_0^c B_i^{b'} \partial_i B_0^{c'} \\
& + \frac{1}{a^2} (\partial_i B_j^c B_i^{b'} B_j^{c'} + B_j^c \partial_i B_i^{b'} B_j^{c'} + B_j^c B_i^{b'} \partial_i B_j^{c'})] = 0
\end{aligned} \tag{15.4.4}$$

which reduces to Eq.(7) of [135] in the Abelian case.

Combining Eq.(15.4.4) with Eq.(15.4.2) we get

$$\begin{aligned}
& (aM)^2 \dot{B}_0^a - M^2 \partial_i B_i^a + 3H (a^2 M^2 B_0^a + g_c \varepsilon^{abc} [- (\partial_j B_j^b) B_0^c - 2B_j^b \partial_j B_0^c - \dot{B}_j^b B_j^c + g_c \varepsilon^{cb'c'} B_j^b B_0^{b'} B_j^{c'}]) \\
& + g_c \varepsilon^{abc} [2H (\partial_i B_i^b B_0^c + B_i^b \partial_i B_0^c + \dot{B}_j^b B_j^c) - \partial_j B_0^b \dot{B}_j^c \\
& - \ddot{B}_j^b B_j^c + \partial_j \dot{B}_0^b B_j^c - \partial^2 B_0^b B_0^c - \partial_i \dot{B}_i^b B_0^c + \frac{1}{a^2} (B_i^b \partial^2 B_i^c + \partial_i B_j^b \partial_j B_i^c + B_j^b \partial_j \partial_i B_i^c + \partial^2 B_j^b B_j^c \\
& - \partial_i \partial_j B_i^b B_j^c)] + g_c^2 \varepsilon^{abc} \varepsilon^{bb'c'} [a^2 (\dot{B}_0^c B_0^{b'} B_0^{c'} + B_0^c \dot{B}_0^{b'} B_0^{c'} + B_0^c B_0^{b'} \dot{B}_0^{c'}) + 2H B_i^c B_0^{b'} B_i^{c'} \\
& - \dot{B}_i^c B_0^{b'} B_i^{c'} - B_i^c \dot{B}_0^{b'} B_i^{c'} - B_i^c B_0^{b'} \dot{B}_i^{c'} - \partial_i B_0^c B_i^{b'} B_0^{c'} - B_0^c \partial_i B_i^{b'} B_0^{c'} - B_0^c B_i^{b'} \partial_i B_0^{c'} \\
& + \frac{1}{a^2} (\partial_i B_j^c B_i^{b'} B_j^{c'} + B_j^c \partial_i B_i^{b'} B_j^{c'} + B_j^c B_i^{b'} \partial_i B_j^{c'})] = 0
\end{aligned} \tag{15.4.5}$$

Plugging this into Eq.(15.4.3) we get

$$\begin{aligned}
& \ddot{B}_n^a + H \dot{B}_n^a - \frac{1}{a^2} \partial_j \partial_j B_n^a + M^2 B_n^a + 2H \partial_n B_0^a \\
& - \frac{1}{(aM)^2} \partial_n [-3H (g_c \varepsilon^{abc} [- (\partial_j B_j^b) B_0^c - 2B_j^b \partial_j B_0^c - \dot{B}_j^b B_j^c + g_c \varepsilon^{cb'c'} B_j^b B_0^{b'} B_j^{c'}]) \\
& + g_c \varepsilon^{abc} [2H (\partial_i B_i^b B_0^c + B_i^b \partial_i B_0^c + \dot{B}_j^b B_j^c) - \partial_j B_0^b \dot{B}_j^c \\
& - \ddot{B}_j^b B_j^c + \partial_j \dot{B}_0^b B_j^c - \partial^2 B_0^b B_0^c - \partial_i \dot{B}_i^b B_0^c + \frac{1}{a^2} (B_i^b \partial^2 B_i^c + \partial_i B_j^b \partial_j B_i^c + B_j^b \partial_j \partial_i B_i^c + \partial^2 B_j^b B_j^c \\
& - \partial_i \partial_j B_i^b B_j^c)] + g_c^2 \varepsilon^{abc} \varepsilon^{bb'c'} [a^2 (\dot{B}_0^c B_0^{b'} B_0^{c'} + B_0^c \dot{B}_0^{b'} B_0^{c'} + B_0^c B_0^{b'} \dot{B}_0^{c'}) + 2H B_i^c B_0^{b'} B_i^{c'} \\
& - \dot{B}_i^c B_0^{b'} B_i^{c'} - B_i^c \dot{B}_0^{b'} B_i^{c'} - B_i^c B_0^{b'} \dot{B}_i^{c'} - \partial_i B_0^c B_i^{b'} B_0^{c'} - B_0^c \partial_i B_i^{b'} B_0^{c'} - B_0^c B_i^{b'} \partial_i B_0^{c'} \\
& + \frac{1}{a^2} (\partial_i B_j^c B_i^{b'} B_j^{c'} + B_j^c \partial_i B_i^{b'} B_j^{c'} + B_j^c B_i^{b'} \partial_i B_j^{c'})] \\
& + g_c \varepsilon^{abc} [H B_0^b B_n^c + \dot{B}_0^b B_n^c + B_0^b \dot{B}_n^c] - g_c \frac{\varepsilon^{abc}}{a^2} [(\partial_j B_j^b) B_n^c + B_j^b \partial_j B_n^c] \\
& + g_c \varepsilon^{abc} [(\partial_n B_0^b) B_0^c - \dot{B}_n^b B_0^c] - g_c \frac{\varepsilon^{abc}}{a^2} [(\partial_n B_j^b) B_j^c - (\partial_j B_n^b) B_j^c] \\
& + g_c^2 \varepsilon^{abc} \varepsilon^{bb'c'} [B_0^c B_n^{b'} B_0^{c'}] - \frac{g_c^2}{a^2} \varepsilon^{abc} \varepsilon^{bb'c'} [B_j^c B_n^{b'} B_j^{c'}] = 0.
\end{aligned} \tag{15.4.6}$$

Let us consider the background part of the vector fields, i.e. $\partial_i B_\mu^a = 0$. Then from Eq.(15.4.2)

$$a^2 M^2 B_0^a + g_c \varepsilon^{abc} \left[-\dot{B}_j^b B_j^c + g_c \varepsilon^{cb'c'} B_j^b B_0^{b'} B_j^{c'} \right] = 0. \quad (15.4.7)$$

Before proceeding with the derivation of the equations of motion for the background and the field perturbations, it is necessary to make some comments about Eqs. (15.4.6) and (15.4.7). One approximation that we have been using in the computation of the cosmological correlators is allowing the A_i^a fields to undergo slow-roll during inflation. One possible way to achieve this is by restricting the parameter space of the background gauge fields through the request that their temporal components should be much smaller than the spatial ones, $B_0^b \ll |B_i^b|/a(t)$, $b = 1, 2, 3$, and, in addition to that, assuming $B_0^b \simeq B_0^c$, $b, c = 1, 2, 3$. With these assumptions, the temporal component can be factored out in Eq. (15.4.7), using the approximation $\dot{B}_i^a \simeq H B_i^a$ (valid in a slow-roll regime). A solution to (15.4.7) is then given by $B_0 = 0$. Adopting this solution and plugging it in Eq (15.4.6), it is easy to show that a slow-roll equation of motion for the physical fields

$$\ddot{A}_i^a + 3H\dot{A}_i^a + m_0^2 A_i^a = 0 \quad (15.4.8)$$

follows from (15.4.6) if we set $M^2 = m_0^2 - 2H^2$ with $\dot{H} \ll m_0^2$ and if

$$\left(\frac{g_c A^1}{m_0} \right)^2 \ll \left| \frac{(A^1)^2}{(A^2)^2 + (A^3)^2 - (A^3)^2 \cos^2 \theta_{13} - (A^2)^2 \cos^2 \theta_{12}} \right|, \quad (15.4.9)$$

$$\left(\frac{g_c A^2}{m_0} \right)^2 \ll \left| \frac{(A^2)^2}{(A^1)^2 + (A^3)^2 - (A^3)^2 \cos^2 \theta_{23} - (A^1)^2 \cos^2 \theta_{12}} \right|, \quad (15.4.10)$$

$$\left(\frac{g_c A^3}{m_0} \right)^2 \ll \left| \frac{(A^3)^2}{(A^1)^2 + (A^2)^2 - (A^2)^2 \cos^2 \theta_{23} - (A^1)^2 \cos^2 \theta_{13}} \right|, \quad (15.4.11)$$

are satisfied. In the equations above, we defined $A^a \equiv |\vec{A}^a|$ and $\cos \theta_{ab} \equiv \hat{A}^a \cdot \hat{A}^b$, a and b running over the gauge indices. The quantities appearing on the right-hand sides of Eqs.(15.4.9) through (15.4.11) can be either large or small w.r.t. one, depending on the specific background configuration, i.e. on the moduli of the gauge fields and the angles θ_{ab} .

Suppose now the conditions described above are all met, then from Eq (15.4.6), in terms of the comoving fields, we have

$$\ddot{B}_i^a + H\dot{B}_i^a + M^2 B_i^a = 0. \quad (15.4.12)$$

Let us now derive the equations for the perturbations. Eq.(15.4.2) becomes

$$\begin{aligned} & \partial_j \delta \dot{B}_j^a - \partial^2 \delta B_0^a + a^2 M^2 \delta B_0^a + g_c \varepsilon^{abc} \left[-\partial_j \delta B_j^b B_0^c - 2B_j^b \partial_j \delta B_0^c - \delta \dot{B}_j^b B_j^c - \dot{B}_j^b \delta B_j^c \right. \\ & \left. + g_c \varepsilon^{cb'c'} \left(\delta B_j^b B_0^{b'} B_j^{c'} + B_j^b \delta B_0^{b'} B_j^{c'} + B_j^b B_0^{b'} \delta B_j^{c'} \right) \right] = 0 \end{aligned} \quad (15.4.13)$$

Eq.(15.4.1) for the field perturbations gives

$$\delta \ddot{B}_i^a + H \delta \dot{B}_i^a - \frac{1}{a^2} \partial_j \partial_j \delta B_i^a + M^2 \delta B_i^a + \frac{1}{a^2} \partial_i \partial_j \delta B_j^a - H \partial_i \delta B_0^a - H \partial_i \delta B_0^a$$

$$\begin{aligned}
& -\frac{g_c}{a^2}\varepsilon^{abc}\left[(\partial_j\delta B_j^b)B_i^c+B_j^b\partial_j\delta B_i^c\right]-\frac{g_c}{a^2}\varepsilon^{abc}\left[(\partial_i\delta B_j^b)B_j^c-(\partial_j\delta B_i^b)B_j^c\right] \\
& -\frac{g_c^2}{a^2}\varepsilon^{abc}\varepsilon^{bb'c'}\left[\delta B_j^cB_i^{b'}B_j^{c'}+B_j^c\delta B_i^{b'}B_j^{c'}+B_j^cB_i^{b'}\delta B_j^{c'}\right] \\
& +g_c\varepsilon^{abc}\left[H\left(B_0^b\delta B_i^c+\delta B_0^bB_i^c\right)+\delta\dot{B}_0^bB_i^c+\dot{B}_0^b\delta B_i^c+\delta B_0^b\dot{B}_i^c+B_0^b\delta\dot{B}_i^c\right] \\
& +g_c\varepsilon^{abc}\left[\partial_i\delta B_0^b\delta B_0^c-\delta\dot{B}_i^bB_0^c-\dot{B}_i^b\delta B_0^c\right] \\
& +g_c^2\varepsilon^{abc}\varepsilon^{bb'c'}\left[\delta B_0^cB_i^{b'}B_0^{c'}+B_0^c\delta B_i^{b'}B_0^{c'}+B_0^cB_i^{b'}\delta B_0^{c'}\right]=0
\end{aligned} \tag{15.4.14}$$

Finally from Eq.(15.4.6) we get

$$\delta\ddot{B}_i^a+H\delta\dot{B}_i^a-\frac{1}{a^2}\partial^2\delta B_i^a+M^2\delta B_i^a+2H\partial_i\delta B_0^a+(\sim g_c\text{terms})=0. \tag{15.4.15}$$

When calculating n-point functions for the gauge bosons, the eigenfunctions we need are provided by free-field solutions, i.e. by solutions of Eq.(15.4.15) with g_c being set to zero. This is exactly the Abelian limit, in fact in this case Eq.(15.4.15) corresponds to (18) of [135] and can be decomposed into a transverse and a longitudinal part

$$\left[\partial_0^2+H\partial_0+M^2+\left(\frac{k}{a}\right)^2\right]\delta\vec{B}^T=0 \tag{15.4.16}$$

$$\left[\partial_0^2+\left(1+\frac{2k^2}{k^2+(aM)^2}\right)H\partial_0+M^2+\left(\frac{k}{a}\right)^2\right]\delta\vec{B}^{\parallel}=0 \tag{15.4.17}$$

where the time derivatives are intended w.r.t. cosmic time.

15.5. Calculation of the number of e-foldings of single-(scalar)field driven inflation in the presence of a vector multiplet

Let us consider the Lagrangian in Eq.(8.11) with $f=1$ and $M^2=m_0^2-2H^2$. Let us assume that the $SU(2)$ gauge multiplet undergoes slow-roll as well as the scalar field but the latter provides the dominant part of the energy density of the universe. This last hypothesis is necessary in order to produce isotropic inflation (i.e. in order for the anisotropy in the expansion that the vector fields introduce to be negligible w.r.t. the isotropic contribution from the scalar field). The expression of the number of e-foldings is

$$N=N_{scalar}+N_{vector}=N_{scalar}+\frac{1}{4m_P^2}\sum_{a=1,2,3}\vec{A}^a\cdot\vec{A}^a. \tag{15.5.1}$$

The previous expression can be easily derived from the equations of motion of the system neglecting terms that are proportional to the $SU(2)$ coupling constant g_c and assuming slow-roll conditions for both the scalar the gauge fields.

The starting point is represented by Einstein equations

$$H^2=\frac{8\pi G}{3}(\rho_{scalar}+\rho_{vector}). \tag{15.5.2}$$

where we split the energy density into a scalar and a vector contribution. In slow-roll approximation, $\rho_{scalar}\sim V(\phi)$. Let us calculate ρ_{vector} . The energy momentum tensor

for the gauge bosons

$$T_{\mu\nu}^{vector} = 2\frac{\delta L}{\delta g^{\mu\nu}} - g_{\mu\nu}L \quad (15.5.3)$$

where, as a remainder, $L = -(1/4)g^{\mu\alpha}g^{\nu\beta}F_{\mu\nu}^a F_{\alpha\beta}^a + (M^2/2)g^{\mu\nu}B_\mu^a B_\nu^a$. So we get

$$\begin{aligned} T_{00}^{vector} &= \frac{\dot{B}_i^a \dot{B}_i^a}{2a^2} + \frac{m_0^2}{2a^2} B_i^a B_i^a + \frac{m_0^2}{2} B_0^a B_0^a - \frac{H}{a^2} \dot{B}_i^a B_i^a + \frac{H^2}{2a^2} B_i^a B_i^a \\ &+ \frac{g_c}{a^2} \varepsilon^{abc} \dot{B}_i^a B_0^b B_i^c + \frac{g_c^2}{2a^2} \varepsilon^{abc} \varepsilon^{ab'c'} B_0^b B_i^c B_0^{b'} B_i^{c'} \\ &+ \frac{g_c^2}{4a^4} \varepsilon^{abc} \varepsilon^{ab'c'} B_i^b B_j^c B_i^{b'} B_j^{c'} \end{aligned} \quad (15.5.4)$$

where sums are taken over all repeated indices. Let us write this in terms of the physical fields

$$\begin{aligned} T_{00}^{vector} &= \frac{\dot{A}_i^a \dot{A}_i^a}{2} + \frac{m_0^2}{2} (A_i^a A_i^a + A_0^a A_0^a) + g_c \varepsilon^{abc} (H A_i^a + \dot{A}_i^a) A_0^b A_i^c + \frac{g_c^2}{2} \varepsilon^{abc} \varepsilon^{ab'c'} A_0^b A_i^c A_0^{b'} A_i^{c'} \\ &+ \frac{g_c^2}{4} \varepsilon^{abc} \varepsilon^{ab'c'} A_i^b A_j^c A_i^{b'} A_j^{c'} \end{aligned} \quad (15.5.5)$$

If we neglect the non-Abelian contribution and we set $A_0^a = 0$, we are left with the Abelian result [123]

$$T_{00}^{vector} = \frac{\dot{A}_i^a \dot{A}_i^a}{2} + \frac{m_0^2}{2} A_i^a A_i^a \quad (15.5.6)$$

The equation of motion for the background vector multiplet \vec{A}^a can be derived from Eq.(15.4.12)

$$\ddot{A}_i^a + 3H\dot{A}_i^a + m_0^2 A_i^a = 0. \quad (15.5.7)$$

which is equal to the equation of a light scalar field of mass m_0 , if $m_0 \ll H$. If the conditions for accelerated expansions are met, Eq.(15.5.7) reduces to

$$3H\dot{A}_i^a + m_0^2 A_i^a \sim 0. \quad (15.5.8)$$

We are now ready to derive Eq.(15.5.1). Let us start from the definition of N and keep in mind Eq.(15.5.2), where we are assuming the existence of a scalar fields ϕ in de-Sitter with a separable potential governed by the usual (background) equation

$$\ddot{\phi} + 3H\dot{\phi} + V' = 0 \quad (15.5.9)$$

and slowly rolling down their potential. Then we have

$$\begin{aligned} N &= \int_{t^*}^t H dt' = \int_{t^*}^t H^2 \frac{dt'}{H} = 8\pi G \int_{t^*}^t \frac{V(\phi)}{3H} dt' + 8\pi G \int_{t^*}^t \frac{V(A)}{3H} dt' \\ &= 8\pi G \int_{t^*}^t \frac{V(\phi)}{3H} \frac{dt}{d\phi} d\phi + 8\pi G \sum_a \int_{t^*}^t \left(\frac{m_0^2}{2} \right) \frac{A_i^a A_i^a}{3H} \frac{dA^a dt'}{dA^a} \end{aligned} \quad (15.5.10)$$

where $A^a \equiv \vec{A}^a \cdot \vec{A}^a$. So

$$N = 8\pi G \int_{\phi(t^*)}^{\phi(t)} \frac{V(\phi)}{3H\dot{\phi}} d\phi + 8\pi G \sum_a \int_{A^a(t^*)}^{A^a(t)} \left(\frac{m_0^2}{4} \right) \frac{A_i^a A_i^a}{3H\dot{A}_j^a A_j^a} dA^a$$

$$\begin{aligned}
&= -\frac{1}{m_P^2} \int_{\phi(t^*)}^{\phi(t)} \frac{V}{V'} d\phi + \frac{1}{m_P^2} \sum_a \int_{A^a(t^*)}^{A^a(t)} \left(\frac{m_0^2}{4} \right) \frac{A_i^a A_i^a}{(-m_0^2) A_j^a A_j^a} dA^a \\
&= -\frac{1}{m_P^2} \int_{\phi(t^*)}^{\phi(t)} \frac{V}{V'} d\phi - \left(\frac{1}{4m_P^2} \right) \sum_a \int_{A^a(t^*)}^{A^a(t)} dA^a
\end{aligned} \tag{15.5.11}$$

after using the slow-roll conditions. Eq.(15.5.1) is thus recovered.

In the final expression for the bispectrum then we can substitute

$$N_a^i \equiv \frac{dN}{dA_i^a} = \left(\frac{1}{2m_P^2} \right) A_i^a \tag{15.5.12}$$

where the derivatives are as usual calculated at the initial time η^* .

The upper limits in integrals such as the ones in Eq.(15.5.11) depend on the chosen path in field space and so they also depend on the initial field configuration. It is important to notice though, as also stated in [148], that if the final time is chosen to be approaching (or later than) the end of inflation, the fields are supposed to have reached their equilibrium values and so N becomes independent of the field values at the final time t . Eq.(15.5.12) is thus recovered.

15.6. Complete expressions for the functions appearing in the bispectrum from quartic interactions

The anisotropy coefficients I_n in Eq. (9.49) are listed below

$$\begin{aligned}
I_{EEE} &\equiv \varepsilon^{aa'b'} \varepsilon^{ac'e} \left[6 \left(\vec{N}^{a'} \cdot \vec{N}^{c'} \right) \left(\vec{N}^{b'} \cdot \vec{A}^e \right) \right. \\
&\quad + \left(\vec{N}^{b'} \cdot \vec{A}^e \right) \left[\left(-2 \left(\hat{k}_3 \cdot \vec{N}^{a'} \right) \left(\hat{k}_3 \cdot \vec{N}^{c'} \right) - 2 \left(\hat{k}_1 \cdot \vec{N}^{a'} \right) \left(\hat{k}_1 \cdot \vec{N}^{c'} \right) \right. \right. \\
&\quad + \left. \left. \left(\hat{k}_1 \cdot \vec{N}^{a'} \right) \left(\hat{k}_3 \cdot \vec{N}^{c'} \right) \hat{k}_1 \cdot \hat{k}_3 + \left(\hat{k}_3 \cdot \vec{N}^{a'} \right) \left(\hat{k}_1 \cdot \vec{N}^{c'} \right) \hat{k}_1 \cdot \hat{k}_3 \right) + (1 \rightarrow 2) + (3 \rightarrow 2) \right] \\
&\quad - \left[\left(2 \left(\vec{N}^{a'} \cdot \vec{N}^{c'} \right) \left(\hat{k}_2 \cdot \vec{N}^{b'} \right) \left(\hat{k}_2 \cdot \vec{A}^e \right) \right) + (2 \rightarrow 1) + (2 \rightarrow 3) \right] \\
&\quad + \left[\left(\hat{k}_2 \cdot \vec{A}^e \left[2 \left(\hat{k}_3 \cdot \vec{N}^{a'} \right) \left(\hat{k}_2 \cdot \vec{N}^{b'} \right) \left(\hat{k}_3 \cdot \vec{N}^{c'} \right) + 2 \left(\hat{k}_1 \cdot \vec{N}^{a'} \right) \left(\hat{k}_2 \cdot \vec{N}^{b'} \right) \left(\hat{k}_1 \cdot \vec{N}^{c'} \right) \right. \right. \right. \\
&\quad \left. \left. - \left(\hat{k}_1 \cdot \vec{N}^{a'} \right) \left(\hat{k}_2 \cdot \vec{N}^{b'} \right) \left(\hat{k}_3 \cdot \vec{N}^{c'} \right) \hat{k}_1 \cdot \hat{k}_3 - \left(\hat{k}_1 \cdot \vec{N}^{a'} \right) \left(\hat{k}_2 \cdot \vec{N}^{b'} \right) \left(\hat{k}_3 \cdot \vec{N}^{c'} \right) \hat{k}_1 \cdot \hat{k}_3 \right] \right. \\
&\quad \left. + (2 \leftrightarrow 1) + (3 \leftrightarrow 2) \right] \tag{15.6.1}
\end{aligned}$$

$$\begin{aligned}
I_{lll} &\equiv \varepsilon^{aa'b'} \varepsilon^{ac'e} \left[\left(\left(\hat{k}_1 \cdot \vec{N}^{a'} \right) \left(\hat{k}_3 \cdot \vec{N}^{b'} \right) \left(\hat{k}_2 \cdot \vec{N}^{c'} \right) \left(\hat{k}_1 \cdot \hat{k}_2 \right) \left(\hat{k}_3 \cdot \vec{A}^e \right) \right. \right. \\
&\quad \left. \left. - \left(\hat{k}_3 \cdot \vec{N}^{a'} \right) \left(\hat{k}_2 \cdot \vec{N}^{b'} \right) \left(\hat{k}_1 \cdot \vec{N}^{c'} \right) \left(\hat{k}_1 \cdot \hat{k}_2 \right) \left(\hat{k}_3 \cdot \vec{A}^e \right) \right) + (1 \leftrightarrow 3) + (2 \leftrightarrow 3) \right] \tag{15.6.2}
\end{aligned}$$

$$\begin{aligned}
I_{llE} &\equiv \varepsilon^{aa'b'} \varepsilon^{ac'e} \left[\left(\vec{N}^{b'} \cdot \vec{A}^e \right) \left(\left(\hat{k}_1 \cdot \vec{N}^{a'} \right) \left(\hat{k}_2 \cdot \vec{N}^{c'} \right) + \left(\hat{k}_2 \cdot \vec{N}^{a'} \right) \left(\hat{k}_1 \cdot \vec{N}^{c'} \right) \right) \hat{k}_1 \cdot \hat{k}_2 \right. \\
&\quad + \left[\left(2 \left(\hat{k}_2 \cdot \vec{A}^e \right) \left(\hat{k}_1 \cdot \vec{N}^{a'} \right) \left(\hat{k}_2 \cdot \vec{N}^{b'} \right) \left(\hat{k}_1 \cdot \vec{N}^{c'} \right) \right) + (1 \leftrightarrow 2) \right] \\
&\quad - \left[\left(\left(\left(\hat{k}_1 \cdot \vec{N}^{a'} \right) \left(\hat{k}_2 \cdot \vec{N}^{c'} \right) + \left(\hat{k}_2 \cdot \vec{N}^{a'} \right) \left(\hat{k}_1 \cdot \vec{N}^{c'} \right) \right) \left(\hat{k}_3 \cdot \vec{N}^{b'} \right) \left(\hat{k}_3 \cdot \vec{A}^e \right) \hat{k}_1 \cdot \hat{k}_2 \right) \right. \\
&\quad \left. + (1 \leftrightarrow 3) + (2 \leftrightarrow 3) \right] \tag{15.6.3}
\end{aligned}$$

$$\begin{aligned}
I_{EEl} &\equiv \varepsilon^{aa'b'} \varepsilon^{ac'e} \left[4 \left(\vec{N}^{b'} \cdot \vec{A}^e \right) \left(\hat{k}_3 \cdot \vec{N}^{a'} \right) \left(\hat{k}_3 \cdot \vec{N}^{c'} \right) \right. \\
&\quad + \left[\left(\left(\hat{k}_2 \cdot \vec{N}^{b'} \right) \left(\hat{k}_2 \cdot \vec{A}^e \right) \left(\left(\hat{k}_1 \cdot \vec{N}^{a'} \right) \left(\hat{k}_3 \cdot \vec{N}^{c'} \right) + \left(\hat{k}_3 \cdot \vec{N}^{a'} \right) \left(\hat{k}_1 \cdot \vec{N}^{c'} \right) \right) \hat{k}_1 \cdot \hat{k}_3 \right) \right.
\end{aligned}$$

$$\begin{aligned}
& + (2 \leftrightarrow 1) + (2 \leftrightarrow 3)] \\
& - \left[\left(2 \left(\hat{k}_2 \cdot \vec{A}^e \right) \left(\hat{k}_2 \cdot \vec{N}^{a'} \right) \left(\hat{k}_3 \cdot \vec{N}^{b'} \right) \left(\hat{k}_2 \cdot \vec{N}^{c'} \right) \right) + (1 \leftrightarrow 2) + (2 \leftrightarrow 3) + (1 \leftrightarrow 3) \right] \\
& - \left[\left(\left(\vec{N}^{b'} \cdot \vec{A}^e \right) \hat{k}_1 \cdot \hat{k}_3 \left(\left(\hat{k}_1 \cdot \vec{N}^{a'} \right) \left(\hat{k}_3 \cdot \vec{N}^{c'} \right) + \left(\hat{k}_3 \cdot \vec{N}^{a'} \right) \left(\hat{k}_1 \cdot \vec{N}^{c'} \right) \right) \right) + (1 \leftrightarrow 2) \right] \\
& + \left[\left(\vec{N}^{a'} \cdot \vec{N}^{b'} \right) \left(\hat{k}_3 \cdot \vec{N}^{c'} \right) \left(\hat{k}_3 \cdot \vec{A}^e \right) + \left(\vec{N}^{c'} \cdot \vec{N}^{b'} \right) \left(\hat{k}_3 \cdot \vec{N}^{a'} \right) \left(\hat{k}_3 \cdot \vec{A}^e \right) \right] \quad (15.6.4)
\end{aligned}$$

where $i \rightarrow j$ means ‘replace \hat{k}_i with \hat{k}_j ’, whereas $i \leftrightarrow j$ means ‘exchange \hat{k}_i with \hat{k}_j ’.

The isotropic functions F_n in (9.49) are given by

$$F_{EEE} = -\frac{1}{24k^6 k_1^2 k_2^2 k_3^2 x^{*2}} [A_{EEE} + (B_{EEE} \cos x^* + C_{EEE} \sin x^*) E_i x^*], \quad (15.6.5)$$

$$F_{lll} = n^6(x^*) F_{EEE}, \quad (15.6.6)$$

$$F_{llE} = n^4(x^*) F_{EEE}, \quad (15.6.7)$$

$$F_{EEl} = n^2(x^*) F_{EEE}, \quad (15.6.8)$$

where

$$\begin{aligned}
A_{EEE} \equiv & kx^{*2} \left(-k^2(k_1^3 + k_2^3 + k_3^3 - 4k_1 k_2 k_3) - k^3(k_2 k_3 + k_1 k_2 + k_1 k_3) \right. \\
& \left. + k_1 k_2 k_3(k_1^2 + k_2^2 + k_3^2 - k_2 k_3 - k_1 k_2 - k_1 k_3)x^{*2} \right) \quad (15.6.9)
\end{aligned}$$

$$B_{EEE} \equiv (k_1^3 + k_2^3 + k_3^3)x^{*3} (-k^3 + k_1 k_2 k_3 x^{*2}) \quad (15.6.10)$$

$$C_{EEE} \equiv -k(k_1^3 + k_2^3 + k_3^3)x^{*2} (-k^2 + (k_2 k_3 + k_1 k_2 + k_1 k_3)x^{*2}) \quad (15.6.11)$$

In the previous equations we set $k \equiv k_1 + k_2 + k_3$.

15.7. More details on computing the analytic expressions of vector-exchange diagrams

We report the expressions of the functions $A, B \dots P$ introduced in Eq. (9.57)

$$A \equiv \left(-16k^2 + k_1 k_2 x^{*2} \right) \cos \left[\frac{(k_1 + k_2) x^*}{4k} \right] - 4k(k_1 + k_2) x^* \sin \left[\frac{(k_1 + k_2) x^*}{4k} \right], \quad (15.7.1)$$

$$B \equiv A [k_1 \rightarrow k_3, k_2 \rightarrow k_4], \quad (15.7.2)$$

$$C \equiv 4k(k_1 + k_2) x^* \cos \left[\frac{(k_1 + k_2) x^*}{4k} \right] + \left(-16k^2 + k_1 k_2 x^{*2} \right) \sin \left[\frac{(k_1 + k_2) x^*}{4k} \right], \quad (15.7.3)$$

$$D \equiv C [k_1 \rightarrow k_3, k_2 \rightarrow k_4], \quad (15.7.4)$$

$$\begin{aligned}
E \equiv & \left(8k^2(k_{\hat{1}2} + k_3 + k_4) - k_{\hat{1}2} k_3 k_4 x^{*2} \right) \cos \left[\frac{(k_{\hat{1}2} + k_3 + k_4) x^*}{4k} \right] \\
& + 2k(k_{\hat{1}2} + k_3 + k_4)^2 x^* \sin \left[\frac{(k_{\hat{1}2} + k_3 + k_4) x^*}{4k} \right], \quad (15.7.5)
\end{aligned}$$

$$F \equiv E [k_3 \rightarrow k_1, k_4 \rightarrow k_2], \quad (15.7.6)$$

$$\begin{aligned}
G \equiv & 2k(k_{\hat{1}2} + k_1 + k_2)^2 x^* \cos \left[\frac{(k_{\hat{1}2} + k_1 + k_2) x^*}{4k} \right] \\
& + \left(-8k^2(k_{\hat{1}2} + k_1 + k_2) - k_{\hat{1}2} k_1 k_2 x^{*2} \right) \sin \left[\frac{(k_{\hat{1}2} + k_1 + k_2) x^*}{4k} \right], \quad (15.7.7)
\end{aligned}$$

$$L \equiv \left(k_1^3 + k_{12}^3 + k_{12}^2 k_2 + k_{12} k_2^2 + k_2^3 (k_{12} + k_2) + k_1 (k_{12}^2 + k_2^2) \right) x^{*2} si \left[\frac{(k_{12} + k_1 + k_2) x^*}{4k} \right],$$

$$M \equiv \left(k_1^3 + k_{12}^3 + k_{12}^2 k_2 + k_{12} k_2^2 + k_2^3 (k_{12} + k_2) + k_1 (k_{12}^2 + k_2^2) \right) x^{*2} ci \left[\frac{(k_{12} + k_1 + k_2) x^*}{4k} \right],$$

$$N \equiv M [k_3 \rightarrow k_1, k_4 \rightarrow k_2], \quad (15.7.8)$$

$$P \equiv L [k_1 \rightarrow k_3, k_2 \rightarrow k_4]. \quad (15.7.9)$$

The anisotropy coefficients introduced in Eq. (9.66) have the following expressions

$$t_1 \equiv k_1 k_3 (\hat{k}_1 \cdot \hat{k}_{12}) (\hat{k}_1 \cdot \hat{k}_2) (\hat{k}_3 \cdot \hat{k}_4) (\hat{k}_3 \cdot \hat{k}_{12}) \quad (15.7.10)$$

$$t_2 \equiv k_1 k_4 (\hat{k}_1 \cdot \hat{k}_{12}) (\hat{k}_1 \cdot \hat{k}_2) (\hat{k}_3 \cdot \hat{k}_4) (\hat{k}_4 \cdot \hat{k}_{12}), \quad (15.7.11)$$

$$t_3 \equiv k_2 k_3 (\hat{k}_2 \cdot \hat{k}_{12}) (\hat{k}_1 \cdot \hat{k}_2) (\hat{k}_3 \cdot \hat{k}_4) (\hat{k}_3 \cdot \hat{k}_{12}), \quad (15.7.12)$$

$$t_4 \equiv k_2 k_4 (\hat{k}_2 \cdot \hat{k}_{12}) (\hat{k}_1 \cdot \hat{k}_2) (\hat{k}_3 \cdot \hat{k}_4) (\hat{k}_4 \cdot \hat{k}_{12}), \quad (15.7.13)$$

$$t_5 \equiv k_1 k_2 (\hat{k}_1 \cdot \hat{k}_{13}) (\hat{k}_1 \cdot \hat{k}_3) (\hat{k}_2 \cdot \hat{k}_4) (\hat{k}_2 \cdot \hat{k}_{13}), \quad (15.7.14)$$

$$t_6 \equiv k_1 k_4 (\hat{k}_1 \cdot \hat{k}_{13}) (\hat{k}_1 \cdot \hat{k}_3) (\hat{k}_2 \cdot \hat{k}_4) (\hat{k}_4 \cdot \hat{k}_{13}), \quad (15.7.15)$$

$$t_7 \equiv k_2 k_3 (\hat{k}_3 \cdot \hat{k}_{13}) (\hat{k}_1 \cdot \hat{k}_3) (\hat{k}_2 \cdot \hat{k}_4) (\hat{k}_2 \cdot \hat{k}_{13}), \quad (15.7.16)$$

$$t_8 \equiv k_3 k_4 (\hat{k}_3 \cdot \hat{k}_{13}) (\hat{k}_1 \cdot \hat{k}_3) (\hat{k}_2 \cdot \hat{k}_4) (\hat{k}_4 \cdot \hat{k}_{13}), \quad (15.7.17)$$

$$t_9 \equiv k_1 k_2 (\hat{k}_1 \cdot \hat{k}_{14}) (\hat{k}_1 \cdot \hat{k}_4) (\hat{k}_2 \cdot \hat{k}_3) (\hat{k}_2 \cdot \hat{k}_{14}), \quad (15.7.18)$$

$$t_{10} \equiv k_1 k_3 (\hat{k}_1 \cdot \hat{k}_{14}) (\hat{k}_1 \cdot \hat{k}_4) (\hat{k}_2 \cdot \hat{k}_3) (\hat{k}_3 \cdot \hat{k}_{14}), \quad (15.7.19)$$

$$t_{11} \equiv k_1 k_3 (\hat{k}_1 \cdot \hat{k}_{12}) (\hat{k}_1 \cdot \hat{k}_2) (\hat{k}_3 \cdot \hat{k}_4) (\hat{k}_3 \cdot \hat{k}_{12}), \quad (15.7.20)$$

$$t_{12} \equiv k_1 k_3 (\hat{k}_1 \cdot \hat{k}_{12}) (\hat{k}_1 \cdot \hat{k}_2) (\hat{k}_3 \cdot \hat{k}_4) (\hat{k}_3 \cdot \hat{k}_{12}). \quad (15.7.21)$$

Let us now list all the scalar products appearing in the equations above

$\hat{k}_1 \cdot \hat{k}_{12} = \frac{k_{12}^2 + k_1^2 + k_2^2}{2k_1 k_{12}}$	$\hat{k}_1 \cdot \hat{k}_{13} = \frac{2k_1^2 + k_4^2 - k_{12}^2 - k_{14}^2}{2k_1 k_{13}}$	$\hat{k}_1 \cdot \hat{k}_{14} = \frac{k_{14}^2 + k_1^2 - k_4^2}{2k_1 k_{14}}$
$\hat{k}_2 \cdot \hat{k}_{12} = \frac{k_{12}^2 + k_1^2 - k_2^2}{2k_2 k_{12}}$	$\hat{k}_3 \cdot \hat{k}_{13} = \frac{2k_3^2 + k_4^2 - k_{12}^2 - k_{14}^2 + k_2^2}{2k_3 k_{13}}$	$\hat{k}_4 \cdot \hat{k}_{14} = \frac{k_{14}^2 - k_1^2 + k_4^2}{2k_4 k_{14}}$
$\hat{k}_3 \cdot \hat{k}_{12} = \frac{k_{12}^2 + k_{14}^2 - 2k_2^2 - k_1^2 - k_3^2}{2k_2 k_{13}}$	$\hat{k}_2 \cdot \hat{k}_{13} = \frac{k_{12}^2 + k_{14}^2 - k_1^2 - 2k_2^2 - k_3^2}{2k_1 k_3}$	$\hat{k}_2 \cdot \hat{k}_4 = \frac{k_1^2 + k_3^2 - k_{12}^2 - k_{14}^2}{2k_2 k_4}$
$\hat{k}_4 \cdot \hat{k}_{12} = \frac{k_{14}^2 + k_{12}^2 - k_1^2 - k_3^2 - 2k_4^2}{2k_4 k_{13}}$	$\hat{k}_4 \cdot \hat{k}_{13} = \frac{k_{14}^2 + k_{12}^2 - k_1^2 - k_3^2 - 2k_4^2}{2k_4 k_{13}}$	$\hat{k}_2 \cdot \hat{k}_{14} = \frac{k_3^2 - k_{14}^2 - k_2^2}{2k_2 k_{14}}$
$\hat{k}_1 \cdot \hat{k}_2 = \frac{k_1^2 + k_2^2 - k_{12}^2}{2k_1 k_2}$	$\hat{k}_1 \cdot \hat{k}_3 = \frac{k_4^2 + k_{12}^2 - k_{12}^2 - k_{14}^2}{2k_1 k_3}$	$\hat{k}_1 \cdot \hat{k}_4 = \frac{k_{14}^2 - k_1^2 - k_4^2}{2k_1 k_4}$
$\hat{k}_3 \cdot \hat{k}_4 = \frac{k_{12}^2 - k_4^2 - k_3^2}{2k_4 k_3}$	$\hat{k}_3 \cdot \hat{k}_{14} = \frac{k_2^2 - k_3^2 - k_{14}^2}{2k_3 k_{14}}$	$\hat{k}_2 \cdot \hat{k}_3 = \frac{k_{14}^2 - k_2^2 - k_3^2}{2k_2 k_3}$

where $k_i \equiv |\vec{k}_i|$, $\hat{k}_i \equiv \vec{k}_i/k_i$, $\vec{k}_{ij} \equiv \vec{k}_i + \vec{k}_j$, $k_{ij} \equiv |\vec{k}_i + \vec{k}_j|$, i and j running over the four external wave vectors.

15.8. Complete expressions for functions appearing in point-interaction diagrams

We provide here the expressions for the coefficients appearing in Eq. (9.89)

$$Q_{EEEE} \equiv x^{*3} [-k(k^3 k_1^4 + k^3 k_2^4 - k_3^3 + k^3 k_3^4 + k^5 k_3 k_4 - 3k^3 k_2^2 k_3 k_4 + k^3 k_3^3 k_4$$

$$\begin{aligned}
& -k_4^3 + k^3 k_3 k_4^3 + k^3 k_4^4 + k^3 k_2 (k_3^3 - k k_3 k_4 - 3k_3^2 k_4 - 3k_3 k_4^2 + k_4^3 \\
& + k^2 (k_3 + k_4)) + k_2^3 (-1 + k^3 (k_3 + k_4)) - 3k^3 k_1^2 (k_3 k_4 + k_2 (k_3 + k_4)) \\
& + k_1^3 (-1 + k^3 (k_2 + k_3 + k_4)) + k^3 k_1 (k_2^3 + k_3^3 - 3k_3^2 k_4 - 3k_3 k_4^2 + k_4^3 \\
& - 3k_2^2 (k_3 + k_4) + k^2 (k_2 + k_3 + k_4) - 3k_2 (k_3^2 + 3k_3 k_4 + k_4^2) - k (k_3 k_4 \\
& + k_2 (k_3 + k_4))) + x^{*5} [k^2 (k_1^3 k_2 k_3 k_4 + k_2 k_3 k_4 (k_2^3 + k_3^3 - 3k_2 k_3 k_4 + k_4^3) \\
& + k_1^4 (k_3 k_4 + k_2 (k_3 + k_4)) - 3k_1^2 (k_3^2 k_4^2 + k_2 k_3 k_4 (k_3 + k_4) + k_2^2 (k_3^2 + k_3 k_4 \\
& + k_4^2)) + k_1 (k_2^3 k_3 k_4 + k_2^4 (k_3 + k_4) - 3k_2^2 k_3 k_4 (k_3 + k_4) + k_3 k_4 (k_3^3 + k_4^3) \\
& + k_2 (k_3^4 + k_3^3 k_4 - 3k_3^2 k_4^2 + k_4^4 + k_3 k_4 (k^2 + k_4^2)))] - 3x^{*7} k_1^2 k_2^2 k_3^2 k_4^2, \tag{15.8.1}
\end{aligned}$$

$$A_{EEEE} \equiv k (k_1^3 + k_2^3 + k_3^3 + k_4^3) x^{*3}, \tag{15.8.2}$$

$$B_{EEEE} \equiv k^4 - k^2 (k_3 k_4 + k_2 (k_3 + k_4 + k_1 (k_2 + k_3 + k_4) x^{*2} + k_1 k_2 k_3 k_4 x^{*4})), \tag{15.8.3}$$

$$C_{EEEE} \equiv k x^* (k^3 - (k_2 k_3 k_4 + k_1 (k_3 k_4 + k_2 (k_3 + k_4))) x^{*2}), \tag{15.8.4}$$

$$D_{EEEE} \equiv -k x^* (k^3 - (k_2 k_3 k_4 + k_1 (k_3 k_4 + k_2 (k_3 k_4))) x^{*2}), \tag{15.8.5}$$

$$E_{EEEE} \equiv k^4 - k^2 (k_3 k_4 + k_2 (k_3 + k_4) + k_1 (k_2 + k_3 + k_4)) x^{*2} + k_1 k_2 k_3 k_4 x^{*4}, \tag{15.8.6}$$

$$F_{EEEE} \equiv k (k_1^3 + k_2^3 + k_3^3 + k_4^3) x^{*3}. \tag{15.8.7}$$

16. References

- [1] For a review, see D. H. Lyth and A. Riotto, *Phys. Rept.* **314**, 1 (1999).
- [2] G. F. Smoot *et al.*, *Astrophys. J.* **396**, L1 (1992)
- [3] C. L. Bennett *et al.*, *Astrophys. J.* **464**, L1 (1996).
- [4] K. M. Gorski, A. J. Banday, C. L. Bennett, G. Hinshaw, A. Kogut, G. F. Smoot and E. L. Wright, *Astrophys. J.* **464** (1996) L11.
- [5] D. N. Spergel *et al.* [WMAP Collaboration], *Astrophys. J. Suppl.* **170**, 377 (2007) [arXiv:astro-ph/0603449].
- [6] P. Creminelli, A. Nicolis, L. Senatore, M. Tegmark and M. Zaldarriaga, *JCAP* **0605**, 004 (2006) [arXiv:astro-ph/0509029].
- [7] E. Komatsu *et al.* [WMAP Collaboration], arXiv:0803.0547 [astro-ph].
- [8] K. M. Smith, L. Senatore and M. Zaldarriaga, *JCAP* **0909**, 006 (2009) [arXiv:0901.2572 [astro-ph]].
- [9] L. Senatore, K. M. Smith and M. Zaldarriaga, arXiv:0905.3746 [astro-ph.CO].
- [10] N. Bartolo, E. Komatsu, S. Matarrese and A. Riotto, *Phys. Rept.* **402**, 103 (2004).
- [11] E. Komatsu *et al.*, arXiv:0902.4759 [astro-ph.CO].
- [12] V. Acquaviva, N. Bartolo, S. Matarrese and A. Riotto, *Nucl. Phys. B* **667**, 119 (2003) [arXiv:astro-ph/0209156].
- [13] J. M. Maldacena, *JHEP* **0305**, 013 (2003) [arXiv:astro-ph/0210603].
- [14] D. Seery, J. E. Lidsey and M. S. Sloth, *JCAP* **0701**, 027 (2007) [arXiv:astro-ph/0610210].
- [15] A. D. Linde, *Phys. Lett. B* **158**, 375 (1985).
- [16] L. A. Kofman, *Phys. Lett. B* **173**, 400 (1986).
- [17] D. Polarski and A. A. Starobinsky, *Phys. Rev. D* **50**, 6123 (1994) [arXiv:astro-ph/9404061].
- [18] J. Garcia-Bellido and D. Wands, *Phys. Rev. D* **53**, 5437 (1996) [arXiv:astro-ph/9511029].
- [19] V. F. Mukhanov and P. J. Steinhardt, *Phys. Lett. B* **422**, 52 (1998) [arXiv:astro-ph/9710038].
- [20] D. Langlois, *Phys. Rev. D* **59**, 123512 (1999) [arXiv:astro-ph/9906080].
- [21] C. Gordon, D. Wands, B. A. Bassett and R. Maartens, *Phys. Rev. D* **63**, 023506 (2001) [arXiv:astro-ph/0009131].
- [22] S. Mollerach, *Phys. Rev. D* **42**, 313 (1990).
- [23] K. Enqvist and M. S. Sloth, *Nucl. Phys. B* **626**, 395 (2002) [arXiv:hep-ph/0109214].
- [24] D. H. Lyth and D. Wands, *Phys. Lett. B* **524**, 5 (2002) [arXiv:hep-ph/0110002].
- [25] D. H. Lyth, C. Ungarelli and D. Wands, *Phys. Rev. D* **67**, 023503 (2003) [arXiv:astro-ph/0208055].
- [26] T. Moroi and T. Takahashi, *Phys. Lett. B* **522**, 215 (2001) [Erratum-ibid. *B* **539**, 303 (2002)] [arXiv:hep-ph/0110096].
- [27] N. Bartolo, S. Matarrese and A. Riotto, *Phys. Rev. D* **69**, 043503 (2004) [arXiv:hep-ph/0309033].
- [28] J. Garriga and V. F. Mukhanov, *Phys. Lett. B* **458**, 219 (1999) [arXiv:hep-th/9904176].
- [29] C. Armendariz-Picon, T. Damour and V. F. Mukhanov, *Phys. Lett. B* **458**, 209 (1999) [arXiv:hep-th/9904075].
- [30] M. Alishahiha, E. Silverstein and D. Tong, *Phys. Rev. D* **70** (2004) 123505 [arXiv:hep-th/0404084].
- [31] X. Chen, M. x. Huang, S. Kachru and G. Shiu, *JCAP* **0701**, 002 (2007) [arXiv:hep-th/0605045].
- [32] N. Arkani-Hamed, P. Creminelli, S. Mukohyama and M. Zaldarriaga, *JCAP* **0404**, 001 (2004) [arXiv:hep-th/0312100].
- [33] N. Bartolo, S. Matarrese and A. Riotto, *Phys. Rev. D* **65**, 103505 (2002) [arXiv:hep-ph/0112261].
- [34] N. Bartolo, S. Matarrese and A. Riotto, *JCAP* **0508**, 010 (2005) [arXiv:astro-ph/0506410].
- [35] D. Seery and J. E. Lidsey, *JCAP* **0701**, 008 (2007) [arXiv:astro-ph/0611034].
- [36] F. Vernizzi and D. Wands, *JCAP* **0605**, 019 (2006) [arXiv:astro-ph/0603799].
- [37] X. Chen and Y. Wang, arXiv:0911.3380 [hep-th].
- [38] C. T. Byrnes, M. Sasaki and D. Wands, *Phys. Rev. D* **74**, 123519 (2006) [arXiv:astro-ph/0611075].
- [39] D. Seery, M. S. Sloth and F. Vernizzi, *JCAP* **0903**, 018 (2009) [arXiv:0811.3934 [astro-ph]].
- [40] M. Sasaki, J. Valiviita and D. Wands, *Phys. Rev. D* **74**, 103003 (2006) [arXiv:astro-ph/0607627].
- [41] X. Chen, M. x. Huang and G. Shiu, *Phys. Rev. D* **74**, 121301 (2006) [arXiv:hep-th/0610235].

- [42] F. Arroja and K. Koyama, Phys. Rev. D **77**, 083517 (2008) [arXiv:0802.1167 [hep-th]].
- [43] F. Arroja, S. Mizuno, K. Koyama and T. Tanaka, Phys. Rev. D **80**, 043527 (2009) [arXiv:0905.3641 [hep-th]].
- [44] X. Chen, B. Hu, M. x. Huang, G. Shiu and Y. Wang, JCAP **0908**, 008 (2009) [arXiv:0905.3494 [astro-ph.CO]].
- [45] X. Gao and B. Hu, JCAP **0908**, 012 (2009) [arXiv:0903.1920 [astro-ph.CO]].
- [46] S. Mizuno, F. Arroja and K. Koyama, arXiv:0907.2439 [hep-th].
- [47] T. Okamoto and W. Hu, Phys. Rev. D **66**, 063008 (2002) [arXiv:astro-ph/0206155].
- [48] N. Kogo and E. Komatsu, Phys. Rev. D **73**, 083007 (2006) [arXiv:astro-ph/0602099].
- [49] J. S. Schwinger, J. Math. Phys. **2**, 407 (1961).
- [50] E. Calzetta and B. L. Hu, Phys. Rev. D **35**, 495 (1987).
- [51] R. D. Jordan, Phys. Rev. D **33**, 444 (1986).
- [52] S. Weinberg, Phys. Rev. D **72**, 043514 (2005) [arXiv:hep-th/0506236].
- [53] S. Weinberg, Phys. Rev. D **74**, 023508 (2006) [arXiv:hep-th/0605244].
- [54] D. Seery, JCAP **0711**, 025 (2007) [arXiv:0707.3377 [astro-ph]].
- [55] D. Seery, JCAP **0802**, 006 (2008) [arXiv:0707.3378 [astro-ph]].
- [56] E. Dimastrogiovanni and N. Bartolo, JCAP **0811**, 016 (2008) [arXiv:0807.2790 [astro-ph]].
- [57] P. Vielva and J. L. Sanz, arXiv:0910.3196 [astro-ph.CO].
- [58] V. Desjacques and U. Seljak, arXiv:0907.2257 [astro-ph.CO].
- [59] See <http://planck.esa.int/>.
- [60] A. de Oliveira-Costa, M. Tegmark, M. Zaldarriaga and A. Hamilton, Phys. Rev. D **69**, 063516 (2004) [arXiv:astro-ph/0307282].
- [61] P. Vielva, E. Martinez-Gonzalez, R. B. Barreiro, J. L. Sanz and L. Cayon, Astrophys. J. **609**, 22 (2004) [arXiv:astro-ph/0310273].
- [62] H. K. Eriksen, F. K. Hansen, A. J. Banday, K. M. Gorski and P. B. Lilje, Astrophys. J. **605**, 14 (2004) [Erratum-ibid. **609**, 1198 (2004)] [arXiv:astro-ph/0307507].
- [63] C. L. Bennett *et al.*, Astrophys. J. **464**, L1 (1996) [arXiv:astro-ph/9601067].
- [64] D. N. Spergel *et al.* [WMAP Collaboration], Astrophys. J. Suppl. **148**, 175 (2003) [arXiv:astro-ph/0302209].
- [65] G. Efstathiou, Mon. Not. Roy. Astron. Soc. **348**, 885 (2004) [arXiv:astro-ph/0310207].
- [66] K. Land and J. Magueijo, Phys. Rev. Lett. **95**, 071301 (2005) [arXiv:astro-ph/0502237].
- [67] M. Cruz, L. Cayon, E. Martinez-Gonzalez, P. Vielva and J. Jin, Astrophys. J. **655**, 11 (2007) [arXiv:astro-ph/0603859].
- [68] F. K. Hansen, P. Cabella, D. Marinucci and N. Vittorio, Astrophys. J. **607**, L67 (2004) [arXiv:astro-ph/0402396].
- [69] F. K. Hansen, A. J. Banday and K. M. Gorski, Mon. Not. Roy. Astron. Soc. **354**, 641 (2004) [arXiv:astro-ph/0404206].
- [70] N. E. Groeneboom, L. Ackerman, I. K. Wehus and H. K. Eriksen, arXiv:0911.0150 [astro-ph.CO].
- [71] D. Hanson and A. Lewis, Phys. Rev. D **80**, 063004 (2009) [arXiv:0908.0963 [astro-ph.CO]].
- [72] R. M. Wald, Phys. Rev. D **28**, 2118 (1983).
- [73] J. D. Barrow and S. Hervik, Phys. Rev. D **73**, 023007 (2006) [arXiv:gr-qc/0511127].
- [74] J. D. Barrow and S. Hervik, Phys. Rev. D **74**, 124017 (2006) [arXiv:gr-qc/0610013].
- [75] E. Di Grezia, G. Esposito, A. Funel, G. Mangano and G. Miele, Phys. Rev. D **68**, 105012 (2003) [arXiv:gr-qc/0305050].
- [76] T. S. Pereira, C. Pitrou and J. P. Uzan, JCAP **0709**, 006 (2007) [arXiv:0707.0736 [astro-ph]].
- [77] C. Pitrou, T. S. Pereira and J. P. Uzan, JCAP **0804**, 004 (2008) [arXiv:0801.3596 [astro-ph]].
- [78] E. Dimastrogiovanni, W. Fischler and S. Paban, JHEP **0807**, 045 (2008) [arXiv:0803.2490 [hep-th]].
- [79] A. E. Gumrukcuoglu, L. Kofman and M. Peloso, Phys. Rev. D **78**, 103525 (2008) [arXiv:0807.1335 [astro-ph]].
- [80] C. Armendariz-Picon and P. B. Greene, Gen. Rel. Grav. **35**, 1637 (2003) [arXiv:hep-th/0301129].

- [81] C. G. Boehmer and D. F. Mota, Phys. Lett. B **663**, 168 (2008) [arXiv:0710.2003 [astro-ph]].
- [82] T. Watanabe, arXiv:0902.1392 [astro-ph.CO].
- [83] G. de Berredo-Peixoto and E. A. de Freitas, Class. Quant. Grav. **26**, 175015 (2009) [arXiv:0902.4025 [gr-qc]].
- [84] C. Germani and A. Kehagias, JCAP **0903**, 028 (2009) [arXiv:0902.3667 [astro-ph.CO]].
- [85] T. Kobayashi and S. Yokoyama, JCAP **0905**, 004 (2009) [arXiv:0903.2769 [astro-ph.CO]].
- [86] T. S. Koivisto and N. J. Nunes, arXiv:0907.3883 [astro-ph.CO].
- [87] C. Germani and A. Kehagias, JCAP **0911**, 005 (2009) [arXiv:0908.0001 [astro-ph.CO]].
- [88] T. S. Koivisto and N. J. Nunes, Phys. Rev. D **80**, 103509 (2009) [arXiv:0908.0920 [astro-ph.CO]].
- [89] T. S. Koivisto, D. F. Mota and C. Pitrou, JHEP **0909**, 092 (2009) [arXiv:0903.4158 [astro-ph.CO]].
- [90] S. Yokoyama and J. Soda, JCAP **0808**, 005 (2008) [arXiv:0805.4265 [astro-ph]].
- [91] C. A. Valenzuela-Toledo, Y. Rodriguez and D. H. Lyth, Phys. Rev. D **80**, 103519 (2009) [arXiv:0909.4064 [astro-ph.CO]].
- [92] K. Dimopoulos, M. Karciauskas, D. H. Lyth and Y. Rodriguez, JCAP **0905**, 013 (2009) [arXiv:0809.1055 [astro-ph]].
- [93] C. A. Valenzuela-Toledo and Y. Rodriguez, arXiv:0910.4208 [astro-ph.CO].
- [94] M. Karciauskas, K. Dimopoulos and D. H. Lyth, Phys. Rev. D **80**, 023509 (2009) [arXiv:0812.0264 [astro-ph]].
- [95] K. Dimopoulos, M. Karciauskas and J. M. Wagstaff, arXiv:0907.1838 [hep-ph].
- [96] K. Dimopoulos, M. Karciauskas and J. M. Wagstaff, arXiv:0909.0475 [hep-ph].
- [97] N. Bartolo, E. Dimastrogiovanni, S. Matarrese and A. Riotto, JCAP **0910**, 015 (2009) [arXiv:0906.4944 [astro-ph.CO]].
- [98] N. Bartolo, E. Dimastrogiovanni, S. Matarrese and A. Riotto, JCAP **0911**, 028 (2009) [arXiv:0909.5621 [astro-ph.CO]].
- [99] V. F. Mukhanov, L. R. W. Abramo, and R. H. Brandenberger, Phys. Rev. Lett. **78**, 1624 (1997).
- [100] L. R. W. Abramo, R. H. Brandenberger, and V. F. Mukhanov, Phys. Rev. **D56**, 3248 (1997).
- [101] D. Boyanovsky, H. J. de Vega and N. G. Sanchez, Phys. Rev. D **71**, 023509 (2005).
- [102] D. Boyanovsky, H. J. de Vega and N. G. Sanchez, Nucl. Phys. B **747**, 25 (2006).
- [103] M. van der Meulen and J. Smit, JCAP **0711**, 023 (2007) [arXiv:0707.0842 [hep-th]].
- [104] S. Weinberg, Phys. Rev. D **72**, 043514 (2005) [arXiv:hep-th/0506236].
- [105] S. Weinberg, Phys. Rev. D **74**, 023508 (2006) [arXiv:hep-th/0605244].
- [106] M. S. Sloth, Nucl. Phys. B **748**, 149 (2006).
- [107] M. S. Sloth, Nucl. Phys. B **775**, 78 (2007).
- [108] D. Seery, JCAP **0711** (2007) 025 [arXiv:0707.3377 [astro-ph]].
- [109] D. Seery, JCAP **0802** (2008) 006 [arXiv:0707.3378 [astro-ph]].
- [110] D. H. Lyth, JCAP **0712**, 016 (2007) [arXiv:0707.0361 [astro-ph]].
- [111] N. Bartolo, S. Matarrese, M. Pietroni, A. Riotto and D. Seery, JCAP **0801** (2008) 015 [arXiv:0711.4263 [astro-ph]].
- [112] K. Enqvist, S. Nurmi, D. Podolsky and G. I. Rigopoulos, JCAP **0804**, 025 (2008) [arXiv:0802.0395 [astro-ph]].
- [113] N. Bartolo, E. Dimastrogiovanni and A. Vallinotto, (to appear).
- [114] L. Leblond and S. Shandera, JCAP **0808**, 007 (2008) [arXiv:0802.2290 [hep-th]].
- [115] S. Shandera, Phys. Rev. D **79**, 123518 (2009) [arXiv:0812.0818 [astro-ph]].
- [116] C. T. Byrnes, K. Koyama, M. Sasaki and D. Wands, JCAP **0711**, 027 (2007) [arXiv:0705.4096 [hep-th]].
- [117] D. H. Lyth and Y. Rodriguez, Phys. Rev. Lett. **95**, 121302 (2005) [arXiv:astro-ph/0504045].
- [118] L. Boubekeur and D. H. Lyth, Phys. Rev. D **73**, 021301 (2006) [arXiv:astro-ph/0504046].
- [119] I. Zaballa, Y. Rodriguez and D. H. Lyth, JCAP **0606** (2006) 013 [arXiv:astro-ph/0603534].
- [120] S. Weinberg, "Cosmology" (Oxford University Press, 2008): Sec. 5.2
- [121] C. P. Burgess, L. Leblond, R. Holman and S. Shandera, arXiv:0912.1608 [hep-th].
- [122] L. H. Ford, Phys. Rev. D **40**, 967 (1989).

- [123] A. Golovnev, V. Mukhanov and V. Vanchurin, JCAP **0806**, 009 (2008) [arXiv:0802.2068 [astro-ph]].
- [124] A. Golovnev, V. Mukhanov and V. Vanchurin, JCAP **0811**, 018 (2008) [arXiv:0810.4304 [astro-ph]].
- [125] A. Golovnev and V. Vanchurin, Phys. Rev. D **79**, 103524 (2009) [arXiv:0903.2977 [astro-ph.CO]].
- [126] C. Armendariz-Picon, JCAP **0407**, 007 (2004) [arXiv:astro-ph/0405267].
- [127] C. G. Boehmer and T. Harko, Eur. Phys. J. C **50**, 423 (2007) [arXiv:gr-qc/0701029].
- [128] T. Koivisto and D. F. Mota, Astrophys. J. **679**, 1 (2008) [arXiv:0707.0279 [astro-ph]].
- [129] T. S. Koivisto and D. F. Mota, JCAP **0808**, 021 (2008) [arXiv:0805.4229 [astro-ph]].
- [130] J. B. Jimenez, R. Lazkoz and A. L. Maroto, arXiv:0904.0433 [astro-ph.CO].
- [131] J. B. Jimenez, R. Lazkoz and A. L. Maroto, Phys. Rev. D **80**, 023004 (2009).
- [132] B. Himmetoglu, C. R. Contaldi and M. Peloso, Phys. Rev. Lett. **102**, 111301 (2009) [arXiv:0809.2779 [astro-ph]].
- [133] B. Himmetoglu, C. R. Contaldi and M. Peloso, Phys. Rev. D **79**, 063517 (2009) [arXiv:0812.1231 [astro-ph]].
- [134] B. Himmetoglu, C. R. Contaldi and M. Peloso, Phys. Rev. D **80**, 123530 (2009) [arXiv:0909.3524 [astro-ph.CO]].
- [135] K. Dimopoulos, Phys. Rev. D **74**, 083502 (2006) [arXiv:hep-ph/0607229].
- [136] K. Dimopoulos and M. Karciuskas, JHEP **0807**, 119 (2008) [arXiv:0803.3041 [hep-th]].
- [137] S. Kanno, M. Kimura, J. Soda and S. Yokoyama, JCAP **0808**, 034 (2008) [arXiv:0806.2422 [hep-ph]].
- [138] L. Ackerman, S. M. Carroll and M. B. Wise, Phys. Rev. D **75**, 083502 (2007) [Erratum-ibid. D **80**, 069901 (2009)] [arXiv:astro-ph/0701357].
- [139] S. M. Carroll, T. R. Dulaney, M. I. Gresham and H. Tam, Phys. Rev. D **79**, 065011 (2009) [arXiv:0812.1049 [hep-th]].
- [140] D. H. Lyth, JCAP **0511**, 006 (2005) [arXiv:astro-ph/0510443].
- [141] L. Alabidi and D. Lyth, JCAP **0608**, 006 (2006) [arXiv:astro-ph/0604569].
- [142] M. P. Salem, Phys. Rev. D **72**, 123516 (2005) [arXiv:astro-ph/0511146].
- [143] F. Bernardeau, L. Kofman and J. P. Uzan, Phys. Rev. D **70**, 083004 (2004) [arXiv:astro-ph/0403315].
- [144] M. a. Watanabe, S. Kanno and J. Soda, Phys. Rev. Lett. **102**, 191302 (2009) [arXiv:0902.2833 [hep-th]].
- [145] B. Himmetoglu, arXiv:0910.3235 [astro-ph.CO].
- [146] T. R. Dulaney and M. I. Gresham, arXiv:1001.2301 [astro-ph.CO].
- [147] D. Seery and J. E. Lidsey, JCAP **0506**, 003 (2005) [arXiv:astro-ph/0503692].
- [148] F. Vernizzi and D. Wands, JCAP **0605**, 019 (2006) [arXiv:astro-ph/0603799].
- [149] J. Martin and J. Yokoyama, JCAP **0801**, 025 (2008) [arXiv:0711.4307 [astro-ph]].
- [150] D. Babich, P. Creminelli and M. Zaldarriaga, JCAP **0408**, 009 (2004) [arXiv:astro-ph/0405356].
- [151] M.E. Peskin and D.V. Schroeder, “An introduction to quantum field theory” (1995): Sec. 9.2
- [152] D. Seery, J. E. Lidsey and M. S. Sloth, JCAP **0701**, 027 (2007) [arXiv:astro-ph/0610210].
- [153] E. Komatsu *et al.*, arXiv:1001.4538 [astro-ph.CO].
- [154] C. L. Bennett *et al.*, arXiv:1001.4758 [astro-ph.CO].