

NEWS & VIEWS

Open Access

Non-destructive OAM measurement via light–matter interaction

Gianluca Ruffato ^{1,2}✉

Abstract

The detection of orbital angular momentum usually relies on optical techniques, which modify the original beam to convert the information carried on its phase into a specific intensity distribution in output. Moreover, the exploitation of high-intensity beams can result destructive for standard optical elements and setups. A recent publication suggests a solution to overcome all those limitations, by probing highly-intense vortex pulses with a structured reference beam in a strong-field photoionization process.

Almost 30 years have passed since the seminal paper of Allen and co-workers¹ ignited a still vivid field of research, which has paved the way to amazing applications in an unexpected variety of areas, including life science, soft and condensed matter, and information and communication technology. In particular, orbital angular momentum (OAM) beams have provided a new degree of freedom to encode information² or increase the Hilbert space in quantum communications³, while their uncommon phase and intensity distributions have enabled advanced techniques in micromanipulation⁴, imaging⁵, and light–matter interaction⁶. Concurrently, those research and applications have inspired the development of new optical components and innovative methods for the controlled generation, manipulation, and measurement of optical angular momentum.

In particular, the measurement of OAM can rely on a rich portfolio of optical elements and architectures with different levels of complexity, integration, and miniaturization, such as interferometric⁷ or diffractive⁸ techniques, holographic methods⁹, conformal mappings^{10,11} multi-plane light conversion¹², and integrated photonics¹³, to name the most used. However, all those solutions inevitably modify the input beam to obtain an output intensity

distribution, which is correlated to the input amount of OAM. In addition, the required physical processing of the input wavefront prevents the application of those methods to high-intensity beams, which may damage or even destroy the optical elements.

A recent experiment coordinated by Professor Y. Liu¹⁴ suggests a possible way-out to detect the OAM of an intense optical vortex, by analyzing the spatial distribution of the electrons emitted in a strong-field photoionization process¹⁵. In such a way, the information on the input field is transferred to secondary carriers, i.e., photoelectrons, with the sacrifice of a negligible amount of photons from the original beam. Thus, from the analysis of the photoelectron momentum distribution or angle-resolved yields, it is possible to infer the amount of OAM carried by the probed optical vortex without affecting significantly its original phase and intensity distribution.

In this work, a two-color co-rotating circular laser field is synthesized by the combination of circularly polarized probing pulses at 800 nm with co-polarized 400-nm optical vortices produced by the cascade of a frequency-doubling non-linear crystal (β -BBO) with a spiral phase plate for OAM generation. The synthesized field is made to interact with a supersonic Argon atom gas jet in a vacuum chamber, and a time-of-flight spectrometer with a position-sensitive detector is used for the three-dimensional reconstruction of photoelectron momentum distribution¹⁶ (Fig. 1). In particular, by inserting a slit in order to spatially sculpture the probing field via

Correspondence: Gianluca Ruffato (gianluca.ruffato@unipd.it)


¹Department of Physics and Astronomy 'G. Galilei', University of Padova, Padova, Italy

²Quantum Technologies Research Center (QTech), University of Padova, Padova, Italy

© The Author(s) 2022



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.



diffraction, the authors show how the consequent spatial structuring of the synthesized field induces an axial asymmetry in the photoelectron distribution, which can be correlated to the OAM amount of the unknown vortex field. Moreover, when the slit is rotated in time, the yields of photoelectrons confined within a defined range exhibits a periodic trend, with the peaks' number and position depending on the value and sign of the probed OAM.

This solution prevents the OAM state from collapsing as in previous methods, so that the measured vortex pulses can be then exploited in subsequent applications, while avoiding at the same time the passage through physical optical elements, which could result in irreparable damage by the intense field.

While the authors provide an experimental proof only for low-order states, a universal scheme to detect any amount of OAM is also suggested using the same set-up. The research opens to new scenarios for the analysis of highly-intense vortex fields, suggesting significant implications to the study of the optical angular momentum of ultrafast and intense optical vortices. The availability of an in situ non-destructive analysis method is expected to enable new applications and further extend the study of intense optical vortices in light-matter interaction and structured light dynamics. Future analyses can also include more complex

laser fields, such as vector¹⁷ or skyrmionic¹⁸ beams, while possible applications can be envisaged to the generation and analysis of structured matter waves, e.g., electron beams¹⁹.

Published online: 10 March 2022

References

- Allen, L. et al. Orbital angular momentum of light and the transformation of Laguerre-Gaussian laser modes. *Phys. Rev. A* **45**, 8185–8189 (1992).
- Wang, J. Twisted optical communications using orbital angular momentum. *Sci. China Phys. Mech. Astron.* **62**, 34201 (2019).
- Mirhosseini, N. et al. High-dimensional quantum cryptography with twisted light. *N. J. Phys.* **17**, 033033 (2015).
- Padgett, M. & Bowman, R. Tweezers with a twist. *Nat. Photonics* **5**, 343–348 (2011).
- Ritsch-Marte, M. Orbital angular momentum light in microscopy. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.* **375**, 20150437 (2017).
- Franke-Arnold, S. Optical angular momentum and atoms. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.* **375**, 20150435 (2017).
- Leach, J. et al. Interferometric methods to measure orbital and spin, or the total angular momentum of a single photon. *Phys. Rev. Lett.* **92**, 013601 (2004).
- Hickmann, J. M. et al. Unveiling a truncated optical lattice associated with a triangular aperture using light's orbital angular momentum. *Phys. Rev. Lett.* **105**, 053904 (2010).
- Gibson, G. et al. Free-space information transfer using light beams carrying orbital angular momentum. *Opt. Express* **12**, 5448–5456 (2004).
- Berkhout, G. C. G. et al. Efficient sorting of orbital angular momentum states of light. *Phys. Rev. Lett.* **105**, 153601 (2010).

11. Wen, Y. H. et al. Spiral transformation for high-resolution and efficient sorting of optical vortex modes. *Phys. Rev. Lett.* **120**, 193904 (2018).
12. Fontaine, N. K. et al. Laguerre-Gaussian mode sorter. *Nat. Commun.* **10**, 1865 (2019).
13. Su, T. H. et al. Demonstration of free space coherent optical communication using integrated silicon photonic orbital angular momentum devices. *Opt. Express* **20**, 9396–9402 (2012).
14. Fang, Y. Q. et al. Probing the orbital angular momentum of intense vortex pulses with strong-field ionization. *Light. Sci. Appl.* **11**, 34 (2022).
15. Ullrich, J. et al. Recoil-ion momentum spectroscopy. *J. Phys. B At. Mol. Optical Phys.* **30**, 2917–2974 (1997).
16. Fang, Y. Q. et al. Photoelectronic mapping of the spin-orbit interaction of intense light fields. *Nat. Photonics* **15**, 115–120 (2021).
17. Rosales-Guzmán, C., Ndagano, B. & Forbes, A. A review of complex vector light fields and their applications. *J. Opt.* **20**, 123001 (2018).
18. Gao, S. J. et al. Paraxial skyrmionic beams. *Phys. Rev. A* **102**, 053513 (2020).
19. Uchida, M. & Tonomura, A. Generation of electron beams carrying orbital angular momentum. *Nature* **464**, 737–739 (2010).