

THE RATIONALITY OF QUATERNIONIC DARMON POINTS OVER GENUS FIELDS OF REAL QUADRATIC FIELDS

MATTEO LONGO AND STEFANO VIGNI

ABSTRACT. Darmon points on p -adic tori and Jacobians of Shimura curves over \mathbb{Q} were introduced in [19] and [20] as generalizations of Darmon's Stark–Heegner points. In this article we study the algebraicity over extensions of a real quadratic field K of the projections of Darmon points to elliptic curves. More precisely, we prove that linear combinations of Darmon points on elliptic curves weighted by certain genus characters of K are rational over the predicted genus fields of K . This extends to an arbitrary quaternionic setting the main theorem on the rationality of Stark–Heegner points obtained by Bertolini and Darmon in [4], and at the same time gives evidence for the rationality conjectures formulated in the joint paper with Rotger [20] and by M. Greenberg in [13]. In light of this result, quaternionic Darmon points represent the first instance of a systematic supply of points of Stark–Heegner type other than Darmon's original ones for which explicit rationality results are known.

1. INTRODUCTION

Given an elliptic curve E over \mathbb{Q} of conductor N , a prime number p dividing N exactly and a real quadratic field K in which p is inert and all primes dividing N/p split (Stark–Heegner hypothesis), Darmon introduced in [7] the notion of Stark–Heegner points on E . These local points live in $E(K_p)$, where K_p is the completion of K at p , but are conjectured to be rational over (narrow) ring class fields of K and to satisfy a Shimura reciprocity law under Galois actions. In fact, the arithmetic properties of Stark–Heegner points are expected to be as rich as those enjoyed by classical Heegner points, which are defined via the theory of complex multiplication and are known to be rational over ring class fields of imaginary quadratic fields.

Stark–Heegner points were later lifted to certain quotients of classical modular Jacobians by Dasgupta in [8]; this was done by proving a rigid analytic uniformization result for modular Jacobians which boils down to an equality of \mathcal{L} -invariants and turns out to be a strong form of a theorem of Greenberg and Stevens ([14]). It is important to observe that both Darmon's and Dasgupta's strategies rely heavily on the theory of modular symbols (and so on the presence of cusps on classical modular curves), and thus they do not extend to more general situations where the Stark–Heegner condition is not satisfied but considerations on the signs of the functional equations of the relevant L -functions predict the existence of points of Stark–Heegner type.

To overcome this difficulty, in [13] M. Greenberg replaced modular symbols by a general cohomological study of Shimura varieties over totally real fields and reinterpreted Darmon's theory in terms of group cohomology. The outcome of his work is, in the simplest case where the base field is \mathbb{Q} , a conjectural construction of local Stark–Heegner points on E which are expected to be rational over appropriate ring class fields of a real quadratic field K . In this context, the field K satisfies a modified Stark–Heegner hypothesis forcing the number of primes dividing N/p and inert in K to be even.

2000 *Mathematics Subject Classification.* 14G05, 11G05.

Key words and phrases. Darmon points, elliptic curves, genus fields.

In [19] we gave an explicit rigid analytic uniformization of the maximal toric quotient of the Jacobian of a Shimura curve over \mathbb{Q} at a prime dividing exactly the level, a result that can be viewed as complementary to the classical theorem of Čerednik and Drinfeld which provides rigid analytic uniformizations at primes dividing the discriminant ([5]). More precisely, denote $J_0^D(Mp)$ the Jacobian variety of the Shimura curve of discriminant $D > 1$ and level Mp . Combining techniques in group cohomology and p -adic integration in the spirit of the works of Dasgupta and Greenberg, we exhibited a p -adic torus which is isogenous to the product of two copies of the p -new quotient of $J_0^D(Mp)$. As a corollary, we offered a proof of the isogeny conjecture formulated in [13, Conjecture 2] (an alternative proof of which has been given by Dasgupta and Greenberg in [9]): this made Greenberg’s construction of local points on elliptic curves over \mathbb{Q} unconditional, but left the rationality conjecture [13, Conjecture 3] wide open.

As an application of our work on rigid analytic uniformizations, in [20] we introduced a new family of Stark–Heegner type points on p -adic tori and Jacobians of Shimura curves, which we called *Darmon points*. These are, at the same time, lifts of Greenberg’s points on elliptic curves and generalizations of Dasgupta’s Stark–Heegner points on modular Jacobians, and we formulated conjectures about their rationality over (narrow) ring class fields of real quadratic fields and their Galois properties. Moreover, assuming these rationality conjectures and applying the methods of [17], we proved the conjecture of Birch and Swinnerton-Dyer for elliptic curves over ring class fields of real quadratic fields in the case of analytic rank 0.

At present, the central open problems in the theories of Stark–Heegner and Darmon points are the rationality conjectures. The first result towards the rationality of Stark–Heegner points was obtained by Bertolini and Darmon in [4], where they proved that linear combinations of Stark–Heegner points weighted by genus characters of K are rational over the predicted genus fields of K . To achieve this, the authors showed that linear combinations of this kind can be expressed in terms of Heegner points and applied their previous results on Hida families of modular forms ([3]).

In this paper we prove the analogue of the results of [4] for projections to elliptic curves of the quaternionic Darmon points introduced in [20]. In other words, we show that linear combinations with genus character coefficients of Darmon points on elliptic curves are rational over the fields that were predicted in [20]. In light of this feature, quaternionic Darmon points represent the first instance of a systematic supply of points of Stark–Heegner type other than Darmon’s original ones for which explicit rationality results are known. Thus our main theorem provides some evidence for both the rationality conjecture formulated in [20] and the conjecture proposed by Greenberg in [13, Conjecture 3] when the totally real base field is \mathbb{Q} . The present article should be viewed as a natural step further in the line of investigation begun in [19] and carried on in [20].

While the reader familiar with the work of Bertolini and Darmon will notice that our strategy is inspired by [4] (e.g., the main result of [3] plays a crucial role in the last part of our arguments), the methods we use are of a different nature and build on the constructions of [19] in an essential way. In fact, in this article we develop a series of new techniques and obtain various auxiliary results which may be of independent interest. Finally, we remark that, as in [4], the scheme of proof presents formal analogies with that of Kronecker’s solution to Pell’s equation (see [4, Introduction] for an illustration of this parallelism).

Now we describe our results more in detail. Fix an elliptic curve E over \mathbb{Q} of square-free conductor N and a real quadratic field K of discriminant d_K prime to N , whose ring of integers we denote \mathcal{O}_K . Suppose that there is a factorization $N = MDp$ such that

- all the primes dividing M (respectively, Dp) are split (respectively, inert) in K ;
- the number of primes dividing D is *even* and ≥ 2 .

Let B denote the indefinite quaternion algebra over \mathbb{Q} of discriminant D and choose Eichler orders $R_0^D(Mp) \subset R_0^D(M)$ of B of level Mp and M , respectively. Write $\Gamma_0^D(Mp)$ and $\Gamma_0^D(M)$ for the groups of elements of norm 1 in $R_0^D(Mp)$ and $R_0^D(M)$, respectively, and consider the Ihara group

$$\Gamma := \left\{ \gamma \in R_0^D(M) \otimes_{\mathbb{Z}} \mathbb{Z}[1/p] \mid \text{norm}(\gamma) = 1 \right\}.$$

For any sign \pm fix an isomorphism $H^1(E(\mathbb{C}), \mathbb{Z})^{\pm} \simeq \mathbb{Z}$, where the superscript denotes the \pm -eigenspace for complex conjugation. Following [19, §7.7], we define a canonical element

$$\mu_E^{\pm} \in H^1(\Gamma, \mathcal{M}_0(\mathbb{Q}))$$

where $\mathcal{M}_0(\mathbb{Q})$ is the \mathbb{Q} -vector space of \mathbb{Q} -valued measures on $\mathbb{P}^1(\mathbb{Q}_p)$ with total mass 0. Its image in $H^1(\Gamma_0^D(Mp), \mathbb{Q})$, obtained by applying Shapiro's lemma, spans the one-dimensional \mathbb{Q} -vector space on which the archimedean involution W_{∞} acts via \pm and the Hecke algebra of level N acts via the character associated with the newform f attached to E by modularity (see §2.2 for details on these actions).

Fix a sign \pm . Using techniques introduced in [19] and [20], in §2.4 we use μ_E^{\pm} to attach a Darmon point $P_{\psi}^{\pm} \in E(K_p)$ to (the $\Gamma_0^D(M)$ -conjugacy class of) each optimal embedding ψ of \mathcal{O}_K into $R_0^D(M)$: these points are the object of our investigation in the present paper.

Let G_K^{\pm} be the narrow class group of K (i.e., the Galois group over K of the narrow Hilbert class field of K). For a genus character χ of K (i.e., an unramified quadratic character of G_K^{\pm}) we define the point

$$P_{\chi} := \sum_{\sigma \in G_K^{\pm}} P_{\psi_{\sigma}}^{\epsilon} \in E(K_p),$$

where the Galois action $\psi \mapsto \psi_{\sigma}$ on (conjugacy classes of) optimal embeddings is described in §2.5 and the choice of $\epsilon \in \{\pm\}$ depends on χ .

The main result of this paper, which is Theorem 5.1 in the text, is the following

Theorem 1.1. *Let χ be a genus character of K associated with a pair (χ_1, χ_2) of Dirichlet characters such that $\chi_i(-MD) = -w_{MD}$ for $i = 1, 2$, where w_{MD} is the eigenvalue of the Atkin-Lehner involution W_{MD} acting on f .*

- (1) *There exists $n \in \mathbb{Z}$ such that $nP_{\chi} \in E(H_{\chi})$ where H_{χ} is the genus field of K cut out by χ .*
- (2) *The point nP_{χ} is of infinite order if and only if $L'(E/K, \chi, 1) \neq 0$.*

Now we outline the strategy of proof of this theorem. In doing this, we state two results (Theorems 1.2 and 1.4) which may be of independent interest.

Write \mathbb{D} for the module of \mathbb{Z}_p -valued measures on $\mathbb{Y} := \mathbb{Z}_p^2$ which are supported on the subset \mathbb{X} of primitive elements (i.e., those $(x, y) \in \mathbb{Y}$ which are not divisible by p). In Section 3 we exploit a control theorem for Hida families in the quaternionic setting (analogous to the one proved by Greenberg and Stevens in [14]) to construct a canonical element

$$\tilde{\mu}_E^{\pm} \in H^1(\Gamma_0^D(M), \mathbb{D}) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$$

such that the image of $\tilde{\mu}_E^{\pm}$ via the map $\pi : \mathbb{X} \rightarrow \mathbb{P}^1(\mathbb{Q}_p)$ taking (x, y) to x/y coincides with μ_E^{\pm} . We fix a representative $\tilde{\mu}_E^{\pm}$ of this class, which can be chosen so that $\pi_*(\tilde{\mu}_E^{\pm}) = \mu_E^{\pm}$. In order to

describe $\tilde{\mu}_E^\pm$ more precisely, embed \mathbb{Z} into the weight space

$$(1) \quad \mathcal{X} := \mathrm{Hom}(\mathbb{Z}_p^\times, \mathbb{Z}_p^\times) \simeq \mathbb{Z}/(p-1)\mathbb{Z} \times (1 + p\mathbb{Z}_p)$$

by sending k to the map $(x \mapsto x^{k-2})$. Hida theory associates with f a neighbourhood \mathbb{U} of 2 in \mathcal{X} (which we assume small enough so that all the results we will state below hold for \mathbb{U}) and a formal q -expansion

$$(2) \quad f_\infty = \sum_{n=1}^{\infty} a_n(\kappa) q^n$$

where $a_1 = 1$ and a_n is a rigid analytic function on \mathbb{U} such that $\sum_{n=1}^{\infty} a_n(k) q^n$ for an even integer $k \geq 2$ is the q -expansion of a p -stabilized weight k eigenform f_k on $\Gamma_0(N)$ and $f_2 = f$. The form f_k is associated, via the Jacquet–Langlands correspondence, with a modular form f_k^{JL} of weight k and level structure $\Gamma_0^D(Mp)$ (so f_k^{JL} is well defined only up to scalars). Thanks to a result of Matsushima and Shimura ([21]), f_k^{JL} corresponds to an element ϕ_k in the first cohomology group of $\Gamma_0^D(Mp)$ with values in the dual of the \mathbb{C} -vector space of homogeneous polynomials of degree $k-2$. Then the property characterizing $\tilde{\mu}_E^\pm$ asserts that integrating homogeneous polynomials of degree $k-2$ against $\tilde{\mu}_{E,\gamma}^\pm$ over $\mathbb{Z}_p \times \mathbb{Z}_p^\times$ (with $\gamma \in \Gamma_0^D(Mp)$) defines a multiple of the projection of ϕ_k to the \pm -eigenspace for W_∞ . See §3.3 for details.

Now let

$$(3) \quad \Phi_{\mathrm{Tate}} : \bar{K}_p / \langle q \rangle \longrightarrow E(\bar{K}_p)$$

denote Tate's p -adic uniformization, where $q \in p\mathbb{Z}_p$ is Tate's period for E at p and \bar{K}_p is an algebraic closure of K_p . Let \log_q be the branch of the p -adic logarithm satisfying $\log_q(q) = 0$ and define $\log_E(P) := \log_q(\Phi_{\mathrm{Tate}}^{-1}(P))$ for all $P \in E(K_p)$.

As a piece of notation, let z_ψ denote the fixed point of $\psi(K^\times)$ acting on $\mathbb{P}^1(\bar{\mathbb{Q}}_p)$ such that $\psi(\alpha)(z_\psi, 1) = \alpha(z_\psi, 1)$ for all $\alpha \in K^\times$ (see §2.3 for precise definitions). Let also ε_K be a generator of the group of norm 1 elements in \mathcal{O}_K^\times such that $\varepsilon_K > 1$ with respect to a fixed embedding $K \hookrightarrow \mathbb{R}$ and define $\gamma_\psi := \psi(\varepsilon_K)$.

The first auxiliary result proved in the paper is the following

Theorem 1.2. *There is an integer $m \neq 0$ such that*

$$\log_E(P_\psi^\pm) = m \cdot \int_{\mathbb{X}} \log_q(x - z_\psi y) d\tilde{\mu}_{E,\gamma_\psi}^\pm.$$

This theorem, which corresponds to Corollary 3.6, allows us to explicitly compute (at least in principle) Darmon points on elliptic curves. Actually, in Theorem 3.4 we prove a more general version for Darmon points on the p -adic torus T/L by means of which we obtained in [19] a parametrization of E (cf. §2.7).

Remark 1.3. The integer m appearing in Theorem 1.2 can be made explicit in terms of the exponent of the abelianization of Γ and the degree of an isogeny between the torus T/L and the elliptic curve E (see §2.7).

In Section 4 we use the cohomology class $\tilde{\mu}_E^\pm$ to introduce a p -adic L -function $L_p(f_\infty/K, \chi, k)$ attached to f_∞ and a genus character χ of K , whose p -adic variable k ranges in \mathbb{U} . The main result proved by Popa in [25] can then be combined with the interpolation properties of $\tilde{\mu}_E^\pm$ to show that, for a classical modular form f_k of even weight $k \geq 4$ and trivial character in the Hida family, the value of $L_p(f_\infty/K, \chi, k)$ is equal, up to multiplication by an explicit non-zero

constant, to the special value of the complex L -function of f_k^\sharp twisted by χ , where the form f_k^\sharp is the p -stabilization of f_k (see Theorem 4.24).

Now recall that a genus character χ of K corresponds to an unordered pair of quadratic Dirichlet characters χ_1 and χ_2 of discriminants d_1 and d_2 , respectively, such that $d_K = d_1 d_2$, and for $j = 1, 2$ let $L_p(f_\infty, \chi_j, k, s)$ denote the Mazur–Kitagawa two-variable p -adic L -function associated with f_∞ and χ_j .

Our last result we highlight in this introduction is the following factorization formula, corresponding to Theorem 4.33 in the main body of the article.

Theorem 1.4. *Let χ be a genus character of K and let (χ_1, χ_2) be the associated pair of Dirichlet characters. There exists a p -adic analytic function η on \mathbb{U} such that $\eta(k) \neq 0$ and*

$$L_p(f_\infty/K, \chi, k) = \eta(k) L_p(f_\infty, \chi_1, k, k/2) L_p(f_\infty, \chi_2, k, k/2)$$

for all $k \in \mathbb{U}$. Moreover, $\eta(2) \in (\mathbb{Q}^\times)^2$.

Here is a brief explanation of how Theorems 1.2 and 1.4 are used to prove Theorem 1.1.

- (1) Theorem 1.4 provides a link between the values at $k = 2$ of the second derivatives of $L_p(f_\infty/K, \chi, k)$ and of the restriction of $L_p(f_\infty, \chi_1, k, s)$ to the line $s = k/2$ (here we need to order the pair (χ_1, χ_2) so that the sign of the functional equation of the L -function $L(E, \chi_1, s)$ is -1 , which can be done thanks to our assumptions).
- (2) On the one hand, the second derivative of $L_p(f_\infty/K, \chi, k)$ evaluated at $k = 2$ is essentially computed by the integrals appearing on the right hand side of Theorem 1.2, thus providing a link with Darmon points (this is proved in Theorem 4.31).
- (3) On the other hand, [3, Theorem 5.4] shows that the value at $k = 2$ of the second derivative of $L_p(f_\infty, \chi_1, k, k/2)$ encodes the logarithm of certain linear combinations of classical Heegner points.

The three observations above establish a relation between Heegner points and Darmon points, yielding the desired rationality result for P_χ .

Remark 1.5. The hypothesis that M be square-free is imposed only to be able to apply the results of [25], which are proved under this assumption. Granting an extension of [25] to more general situations, our arguments work equally well under the weaker condition that M be a product of odd powers of distinct primes (and this restriction comes into play exclusively in §4.5 for some calculations with local Hilbert symbols).

2. DARMON POINTS ON p -ADIC TORI AND ELLIPTIC CURVES

2.1. Quaternion algebras and Shimura curves. Fix throughout the paper embeddings

$$(4) \quad \bar{\mathbb{Q}} \hookrightarrow \bar{\mathbb{Q}}_p, \quad \bar{\mathbb{Q}}_p \hookrightarrow \mathbb{C}.$$

Let $D > 1$ be a square-free product of an *even* number of primes and let $M \geq 1$ be a square-free integer prime to D . We also fix a prime number p not dividing $6MD$. Write B for the (indefinite) quaternion algebra over \mathbb{Q} of discriminant D .

Let $K = \mathbb{Q}(\sqrt{d_K})$ be a real quadratic field with fundamental discriminant d_K such that all the primes dividing Dp are inert in K and all the primes dividing M are split in K , and fix an embedding $K \hookrightarrow \mathbb{R}$. For the rest of the paper, fix an auxiliary real quadratic field F such that all the primes dividing D are inert in F while p is split in F . In particular, $F \neq K$. The condition on the primes dividing D implies that F is a splitting field for B , so we can (and do) choose an isomorphism $i_F : B \otimes_{\mathbb{Q}} F \xrightarrow{\sim} M_2(F)$ of F -algebras. Since p splits in F , the embedding $\bar{\mathbb{Q}} \hookrightarrow \bar{\mathbb{Q}}_p$

fixed in (4) and the isomorphism i_F induce an isomorphism $i_p : B \otimes_{\mathbb{Q}} \mathbb{Q}_p \xrightarrow{\cong} M_2(\mathbb{Q}_p)$ of \mathbb{Q}_p -algebras. Moreover, fix an embedding $F \hookrightarrow \mathbb{R}$; this induces via i_F an isomorphism $i_{\infty} : B \otimes_{\mathbb{Q}} \mathbb{R} \xrightarrow{\cong} M_2(\mathbb{R})$ of \mathbb{R} -algebras. Finally, for all primes $\ell \nmid Dp$ fix isomorphisms $i_{\ell} : B \otimes_{\mathbb{Q}} \mathbb{Q}_{\ell} \xrightarrow{\cong} M_2(\mathbb{Q}_{\ell})$ of \mathbb{Q}_{ℓ} -algebras.

Remark 2.1. When describing local arguments at p , the fact that p splits in F allows us to work with \mathbb{Q}_p and not with quadratic extensions of it, which simplifies the computations.

Let R^{\max} be the maximal order of B satisfying the following condition: if ℓ is a prime such that $\ell \nmid D$ then $i_{\ell}(R^{\max} \otimes \mathbb{Z}_{\ell}) = M_2(\mathbb{Z}_{\ell})$. We fix an Eichler order $R_0^D(M)$ of level M contained in R^{\max} by requiring that for all primes $\ell \nmid M$ the order $i_{\ell}(R_0^D(M) \otimes \mathbb{Z}_{\ell})$ is equal to the order $R_0^{\text{loc}}(\ell)$ of $M_2(\mathbb{Z}_{\ell})$ consisting of those matrices $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ with $c \equiv 0 \pmod{\ell}$. Furthermore, we also fix an Eichler order $R_0^D(Mp) \subset R_0^D(M)$ of level Mp by requiring that, in addition, $i_p(R_0^D(Mp) \otimes \mathbb{Z}_p)$ is equal to the order $R_0^{\text{loc}}(p)$ of $M_2(\mathbb{Z}_p)$ consisting of those matrices $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ with $c \equiv 0 \pmod{p}$. We denote by $\Gamma_0^D(M)$ and $\Gamma_0^D(Mp)$ the groups of elements in $R_0^D(M)$ and $R_0^D(Mp)$, respectively, of reduced norm 1.

Finally, let $X_0^D(M)$ (respectively, $X_0^D(Mp)$) denote the (compact) Shimura curve whose corresponding Riemann surface is defined as the analytic quotient $\Gamma_0^D(M) \backslash \mathcal{H}$ (respectively, $\Gamma_0^D(Mp) \backslash \mathcal{H}$), where \mathcal{H} is the complex upper half plane and the elements of B^{\times} with positive norm act on it by Möbius (i.e., fractional linear) transformations via i_{∞} .

2.2. Hecke algebras. For any subgroup G of B^{\times} consisting of elements having reduced norm 1 and any subsemigroup S of B^{\times} such that (G, S) is a Hecke pair in the sense of [1, §1.1] we denote by $\mathcal{H}(G, S)$ the Hecke algebra (over \mathbb{Z}) of the pair (G, S) whose elements are combinations with integer coefficients of double cosets $T(s) := GsG$ for $s \in S$.

Let norm be the reduced norm of B , let $g \mapsto g^* := \text{norm}(g)g^{-1}$ be the main involution of B^{\times} and for any subset S of B^{\times} let S^* denote the image of S under the main involution.

If M is a left $\mathbb{Z}[S^*]$ -module, the group $H^r(G, M)$ has a natural right action of $\mathcal{H}(G, S)$ defined on the level of cochains $c \in Z^r(G, M)$ as follows: for $s \in S$ set

$$(c|T(s))(\gamma_1, \dots, \gamma_r) := \sum_i s_i^* \cdot c(t_i(\gamma_1), \dots, t_i(\gamma_r))$$

where $GsG = \coprod_i Gs_i$ (finite disjoint decomposition) and $t_i : G \rightarrow G$ is defined by the equations $Gs_i\gamma = Gs_j$ (for some j) and $g_i\gamma = t_i(\gamma)g_j$. We also define a right action of $\mathcal{H}(G, S)$ on $H_r(G, M)$ by the formula

$$c|T(s) := \sum_{i,j} s_i(m_j) \otimes [t_i(\gamma_{1,j}) | \dots | t_i(\gamma_{r,j})]$$

for $c = \sum_j m_j \otimes [\gamma_{1,j} | \dots | \gamma_{r,j}] \in Z_r(G, M)$.

Remark 2.2. The above formalism of Hecke operators slightly differs from the one adopted in [19, §2.1], where we considered left actions on homology and cohomology groups instead of right actions.

For primes ℓ let $\Sigma_{\ell}^{\text{loc}}$ denote the semigroup of elements in $R^{\max} \otimes \mathbb{Z}_{\ell}$ with non-zero norm. For every prime $\ell \nmid Mp$ let $\Sigma_0^{\text{loc}}(\ell) \subset \Sigma_{\ell}^{\text{loc}}$ be the inverse image under i_{ℓ} of the semigroup of matrices $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{GL}_2(\mathbb{Q}_{\ell}) \cap M_2(\mathbb{Z}_{\ell})$ with $a \in \mathbb{Z}_{\ell}^{\times}$ and $c \equiv 0 \pmod{\ell}$. Then define the semigroup

$$S_0^D(Mp) := B^+ \cap \left(\prod_{\ell \nmid Mp} \Sigma_0^{\text{loc}}(\ell) \times \prod_{\ell \nmid Mp} \Sigma_{\ell}^{\text{loc}} \right)$$

where B^+ is the subgroup of elements in B^\times of positive norm. Then we can form the commutative Hecke algebra $\mathcal{H}(Mp) = \mathcal{H}(\Gamma_0^D(Mp), S_0^D(Mp))$ generated over \mathbb{Z} by the standard elements $T_n = \sum T(\alpha)$ for $n \geq 1$ (the sum being over all the elements $\alpha \in S$ of norm n) and $T_{n,n} = T(n)$ for $n \geq 1$, $n \nmid MDp$. Analogously, we can define the Hecke algebra $\mathcal{H}(M)$.

As in [19], consider the Ihara group

$$\Gamma := \left\{ \gamma \in R_0^D(M) \otimes_{\mathbb{Z}} \mathbb{Z}[1/p] \mid \text{norm}(\gamma) = 1 \right\} \hookrightarrow \text{SL}_2(\mathbb{Q}_p)$$

where $\text{norm} : B \rightarrow \mathbb{Q}$ denotes the reduced norm map and the injection is induced by the composition of the canonical injection $B \hookrightarrow B \otimes_{\mathbb{Q}} \mathbb{Q}_p$ with the isomorphism i_p fixed in (4). If

$$S_p := B^+ \cap \left(\text{M}_2(\mathbb{Q}_p) \times \prod_{\ell|M} \Sigma_0^{\text{loc}}(\ell) \times \prod_{\ell \nmid Mp} \Sigma_{\ell}^{\text{loc}} \right)$$

then we may also consider the Hecke algebra $\mathcal{H}(\Gamma, S_p)$. See [19, §2.2 and §2.3] for details.

Now we introduce Atkin–Lehner involutions. For every prime $\ell \nmid MDp$ choose an element $\omega_{\ell} \in R_0^D(Mp)$ of norm ℓ ; if $\ell \mid Mp$ then we can (and do) choose ω_{ℓ} in such a way that $i_{\ell}(\omega_{\ell}) = \begin{pmatrix} 0 & -1 \\ \ell & 0 \end{pmatrix}$ up to an element of $R_0^{\text{loc}}(\ell)$ of norm 1. It is known that ω_{ℓ} normalizes both $\Gamma_0^D(Mp)$ and $\Gamma_0^D(M)$ when $\ell \mid MD$ (cf. [24, §2]), and it turns out that the same is true of ω_p . Finally, choose $\omega_{\infty} \in R_0^D(Mp)$ of norm -1 ; of course, ω_{∞} normalizes both $\Gamma_0^D(Mp)$ and $\Gamma_0^D(M)$.

Suppose that a semigroup $S \subset B^\times$ contains ω_{∞} and ω_{ℓ} for all primes $\ell \mid MDp$. Then define $W_{\infty} := T(\omega_{\infty})$ and $W_{\ell} := T(\omega_{\ell})$ in $\mathcal{H}(G, S)$; further, for every integer $m \mid MDp$ set $W_m := \prod_{\ell \mid m} W_{\ell}$. Note that $W_m = T(\omega_m)$ where $\omega_m := \prod_{\ell \mid m} \omega_{\ell}$. In particular, W_p belongs to $\mathcal{H}(\Gamma, S_p)$ and, in fact, $W_p = U_p$ in this Hecke algebra (see [19, §2.3] for details).

2.3. Quadratic forms. Let τ denote the generator of $\text{Gal}(F(\sqrt{d_K})/F)$. If $\psi : K \hookrightarrow B$ is an embedding of \mathbb{Q} -algebras then we may consider the quadratic form $Q_{\psi}(x, y) \in P_2(F)$ associated with ψ , which is defined by

$$Q_{\psi}(x, y) := cx^2 - 2axy - by^2 \quad \text{where } i_F(\psi(\sqrt{d_K})) = \begin{pmatrix} a & b \\ c & -a \end{pmatrix}.$$

We explicitly observe that $c \neq 0$. In fact, if $c = 0$ then $d = a^2$, whence $K \subset F$, a possibility which is ruled out by our choice of F .

We can factor the quadratic form $Q_{\psi}(x, y)$ as

$$Q_{\psi}(x, y) = c(x - z_{\psi}y)(x - \bar{z}_{\psi}y)$$

where $z_{\psi}, \bar{z}_{\psi} \in F(\sqrt{d_K}) - F$ are the roots of the equation $cz^2 - 2az - b = 0$ and $\tau(z_{\psi}) = \bar{z}_{\psi}$. The two roots z_{ψ} and \bar{z}_{ψ} are the only fixed points for the action of $\psi(K^\times)$ on $\mathbb{P}^1(F(\sqrt{d_K}))$ by Möbius transformations via i_F . We may also order z_{ψ} and \bar{z}_{ψ} by requiring that

$$\psi(\alpha) \begin{pmatrix} z_{\psi} \\ 1 \end{pmatrix} = \alpha \begin{pmatrix} z_{\psi} \\ 1 \end{pmatrix}, \quad \psi(\alpha) \begin{pmatrix} \bar{z}_{\psi} \\ 1 \end{pmatrix} = \tau(\alpha) \begin{pmatrix} \bar{z}_{\psi} \\ 1 \end{pmatrix}$$

for all $\alpha \in K$.

Let $\mathcal{H}_p := \mathbb{C}_p - \mathbb{Q}_p$ denote Drinfeld's p -adic half plane. The completion K_p of K at the prime (p) is the (unique, up to isomorphism) unramified quadratic extension of \mathbb{Q}_p . Since p is split in F , the completion of $F(\sqrt{d_K})$ with respect to the valuation induced by the embedding $\mathbb{Q} \hookrightarrow \mathbb{Q}_p$

chosen in (4) is again K_p . Thus z_ψ and \bar{z}_ψ can also be seen as points in $K_p - \mathbb{Q}_p$. Since there are canonical isomorphisms

$$\mathrm{Gal}(F(\sqrt{d_K})/F) \simeq \mathrm{Gal}(K/\mathbb{Q}) \simeq \mathrm{Gal}(K_p/\mathbb{Q}_p),$$

in the rest of the article we shall view τ as the generator of $\mathrm{Gal}(K_p/\mathbb{Q}_p)$ as well.

2.4. Darmon points on tori. For any abelian group A denote by $\mathcal{M}(A)$ the group of A -valued measures on $\mathbb{P}^1(\mathbb{Q}_p)$ and by $\mathcal{M}_0(A)$ those measures in $\mathcal{M}(A)$ with total mass 0. There is a canonical left action of $\mathrm{GL}_2(\mathbb{Q}_p)$ on $\mathcal{M}(A)$ and $\mathcal{M}_0(A)$ defined by the integration formula

$$\int_{\mathbb{P}^1(\mathbb{Q}_p)} \varphi(t) d(\gamma \cdot \nu)(t) := \int_{\mathbb{P}^1(\mathbb{Q}_p)} \varphi\left(\frac{at+b}{ct+d}\right) d\nu$$

for all step functions $\varphi : \mathbb{P}^1(\mathbb{Q}_p) \rightarrow \mathbb{C}_p$ and all $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{GL}_2(\mathbb{Q}_p)$. Then B^\times acts on $\mathcal{M}(A)$ and $\mathcal{M}_0(A)$ via i_p and the embedding $B \hookrightarrow B \otimes \mathbb{Q}_p$.

The group $H^1(\Gamma, \mathcal{M}_0(\mathbb{Q}))$ is endowed with actions of the Hecke algebra $\mathcal{H}(\Gamma, S_p)$ considered in §2.2 and of the involutions W_m and W_∞ . For any choice of sign \pm , define $H^1(\Gamma, \mathcal{M}_0(\mathbb{Q}))^{f, \pm}$ to be the subspace of $H^1(\Gamma, \mathcal{M}_0(\mathbb{Q}))$ consisting of those ξ such that $\xi|T = \theta_f(T)\xi$ for all $T \in \mathcal{H}(Mp)$ and $\xi|W_\infty = \pm\xi$ where $\theta_f : \mathcal{H}(Mp) \rightarrow \mathbb{Z}$ is the morphism associated with f . By [13, Proposition 25], the \mathbb{Q} -vector space $H^1(\Gamma, \mathcal{M}_0(\mathbb{Q}))^{f, \pm}$ has dimension 1. In [19, Section 4] we constructed an explicit generator for this vector space. Since we will need this description in the following computations, we will briefly review setting and results of [19].

Recall the identification between $H^1(\Gamma, \mathcal{M}_0(A))$ with $H^1(\Gamma, \mathcal{F}_{\mathrm{har}}(A))$ where, for any abelian group A , the symbol $\mathcal{F}_{\mathrm{har}}(A)$ stands for the abelian group of A -valued harmonic cocycles (see, e.g., [13, Lemma 27] and [19, §4.1]). Write \mathcal{T} for the Bruhat–Tits tree of $\mathrm{PGL}_2(\mathbb{Q}_p)$ (see [26, Ch. II, §1]), whose set of vertices (respectively, oriented edges) will be denoted by \mathcal{V} (respectively, \mathcal{E}). The groups B^\times and $(B \otimes \mathbb{Q}_p)^\times$ act on the left on \mathcal{T} via i_p . Let $v_* \in \mathcal{V}$ be the distinguished vertex corresponding to the maximal order $M_2(\mathbb{Z}_p)$ of $M_2(\mathbb{Q}_p)$, and say that a vertex $v \in \mathcal{V}$ is *even* (respectively, *odd*) if its distance from v_* is even (respectively, odd). Moreover, denote by \hat{v}_* the vertex corresponding to $\begin{pmatrix} \mathbb{Z}_p & p^{-1}\mathbb{Z}_p \\ p\mathbb{Z}_p & \mathbb{Z}_p \end{pmatrix}$, fix the edge $e_* = (v_*, \hat{v}_*) \in \mathcal{E}$ and say that an edge $e = (v_1, v_2) \in \mathcal{E}$ is *even* (respectively, *odd*) if v_1 is even (respectively, odd). Write \mathcal{E}^+ for the set of even vertices of \mathcal{T} . Finally, if $e = (v_1, v_2)$ write \bar{e} for the reversed edge (v_2, v_1) .

Choose a harmonic (in the sense of [19, Definition 4.6]) system of representatives $\mathcal{Y}_{\mathrm{har}}$ for the cosets in $\Gamma_0^D(Mp) \backslash \Gamma$, which can be written as $\mathcal{Y}_{\mathrm{har}} = \{\gamma_e\}_{e \in \mathcal{E}^+}$ with $\gamma_e \in \Gamma$ such that $\gamma_e(e) = e_*$. Set $H_E := H_1(E(\mathbb{C}), \mathbb{Z})$ and write H_E^\pm for the \pm -eigenspace for complex conjugation acting on H_E . Fix an isomorphism $H_E^\pm \simeq \mathbb{Z}$. Keeping in mind that H_E^\pm is (canonically isomorphic to) a quotient of $\Gamma_0^D(Mp)$, define, as in [19, Definition 4.2], the *universal 1-cochain* $\mu_{\mathrm{univ}}^{\mathcal{Y}} : \Gamma \rightarrow \mathcal{F}_{\mathrm{har}}(\mathbb{Z})$ associated with \mathcal{Y} by the following rules:

- if $\gamma \in \Gamma$ and $e \in \mathcal{E}^+$ let $g_{\gamma, e} \in \Gamma_0^D(Mp)$ be defined by the equation $\gamma_e \gamma = g_{\gamma, e} \gamma_{\gamma^{-1}(e)}$, then set $\mu_{\mathrm{univ}}^{\mathcal{Y}}(\gamma)(e) := [g_{\gamma, e}] \in H_E^\pm \simeq \mathbb{Z}$;
- if $\gamma \in \Gamma$ and $e \notin \mathcal{E}^+$ then set $\mu_{\mathrm{univ}}^{\mathcal{Y}}(\gamma)(e) := -\mu_{\mathrm{univ}}^{\mathcal{Y}}(\gamma)(\bar{e})$.

Harmonic systems always exist, by [19, Proposition 4.8]. The canonical generator $\boldsymbol{\mu}_f^\pm$ of $H^1(\Gamma, \mathcal{M}_0(\mathbb{Q}))$ is obtained by fixing a prime number $r \nmid MDp$, applying the Hecke operator $t_r := T_r - (r+1)$ to the class of $\mu_{\mathrm{univ}}^{\mathcal{Y}}$ in $H^1(\Gamma, \mathcal{F}_{\mathrm{har}}(\mathbb{Z}))$ and then using the isomorphism

$$H^1(\Gamma, \mathcal{M}_0(\mathbb{Q})) \simeq H^1(\Gamma, \mathcal{M}_0(\mathbb{Z})) \otimes_{\mathbb{Z}} \mathbb{Q}.$$

As shown in [19, Lemma 4.11], the resulting class μ_f^\pm independent of the choice of \mathcal{Y}_{har} as above. We also fix a representative μ_f^\pm of μ_f^\pm .

Let $T := \mathbb{G}_m$ denote the multiplicative group (viewed as a functor on commutative \mathbb{Q} -algebras). As explained in [13, Section 6], the measure μ_f^\pm can be used to define a lattice L in $T(\mathbb{C}_p)$ as the image of the composition

$$(5) \quad H_2(\Gamma, \mathbb{Z}) \longrightarrow H_1(\Gamma, \text{Div}^0(\mathcal{H}_p)) \longrightarrow T(\mathbb{C}_p).$$

Here the first map is extracted from the long exact sequence in homology associated with the short exact sequence

$$0 \longrightarrow \text{Div}^0(\mathcal{H}_p) \longrightarrow \text{Div}(\mathcal{H}_p) \xrightarrow{\text{deg}} \mathbb{Z} \longrightarrow 0$$

and the second is the integration map described in [19, §5.1]. In fact, the lattice L is contained in $T(\mathbb{Q}_p)$ and is Hecke-stable ([19, Proposition 6.1]). Fix $z \in K_p - \mathbb{Q}_p$ and let $\mathbf{d}_z \in H^2(\Gamma, T(\mathbb{C}_p))$ be the cohomology class represented by the 2-cocycle

$$(6) \quad d_z : \Gamma \times \Gamma \longrightarrow T(K_p), \quad (\gamma_1, \gamma_2) \longmapsto \int_{\mathbb{P}^1(\mathbb{Q}_p)} \frac{s - \gamma_1^{-1}(z)}{s - z} d\mu_{f, \gamma_2}^\pm(s).$$

(See [19, §5.1] for the definition of the multiplicative integral.) The class \mathbf{d}_z does not depend on the choice of the representative μ_f^\pm of μ_f^\pm . Write \bar{d}_z for the composition of d_z with the canonical projection onto $T(K_p)/L$. By construction, L is the smallest subgroup of $T(\mathbb{Q}_p)$ such that \mathbf{d}_z becomes trivial in $H^2(\Gamma, T(\mathbb{C}_p)/L)$, so there exists

$$\beta_z : \Gamma \longrightarrow T/L$$

such that

$$\bar{d}_z(\gamma_1, \gamma_2) = \beta_z(\gamma_1 \gamma_2) \cdot \beta_z(\gamma_1)^{-1} \cdot \beta_z(\gamma_2)^{-1}$$

for all $\gamma_1, \gamma_2 \in \Gamma$ (the action of Γ on H_E^\pm used in [19] is the trivial one). Note that β_z is well defined only up to elements in $\text{Hom}(\Gamma, T/L)$. To deal with this ambiguity, recall that Γ^{ab} is a finite group ([20, Proposition 2.1]). Thus, if t is the exponent of Γ^{ab} then $t \cdot \beta_z$ is well defined (i.e., it depends only on the choice of a representative for μ_f^\pm).

Let H_K^+ denote the narrow Hilbert class field of K and set $G_K^+ := \text{Gal}(H_K^+/K)$. Write $\text{Pic}^+(\mathcal{O}_K)$ for the narrow class group of the ring of integers \mathcal{O}_K of K . The reciprocity map of global class field theory induces an isomorphism

$$(7) \quad \text{rec} : \text{Pic}^+(\mathcal{O}_K) \xrightarrow{\cong} G_K^+.$$

Let $\psi : K \hookrightarrow B$ be an optimal embedding of \mathcal{O}_K into $R_0^D(M)$, i.e. an embedding of K into B such that $\psi(\mathcal{O}_K) = \psi(K) \cap R_0^D(M)$, and write $\text{Emb}(\mathcal{O}_K, R_0^D(M))$ for the set of such embeddings. The group $\Gamma_0^D(M)$ acts on $\text{Emb}(\mathcal{O}_K, R_0^D(M))$ by conjugation.

Recall the points z_ψ and \bar{z}_ψ defined in §2.3. By Dirichlet's unit theorem, the abelian group of units in \mathcal{O}_K^\times of norm 1 is free of rank one. Choose a generator ε_K of this group such that $\varepsilon_K > 1$ with respect to the fixed embedding $K \hookrightarrow \mathbb{R}$ of §2.1 and set $\gamma_\psi := \psi(\varepsilon_K) \in \Gamma_0^D(M)$.

Definition 2.3. The *Darmon points* on $T(K_p)/L$ are the points

$$\mathcal{P}_\psi := t \cdot \beta_{z_\psi}(\gamma_\psi) \in T(K_p)/L$$

where ψ varies in $\text{Emb}(\mathcal{O}_K, R_0^D(M))$.

Let ψ be as before. While, thanks to the multiplicative factor t , the point \mathcal{P}_ψ does not depend on the choice of a map β_{z_ψ} splitting the 2-cocycle d_{z_ψ} modulo L , it might *a priori* depend on the choice of the representative μ_f^\pm of $\boldsymbol{\mu}_f^\pm$. In fact, this is not the case: a direct calculation explained in [20] shows that the point \mathcal{P}_ψ is independent of the choice of a representative of $\boldsymbol{\mu}_f^\pm$ and depends only on the $\Gamma_0^D(M)$ -conjugacy class of ψ . Although, in light of this result, the symbol $\mathcal{P}_{[\psi]}$ would be more appropriate, in order not to burden our notation we will continue to write \mathcal{P}_ψ for the Darmon points. However, the reader should always bear in mind that $\mathcal{P}_\psi = \mathcal{P}_{\psi'}$ whenever ψ and ψ' are $\Gamma_0^D(M)$ -conjugate.

2.5. Galois action on optimal embeddings. As in [20, §4.2], for every prime ℓ dividing MD fix orientations of $R_0^D(M)$ and \mathcal{O}_K at ℓ , i.e., ring homomorphisms

$$\mathfrak{D}_\ell : R_0^D(M) \longrightarrow \mathbb{F}_{\ell^\delta}, \quad \mathfrak{o}_\ell : \mathcal{O}_K \longrightarrow \mathbb{F}_{\ell^\delta}$$

where $\delta = 1$ if $\ell|M$ and $\delta = 2$ if $\ell|D$. There are exactly two orientations of \mathcal{O}_K at every $\ell|MD$.

Definition 2.4. 1) Two embeddings $\psi, \psi' \in \text{Emb}(\mathcal{O}_K, R_0^D(M))$ are said to have *the same orientation* at a prime $\ell|MD$ if $\mathfrak{D}_\ell \circ (\psi|_{\mathcal{O}_K}) = \mathfrak{D}_\ell \circ (\psi'|_{\mathcal{O}_K})$ and are said to have *opposite orientations* at ℓ otherwise.

2) An embedding $\psi \in \text{Emb}(\mathcal{O}_K, R_0^D(M))$ is said to be *oriented* if $\mathfrak{D}_\ell \circ (\psi|_{\mathcal{O}_K}) = \mathfrak{o}_\ell$ for all primes $\ell|MD$.

We denote the set of oriented optimal embeddings of \mathcal{O}_K into $R_0^D(M)$ by $\mathcal{E}(\mathcal{O}_K, R_0^D(M))$. Note that the action of $\Gamma_0^D(M)$ on $\text{Emb}(\mathcal{O}_K, R_0^D(M))$ by conjugation restricts to an action on $\mathcal{E}(\mathcal{O}_K, R_0^D(M))$. Moreover, if $\psi \in \mathcal{E}(\mathcal{O}_K, R_0^D(M))$ then

$$\psi^* := \omega_\infty \psi \omega_\infty^{-1}$$

is in $\mathcal{E}(\mathcal{O}_K, R_0^D(M))$ too.

The next lemma, the proof of which we omit, collects some basic properties of (oriented) optimal embeddings.

Lemma 2.5. *Take $\psi, \psi' \in \text{Emb}(\mathcal{O}_K, R_0^D(M))$.*

- (1) *If ψ and ψ' have opposite orientations at a prime dividing MD then at most one of them is oriented in the sense of Definition 2.4.*
- (2) *If ℓ is a prime dividing MD then ψ and $\omega_\ell \psi \omega_\ell^{-1}$ have opposite orientations at ℓ and the same orientation at all the other primes dividing MD .*
- (3) *The embeddings ψ and $\psi \circ \tau$ have opposite orientations at all the primes dividing MD .*

The principal ideal $(\sqrt{d_K})$ is a proper \mathcal{O}_K -ideal of K , so we can consider its class \mathfrak{D}_K in $\text{Pic}^+(\mathcal{O}_K)$; define $\sigma_K := \text{rec}(\mathfrak{D}_K) \in G_K^+$. From here to the end of this subsection, for notational convenience set $R := R_0^D(M)$. We want to recall the actions of $\text{Pic}^+(\mathcal{O}_K)$ and G_K^+ on the sets of $\Gamma_0^D(M)$ -conjugacy classes of optimal and oriented optimal embeddings of \mathcal{O}_K into R . For details, see [20, Proposition 4.2]. Let $\mathfrak{a} \subset \mathcal{O}_K$ be an ideal representing a class $[\mathfrak{a}] \in \text{Pic}^+(\mathcal{O}_K)$ and let $\psi \in \text{Emb}(\mathcal{O}_K, R)$. Since the quaternion algebra B is indefinite, the left R -ideal $R\psi(\mathfrak{a})$ is principal; let $a \in R$ be a generator of this ideal with positive reduced norm, which is unique up to elements in $\Gamma_0^D(M)$. The right action of $\psi(\mathcal{O}_K)$ on $R\psi(\mathfrak{a})$ shows that $\psi(\mathcal{O}_K)$ is contained in the right order of $R\psi(\mathfrak{a})$, which is equal to $a^{-1}Ra$. It can be checked that this recipe induces a well-defined action of $\text{Pic}^+(\mathcal{O}_K)$ on conjugacy classes of embeddings given by

$$[\mathfrak{a}] \star [\psi] := [a\psi a^{-1}] \in \text{Emb}(\mathcal{O}_K, R)/\Gamma_0^D(M).$$

Notice that if $\mathfrak{a} = (\sqrt{d_K})$ then we can take $a = \omega_\infty \cdot \psi(\sqrt{d_K})$, whence

$$(8) \quad \mathfrak{D}_K \star [\psi] = [\omega_\infty \psi \omega_\infty^{-1}] = [\psi^*].$$

Recalling the reciprocity map of (7), for all $\sigma \in G_c^+$ and $[\psi] \in \text{Emb}(\mathcal{O}_K, R)/\Gamma_0^D(M)$ define

$$\sigma \star [\psi] := \text{rec}^{-1}(\sigma) \star [\psi].$$

In particular, it follows from (8) that $\sigma_K \star [\psi] = [\psi^*]$ for all $\psi \in \text{Emb}(\mathcal{O}_K, R)$. Finally, for every $\sigma \in G_c^+$ and every $\psi \in \text{Emb}(\mathcal{O}_K, R)$ choose an element

$$\sigma \star \psi \in \sigma \star [\psi].$$

If now ψ is an oriented optimal embedding (with respect to the orientations of \mathcal{O}_K and R fixed above) then, with notation as before, the Eichler order $a^{-1}Ra$ inherits an orientation from the one of R . Actually, it can be checked that we get an induced action of $\text{Pic}^+(\mathcal{O}_K)$ (and G_K^+) on the set $\mathcal{E}(\mathcal{O}_K, R)/\Gamma_0^D(M)$, and this action turns out to be free and transitive. In light of this, it is convenient to describe a (non-canonical) bijection between $\mathcal{E}(\mathcal{O}_K, R)/\Gamma_0^D(M)$ and G_K^+ . To this end, fix once and for all an auxiliary embedding $\psi_0 \in \mathcal{E}(\mathcal{O}_K, R)$ and define $E : G_K^+ \rightarrow \mathcal{E}(\mathcal{O}_K, R)/\Gamma_0^D(M)$ by $E(\sigma) := \sigma \star [\psi_0]$. Finally, set

$$G := E^{-1} : \mathcal{E}(\mathcal{O}_K, R)/\Gamma_0^D(M) \longrightarrow G_K^+,$$

which is a bijection satisfying

$$(9) \quad G([\psi^*]) = \sigma_K \cdot G([\psi])$$

for all $\psi \in \mathcal{E}(\mathcal{O}_K, R)$. Now choose for every $\sigma \in G_K^+$ an embedding $\psi_\sigma \in E(\sigma)$, so that the family $\{\psi_\sigma\}_{\sigma \in G_K^+}$ is a set of representatives of the $\Gamma_0^D(M)$ -conjugacy classes of oriented optimal embeddings of \mathcal{O}_K into R . If $\gamma, \gamma' \in R$ write $\gamma \sim \gamma'$ to indicate that γ and γ' are in the same $\Gamma_0^D(M)$ -conjugacy class, and adopt a similar notation for (oriented) optimal embeddings of \mathcal{O}_K into R . For all $\sigma, \sigma' \in G_K^+$ one has

$$\sigma \star [\psi_{\sigma'}] = \sigma \star E(\sigma') = \sigma \star (\sigma' \star [\psi_0]) = (\sigma\sigma') \star [\psi_0] = E(\sigma\sigma'),$$

which means that

$$\sigma \star \psi_{\sigma'} \sim \psi_{\sigma\sigma'}.$$

Furthermore, since $G([\psi_\sigma^*]) = \sigma \cdot G([\psi_\sigma]) = \sigma_K \sigma$, by equality (9), we deduce that

$$\psi_\sigma^* \sim \psi_{\sigma_K \sigma}$$

for all $\sigma \in G_K^+$.

To conclude this subsection, we prove a lemma which will be useful later on. As above, for every prime ℓ dividing MD fix an orientation $\mathfrak{D}_\ell : R \rightarrow \mathbb{F}_{\ell^s}$ of the Eichler order R at ℓ .

Lemma 2.6. *Let $\psi, \psi' \in \text{Emb}(\mathcal{O}_K, R)$. If ψ and ψ' have the same orientation at all primes dividing MD then there exists a unique $\sigma \in G_K^+$ such that $\psi \sim \sigma \star \psi'$.*

Proof. For every prime $\ell | MD$ choose the unique orientation of \mathcal{O}_c at ℓ such that ψ and ψ' become oriented optimal embeddings. Since the action of G_K^+ on $\Gamma_0^D(M)$ -conjugacy classes of oriented optimal embeddings is free and transitive, there exists a unique $\sigma \in G_K^+$ such that $[\psi] = \sigma \star [\psi']$, and the claim follows. \square

Remark 2.7. The element $\sigma \in G_K^+$ whose existence is established in Lemma 2.6 depends on the chosen orientations \mathfrak{D}_ℓ of R .

2.6. Real conjugation on Darmon points. In the sequel we will need to understand the action of the local Galois group $\text{Gal}(K_p/\mathbb{Q}_p)$ on Darmon points. As in §2.3, denote by τ the generator of $\text{Gal}(K_p/\mathbb{Q}_p)$. We start with the following

Lemma 2.8. $\mathcal{P}_{\psi \circ \tau} = \tau(\mathcal{P}_\psi)^{-1}$.

Proof. First recall that \bar{z}_ψ satisfies the condition $\psi(\alpha)\begin{pmatrix} \bar{z}_\psi \\ 1 \end{pmatrix} = \bar{\alpha}\begin{pmatrix} \bar{z}_\psi \\ 1 \end{pmatrix}$, so $z_{\psi \circ \tau} = \bar{z}_\psi$. Since $(\psi \circ \tau)(u) = \psi(u^{-1}) = \gamma_\psi^{-1}$, it follows that $\mathcal{P}_{\psi \circ \tau} = t \cdot \beta_{\bar{z}_\psi}(\gamma_\psi^{-1})$. Observe that the lattice L , being contained in $T(\mathbb{Q}_p)$, inherits the trivial action of τ . With notation as before, if β_z splits \bar{d}_z then $\tau \circ \beta_z$ splits $\bar{d}_{\tau(z)}$, so it follows that

$$(10) \quad \mathcal{P}_{\psi \circ \tau} = t \cdot \beta_{\bar{z}_\psi}(\gamma_\psi^{-1}) = t \cdot \tau(\beta_{z_\psi}(\gamma_\psi^{-1})) = \tau(t \cdot \beta_{z_\psi}(\gamma_\psi^{-1})).$$

It remains to show that $\beta_{z_\psi}(\gamma_\psi^{-1}) = \beta_{z_\psi}(\gamma_\psi)^{-1}$. To prove this equality, note that, since $\gamma_\psi(z_\psi) = z_\psi$ and $\mu_{f,\gamma}^\pm$ has total mass 0 for all $\gamma \in \Gamma$, one has

$$d_{z_\psi}(\gamma_\psi, \gamma) = \int_{\mathbb{P}^1(\mathbb{Q}_p)} \frac{s - \gamma_\psi^{-1}(z_\psi)}{s - z_\psi} d\mu_{f,\gamma}^\pm = 1$$

for all $\gamma \in \Gamma$. Thus we obtain that

$$(11) \quad 1 = d_{z_\psi}(\gamma_\psi, \gamma_\psi^{-1}) = \beta_{z_\psi}(1) \cdot \beta_{z_\psi}(\gamma_\psi)^{-1} \cdot \beta_{z_\psi}(\gamma_\psi^{-1})^{-1}.$$

On the other hand, one has

$$1 = d_{z_\psi}(\gamma_\psi, 1) = \beta_{z_\psi}(\gamma_\psi) \cdot \beta_{z_\psi}(\gamma_\psi)^{-1} \cdot \beta_{z_\psi}(1)^{-1},$$

from which we also get $\beta_{z_\psi}(1) = 1$. Substituting in (11) gives $\beta_{z_\psi}(\gamma_\psi^{-1}) = \beta_{z_\psi}(\gamma_\psi)^{-1}$, as claimed. Equality (10) then becomes

$$\mathcal{P}_{\psi \circ \tau} = \tau(t \cdot \beta_{z_\psi}(\gamma_\psi)^{-1}) = \tau(\mathcal{P}_\psi)^{-1},$$

completing the proof. \square

By part (3) of Lemma 2.5, if $\psi \in \text{Emb}(\mathcal{O}_K, R_0^D(M))$ then ψ and $\psi \circ \tau$ have opposite orientations at every prime dividing MD . Since conjugation by ω_{MD} reverses the orientation at every prime (by (2) of the same lemma) and ω_{MD} normalizes $\Gamma_0^D(M)$, Lemma 2.6 ensures that there exists a unique $\sigma_\psi \in G_K^+$ such that

$$(12) \quad \psi \circ \tau \sim \omega_{MD}(\sigma_\psi \star \psi) \omega_{MD}^{-1}.$$

Now we can prove

Proposition 2.9. *Let w_{MD} denote the eigenvalue of W_{MD} on H_E^\pm . Then*

$$\tau(\mathcal{P}_\psi) = \mathcal{P}_{\sigma_\psi \star \psi}^{-w_{MD}}$$

where $\sigma = \sigma_\psi$ is the element of G_K^+ appearing in (12).

Proof. We first notice that the integration map appearing in (5) is equivariant for the action of W_{MD} . This can be shown as in [19, Proposition 5.1], where the same result is proved for W_p and W_∞ , so we just sketch the arguments in this case. Observe that $\omega_{MD} \in R_0^D(Mp)$ satisfies $\Gamma \omega_{MD} = \omega_{MD} \Gamma$ and $\Gamma_0^D(Mp) \omega_{MD} = \omega_{MD} \Gamma_0^D(Mp)$. Let \mathcal{Y}_{rad} be a fixed radial system of representatives for $\Gamma_0^D(Mp) \backslash \Gamma$ used to compute a representative μ_f^\pm of $\mu_{f,\gamma}^\pm$ (so $\mu_{f,\gamma}^\pm = t_r \mu_{\text{univ},\gamma}^{\mathcal{Y}_{\text{rad}}}$). Introduce the system

$$\mathcal{Y}'_{\text{rad}} := \{\omega_{MD}^{-1} \gamma_{\omega(e)} \omega_{MD}\}_{e \in \mathcal{E}^+}.$$

As noticed in [19, §5.2], $\mathcal{Y}'_{\text{rad}}$ is again radial and thus can be used to define a representative of μ_f^\pm . Since the Hecke action is defined at the cochain level, one easily verifies that

$$(13) \quad \int_{\mathbb{P}^1(\mathbb{Q}_p)} \frac{t - \omega_{MD}\gamma_1^{-1}(z_\psi)}{t - \omega_{MD}z_\psi} dt_r \mu_{\text{univ}, \omega_{MD}\gamma_2\omega_{MD}^{-1}}^{\mathcal{Y}'_{\text{rad}}} = W_{MD} \cdot \int_{\mathbb{P}^1(\mathbb{Q}_p)} \frac{t - \gamma_1^{-1}(z_\psi)}{t - z_\psi} dt_r \mu_{\text{univ}, \gamma_2}^{\mathcal{Y}'_{\text{rad}}}$$

for all $\gamma_1, \gamma_2 \in \Gamma$. To simplify notations, set $\mu'_\gamma := t_r \mu_{\text{univ}, \gamma}^{\mathcal{Y}'_{\text{rad}}}$ and $\psi' := \omega_{MD}(\sigma \star \psi)\omega_{MD}^{-1}$, with $\sigma = \sigma_\psi$ as in (12). Then $z_{\psi'} = \omega_{MD}(z_\psi)$ and $\bar{z}_{\psi'} = \omega_{MD}(\bar{z}_\psi)$. By (13), if β'_{z_ψ} splits the 2-cocycle

$$(\gamma_1, \gamma_2) \mapsto \int_{\mathbb{P}^1(\mathbb{Q}_p)} \frac{t - \gamma_1^{-1}(z_\psi)}{t - z_\psi} d\mu'_{\gamma_2}$$

then $\gamma \mapsto w_{MD}\beta'_{z_\psi}(\omega_{MD}^{-1}\gamma\omega_{MD})$ splits the 2-cocycle $d_{\omega_{MD}z_\psi} = d_{z_{\psi'}}$ defined in (6) (here use the fact that W_{MD} acts on T as multiplication by w_{MD}). It follows that

$$w_{MD} \cdot \beta'_{z_\psi}(\omega_{MD}^{-1}\gamma\omega_{MD}) = \beta_{z_{\psi'}}(\gamma) + \varphi(\gamma)$$

where $\varphi : \Gamma \rightarrow T(K_p)/L_T$ is a homomorphism. Therefore we get

$$(14) \quad \mathcal{P}_{\psi'} = t \cdot \beta_{z_{\psi'}}(\gamma_{\psi'}) = w_{MD}t \cdot \beta'_{z_\psi}(\omega_{MD}^{-1}\gamma_{\psi'}\omega_{MD}) = w_{MD}t \cdot \beta'_{z_\psi}(\gamma_{\sigma\star\psi}).$$

As remarked in §2.4, $t \cdot \beta'_{z_\psi}(\gamma_{\sigma\star\psi}) = t \cdot \beta_{z_\psi}(\gamma_{\sigma\star\psi})$, hence (14) implies that

$$(15) \quad \mathcal{P}_{\psi'} = \mathcal{P}_{\sigma\star\psi}^{w_{MD}}.$$

From Lemma 2.8 and (12) we obtain

$$(16) \quad \tau(\mathcal{P}_{\psi'}) = \mathcal{P}_{\psi'}^{-1},$$

and then combining (15) with (16) yields the result. \square

2.7. Darmon points on elliptic curves. Define a p -adic logarithm on the quotient $T(\mathbb{C}_p)/L$ as follows. As in the introduction, let $q \in p\mathbb{Z}_p$ be Tate's period for E at p and let \log_q be the branch of the p -adic logarithm such that $\log_q(q) = 0$. Since $T(\mathbb{C}_p) \simeq \mathbb{C}_p^\times$, the fact that \log_q induces a map on $T(\mathbb{C}_p)/L$ is an immediate consequence of the following

Lemma 2.10. $\log_q(L) = 0$.

Proof. By [19, Theorem 7.16], the lattices L and $\langle q \rangle$ are homothetic, i.e., $L \cap \langle q \rangle$ has finite index both in L and in $\langle q \rangle$. If $n := [L : L \cap \langle q \rangle]$ then $L^n \subset \langle q \rangle$, so $\log_q(L^n) = 0$ by the choice of \log_q . Since \log_q is a homomorphism from \mathbb{C}_p^\times to \mathbb{C}_p , the lemma is proved. \square

From now on, by an abuse of notation we will use the symbol \log_q for both maps induced by \log_q on $T(\bar{K}_p)/L$ and $\bar{K}_p^\times/\langle q \rangle$; these two maps are K_p -valued on the K_p -rational points.

As in the proof of the lemma above, set $n := [L : L \cap \langle q \rangle]$. Raising to the n -th power induces a Galois-equivariant isogeny of p -adic tori

$$(17) \quad T(\bar{K}_p)/L \longrightarrow \bar{K}_p^\times/\langle q \rangle$$

defined over K_p , and then composing (17) with Tate's analytic uniformization Φ_{Tate} introduced in (3) yields a Galois-equivariant isogeny

$$\varphi_E : T(\bar{K}_p)/L \longrightarrow E(\bar{K}_p).$$

As in [4, p. 346], define a homomorphism

$$\log_E : E(K_p) \longrightarrow K_p$$

by the rule $\log_E(P) := \log_q(\Phi_{\text{Tate}}^{-1}(P))$ for all $P \in E(K_p)$. It follows that

$$(18) \quad \log_E(\varphi_E(x)) = n \log_q(x) \quad \text{for all } x \in T(K_p)/L.$$

For every $\psi \in \text{Emb}(\mathcal{O}_c, R_0^D(M))$ define

$$P_\psi^\pm := \varphi_E(\mathcal{P}_\psi) \in E(K_p).$$

Remark 2.11. Since the same is true of \mathcal{P}_ψ , the point P_ψ^\pm is independent of the representative of μ_f^\pm , and $P_\psi^\pm = P_{\psi'}^\pm$ if ψ and ψ' are $\Gamma_0^D(M)$ -conjugate. However, as made explicit by the notation, P_ψ^\pm does depend on the choice of sign \pm .

3. EXPLICIT FORMULAS FOR DARMON POINTS

The aim of this section is to use a control theorem of Greenberg–Stevens type in the quaternionic setting (Theorem 3.1) to obtain a canonical lift $\tilde{\mu}_f^\pm$ of μ_f^\pm . The role of this explicit lift is twofold: on the one hand, it is used to perform computations with Darmon points (cf. Corollary 3.6); on the other hand, in Section 4 it allows us to define a p -adic L -function which, thanks to Popa’s work ([25]), encodes (for suitable integers k) special values of the L -functions of the Hida forms f_k over K twisted by genus characters of K .

3.1. Modular forms on quaternion algebras. For any ring R and any integer $n \geq 0$ let $P_n(R) := \text{Sym}^n(R)$ denote the R -module of homogeneous polynomials $P(x, y)$ in two variables x, y of degree $n \geq 0$ with coefficients in R . It is equipped with a right action of the group $\text{GL}_2(R)$ given by the rule

$$(P|\gamma)(x, y) := P(ax + by, cx + dy) \quad \text{for } \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix}.$$

The R -linear dual $V_n(R)$ of $P_n(R)$ is then endowed with a left action of $\text{GL}_2(R)$ by the formula

$$(\gamma \cdot \phi)(P) := \phi(P|\gamma).$$

Finally, if R is a field extension of F then $P_n(R)$ (respectively, $V_n(R)$) is equipped with a right (respectively, left) action of B^\times via the map i_F and the rules above.

Let g be a cusp form of level $\Gamma_0^D(Mp)$ and weight k , suppose that g is an eigenform for all the Hecke operators in the usual Hecke algebra \mathfrak{h}_k and write $\lambda : \mathfrak{h}_k \rightarrow \mathbb{C}$ for the corresponding morphism. Let L be a subfield of \mathbb{C} containing the image of λ . If ι denotes complex conjugation, define $H^1(X_0^D(Mp), V_{k-2}(L))^{g, \pm}$ to be the L -subspace of $H^1(X_0^D(Mp), V_{k-2}(L))$ consisting of the elements ξ such that $\xi|T = \lambda(T)\xi$ for all $T \in \mathfrak{h}_k$ and $\xi|\iota = \pm\xi$. By [21], we know that

$$\dim_L \left(H^1(X_0^D(Mp), V_{k-2}(L))^{g, \pm} \right) = 1.$$

Recall that there is a canonical isomorphism

$$H^1(X_0^D(Mp), V_{k-2}(\mathbb{C})) \simeq H^1(\Gamma_0^D(Mp), V_{k-2}(\mathbb{C}))$$

which is equivariant for the action of the involution ι . Then the complex vector space $H^1(\Gamma_0^D(Mp), V_{k-2}(\mathbb{C}))^{g, \pm}$ is spanned by the projection on the \pm -eigenspace for ι of the class represented by the 1-cocycle $\gamma \mapsto v(g)_\gamma$ with

$$v(g)_\gamma(P) := \int_\tau^{\gamma(\tau)} g(z)P(z, 1)dz$$

(this class does not depend on the choice of $\tau \in \mathcal{H}$). For details, see [27, §8.2].

3.2. Hida families. Recall the group \mathcal{X} in (1) and the Hida family $f_\infty = \sum_{n=1}^{\infty} a_n(\kappa)q^n$ of f in (2), where the p -adic analytic functions a_n are all defined in a p -adic neighbourhood \mathbb{U}_f of 2 in \mathcal{X} . For a fixed even integer $k \geq 2$ in \mathbb{U}_f write f_k for the normalized eigenform of weight k on $\Gamma_0(MDp)$ whose q -expansion is obtained by setting $\kappa = k$ in f_∞ ; in particular, $f_2 = f$ (see [3, §1.2] for details). Shrinking it if necessary, we can assume that \mathbb{U}_f is contained in the residue class of 2 modulo $p - 1$. For any even integer $k \in \mathbb{U}_f$ denote by F_k the finite extension of \mathbb{Q}_p generated by the Fourier coefficients of f_k . In the notations of §3.1, define the one-dimensional F_k -vector space

$$\mathbb{W}_{f_k}^\pm := H^1(\Gamma_0^D(Mp), V_{k-2}(F_k))^{f_k, \pm},$$

then fix a generator ϕ_k^\pm of $\mathbb{W}_{f_k}^\pm$ as well as a representative ϕ_k^\pm of ϕ_k^\pm .

3.3. The control theorem. Let $\mathbb{X} := (\mathbb{Z}_p^2)'$ denote the set of primitive vectors in $\mathbb{Y} := \mathbb{Z}_p^2$. For any \mathbb{Z}_p -module G let $\tilde{\mathbb{D}}(G)$ be the group of G -valued measures on \mathbb{Y} and let $\mathbb{D}(G)$ be the subgroup consisting of those measures which are supported on \mathbb{X} . The left action of $\Sigma := \mathrm{GL}_2(\mathbb{Q}_p) \cap \mathrm{M}_2(\mathbb{Z}_p)$ on $\tilde{\mathbb{D}}(G)$ considered in [19] is given by the integration formula

$$\int_{\mathbb{Y}} \varphi(x, y) d(\gamma \cdot \nu)(x, y) = \int_{\mathbb{Y}} \varphi(\gamma \cdot (x, y)) d\nu(x, y)$$

with $\gamma \cdot (x, y) := (ax + by, cx + dy)$ for all step functions $\varphi : \mathbb{Y} \rightarrow G$, all measures $\nu \in \tilde{\mathbb{D}}(G)$ and all matrices $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Sigma$. This action also induces an action of Σ on $\mathbb{D}(G)$ (see [19, Lemma 7.4]). Denote by $\pi : \mathbb{X} \rightarrow \mathbb{P}^1(\mathbb{Q}_p)$ the fibration defined by $(a, b) \mapsto a/b$, so that we obtain a canonical map

$$\pi_* : H^1(\Gamma_0^D(M), \mathbb{D}(\mathbb{Q}_p)) \longrightarrow H^1(\Gamma_0^D(M), \mathcal{M}(\mathbb{Q}_p)).$$

To simplify the notation, write $\mathbb{D} := \mathbb{D}(\mathbb{Z}_p)$. The group $H^1(\Gamma_0^D(M), \mathbb{D})$ comes equipped with an action of the Hecke algebra $\mathcal{H}(M)$, which allows us to define its *ordinary submodule* as

$$\mathbb{W} := \bigcap_{n=1}^{\infty} H^1(\Gamma_0^D(M), \mathbb{D}) | T_p^n$$

where $T_p \in \mathcal{H}(M)$ is the p -th Hecke operator. One can show that T_p acts invertibly on \mathbb{W} and that $H^1(\Gamma_0^D(M), \mathbb{D})$ decomposes as a direct sum of \mathbb{W} and a submodule on which T_p acts nilpotently. Finally, let

$$\Lambda := \mathbb{Z}_p[[1 + p\mathbb{Z}_p]] \simeq \mathbb{Z}_p[[T]]$$

be the Iwasawa algebra. Following the constructions in [14, Sections 1 and 5], one endows \mathbb{W} with a structure of (finitely generated) Λ -module.

The following Control Theorem is probably known to experts. To state this result, let us introduce a minimal set of notations. For details, proofs, and a dictionary between classical Hida families and their quaternionic counterparts, see [18, Sections 5 and 6].

As in [18, §5.3], let \mathcal{R} be the integral closure of Λ in the primitive component to which f belongs. The ring \mathcal{R} is a complete local noetherian domain, finitely generated as a Λ -module and equipped with a canonical map $\eta : \mathfrak{h}_\infty^{\mathrm{ord}} \rightarrow \mathcal{R}$ of Λ -algebras (where $\mathfrak{h}_\infty^{\mathrm{ord}}$ is Hida's ordinary Hecke algebra: see [18, §5.1]). If $k \in \mathbb{U}_f$ is an even positive integer there exists a continuous morphism $\vartheta_k : \mathcal{R} \rightarrow \bar{\mathbb{Q}}_p$ whose restriction to group-like elements in Λ is the character $x \mapsto x^{k-2}$ and such that $f_k = \sum_{n=1}^{\infty} (\vartheta_k \circ \eta)(T_n)q^n$, where $T_n \in \mathfrak{h}_\infty^{\mathrm{ord}}$ is the n -th Hecke operator. Set $\mathfrak{p}_k := \ker(\vartheta_k)$; if $\mathcal{R}_{\mathfrak{p}_k}$ is the localization of \mathcal{R} at \mathfrak{p}_k , we have $\mathcal{R}_{\mathfrak{p}_k}/\mathfrak{p}_k \mathcal{R}_{\mathfrak{p}_k} \simeq F_k$. Define

$$\mathbb{W}_{\mathfrak{p}_k}^\pm := \mathbb{W}^\pm \otimes_\Lambda \mathcal{R}_{\mathfrak{p}_k}$$

where \mathbb{W}^\pm denotes the \pm -eigenmodule for the action of W_∞ on \mathbb{W} .

Let $\nu \in \mathbb{W}$ and fix a representative ν of ν and a field extension L of \mathbb{Q}_p . The function

$$\gamma \longmapsto \rho_k(\nu_\gamma) := \left(P(x, y) \mapsto \int_{\mathbb{Z}_p \times \mathbb{Z}_p^\times} P(x, y) d\nu_\gamma \right)$$

defined on $\Gamma_0^D(Mp)$ with values in $V_{k-2}(L)$ is a 1-cocycle. Its class in $H^1(\Gamma_0^D(Mp), V_{k-2}(L))$, denoted $\rho_k(\nu)$, does not depend on the choice of ν .

Theorem 3.1. *Let $k \in \mathbb{U}_f$ be an even positive integer. For any choice of sign \pm the $\mathcal{R}_{\mathfrak{p}_k}$ -module $\mathbb{W}_{\mathfrak{p}_k}^\pm$ is free of rank 1 and ρ_k induces an isomorphism $\rho_k : \mathbb{W}_{\mathfrak{p}_k}^\pm / \mathfrak{p}_k \mathbb{W}_{\mathfrak{p}_k}^\pm \xrightarrow{\cong} \mathbb{W}_{f_k}^\pm$.*

The proof of Theorem 3.1 follows the arguments in [14, §5 and §6] closely, and the articles [2] and [9] provide generalizations of the results of [14] to the quaternionic setting we are working in. However, for lack of precise references for the statement we need, a full proof of a more general version of it will be given in a subsequent paper.

Proposition 3.2. *There exists $\tilde{\mu}_f^\pm \in \mathbb{W}^\pm \otimes_{\mathbb{Z}_p} \mathbb{Q}_p$ with the following properties:*

- (1) $\pi_*(\tilde{\mu}_f^\pm) = \text{res}_{\Gamma_0^D(M)}^\Gamma(\mu_f^\pm)$ in $H^1(\Gamma_0^D(M), \mathcal{M}_0(\mathbb{Q}_p))$;
- (2) *at the cost of replacing \mathbb{U}_f with a smaller neighbourhood, for every $k \in \mathbb{U}_f$ there exists $\lambda_B^\pm(k) \in F_k^\times$ such that $\phi_k^\pm = \lambda_B^\pm(k) \rho_k(\tilde{\mu}_f^\pm)$ in $\mathbb{W}_{f_k}^\pm$.*

Proof. In light of Theorem 3.1, we can (and do) choose $\Phi^\pm \in \mathbb{W}^\pm$ such that $\rho_2(\Phi^\pm) = \lambda_1^\pm \phi_2^\pm$ with $\lambda_1^\pm \in \mathbb{Q}_p^\times$. Fix a representative Φ^\pm of Φ^\pm and note that $\rho_2(\Phi^\pm)$ is represented by the cocycle $\gamma \mapsto (\pi_*(\Phi_\gamma^\pm))(\mathbb{Z}_p)$ on $\Gamma_0^D(Mp)$. By [13, Proposition 25], there is a canonical isomorphism

$$\mathcal{S} : H^1(\Gamma, \mathcal{M}_0(\mathbb{Q}_p))^{f, \pm} \xrightarrow{\cong} \mathbb{W}_f^\pm,$$

and thus there exists $\lambda_2^\pm \in \mathbb{Q}_p^\times$ such that $\mathcal{S}(\mu_f^\pm) = \lambda_2^\pm \phi_2^\pm$. Moreover, observe that $\mathcal{S}(\mu_f^\pm)$ is represented by the cocycle $\gamma \mapsto \mu_{f, \gamma}^\pm(\mathbb{Z}_p)$ on $\Gamma_0^D(Mp)$.

Let $\lambda^\pm := \lambda_2^\pm / \lambda_1^\pm$, define $c_\gamma := \lambda^\pm \pi_*(\Phi_\gamma^\pm) - \mu_{f, \gamma}^\pm$ for $\gamma \in \Gamma_0^D(M)$ and set $\mathbf{c} := [c]$ in $H^1(\Gamma_0^D(M), \mathcal{M}(\mathbb{Q}_p))$. Using the surjectivity of π_* proved in [19, Proposition 7.7], choose $\tilde{\mathbf{c}} \in \mathbb{W}^\pm \otimes_{\mathbb{Z}} \mathbb{Q}$ such that $\pi_*(\tilde{\mathbf{c}}) = \mathbf{c}$. (Note that one can choose $\tilde{\mathbf{c}}$ in the ordinary submodule because the arrows between Tate modules at the end of the proof of [19, Proposition 7.7] respect ordinary summands; the same remark applies to the \pm -eigenmodules for W_∞). Since $[\gamma \mapsto c_\gamma(\mathbb{Z}_p)] = 0$ in \mathbb{W}_f^\pm , one has $\rho_2(\tilde{\mathbf{c}}) = 0$. Therefore the class $\tilde{\mu}_f^\pm := \lambda^\pm \Phi^\pm - \tilde{\mathbf{c}}$ satisfies $\rho_2(\tilde{\mu}_f^\pm) = \mathcal{S}(\mu_f^\pm)$ and $\pi_*(\tilde{\mu}_f^\pm) = \text{res}_{\Gamma_0^D(M)}^\Gamma(\mu_f^\pm)$, which proves part (1).

As for part (2), by Theorem 3.1 it is enough to observe that, since the intersection of infinitely many prime ideals of height 1 of \mathcal{R} is trivial, $\rho_k(\tilde{\mu}_f^\pm)$ is 0 only for finitely many integers k , among which the integer 2 does not appear because $\rho_2(\tilde{\mu}_f^\pm) = \mathcal{S}(\mu_f^\pm)$. \square

Definition 3.3. The open set \mathbb{U}_f of Proposition 3.2 is the *domain of analyticity* of $\tilde{\mu}_f^\pm$.

3.4. Explicit formulas for Darmon points. As explained in [19, §7.3], we can choose a representative $\tilde{\mu}_f^\pm$ of $\tilde{\mu}_f^\pm$ in Proposition 3.2 so that $\pi_*(\tilde{\mu}_{f, \gamma}^\pm) = \mu_{f, \gamma}^\pm$ for all $\gamma \in \Gamma_0^D(M)$ (here the map π_* is viewed as defined on cocycles). From now on we fix such a choice of $\tilde{\mu}_f^\pm$.

Let $\log : \mathbb{C}_p^\times \rightarrow \mathbb{C}_p$ be any branch of the p -adic logarithm; such a branch is determined by choosing $\theta \in \mathbb{C}_p^\times$ with $|\theta|_p < 1$ (where $|\cdot|_p$ is the p -adic absolute value on \mathbb{C}_p normalized so that

$|p|_p = 1/p$), setting $\log(\theta) := 0$ and requiring that \log be a homomorphism. In particular, since $x - z_\psi y \in (\mathcal{O}_K \otimes \mathbb{Z}_p)^\times$ when $(x, y) \in \mathbb{X}$, the value $\log(x - z_\psi y)$ for $(x, y) \in \mathbb{X}$ does not depend on the choice of \log (see also the discussion in the proof of [4, Theorem 2.5]).

To simplify the notations, let $G_0 := \Gamma_0^D(Mp)$ and $G_1 := \Gamma_0^D(M)$. Define $G_2 = \hat{\Gamma}_0^D(M) := \omega_p^{-1}G_1\omega_p$. Recall that Γ is the amalgamated product $\Gamma = G_1 *_{G_0} G_2$ with respect to the natural inclusion $G_0 \subset G_1$ and the injection $\varsigma : G_0 \hookrightarrow G_2$ sending x to $\omega_p x \omega_p^{-1}$ (cf. [19, §4.1]).

Theorem 3.4. $\log(\mathcal{P}_\psi) = -t \cdot \int_{\mathbb{X}} \log(x - z_\psi y) d\tilde{\mu}_{f, \gamma_\psi}^\pm \pmod{\log(L)}$.

Proof. Define the 1-cochain $\rho_\gamma := -\int_{\mathbb{X}} \log(x - z_\psi y) d\tilde{\mu}_{f, \gamma}^\pm$ for $\gamma \in G_1$ and let $\hat{\rho}_\gamma$ be defined from ρ_γ for $\gamma \in G_2$ as in [19, §7.5]. Recall that $\pi_*(\tilde{\mu}_f^\pm) = \mu_f^\pm$. A combination of this equality with the calculations in [19, §7.6] shows that $\log(\mathbf{d}_z) = \Delta(\boldsymbol{\rho} - \hat{\boldsymbol{\rho}})$, where Δ is the connecting map appearing in the Mayer–Vietoris exact sequence and $\boldsymbol{\rho} - \hat{\boldsymbol{\rho}} \in H^1(G_0, K_p)$ is represented by $\rho - \hat{\rho}$. We already know that, modulo $\log(L)$, the 2-cocycle $\log(\mathbf{d}_z)$ is split by $\log(\beta_z)$. Hence

$$\delta(\log(\beta_z)) = \Delta(\rho - \hat{\rho}) \pmod{\log(L)}.$$

Since t annihilates Γ^{ab} , the claim is then a consequence of the following lemma. \square

Lemma 3.5. *Let A be an abelian group with trivial Γ -action and consider the portion of the Mayer–Vietoris exact sequence given by*

$$H^1(G_1, A) \oplus H^1(G_2, A) \longrightarrow H^1(G_0, A) \xrightarrow{\Delta} H^2(G, A) \longrightarrow H^2(G_1, A) \oplus H^2(G_2, A).$$

Assume that

- $c = \Delta(\rho)$ for some $\rho \in H^1(G_0, A)$, so there are 1-cochains $\theta_1 \in C^1(G_1, A)$ and $\theta_2 \in C^1(G_2, A)$ such that $c|_{G_1} = \delta(\theta_1)$ and $c|_{G_2} = \delta(\theta_2)$, where δ is the connecting map in cohomology and $\rho = \theta_1|_{G_0} - \theta_2|_{G_0}$;
- c is trivial in $H^2(G, A)$, so $c = \delta(b)$ for some 1-cochain $b \in C^1(G, A)$.

Then $b|_{G_1} = \theta_1 + \varphi_1|_{G_1}$ and $b|_{G_2} = \theta_2 + \varphi_2|_{G_2}$ where $\varphi_1, \varphi_2 : G \rightarrow A$ are group homomorphisms.

Proof. We only prove the existence of φ_1 , since the result for φ_2 can be shown in the same way up to the non-trivial permutation of the set of indices $\{1, 2\}$. First of all, since $c|_{G_1} = \delta(\theta_1) = \delta(b|_{G_1})$ and all the groups involved act trivially on A , it is immediate to deduce that

$$b|_{G_1} = \theta_1 + \psi_1$$

where $\psi_1 : G_1 \rightarrow A$ is a group homomorphism. We want to prove that ψ_1 extends to a homomorphism $\varphi_1 : G \rightarrow A$. To this end, consider the natural isomorphism $\theta : G_1 \xrightarrow{\sim} G_2$ defined by $x \mapsto \omega_p x \omega_p^{-1}$ and define the homomorphism

$$\tilde{\psi}_1 := \psi_1 \circ \theta^{-1} : G_2 \longrightarrow A$$

It follows that $\tilde{\psi}_1 \circ \varsigma = \psi_1|_{G_0}$. But then the universal property of the amalgamated product (see [26, Ch. I, §1]) ensures that there exists a homomorphism $\varphi_1 : G \rightarrow A$ such that $\varphi_1|_{G_1} = \psi_1$, as was to be shown. \square

Now recall the notation in §2.7.

Corollary 3.6. $\log_E(P_\psi) = -nt \cdot \int_{\mathbb{X}} \log_q(x - z_\psi y) d\tilde{\mu}_{f, \gamma_\psi}^\pm$.

Proof. Immediate from Theorem 3.4 and formula (18). \square

4. SPECIAL VALUES OF L -FUNCTIONS

In this section we use the class $\tilde{\mu}_f^\pm$ to define a p -adic L -function $L_p(f_\infty/K, \chi, \kappa)$ associated with f_∞ , K and a genus character χ of K and defined on the domain of analyticity \mathbb{U}_f of f (cf. Definition 3.3). Using Popa's results in [25], we establish a link between the values of $L_p(f_\infty/K, \chi, \kappa)$ at even positive integers $\kappa = k$ and the special values of the complex L -functions of the forms f_k over K twisted by χ .

4.1. Modular forms and 1-cocycles. First, we extend the notation employed in §3.1. For any cusp form g of level $\Gamma_0^D(Mp)$ and weight $k \geq 2$, any point $\tau \in \mathcal{H}$ and any element $\gamma \in B^\times$ of positive norm define $I(g, \tau, \gamma) \in V_{k-2}(\mathbb{C})$ by the formula

$$I(g, \tau, \gamma)(P(x, y)) := \int_\tau^{\gamma(\tau)} g(z)P(z, 1)dz$$

for all $P \in P_{k-2}(\mathbb{C})$. As in §3.1, let B^\times act on the left on $V_n(\mathbb{C})$ via i_F . A direct computation shows that

$$(19) \quad (b \cdot I(g|b, \tau, \gamma))(P) = \det(i_F(b))^{k-2} I(g, b(\tau), b\gamma b^{-1})(P)$$

for all g and τ as above, all $b, \gamma \in B^\times$ of positive norm and all $P \in P_{k-2}(\mathbb{C})$. The cohomology class in $H^1(\Gamma_0^D(Mp), V_{k-2}(\mathbb{C}))$ represented by the cocycle $\gamma \mapsto I(g, \tau, \gamma)$ does not depend on the choice of τ . Denote $\gamma \mapsto I^\pm(g, \tau, \gamma)$ the \pm -component of $\gamma \mapsto I(g, \tau, \gamma)$ under the action of ι ; the class represented by this cocycle is again independent of the choice of τ .

If f_k^{JL} is a form on $\Gamma_0^D(Mp)$ associated with f_k by the Jacquet–Langlands correspondence, it follows that the cohomology class represented by $\gamma \mapsto I^\pm(f_k^{\text{JL}}, \tau, \gamma)$ is a multiple of the generator ϕ_k^\pm introduced in §3.3. From now on, for a fixed sign \pm we normalize the choice of $f_k^{\text{JL}} = f_{k, \pm}^{\text{JL}}$ in such a way that

$$(20) \quad [\gamma \mapsto I^\pm(f_k^{\text{JL}}, \tau, \gamma)] = \phi_k^\pm$$

in $H^1(\Gamma_0^D(Mp), V_{k-2}(\mathbb{C}))^{f_k, \pm}$, where we identify $\phi_k^\pm \in \mathbb{W}_k^\pm$ with its image in this cohomology group via the embedding in (4).

Let $L = \mathbb{Q}(\sqrt{d_L})$ be a real quadratic field with discriminant d_L such that

- all the primes dividing M are split in L ;
- all the primes dividing D are inert in L ;
- the prime p is unramified in L .

Let S denote either M or Mp and let $\psi : L \hookrightarrow B$ be an optimal embedding of the maximal order \mathcal{O}_L of L into $R_0^D(S)$ (note that such an embedding exists unless $S = Mp$ and p is inert in L). As done in §2.3 for $L = K$, we may consider the element $i_F(\psi(\sqrt{d_L})) = \begin{pmatrix} a & b \\ c & -a \end{pmatrix}$ and the quadratic form

$$Q_\psi(x, y) = cx^2 - 2axy - by^2.$$

In order to uniformize our notation, we assume that $L \neq F$, so that $c \neq 0$. Moreover, for every even integer $k \geq 2$ define

$$Q_\psi^{(k)}(x, y) := Q_\psi(x, y)^{\frac{k-2}{2}}.$$

As before, for $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Sigma := \text{GL}_2(\mathbb{Q}_p) \cap \text{M}_2(\mathbb{Z}_p)$ write $\gamma(x, y) := (ax + by, cx + dy)$. Then

$$(21) \quad Q_{\gamma\psi\gamma^{-1}}(x, y) = \det(\gamma) \cdot Q_\psi(\gamma^{-1}(x, y))$$

for all $\gamma \in \Sigma$.

Fix an embedding $L \hookrightarrow \mathbb{R}$ (coinciding with the one fixed in §2.1 when $L = K$) and a fundamental unit ε_L of L such that $\varepsilon_L > 1$ under this embedding, then set $\gamma_\psi := \psi(\varepsilon_L)$. An easy computation shows that if $k \geq 2$ is the weight of g then

$$(22) \quad I(g, \psi) := I(g, \tau, \gamma_\psi) \left(Q_\psi^{(k)}(x, y) \right)$$

does not depend on τ , justifying the notation. Moreover, one checks that $I(g, \psi)$ depends only on the $\Gamma_0^D(S)$ -conjugacy class of ψ . We define $I^\pm(g, \psi)$ in an analogous manner.

4.2. Preliminary calculations. Before proceeding, in this subsection we collect a number of preliminary calculations that will be used in the proof of the main results of this section, namely Theorems 4.24 and 4.33.

Let $\nu|_{\mathbb{U}}$ denote the restriction of a measure ν on \mathbb{Y} to the compact open subset $\mathbb{U} \subset \mathbb{Y}$. Define the compact open subsets

$$\mathbb{X}_\infty := \mathbb{Z}_p^\times \times p\mathbb{Z}_p, \quad \mathbb{X}_{\text{aff}} := \mathbb{Z}_p \times \mathbb{Z}_p^\times$$

of \mathbb{X} , so that $\mathbb{X} = \mathbb{X}_{\text{aff}} \coprod \mathbb{X}_\infty$.

Lemma 4.1. *With self-explaining notation, for every $\nu \in H^1(\Gamma_0^D(M), \mathbb{D})$ one has*

$$\text{res}(\nu)|_{\mathbb{X}_\infty} = \left(\text{res}(\nu)|_{\mathbb{X}_{\text{aff}}} \right) |U_p^{-1}W_p.$$

Proof. A variation on the proof of [19, Lemma 7.11], where a different convention for the Hecke action is adopted. The details are omitted. \square

Notation 4.2. In the following, we often let $[\gamma \mapsto c_\gamma]$ or even $[c_\gamma]$ denote the class represented in $H^1(G, M)$ by a cocycle $c \in Z^1(G, M)$. Moreover, we write $c \equiv c'$ or (by abuse of notation) $c_\gamma \equiv c'_\gamma$ if the two cocycles c and c' are cohomologous, i.e., differ by a coboundary.

Lemma 4.3. *Let L be a field, let $\nu \in H^1(\Gamma_0^D(Mp), \mathbb{D})$ and fix a representative ν of ν . Then*

$$\left[P \mapsto \int_{\mathbb{Z}_p^\times \times \mathbb{Z}_p^\times} P(x, y) d\nu_\gamma \right] = \left[P \mapsto \int_{\mathbb{X}_{\text{aff}}} P(x, y) d\nu_\gamma - \int_{\mathbb{X}_\infty} (P|\alpha)(x, y) d\nu_{\alpha^{-1}\gamma\alpha} \right]$$

in $H^1(\Gamma_0^D(Mp), V_{k-2}(L))$, where $\alpha \in \Gamma_0^D(M)$ satisfies $\alpha\mathbb{X}_\infty = p\mathbb{Z}_p \times \mathbb{Z}_p^\times$.

Proof. Using the decomposition $\mathbb{Z}_p^\times \times \mathbb{Z}_p^\times = \mathbb{X}_{\text{aff}} - (p\mathbb{Z}_p \times \mathbb{Z}_p^\times)$, write

$$\int_{\mathbb{Z}_p^\times \times \mathbb{Z}_p^\times} P(x, y) d\nu_\gamma = \int_{\mathbb{X}_{\text{aff}}} P(x, y) d\nu_\gamma - \int_{p\mathbb{Z}_p \times \mathbb{Z}_p^\times} P(x, y) d\nu_\gamma.$$

Let α be as in the statement of the lemma. First of all, observe that

$$\int_{p\mathbb{Z}_p \times \mathbb{Z}_p^\times} P(x, y) d\nu_\gamma = \int_{\alpha\mathbb{X}_\infty} P(x, y) d\nu_\gamma = \int_{\mathbb{X}_\infty} (P|\alpha)(x, y) d\alpha^{-1}\nu_\gamma.$$

Since $\nu_{\alpha^{-1}\gamma\alpha} = \alpha^{-1}\nu_\gamma + \alpha^{-1}\gamma\nu_\alpha - \alpha^{-1}\nu_\alpha$, it follows that

$$\int_{\mathbb{X}_\infty} (P|\alpha)(x, y) d\alpha^{-1}\nu_\gamma = \int_{\mathbb{X}_\infty} (P|\alpha)(x, y) d\nu_{\alpha^{-1}\gamma\alpha} + \int_{\mathbb{X}_\infty} (P|\alpha)(x, y) d\alpha^{-1}(\nu_\alpha - \gamma\nu_\alpha).$$

Now we show that the map h on $\Gamma_0^D(Mp)$ such that $h_\gamma(P) := \int_{\mathbb{X}_\infty} (P|\alpha)(x, y) d\alpha^{-1}(\nu_\alpha - \gamma\nu_\alpha)$ is a coboundary. Using the fact that \mathbb{X}_∞ is stable under the action of $\Gamma_0^D(Mp)$, an easy computation shows that

$$\int_{\mathbb{X}_\infty} (P|\alpha)(x, y) d\alpha^{-1}\gamma\nu_\alpha = \int_{\mathbb{X}_\infty} (P|\gamma\alpha)(x, y) d\alpha^{-1}\nu_\alpha.$$

Hence if $v(P) := - \int_{\mathbb{X}_\infty} (P|\alpha)(x, y) d\nu_\alpha$ then $h_\gamma(P) = (\gamma v - v)(P)$, and we are done. \square

Recall the domain of analyticity \mathbb{U}_f introduced in Definition 3.3. In light of Lemma 4.3 and normalization (20), we can restate Proposition 3.2 as follows.

Proposition 4.4. *For any choice of $\tau \in \mathcal{H}$ and every $k \in \mathbb{U}_f$ the equality*

$$\left[P \mapsto \int_{\mathbb{X}_{\text{aff}}} P(x, y) d\tilde{\mu}_{f, \gamma}^\pm \right] = \left[\lambda_B^\pm(k) I^\pm(f_k^{\text{JL}}, \tau, \gamma) \right]$$

holds in $H^1(\Gamma_0^D(Mp), V_{k-2}(\mathbb{C}))$.

Lemma 4.5. *For every $k \in \mathbb{U}_f$, one has*

$$\left[P \mapsto \int_{\mathbb{X}_\infty} P(x, y) d\tilde{\mu}_{f, \gamma}^\pm \right] = \left[P \mapsto a_p(k)^{-1} \int_{\mathbb{X}_{\text{aff}}} P(x, y) d(\tilde{\mu}_{f, \gamma}^\pm | W_p) \right]$$

in $H^1(\Gamma_0^D(Mp), V_{k-2}(F_k))$.

Proof. By Lemma 4.1, there exists a \mathbb{Z}_p -valued measure m on \mathbb{X}_∞ such that

$$\int_{\mathbb{X}_\infty} P(x, y) d\tilde{\mu}_{f, \gamma}^\pm = \int_{\mathbb{X}_{\text{aff}}} P(x, y) d(\tilde{\mu}_{f, \gamma}^\pm | U_p^{-1} W_p) + \int_{\mathbb{X}_\infty} P(x, y) d(\gamma m - m).$$

Since $\tilde{\mu}_f^\pm | U_p^{-1} \equiv a_p(k)^{-1} \tilde{\mu}_f^\pm$, there exists also an F_k -valued measure m' on \mathbb{X}_{aff} such that

$$\int_{\mathbb{X}_{\text{aff}}} P(x, y) d(\tilde{\mu}_{f, \gamma}^\pm | U_p^{-1} W_p) = a_p(k)^{-1} \int_{\mathbb{X}_{\text{aff}}} P(x, y) d(\tilde{\mu}_{f, \gamma}^\pm | W_p) + \int_{\mathbb{X}_{\text{aff}}} (P|\omega_p^*)(x, y) d(\gamma m' - m')$$

where $\omega_p^* := p\omega_p^{-1}$. Therefore, setting $v(P) := \int_{\mathbb{X}_\infty} P dm + \int_{\mathbb{X}_{\text{aff}}} (P|\omega_p^*) dm'$ we get the equality

$$\int_{\mathbb{X}_\infty} P(x, y) d\tilde{\mu}_{f, \gamma}^\pm = a_p(k)^{-1} \int_{\mathbb{X}_{\text{aff}}} P(x, y) d(\tilde{\mu}_{f, \gamma}^\pm | W_p) + (\gamma v - v)(P),$$

which completes the proof. \square

Lemma 4.6. *For any choice of $\tau \in \mathcal{H}$ and every $k \in \mathbb{U}_f$, the equality*

$$\left[P \mapsto \int_{\mathbb{X}_{\text{aff}}} P(x, y) d\tilde{\mu}_{f, \gamma}^\pm | W_p \right] = \left[\lambda_B^\pm(k) I^\pm(f_k^{\text{JL}}, \tau, \gamma) | W_p \right]$$

holds in $H^1(\Gamma_0^D(Mp), V_{k-2}(\mathbb{C}))$.

Lemma 4.7. *For any choice of $\tau \in \mathcal{H}$ and every $k \in \mathbb{U}_f$, the equality*

$$\left[P \mapsto \int_{\mathbb{X}_\infty} P(x, y) d\tilde{\mu}_{f, \gamma}^\pm \right] = \left[P \mapsto \frac{\lambda_B^\pm(k) p^{k-2}}{a_p(k)} I^\pm(f_k^{\text{JL}}, \tau, \omega_p \gamma \omega_p^{-1})(P(x, y) | \omega_p^{-1}) \right]$$

holds in $H^1(\Gamma_0^D(Mp), V_{k-2}(\mathbb{C}))$.

Proof. Combining Lemmas 4.5 and 4.6, we get

$$\left[P \mapsto \int_{\mathbb{X}_{\text{aff}}} P(x, y) d(\tilde{\mu}_{f, \gamma}^\pm | W_p) \right] = \left[P \mapsto \lambda_B^\pm(k) a_p(k)^{-1} I^\pm(f_k^{\text{JL}}, \tau, \omega_p \gamma \omega_p^{-1})(P(x, y) | \omega_p^*) \right].$$

Since $\omega_p^* = p\omega_p^{-1}$, the claim follows. \square

Let $k \geq 4$ be an even integer and let f_k^\sharp be the p -stabilization of f_k as defined, e.g., in [14, p. 410]. Then f_k^\sharp corresponds by Jacquet–Langlands to a form $f_k^{\text{JL},\sharp}$, which can be chosen in such a way that

$$(23) \quad f_k^{\text{JL}} = f_k^{\text{JL},\sharp} - a_p(k)^{-1} f_k^{\text{JL},\sharp}|W_p.$$

Combining (19) and (23), for all polynomials P of degree $k-2$ and any $\tau \in \mathcal{H}$ we get

$$\begin{aligned} I(f_k^{\text{JL}}, \tau, \gamma)(P) &= \int_{\tau}^{\gamma(\tau)} P(z, 1) f_k^{\text{JL},\sharp}(z) dz - a_p(k)^{-1} \int_{\tau}^{\gamma(\tau)} P(z, 1) f_k^{\text{JL},\sharp}(z) |W_p dz \\ &= I(f_k^{\text{JL},\sharp}, \tau, \gamma)(P) - a_p(k)^{-1} I(f_k^{\text{JL},\sharp}|W_p, \tau, \gamma)(P) \\ &= I(f_k^{\text{JL},\sharp}, \tau, \gamma)(P) - p^{k-2} a_p(k)^{-1} I(f_k^{\text{JL},\sharp}, \omega_p(\tau), \omega_p \gamma \omega_p^{-1})(P|\omega_p^{-1}). \end{aligned}$$

In particular, taking \pm -eigenspaces gives

$$(24) \quad I^{\pm}(f_k^{\text{JL}}, \tau, \gamma)(P) = I^{\pm}(f_k^{\text{JL},\sharp}, \tau, \gamma)(P) - p^{k-2} a_p(k)^{-1} I^{\pm}(f_k^{\text{JL},\sharp}, \omega_p(\tau), \omega_p \gamma \omega_p^{-1})(P|\omega_p^{-1}).$$

Proposition 4.8. *Let $k \geq 4$ be an even integer in \mathbb{U}_f and let $\tau \in \mathcal{H}$. Then*

$$\left[P \mapsto \int_{\mathbb{X}} P(x, y) d\tilde{\mu}_{f,\gamma}^{\pm} \right] = \left[\lambda_B^{\pm}(k) \left(1 - \frac{p^{k-2}}{a_p(k)^2} \right) I^{\pm}(f_k^{\text{JL},\sharp}, \tau, \gamma) \right]$$

in $H^1(\Gamma_0^D(Mp), V_{k-2}(\mathbb{C}))$.

Proof. Split the integral on the left hand side

$$(25) \quad \int_{\mathbb{X}} P(x, y) d\tilde{\mu}_{f,\gamma}^{\pm} = \int_{\mathbb{X}_{\text{aff}}} P(x, y) d\tilde{\mu}_{f,\gamma}^{\pm} + \int_{\mathbb{X}_{\infty}} P(x, y) d\tilde{\mu}_{f,\gamma}^{\pm}.$$

We begin by evaluating the first integral on the right hand side of (25). A straightforward computation shows that the class in $H^1(\Gamma_0^D(Mp), V_{k-2}(\mathbb{C}))$ of the cocycle

$$\gamma \mapsto \left(P(x, y) \mapsto \int_{\tau}^{\omega_p \gamma \omega_p^{-1}(\tau)} f_k^{\text{JL},\sharp} P(z, 1) | \omega_p^{-1} dz \right)$$

is independent of the choice of $\tau \in \mathcal{H}$. Then, using Proposition 4.4 and (24), we see that the cocycle appearing as the first term on the right hand side of (25) is equivalent to

$$(26) \quad \gamma \mapsto \left(P \mapsto \lambda_B^{\pm}(k) I^{\pm}(f_k^{\text{JL},\sharp}, \tau, \gamma)(P) - \lambda_B^{\pm}(k) p^{k-2} a_p(k)^{-1} I^{\pm}(f_k^{\text{JL},\sharp}, \tau, \omega_p \gamma \omega_p^{-1})(P|\omega_p^{-1}) \right).$$

Now we compute the second term on the right of (25). Using Lemma 4.7 and (24), we compute

$$\begin{aligned} I^{\pm}(f_k^{\text{JL}}, \tau, \omega_p \gamma \omega_p^{-1})(P|\omega_p^{-1}) \\ = I^{\pm}(f_k^{\text{JL},\sharp}, \tau, \omega_p \gamma \omega_p^{-1})(P|\omega_p^{-1}) - p^{k-2} a_p(k)^{-1} I^{\pm}(f_k^{\text{JL},\sharp}, \omega_p(\tau), \omega_p^2 \gamma \omega_p^{-2})(P|\omega_p^{-2}). \end{aligned}$$

To study the second term on the right hand side of the above equation, we apply (19), the equality $f_k^{\text{JL},\sharp}|W_p^2 = p^{k-2} f_k^{\text{JL},\sharp}$ and the fact that the class of $\gamma \mapsto I^{\pm}(f_k^{\text{JL},\sharp}, \tau, \gamma)$ does not depend on the choice of $\tau \in \mathcal{H}$. Finally, we obtain that $P \mapsto \int_{\mathbb{X}_{\infty}} P(x, y) d\tilde{\mu}_{f,\gamma}^{\pm}$ is equivalent to

$$(27) \quad P \mapsto \lambda_B^{\pm}(k) a_p(k)^{-1} p^{k-2} \left(I^{\pm}(f_k^{\text{JL},\sharp}, \tau, \omega_p \gamma \omega_p^{-1})(P|\omega_p^{-1}) - a_p(k)^{-1} I^{\pm}(f_k^{\text{JL},\sharp}, \tau, \gamma)(P) \right).$$

Combining (26) and (27) yields the result. \square

Corollary 4.9. *Let $k \geq 4$ be an even integer in \mathbb{U}_f and let $\tau \in \mathcal{H}$. Then*

$$\left[P \mapsto \int_{\mathbb{X}} P(x, y) d\tilde{\mu}_{f, \gamma}^{\pm} \right] = \left[\lambda_B^{\pm}(k) \left(1 - \frac{p^{k-2}}{a_p(k)^2} \right) I^{\pm}(f_k^{\text{JL}, \#}, \tau, \gamma) \right]$$

in $H^1(\Gamma_0^D(M), V_{k-2}(\mathbb{C}))$.

Proof. If this were not true then the left hand side and the right hand side would represent different classes in $H^1(\Gamma_0^D(M), V_{k-2}(\mathbb{C}))$ that, by Proposition 4.8, when restricted to $\Gamma_0^D(Mp)$ share the same Hecke eigenvalues at all primes except, possibly, the prime p . By multiplicity one, this cannot occur because $f_k^{\#}$ is a newform. \square

Proposition 4.10. *For any $\tau \in \mathcal{H}$, the equality*

$$\left[\gamma \mapsto \int_{\mathbb{Z}_p^{\times} \times \mathbb{Z}_p^{\times}} d\tilde{\mu}_{f, \gamma}^{\pm} \right] = \left[\gamma \mapsto 2\lambda_B^{\pm}(2) \cdot \int_{\tau}^{\gamma(\tau)} f^{\text{JL}} dz \right]$$

holds in $H^1(\Gamma_0^D(Mp), \mathbb{C})$.

Proof. Combining (19), Proposition 4.4 and Lemma 4.7, it follows that

$$\int_{\mathbb{Z}_p^{\times} \times \mathbb{Z}_p^{\times}} d\tilde{\mu}_{f, \gamma}^{\pm} = \lambda_B^{\pm}(2) \left(\int_{\tau}^{\gamma(\tau)} f^{\text{JL}} dz - \frac{1}{a_p} \int_{\tau}^{\gamma(\tau)} (f^{\text{JL}}|_{W_p}) dz \right)$$

for all $\gamma \in \Gamma_0^D(Mp)$. Now $f^{\text{JL}}|_{W_p} = -a_p f^{\text{JL}}$, and the result follows. \square

Define $w_p := -a_p$, so that $f_k^{\text{JL}, \#}|_{W_p} = w_p p^{(k-2)/2} f_k^{\text{JL}, \#}$ (it is known that the value w_p is common to all the forms f_k in the Hida family: see, e.g., [23, §3.4.4]).

Proposition 4.11. *Let $k \geq 4$ be an even integer in \mathbb{U}_f and let $\tau \in \mathcal{H}$. Then*

$$\left[P \mapsto \int_{\mathbb{Z}_p^{\times} \times \mathbb{Z}_p^{\times}} P(x, y) d\tilde{\mu}_{f, \gamma}^{\pm} \right] = \left[\lambda_B^{\pm}(k) \left(1 - \frac{w_p p^{(k-2)/2}}{a_p(k)} \right)^2 I^{\pm}(f_k^{\text{JL}, \#}, \tau, \gamma) \right]$$

in $H^1(\Gamma_0^D(Mp), V_{k-2}(\mathbb{C}))$.

Proof. Combining Lemma 4.3 with the arguments in the proof of Proposition 4.8, it follows that the cocycle on $\Gamma_0^D(Mp)$ sending γ to $(P \mapsto \int_{\mathbb{Z}_p^{\times} \times \mathbb{Z}_p^{\times}} P d\tilde{\mu}_{f, \gamma}^{\pm})$ is equivalent to the difference between the cocycles sending γ to the maps which take \tilde{P} to

$$\lambda_B^{\pm}(k) I^{\pm}(f_k^{\text{JL}, \#}, \tau, \gamma)(P) - \frac{\lambda_B^{\pm}(k) p^{k-2}}{a_p(k)} I^{\pm}(f_k^{\text{JL}, \#}, \tau, \omega_p \gamma \omega_p^{-1})(P|_{\omega_p^{-1}})$$

and

$$\frac{\lambda_B^{\pm}(k) p^{k-2}}{a_p(k)} I^{\pm}(f_k^{\text{JL}, \#}, \tau, \omega_p \alpha^{-1} \gamma \alpha \omega_p^{-1})(P|\alpha \omega_p^{-1}) - \frac{\lambda_B^{\pm}(k) p^{k-2}}{a_p(k)^2} I^{\pm}(f_k^{\text{JL}, \#}, \tau, \alpha^{-1} \gamma \alpha)(P|\alpha),$$

with α as in the statement of Lemma 4.3. Using (19) and the fact that $f_k^{\text{JL}, \#}$ and $f_k^{\text{JL}, \#}|_{W_p}$ have level $\Gamma_0^D(M)$, it can be checked that the sums above are equal to

$$\lambda_B^{\pm}(k) I^{\pm}(f_k^{\text{JL}, \#}, \tau, \gamma)(P) - \frac{\lambda_B^{\pm}(k)}{a_p(k)} I^{\pm}(f_k^{\text{JL}, \#}|_{W_p}, \tau, \gamma)(P)$$

and

$$\frac{\lambda_B^{\pm}(k)}{a_p(k)} I^{\pm}(f_k^{\text{JL}, \#}|_{W_p}, \tau, \gamma)(P) - \frac{\lambda_B^{\pm}(k) p^{k-2}}{a_p(k)^2} I^{\pm}(f_k^{\text{JL}, \#}, \tau, \gamma)(P),$$

respectively. Taking the difference of these two expressions yields the result. \square

Corollary 4.12. *Let $k \geq 4$ be an even integer in \mathbb{U}_f and let $\tau \in \mathcal{H}$. Then*

$$\left[P \mapsto \int_{\mathbb{Z}_p^\times \times \mathbb{Z}_p^\times} P(x, y) d\tilde{\mu}_{f, \gamma}^\pm \right] = \left[\lambda_B^\pm(k) \left(1 - \frac{w_p p^{(k-2)/2}}{a_p(k)} \right)^2 I^\pm(f_k^{\text{JL}, \#}, \tau, \gamma) \right]$$

in $H^1(\Gamma_0^D(M), V_{k-2}(\mathbb{C}))$.

Proof. In light of Proposition 4.11, proceed as in the proof of Corollary 4.9. \square

4.3. Popa's work. Let \mathbb{A} be the adèle ring of \mathbb{Q} and set $B_{\mathbb{A}} := B \otimes_{\mathbb{Q}} \mathbb{A}$ for the adelization of our quaternion algebra B . Let g be a weight k normalized newform of level $\Gamma_0(SD)$ where S is a square-free positive integer prime to D . Let $R_0^D(S)$ be an Eichler order of B of level S and let $\Gamma_0^D(S)$ denote the group of its elements of norm 1. Let g^{JL} denote a modular form on $\Gamma_0^D(S)$ corresponding to g via the Jacquet–Langlands correspondence; notice that g^{JL} is well defined up to multiplication by non-zero constants. Let φ_g and φ_g^{JL} denote the automorphic forms on $\text{GL}_2(\mathbb{A})$ and $B_{\mathbb{A}}^\times$ associated with g and g^{JL} , respectively, whose definitions can be found, e.g., in [10, §3] and [10, §10], respectively. We just recall, because it will be used in the following computations, that φ_g^{JL} is defined in *loc. cit.* using the decomposition

$$(28) \quad B_{\mathbb{A}}^\times = B^\times (B_\infty^\times)^0 K_0^D(S)$$

where $K_0^D(S) := (R_0^D(S) \otimes_{\mathbb{Z}} \hat{\mathbb{Z}})^\times$ ($\hat{\mathbb{Z}}$ denotes, as usual, the profinite completion of \mathbb{Z}) and $(B_\infty^\times)^0$ is the connected component of the identity in $B_\infty^\times := (B \otimes_{\mathbb{Q}} \mathbb{R})^\times$ (cf. [10, Lemma 10.3] for details).

Fix a real quadratic field $L = \mathbb{Q}(\sqrt{d_L})$ such that all the primes dividing S (respectively, D) are split (respectively, inert) in L . For simplicity, we assume that $L \neq F$ (actually, the case $L = F$ is easier but requires different conventions: note, for example, that one may diagonally embed F into $M_2(F)$ and that the quadratic form associated with this embedding has $c = b = 0$, in the notations of §4.1). Set $G_L^+ := \text{Gal}(H_L^+/L)$ where H_L^+ is the narrow Hilbert class field of L . Fix an unramified character $\chi : G_L^+ \rightarrow \mathbb{C}^\times$. By class field theory, χ can be identified with a finite-order Hecke character $\chi : \mathbb{A}^\times L^\times \backslash \mathbb{A}_L^\times \rightarrow \mathbb{C}^\times$, where $\mathbb{A}_L = L \otimes_{\mathbb{Q}} \mathbb{A}$ is the adèle ring of L . Denote by π_χ the automorphic representation of $\text{GL}_2(\mathbb{A})$ attached to χ via the Jacquet–Langlands correspondence.

Fix an embedding $L \hookrightarrow \mathbb{R}$ and choose the generator ε_L of the group of norm 1 elements in \mathcal{O}_L so that $\varepsilon_L > 1$ with respect to this embedding. In the special case where $L = K$ fix the embedding $K \hookrightarrow \mathbb{R}$ as in §2.1, so that ε_L is the element denoted by ε_1 in §2.4. Write dx for the Haar measure on $L^\times \mathbb{A}^\times \backslash \mathbb{A}_L^\times$ defined in [25, p. 840], which is normalized in such a way that the total mass is equal to $h_L^+ \ln \varepsilon_L$, where h_L^+ is the cardinality of $\text{Pic}^+(\mathcal{O}_L)$. As above, write $\text{Emb}(\mathcal{O}_L, R_0^D(S))$ for the set of optimal embeddings of \mathcal{O}_L into $R_0^D(S)$. Fix also an embedding $\psi_0 \in \text{Emb}(\mathcal{O}_L, R_0^D(S))$ (if $L = K$ choose ψ_0 as in §2.6) and write $\psi_{0, \mathbb{A}}$ for its adelization. Also, write $\hat{\psi}_0$ and $\psi_{0, \infty}$ for the finite and the infinite part of $\psi_{0, \mathbb{A}}$, respectively. At the archimedean places of L we may also consider, as in [25, (5.3.1)], the diagonal embedding $\psi_{\text{diag}} : L \otimes_{\mathbb{Q}} \mathbb{R} \hookrightarrow M_2(\mathbb{R})$ obtained by extending by \mathbb{R} -linearity the embedding $L \hookrightarrow M_2(\mathbb{R})$ given by $x \mapsto \begin{pmatrix} x & 0 \\ 0 & \bar{x} \end{pmatrix}$ where $x \mapsto \bar{x}$ is the involution of L . Let $i_F(\psi_0(\sqrt{d_L})) = \begin{pmatrix} a & b \\ c & -a \end{pmatrix}$ and define $\gamma_\infty := \begin{pmatrix} a + \sqrt{c} & a - \sqrt{c} \\ c & -c \end{pmatrix} \in \text{GL}_2(\mathbb{R})$ with $c > 0$ (note that, since $L \neq F$, we always have $c \neq 0$). Then $\psi_{0, \infty} = \gamma_\infty \psi_{\text{diag}} \gamma_\infty^{-1}$. Define

$$l(\varphi_g^{\text{JL}}, \chi) := \int_{\mathbb{A}^\times L^\times \backslash \mathbb{A}_L^\times} \varphi_g^{\text{JL}}(\psi_{0, \mathbb{A}}(x) \gamma_\infty) \chi^{-1}(x) dx.$$

By [25, Theorem 5.4.1], there is the formula

$$L(1/2, \pi_g \times \pi_\chi) = \frac{4}{\sqrt{d_L}} \cdot \prod_{\ell|D} \frac{\ell+1}{\ell-1} \cdot \frac{\|\varphi_g\|^2}{\|\varphi_g^{\text{JL}}\|^2} \cdot |l(\varphi_g^{\text{JL}}, \chi)|^2.$$

Although $l(\varphi_g^{\text{JL}}, \chi)$ depends on the choice of g^{JL} and φ_g^{JL} , the right hand side of the above expression does not depend on it.

Now we compute the term $l(\varphi_g^{\text{JL}}, \chi)$ more explicitly. Set $\tau_0 := i_F(\gamma_\infty)(i)$. Following [25, §6.1], one shows that

$$l(\varphi_g^{\text{JL}}, \chi) = \frac{1}{(2i)^{k/2} \sqrt{d_L}^{(k-2)/2}} \sum_{[\mathfrak{a}] \in \text{Pic}^+(\mathcal{O}_L)} \chi^{-1}([\mathfrak{a}]) M([\mathfrak{a}])$$

with

$$(29) \quad M([\mathfrak{a}]) := \det(\alpha_{[\mathfrak{a}]})^{\frac{2-k}{2}} \int_{\tau_{[\mathfrak{a}]}}^{(\alpha_{[\mathfrak{a}]}^{-1} \psi_0 \alpha_{[\mathfrak{a}]})^{(\varepsilon_L) \tau_{[\mathfrak{a}]}}} f(z) (Q_{\psi_0}^{(k)} | \alpha_{[\mathfrak{a}]}) (z, 1) dz.$$

The term $\alpha_{[\mathfrak{a}]} \in (B_\infty^\times)^0$ is defined, using decomposition (28), by the equation $\hat{\psi}_0(\hat{\mathfrak{a}}) = b \alpha_{[\mathfrak{a}]}^{-1} \mathbf{k}$ with $b \in B^\times$, $\alpha_{[\mathfrak{a}]}^{-1} \in (B_\infty^\times)^0$, $\mathbf{k} \in K_0^D(S)$ and $\tau_{[\mathfrak{a}]} := \alpha_{[\mathfrak{a}]}^{-1}(\tau_0)$; here $\hat{\mathfrak{a}}$ is any finite idele that corresponds to an ideal \mathfrak{a} representing the class $[\mathfrak{a}]$. Although $\alpha_{[\mathfrak{a}]}$ is not unique, the results will not depend on the actual choice. To simplify notations, we also identify $\alpha_{[\mathfrak{a}]}$ with its image via i_∞ and view $\alpha_{[\mathfrak{a}]}$ as an element of $\text{GL}_2^+(\mathbb{R})$.

Let h_L be the class number of L , so that $h_L^+/h_L = 1$ or 2 , let $\text{Pic}(\mathcal{O}_L)$ be the ideal class group of L and write H_L for the Hilbert class field of L (hence $\text{Pic}(\mathcal{O}_L) \simeq \text{Gal}(H_L/L)$ via the reciprocity map of class field theory). Let \mathfrak{D}_L denote the class of $(\sqrt{d_L})$ in $\text{Pic}^+(\mathcal{O}_L)$. If for every $\sigma \in \text{Gal}(H_L/L)$ we write $[\mathfrak{a}_\sigma] = \text{rec}^{-1}(\sigma)$ for an ideal \mathfrak{a}_σ of \mathcal{O}_L then

$$\text{Pic}(\mathcal{O}_L) = \{[\mathfrak{a}_\sigma]\}_{\sigma \in \text{Gal}(H_L/L)}$$

and

$$\text{Pic}^+(\mathcal{O}_L) = \begin{cases} \{[\mathfrak{a}_\sigma]\}_{\sigma \in \text{Gal}(H_L/L)} & \text{if } h_L^+ = h_L, \\ \{[\mathfrak{a}_\sigma]\}_{\sigma \in \text{Gal}(H_L/L)} \cup \{\mathfrak{D}_L \cdot [\mathfrak{a}_\sigma]\}_{\sigma \in \text{Gal}(H_L/L)} & \text{otherwise.} \end{cases}$$

Here, with a slight abuse of notation, we have adopted the symbol $[\star]$ to denote ideal classes both in $\text{Pic}(\mathcal{O}_L)$ and in $\text{Pic}^+(\mathcal{O}_L)$.

Write $\sigma_L \in G_L^+$ for the image of \mathfrak{D}_L under the reciprocity map. For every $\sigma \in G_L^+$ let $[\mathfrak{a}_\sigma]$ be the corresponding element in $\text{Pic}^+(\mathcal{O}_L)$ via the Artin map, represented by an ideal $\mathfrak{a}_\sigma \subset \mathcal{O}_L$. We extend in the obvious way the notation and results of §2.5 about oriented optimal embeddings to the more general case needed here. In particular, for an oriented optimal embedding $\psi \in \mathcal{E}(\mathcal{O}_L, R_0^D(S))$ define

$$(30) \quad [\mathfrak{a}_\sigma] \star [\psi] := [\mathfrak{a}_\sigma \psi \mathfrak{a}_\sigma^{-1}] \in \mathcal{E}(\mathcal{O}_L, R_0^D(S)) / \Gamma_0^D(S)$$

where $\mathfrak{a}_\sigma \in R_0^D(S)$ has positive reduced norm and satisfies $R_0^D(S) \psi(\mathfrak{a}_\sigma) = R_0^D(S) \mathfrak{a}_\sigma$. Fixing ψ , the rule $\sigma \mapsto [\psi_\sigma]$ with $\psi_\sigma := \mathfrak{a}_\sigma \psi \mathfrak{a}_\sigma^{-1}$ is the inverse of a bijection G between $\mathcal{E}(\mathcal{O}_L, R_0^D(S)) / \Gamma_0^D(S)$ and G_L^+ . The map G satisfies the equation $G([\psi^*]) = \sigma_L \cdot G([\psi])$ for all $\psi \in \mathcal{E}(\mathcal{O}_L, R_0^D(S))$, where $\psi^* := \omega_\infty \psi \omega_\infty^{-1}$. The family $\{\psi_\sigma\}_{\sigma \in G_L^+}$ is a set of representatives of the $\Gamma_0^D(S)$ -conjugacy classes of oriented optimal embeddings of \mathcal{O}_L into $R_0^D(S)$. Finally, for every $\sigma \in G_L^+$ set $\gamma_\sigma := \psi_\sigma(\varepsilon_L) \in \Gamma_0^D(S)$.

Remark 4.13. We explicitly observe a fact that will be used in the proof of Theorem 4.33. Suppose that, in the above discussion, $S = Mp$. Since the quadratic order \mathcal{O}_L is maximal, an oriented optimal embedding of \mathcal{O}_L into $R_0^D(Mp)$ can be naturally viewed as an element of $\mathcal{E}(\mathcal{O}_L, R_0^D(M))$ too. Moreover, the element a_σ described above also satisfies $R_0^D(M)\psi(\mathfrak{a}_\sigma) = R_0^D(M)a_\sigma$. This shows that we can (and do) fix via (30) a family $\{\psi_\sigma\}_{\sigma \in G_L^+}$ which is a set of representatives of the $\Gamma_0^D(S)$ -conjugacy classes of oriented optimal embeddings of \mathcal{O}_L into $R_0^D(S)$ for S equal to either M or Mp .

If the subscript ∞ indicates that an element of B^\times of positive norm is to be viewed as belonging to $(B_\infty^\times)^0$, it follows that $\hat{\psi}_0(\hat{\mathfrak{a}}_\sigma) = a_\sigma(a_\sigma^{-1})_\infty \mathbf{k}$, with respect to decomposition (28), for a suitable $\mathbf{k} \in K_0^D(S)$. Hence for every $\sigma \in G_L^+$ we can take $\alpha_{[\mathfrak{a}_\sigma]} = a_\sigma$, in the above notation. Moreover, if $h_L^+ \neq h_L$ then for every $\sigma \in G_L^+$ we can also take $\alpha_{\mathfrak{a}_L \cdot [\mathfrak{a}_\sigma]}$ to be equal to $\omega_\infty a_\sigma \psi_0(\sqrt{d_L})$.

In light of the expression of the integrals $M([\mathfrak{a}_\sigma])$ given in (29), we summarize the above calculations in the following

Theorem 4.14 (Popa). *Let χ be a complex-valued character of G_L^+ . Then*

$$L(1/2, \pi_g \times \pi_\chi) = \frac{4}{(2i)^k \sqrt{d_L}^{k-1}} \cdot \frac{\|\varphi_g\|^2}{\|\varphi_g^{\text{JL}}\|^2} \cdot \prod_{\ell|D} \frac{\ell+1}{\ell-1} \cdot \left| \sum_{\sigma \in G_L^+} \chi^{-1}(\sigma) \int_{\tau_0}^{\gamma_\sigma(\tau_0)} g^{\text{JL}}(z) Q_{\psi_\sigma}^{(k)}(z, 1) dz \right|^2.$$

With notation as in (22), the next task is to rewrite more explicitly the integral

$$\Theta_\psi := \int_{\tau_0}^{\gamma_\psi(\tau_0)} g^{\text{JL}}(z) Q_\psi^{(k)}(z, 1) dz = I(g^{\text{JL}}, \psi)$$

for any of the $\psi = \psi_\sigma$ and $\gamma_\psi = \gamma_\sigma$. Define

$$(g^{\text{JL}}|_{\omega_\infty})(z) := (Cz + D)^{-k} \cdot \overline{g^{\text{JL}}(\omega_\infty \bar{z})},$$

where $i_\infty(\omega_\infty) = \begin{pmatrix} * & * \\ C & D \end{pmatrix}$.

Lemma 4.15. *Up to rescaling g^{JL} , we may assume that $g^{\text{JL}}|_{\omega_\infty} = g^{\text{JL}}$.*

Proof. A simple computation shows that $g^{\text{JL}}|_{\omega_\infty}$ has the same weight and level as g^{JL} . Since the Hecke eigenvalues of g are real because g is a newform with trivial character, one sees that the same is true of g^{JL} . This can be used to show that $g^{\text{JL}}|_{\omega_\infty}$ has the same eigenvalues as g^{JL} and so, by multiplicity one, there exists a complex number λ such that $g^{\text{JL}}|_{\omega_\infty} = \lambda g^{\text{JL}}$. Since $\omega_\infty^2 \in \Gamma_0^D(M)$, we have that $(g^{\text{JL}}|_{\omega_\infty})|_{\omega_\infty} = g^{\text{JL}}$. (Thus, in particular, $|\lambda| = 1$, but we will not need this here). If $\lambda = 1$ then g^{JL} verifies the claimed condition without rescaling, while if $\lambda = -1$ then ig^{JL} does the job. In the other cases, the modular form $g^{\text{JL}} + g^{\text{JL}}|_{\omega_\infty} = (1 + \lambda)g^{\text{JL}}$ has the required property. \square

From now on assume that g^{JL} verifies the condition in Lemma 4.15. Using this, a simple computation shows that

$$\overline{\Theta_\psi} = \int_{\omega_\infty \bar{\tau}_0}^{\gamma_{\psi^*}(\omega_\infty \bar{\tau}_0)} g^{\text{JL}}(z) Q_{\psi^*}^{(k)}(z, 1) dz = I(g^{\text{JL}}, \psi^*) = \Theta_{\psi^*}$$

(recall that $\psi^* := \omega_\infty \psi \omega_\infty^{-1}$). By the above discussion, the embeddings ψ_σ^* and $\psi_{\sigma_L \sigma}$ are $\Gamma_0^D(S)$ -conjugate. Since Θ_ψ depends only on the $\Gamma_0^D(S)$ -conjugacy class of ψ , we obtain

$$(31) \quad \overline{\Theta_{\psi_\sigma}} = \Theta_{\psi_{\sigma_L \sigma}}.$$

Let χ be a *genus character* of L , i.e., an unramified quadratic character of G_L^+ . Thanks to (31), we have

$$\overline{\sum_{\sigma \in G_L^+} \chi(\sigma) \Theta_{\psi_\sigma}} = \sum_{\sigma \in G_L^+} \chi(\sigma) \overline{\Theta_{\psi_\sigma}} = \sum_{\sigma \in G_L^+} \chi(\sigma) \Theta_{\psi_{\sigma_L \sigma}} = \chi(\sigma_L) \cdot \left(\sum_{\sigma \in G_L^+} \chi(\sigma) \Theta_{\psi_\sigma} \right).$$

Definition 4.16. The *sign* of χ is the sign of $\chi(\sigma_L)$.

A straightforward calculation shows that

$$\overline{\int_{\tau_0}^{\gamma(\tau_0)} g(z) P(z, 1) dz} = - \int_{\omega_\infty \bar{\tau}_0}^{\omega_\infty \gamma \omega_\infty^{-1}(\omega_\infty \bar{\tau}_0)} g(z) (P|_{\omega_\infty^*})(z, 1) dz.$$

In other words, the action of complex conjugation on $I(g, \tau, \gamma)(P)$ coincides, up to a change of sign, with the action of W_∞ (for a more detailed discussion of this relation in the case of elliptic modular forms, see [15, p. 588]). Hence if χ has sign $-\epsilon \in \{\pm\}$ then

$$\sum_{\sigma \in G_L^+} \chi(\sigma) \Theta_{\psi_\sigma} = \sum_{\sigma \in G_L^+} \chi(\sigma) I^\epsilon(g^{\text{JL}}, \psi_\sigma).$$

Summing up, Theorem 4.14 admits the following reformulation for genus characters.

Corollary 4.17. *Let χ be a genus character of L of sign $-\epsilon$. Then*

$$L(1/2, \pi_g \times \pi_\chi) = \frac{\epsilon 4}{(2i)^k \sqrt{d_L}^{k-1}} \cdot \frac{\|\varphi_g\|^2}{\|\varphi_g^{\text{JL}}\|^2} \cdot \prod_{\ell|D} \frac{\ell+1}{\ell-1} \cdot \left(\sum_{\sigma \in G_L^+} \chi(\sigma) I^\epsilon(g^{\text{JL}}, \psi_\sigma) \right)^2.$$

4.4. p -adic L -functions over real quadratic fields. Before studying p -adic L -functions, let us observe a simple fact. Let L be a real quadratic field and let $\psi : L \hookrightarrow B$ be an optimal embedding of \mathcal{O}_L into $R_0^D(N)$, where N denotes either M or Mp . Fix a subset \mathbb{U} of \mathbb{X} such that $\gamma_\psi \mathbb{U} = \mathbb{U}$. Let $k \geq 2$ be an *even* integer, fix $\nu \in \mathbb{W}$ and choose a representative ν of ν .

Lemma 4.18. *Let the notation be as above.*

- (1) *The value $\int_{\mathbb{U}} Q_\psi^{(k)}(x, y) d\nu_{\gamma_\psi}$ does not depend on the choice of a representative ν of ν .*
- (2) *If $\psi' = \alpha \psi \alpha^{-1}$ for some $\alpha \in \Gamma_0^D(N)$ then $\int_{\mathbb{U}} Q_\psi^{(k)}(x, y) d\nu_{\gamma_\psi} = \int_{\alpha \mathbb{U}} Q_{\psi'}^{(k)}(x, y) d\nu_{\gamma_{\psi'}}$.*
- (3) *If $\gamma \mathbb{U} = \mathbb{U}$ for all $\gamma \in \Gamma_0^D(N)$ then $\int_{\mathbb{U}} Q_\psi^{(k)}(x, y) d\nu_{\gamma_\psi}$ depends only on the $\Gamma_0^D(N)$ -conjugacy class of ψ .*

Proof. The proof follows from a direct calculation using the equality

$$(32) \quad Q_\psi(\gamma_\psi(x, y)) = Q_\psi(x, y),$$

which is a consequence of (21). Details are left to the reader. \square

Recall now the fixed real quadratic field K where p is inert. Recall also the domain of analyticity \mathbb{U}_f of $\tilde{\mu}_f^\pm$ introduced in Definition 3.3. Let $\langle x \rangle$ denote the principal unit attached to $x \in \mathbb{Q}_p^\times$, defined as the unique element of $1 + p\mathbb{Z}_p$ such that $x = p^{\text{ord}_p(x)} \zeta_x \langle x \rangle$ where ζ_x is a $(p-1)$ -st root of unity. For $k \in \mathbb{U}_f$ and $(x, y) \in \mathbb{X}$ define

$$Q_\psi^{(k)}(x, y) := Q_\psi(x, y)^{\frac{k-2}{2}} = \exp_p \left(\frac{k-2}{2} \log(\langle Q_\psi(x, y) \rangle) \right)$$

where \exp_p is the p -adic exponential and \log is *any* branch of the p -adic logarithm. Since $Q_\psi(x, y) = \langle Q_\psi(x, y) \rangle$ for $(x, y) \in \mathbb{X}$, this definition does not depend on the choice of \log .

Definition 4.19. Let ψ be an optimal embedding of \mathcal{O}_K into $R_0^D(M)$. The *partial square root p -adic L -function* attached to f_∞ , K and ψ is

$$\mathcal{L}_p^\pm(f_\infty, \psi, k) := \int_{\mathbb{X}} Q_\psi^{(k)}(x, y) d\tilde{\mu}_{f, \gamma_\psi}^\pm(x, y)$$

where k varies in \mathbb{U}_f .

Remark 4.20. Since $Q_\psi(x, y) = \langle Q_\psi(x, y) \rangle$ on \mathbb{X} , we have

$$\mathcal{L}_p^\pm(f_\infty, \psi, k) = \int_{\mathbb{X}} \langle Q_\psi(x, y) \rangle^{\frac{k-2}{2}} d\tilde{\mu}_{f, \gamma_\psi}^\pm(x, y).$$

Moreover, if $k \geq 2$ is an integer in \mathbb{U}_f then $Q_\psi^{(k)}(x, y)$ is equal to the $(k-2)/2$ -fold self product of $Q_\psi(x, y)$. This justifies the restriction of \mathbb{U}_f to the residue class of 2 modulo $p-1$ in \mathcal{X} .

Proposition 4.21. $\mathcal{L}_p^\pm(f_\infty, \psi, k)$ is independent of the choice of a representative $\tilde{\mu}_f^\pm$ of $\tilde{\mu}_f^\pm$ and only depends on the $\Gamma_0^D(M)$ -conjugacy class of ψ .

Proof. Lemma 4.18 shows that the restriction of $\mathcal{L}_p^\pm(f_\infty, \psi, k)$ to the subset of \mathbb{U}_f consisting of points with trivial character has the required properties. The result then follows from a density argument. \square

Proposition 4.22. For every even integer $k \geq 4$ in \mathbb{U}_f one has

$$\mathcal{L}^\pm(f_\infty, \psi, k) = \lambda_B^\pm(k) \left(1 - \frac{p^{k-2}}{a_p(k)^2}\right) I^\pm(f_k^{\text{JL}, \sharp}, \psi).$$

Proof. Corollary 4.9 ensures the existence of $v \in V_{k-2}(\mathbb{C})$ such that

$$\int_{\mathbb{X}} Q_\psi^{(k)}(x, y) d\tilde{\mu}_{f, \gamma_\psi}^\pm(x, y) = \lambda_B^\pm(k) \left(1 - \frac{p^{k-2}}{a_p(k)^2}\right) I^\pm(f_k^{\text{JL}, \sharp}, \psi) + (\gamma v - v) \left(Q_\psi^{(k)}(x, y)\right)$$

for all $\gamma \in \Gamma_0^D(Mp)$. We conclude by evaluating this expression at $\gamma = \gamma_\psi$ and using (32). \square

Definition 4.23. Let χ be a quadratic character of G_K^+ of sign $-\epsilon$ and let k vary in \mathbb{U}_f .

(1) The *square root p -adic L -function* attached to f_∞ and χ is

$$\mathcal{L}_p(f_\infty/K, \chi, k) := \sum_{\sigma \in G_K^+} \chi(\sigma) \mathcal{L}_p^\epsilon(f_\infty, \psi_\sigma, k).$$

(2) The *p -adic L -function* attached to f_∞ and χ is

$$L_p(f_\infty/K, \chi, k) := \mathcal{L}_p(f_\infty/K, \chi, k)^2.$$

Theorem 4.24. Let χ be a genus character of K of sign $-\epsilon$. Then for all even integers $k \geq 4$ in \mathbb{U}_f we have

$$L_p(f_\infty/K, \chi, k) = \lambda_B^\epsilon(k)^2 \left(1 - \frac{p^{k-2}}{a_p(k)^2}\right)^2 \prod_{\ell|D} \frac{\ell-1}{\ell+1} \cdot \frac{(2i)^k \sqrt{d_K}^{k-1}}{\epsilon 4} \cdot \frac{\|\varphi_{f_k}^{\text{JL}, \sharp}\|^2}{\|\varphi_{f_k}^\sharp\|^2} \cdot L_K(1/2, \pi_{f_k}^\sharp \times \pi_\chi).$$

Proof. If $k \geq 2$ then f_k^\sharp is a newform on $\Gamma_0^D(M)$, so the result follows immediately from Corollary 4.17 and Proposition 4.22. \square

Remark 4.25. The above result holds also for $k = 2$ if we set $f_2^\sharp := 0$. Indeed, in this case

$$(33) \quad \mathcal{L}_p(f_\infty/K, \chi, 2) = \int_{\mathbb{X}} d\tilde{\mu}_{f, \gamma_\psi} = \int_{\mathbb{P}^1(\mathbb{Q}_p)} d\pi_*(\mu_{f, \gamma_\psi}) = 0,$$

and the equality in the statement is trivially verified.

4.5. Atkin–Lehner involutions and genus characters. A genus character χ of K cuts out the genus field H_χ of K given by

$$H_\chi = \mathbb{Q}(\sqrt{d_1}, \sqrt{d_2})$$

where $d_K = d_1 d_2$. The extension H_χ/\mathbb{Q} is biquadratic unless χ is trivial (in which case, of course, $H_\chi = K$). Let χ_1, χ_2 and ϵ_K be the characters associated with the quadratic fields $\mathbb{Q}(\sqrt{d_1}), \mathbb{Q}(\sqrt{d_2})$ and K , respectively, so that $\epsilon_K = \chi_1 \cdot \chi_2$. In fact, the genus characters of K are in bijection with the factorizations of d into a product of relatively prime discriminants d_1 and d_2 or, equivalently, with the unordered pairs (χ_1, χ_2) of primitive quadratic Dirichlet characters of coprime conductors satisfying $\epsilon_K = \chi_1 \cdot \chi_2$ (the trivial character corresponds to the factorization $d = 1 \cdot d$). For more details see, e.g., [6, Ch. 14, §G].

Let now ψ be an oriented optimal embedding of \mathcal{O}_K into $R_0^D(M)$ and recall the unique element $\sigma_\psi \in G_K^+$ introduced in (12) such that

$$(34) \quad \psi \circ \tau \sim \omega_{MD}(\sigma_\psi \star \psi) \omega_{MD}^{-1}.$$

To ease the writing, from now until the end of this subsection set $G := G_K^+$. Denote by G^2 the subgroup of G consisting of the squares of the elements of G .

Lemma 4.26. *The class of σ_ψ in G/G^2 does not depend on ψ .*

In light of this result, write $\bar{\sigma}$ for the class of any σ_ψ in G/G^2 .

Proof. To begin with, recall that σ_ψ is characterized as the unique element of G such that

$$[\omega_{MD}^{-1}(\psi \circ \tau) \omega_{MD}] = \sigma_\psi \star [\psi].$$

Suppose that ψ' is another oriented optimal embedding of \mathcal{O}_K into $R_0^D(M)$ and write $[\psi'] = \sigma \star [\psi]$ for a suitable $\sigma \in G$. Then

$$(35) \quad \begin{aligned} (\sigma_\psi \sigma^{-2}) \star [\psi'] &= (\sigma_\psi \sigma^{-2}) \star (\sigma \star [\psi]) = \sigma_\psi \star (\sigma^{-1} \star [\psi]) \\ &= \sigma^{-1} \star (\sigma_\psi \star [\psi]) = \sigma^{-1} \star [\omega_{MD}^{-1}(\psi \circ \tau) \omega_{MD}]. \end{aligned}$$

On the other hand, $\sigma_{\psi'}$ is characterized as the unique element of G such that

$$(36) \quad [\omega_{MD}^{-1}((\sigma \star \psi) \circ \tau) \omega_{MD}] = [\omega_{MD}^{-1}(\psi' \circ \tau) \omega_{MD}] = \sigma_{\psi'} \star [\psi'].$$

If $\mathfrak{a} \subset \mathcal{O}_K$ is an ideal whose class $[\mathfrak{a}] \in \text{Pic}^+(\mathcal{O}_K)$ corresponds to σ under the reciprocity map then σ^{-1} is represented by $\bar{\mathfrak{a}} := \tau(\mathfrak{a})$. With notation as in §2.5, it follows that

$$R_0^D(M)(\psi \circ \tau)(\bar{\mathfrak{a}}) = R_0^D(M)\psi(\mathfrak{a}) = R_0^D(M)a,$$

and of course $(a\psi a^{-1}) \circ \tau = a(\psi \circ \tau)a^{-1}$. Hence

$$(37) \quad \sigma^{-1} \star [\omega_{MD}^{-1}(\psi \circ \tau) \omega_{MD}] = [\omega_{MD}^{-1}((\sigma \star \psi) \circ \tau) \omega_{MD}],$$

and by comparing (35) and (36) we conclude that

$$(38) \quad \sigma_{\psi'} = \sigma_\psi \sigma^{-2}.$$

The lemma is proved. \square

Remark 4.27. Equality (37) is a manifestation of the fact that the Galois extension H_K^+/\mathbb{Q} is generalized dihedral, which is expressed by the formula $\tau\sigma\tau = \sigma^{-1}$ for all $\sigma \in G$.

Now we want to give a description of the quaternion algebra B in terms of the Galois element σ_ψ introduced in (34) (or, rather, of $\sigma_K\sigma_\psi$). This represents the analogue in our real quadratic setting of results for CM points on Shimura curves obtained by Jordan in [16] (see [12, Lemma 5.10] for a detailed proof of a generalization of Jordan's results).

Recall the element $\omega_\infty \in R_0^D(M)$ of norm -1 and consider

$$\mathfrak{D}_K \star [\psi \circ \tau] = [\omega_\infty(\psi \circ \tau)\omega_\infty^{-1}] = [(\psi \circ \tau)^*].$$

Since $\omega_\infty(\psi \circ \tau)\omega_\infty^{-1}$ has the same orientation as $\psi \circ \tau$ at all the primes dividing MD , it follows from Lemma 2.6 that there exists $\sigma_{\mathfrak{a}} \in G$ corresponding to (the class of) an ideal \mathfrak{a} of \mathcal{O}_K via the reciprocity map of class field theory such that

$$(39) \quad [(\psi \circ \tau)^*] = [\omega_{MD}(\sigma_{\mathfrak{a}} \star \psi)\omega_{MD}^{-1}].$$

Proposition 4.28. *There is an isomorphism*

$$(40) \quad B \simeq \left(\frac{d_K, MDN_{K/\mathbb{Q}}(\mathfrak{a})}{\mathbb{Q}} \right)$$

of quaternion algebras over \mathbb{Q} . Moreover, if \mathfrak{b} is an ideal of \mathcal{O}_K satisfying (40) and $\sigma_{\mathfrak{b}} \in G$ corresponds to the class of \mathfrak{b} in $\text{Pic}^+(\mathcal{O}_K)$ then the images of $\sigma_{\mathfrak{a}}$ and $\sigma_{\mathfrak{b}}$ in G/G^2 are equal.

Proof. Set $i := \psi(\sqrt{d_K}) \in R_0^D(M)$. We first show the following

Claim. *There exists $j \in R_0^D(M)$ such that $j^2 = MDN_{K/\mathbb{Q}}(\mathfrak{a})$ and $ij = -ji$.*

Proof of Claim. As in §2.5, let $a \in R_0^D(M)$ be an element of positive reduced norm such that $R_0^D(M)\psi(\mathfrak{a}) = R_0^D(M)a$. In particular, the reduced norm of a is equal to $N_{K/\mathbb{Q}}(\mathfrak{a})$. Then equality (39) means that there exists $\alpha \in \Gamma_0^D(M)$ such that

$$\alpha\omega_\infty(\psi \circ \tau)\omega_\infty^{-1}\alpha^{-1} = \omega_{MD}a\psi a^{-1}\omega_{MD}^{-1}.$$

Define $j := \omega_\infty^{-1}\alpha^{-1}\omega_{MD}a \in R_0^D(M)$, so that

$$(41) \quad \psi \circ \tau = j\psi j^{-1}.$$

Now we want to prove that $j^2 \in \mathbb{Q}^\times$ and $ij = -ji$. To begin with, the relation $ij = -ji$ follows immediately from (41) and the fact that $(\psi \circ \tau)(\sqrt{d_K}) = -\psi(\sqrt{d_K})$. As for the first assertion, we use the canonical injection $B \hookrightarrow M_2(F)$ and denote by the same symbol ψ the injection $K \hookrightarrow M_2(F)$ obtained by composing ψ with this map. Recall that, in our notation, $\psi(\sqrt{d_K}) = \begin{pmatrix} a & b \\ c & -a \end{pmatrix} \in M_2(F)$. Write $j = \begin{pmatrix} x & y \\ z & w \end{pmatrix} \in M_2(F)$. Since $ij = -ji$, we see that $w = -x$, and then it follows that $j^2 = \begin{pmatrix} -\text{norm}(j) & 0 \\ 0 & -\text{norm}(j) \end{pmatrix}$. Thus $j^2 \in \mathbb{Q}^\times$ because $\text{norm}(j) \in \mathbb{Q}$. Finally, since $\text{norm}(j) = -MDN_{K/\mathbb{Q}}(\mathfrak{a}) \in \mathbb{Z}$ we conclude that $j^2 = MDN_{K/\mathbb{Q}}(\mathfrak{a})$. \square

Since $i, j \in R_0^D(M)$, in light of the above Claim we conclude that

$$B \simeq \mathbb{Q} \oplus \mathbb{Q}i \oplus \mathbb{Q}j \oplus \mathbb{Q}ij = \left(\frac{d_K, MDN_{K/\mathbb{Q}}(\mathfrak{a})}{\mathbb{Q}} \right).$$

To complete the proof of the proposition, one can proceed as in [12, Remark 5.11] and note that, by genus theory, if the ideal \mathfrak{b} of \mathcal{O}_K satisfies (40) then $\sigma_{\mathfrak{a}}$ and $\sigma_{\mathfrak{b}}$ have the same image in G/G^2 . \square

With notation as before, it follows that

$$(42) \quad \sigma_\psi = \sigma_K \sigma_{\mathfrak{a}}.$$

Now let λ be a prime number that splits in K and is such that $\lambda \equiv MD \pmod{d_K}$. A calculation with local Hilbert symbols analogous to the one in [25, §5.1] shows that there is an isomorphism

$$(43) \quad B \simeq \left(\frac{d_K, MD\lambda}{\mathbb{Q}} \right)$$

of quaternion algebras over \mathbb{Q} (here we use the fact that M is square-free, although the condition that M be a product of odd powers of distinct primes would suffice). Choose an ideal \mathfrak{b} of \mathcal{O}_K such that $N_{K/\mathbb{Q}}(\mathfrak{b}) = \lambda$ (such a \mathfrak{b} exists because λ splits in K). Combining isomorphism (43) with Proposition 4.28 and equality (42), it follows that the class of $\sigma_K \sigma_{\mathfrak{b}}$ in G/G^2 coincides with the class $\bar{\sigma}$ of σ_ψ .

Suppose that the genus character χ is associated with the pair (χ_1, χ_2) of quadratic characters. It is now straightforward to prove the main result of this subsection.

Proposition 4.29. $\chi(\bar{\sigma}) = \chi_1(-MD)$.

Proof. With notation as before, the congruence $N_{K/\mathbb{Q}}(\mathfrak{b}) \equiv MD \pmod{d_K}$ implies that $N_{K/\mathbb{Q}}(\mathfrak{b}) \equiv MD \pmod{d_1}$. Hence, since $\chi(\sigma_K) = \text{sign}(d_1) = \chi_1(-1)$ (cf. [6, Remark 14.47]), we get that

$$\chi(\bar{\sigma}) = \chi(\sigma_K) \cdot \chi(\sigma_{\mathfrak{b}}) = \chi_1(-MD),$$

as was to be shown. \square

Remark 4.30. Throughout this article we assume that $D > 1$, i.e., that B is *not* isomorphic to the split algebra $M_2(\mathbb{Q})$. However, a moment's thought reveals that the arguments in this subsection remain valid also when $D = 1$. With this in mind, it turns out that if $D = 1$ then Proposition 4.29 reduces to [4, Proposition 1.8], and so our arguments provide an alternative (and more conceptual) proof of this result.

4.6. Derivatives of p -adic L -functions and Darmon points. Let \log_E be as in Corollary 3.6, where ψ is an optimal embedding of \mathcal{O}_K into $R_0^D(M)$. Set $J_\psi^\pm := \log_E(P_\psi^\pm)$ and for a genus character χ of K of sign $-\epsilon \in \{\pm\}$ define

$$J_\chi := \sum_{\sigma \in G_K^+} \chi(\sigma) J_{\psi_\sigma}^\epsilon = \log_E(P_\chi)$$

with $P_\chi := \sum_{\sigma \in G_K^+} \chi(\sigma) P_{\psi_\sigma}^\epsilon \in E(K_p)$. Finally, recall the integers n and t in Corollary 3.6.

Theorem 4.31. *With notation as before, there is an equality*

$$\frac{d}{dk} \mathcal{L}_p(f_\infty/K, \chi, k)_{k=2} = \frac{w_{MD} \chi_1(-MD) - 1}{2nt} J_\chi.$$

Proof. Recall the choice \log_q of the branch of the logarithm made in §2.7. The integrals below do not depend on this choice, as observed before Definition 4.19. However, we will write \log_q instead of a generic \log to directly obtain the result to be shown. First, we have

$$\frac{d}{dk} \mathcal{L}_p^\epsilon(f_\infty, \psi, k)_{k=2} = \int_{\mathbb{X}} \log_q(Q_\psi(x, y)) d\tilde{\mu}_{f, \gamma_\psi}^\epsilon.$$

Since $Q_\psi(x, y) = c(x - z_\psi y)(x - \bar{z}_\psi y)$, we can write

$$\log_q(Q_\psi(x, y)) = \log_q(c) + \log_q(x - z_\psi y) + \log_q(x - \bar{z}_\psi y).$$

Since

$$\int_{\mathbb{X}} \log_q(c) d\tilde{\mu}_{f,\gamma_\psi}^\epsilon = \int_{\mathbb{P}^1(\mathbb{Q}_p)} \log_q(c) d\pi_*(\tilde{\mu}_f^\epsilon)_{\gamma_\psi} = 0$$

(because $\pi_*(\tilde{\mu}_f)$ is cohomologous to μ_f and thus has total mass 0 at every γ), we find that

$$\frac{d}{dk} \mathcal{L}_p^\epsilon(f_\infty, \psi, k)_{k=2} = \frac{1}{2} \int_{\mathbb{X}} \log_q(x - z_\psi y) d\mu_{f,\gamma_\psi}^\epsilon + \frac{1}{2} \int_{\mathbb{X}} \log_q(x - \bar{z}_\psi y) d\mu_{f,\gamma_\psi}^\epsilon.$$

Thanks to Corollary 3.6, the first term on the right hand side is equal to $-J_\psi^\epsilon/2nt$. By the same result, the second term on the right hand side is equal to $-\tau(J_\psi^\epsilon)/2nt$. Proposition 2.9 then shows that

$$(44) \quad \frac{d}{dk} \mathcal{L}_p^\epsilon(f_\infty, \psi, k)_{k=2} = \frac{w_{MD} J_{\sigma_\psi \star \psi}^\epsilon - J_\psi^\epsilon}{2nt}$$

with σ_ψ as in (12). Using formula (38), it is easy to check that the map from G_K^+ to itself sending σ to $\sigma_{\psi\sigma}$ is a bijection. In light of this fact and Proposition 4.29, specializing (44) to $\psi = \psi_\sigma$ and then summing over all $\sigma \in G_K^+$ yields the result. \square

Corollary 4.32. *There is an equality*

$$\frac{d^2}{d^2k} L_p(f_\infty/K, \chi, k)_{k=2} = \begin{cases} J_\chi^2/(nt)^2 & \text{if } \chi_1(-MD) = -w_{MD}; \\ 0 & \text{if } \chi_1(-MD) = w_{MD}. \end{cases}$$

Proof. The second derivative of $L_p(f_\infty/K, \chi, k)$ evaluated at $k = 2$ is equal to the sum

$$2 \left(\frac{d}{dk} \mathcal{L}_p(f_\infty/K, \chi, k)_{k=2} \right)^2 + 2 \mathcal{L}_p(f_\infty/K, \chi, 2) \frac{d^2}{d^2k} \mathcal{L}_p(f_\infty/K, \chi, k)_{k=2}.$$

Since $\mathcal{L}_p(f_\infty/K, \chi, 2) = 0$ by (33), the result follows from Theorem 4.31. \square

4.7. Factorization formulas for p -adic L -functions. For $j = 1, 2$ let $L_p(f_\infty, \chi_j, k, s)$ be the Mazur–Kitagawa p -adic L -function associated with f_∞ and χ_j ; we refer to [3, §1.4] for its definition and main properties. In particular, recall that, by [4, Corollary 3.2], if $\chi_j(-1) = (-1)^{(k-2)/2} \tilde{\epsilon}$ with $\tilde{\epsilon} = \pm 1$ (since K is real, $\epsilon_K(-1) = 1$, so $\chi_1(-1) = \chi_2(-1)$) we have

$$(45) \quad L_p(f_\infty, \chi_j, k, k/2) = \lambda^{\tilde{\epsilon}}(k) \left(1 - \frac{\chi_j(p) p^{(k-2)/2}}{a_p(k)} \right)^2 \cdot \frac{\Gamma(k/2) \tau(\chi_j)}{(-2\pi i)^{(k-2)/2}} \cdot \frac{L(f_k^\sharp, \chi_j, k/2)}{\Omega_{f_k}^{\tilde{\epsilon}}}$$

where $\tau(\chi_j) := \sum_{a=1}^{d_j} \chi_j(a) e^{2\pi i a/d_j}$ is the Gauss sum attached to χ_j , $\Omega_{f_k}^{\tilde{\epsilon}}$ is the complex period of f_k^\sharp defined as in [3, Proposition 1.1] and $\lambda^{\tilde{\epsilon}}(k) \in \mathbb{C}_p$ is defined as in [3, Theorem 1.5]. Finally, recall that the sign of the functional equation of the L -function $L(f, \delta, s)$ of f twisted by a quadratic Dirichlet character δ is equal to $-w_{MDp} \delta(-MDp)$ and that the p -adic L -function $L_p(f, \delta, s)$ has an exceptional zero at $s = 1$ if and only if $\delta(p) = -w_p$.

The following factorization result will play a crucial role in our subsequent arguments.

Theorem 4.33. *There exist a neighbourhood $\mathbb{U} \subset \mathbb{U}_f$ of 2 and a p -adic analytic function η on \mathbb{U} such that*

- (1) $\eta(k) \neq 0$ for all $k \in \mathbb{U}$ and $\eta(2) \in (\mathbb{Q}^\times)^2$;
- (2) for all $k \in \mathbb{U}$ there is a factorization

$$L_p(f_\infty/K, \chi, k) = \eta(k) L_p(f_\infty, \chi_1, k, k/2) L_p(f_\infty, \chi_2, k, k/2).$$

Proof. By comparing Euler factors, we first notice that there is a factorization of complex L -functions

$$L(s, \pi_{f_k} \times \pi_\chi) = L(s, \pi_{f_k} \times \pi_{\chi_1})L(s, \pi_{f_k} \times \pi_{\chi_2}).$$

On the other hand (see, e.g., [11, p. 202]), for $j = 1, 2$ one has

$$L(s, \pi_{f_k^\#} \times \pi_{\chi_j}) = \frac{\Gamma(s + (k-1)/2)}{(2\pi)^{s+(k-1)/2}} L(f_k^\#, \chi_j, s + (k-1)/2).$$

Let the sign of χ be $-\epsilon$. Then it follows from Theorem 4.24 that for all even integers $k \geq 4$ in \mathbb{U}_f there is an equality

$$(46) \quad L_p(f_\infty/K, \chi, k) = \lambda_B^\epsilon(k)^2 \left(1 - \frac{p^{k-2}}{a_p(k)^2}\right)^2 \prod_{\ell|D} \frac{\ell-1}{\ell+1} \cdot \frac{\sqrt{d_K}^{k-1}}{4\epsilon} \cdot \frac{\|\varphi_{f_k^\#}^{\text{JL}}\|^2}{\|\varphi_{f_k^\#}\|^2} \cdot \frac{\Gamma(k/2)^2}{(-i\pi)^k} \\ \times L(f_k^\#, \chi_1, k/2) L(f_k^\#, \chi_2, k/2).$$

Since p is inert in K , we have $\chi_1(p) = -\chi_2(p)$, and then formulas (45) and (46) imply that

$$(47) \quad L_p(f_\infty/K, \chi, k) = \eta(k) L_p(f_\infty, \chi_1, k, k/2) L_p(f_\infty, \chi_2, k, k/2)$$

for all even integers $k \geq 4$, where $\eta(k) \in \mathbb{C}_p$ can be made explicit by comparing (45) and (46).

Now we show that η can be extended to an analytic function with the prescribed properties. For this we need an argument similar to the one in the proof of [3, Proposition 5.2]. Consider the set of primitive Dirichlet characters χ'_1 of conductor prime to MDp such that

- $\chi'_1(-1) = \chi_1(-1) = \chi_2(-1)$;
- $\chi'_1(-MD) = -w_{MD}$;
- $\chi'_1(p) = -a_p = w_p$.

The condition $\chi'_1(-MD) = -w_{MD}$ implies that $L(f, \chi'_1, s)$ vanishes of even order at $s = 1$, while the condition $\chi'_1(p) = -a_p = w_p$ implies that $L_p(f, \chi'_1, 1)$ does not have an exceptional zero, so that $L_p(f, \chi'_1, 1) \in \mathbb{Q}^\times$. Fix a character χ'_1 as above such that $L(f, \chi'_1, 1) \neq 0$ (by the main result of [22], there are infinitely many).

Consider now the set of primitive Dirichlet characters χ'_2 of conductor prime to MDp and the conductor of χ'_1 satisfying

- $\chi'_2(-1) = \chi'_1(-1)$;
- $\chi'_2(\ell) = \chi'_1(\ell)$ for $\ell|pM$;
- $\chi'_2(\ell) = -\chi'_1(\ell)$ for $\ell|D$.

Since D has an even number of prime factors, $\chi'_2(-MD) = \chi'_1(-MD)$ and $L(f, \chi'_2, s)$ also vanishes to even order at $s = 1$. Also, since $\chi'_2(p) = -a_p$, the p -adic L -function $L_p(f, \chi'_2, s)$ does not have an exceptional zero. Hence, again by [22], there are infinitely many characters χ'_2 as above such that $L(f, \chi'_2, 1) \neq 0$ and $L_p(f, \chi'_2, 1) \neq 0$. Choose a pair (χ'_1, χ'_2) as prescribed above and denote by d'_1 and d'_2 the conductors of χ'_1 and χ'_2 , respectively. Let $d_{K'} := d'_1 \cdot d'_2$ and consider the real quadratic field $K' := \mathbb{Q}(\sqrt{d_{K'}})$ and the genus character χ' of K' corresponding to the pair (χ'_1, χ'_2) . In particular, by definition, the sign of χ' is equal to the sign of χ . Recall that $G_{K'}^+ := \text{Gal}(H_{K'}^+/K')$ and fix a set $\{\psi'_\sigma\}$ of representatives of the $\Gamma_0^D(Mp)$ -equivalence classes of oriented optimal embeddings of $\mathcal{O}_{K'}$ into $R_0^D(Mp)$ as described in Remark 4.13, so that this family is also a set of representatives for the $\Gamma_0^D(M)$ -equivalence classes of oriented optimal embeddings

of $\mathcal{O}_{K'}$ into $R_0^D(M)$. Now consider the expression

$$(48) \quad L_p(f_\infty/K', \chi', k) := \left(\sum_{\sigma \in G_{K'}^+} \chi'(\sigma) \int_{\mathbb{Z}_p^\times \times \mathbb{Z}_p^\times} Q_{\psi'_\sigma}^{(k)}(x, y) d\tilde{\mu}_{f, \gamma_{\psi'_\sigma}}^\epsilon \right)^2.$$

Combining the arguments in the proof of Proposition 4.22 with Proposition 4.10 and Corollary 4.12, we see that

$$\int_{\mathbb{Z}_p^\times \times \mathbb{Z}_p^\times} d\tilde{\mu}_{f, \gamma_{\psi'_\sigma}}^\epsilon = 2\lambda_B^\epsilon(2) I^\epsilon(f^{\text{JL}, \#}, \tau, \gamma_{\psi'_\sigma})(1)$$

and

$$\int_{\mathbb{Z}_p^\times \times \mathbb{Z}_p^\times} Q_{\psi'_\sigma}^{(k)}(x, y) d\tilde{\mu}_{f, \gamma_{\psi'_\sigma}}^\epsilon = \lambda_B^\epsilon(k) \left(1 - \frac{w_p p^{(k-2)/2}}{a_p(k)} \right)^2 I^\epsilon(f_k^{\text{JL}, \#}, \tau, \gamma_{\psi'_\sigma}) \left(Q_{\psi'_\sigma}^{(k)}(x, y) \right)$$

for f_k with trivial character. Set $f_2^\# := f$, $f_2^{\text{JL}, \#} := f^{\text{JL}}$ and $\varphi_{f_2^\#}^{\text{JL}, \#} := \varphi_f^{\text{JL}}/2$. For all even $k \geq 2$ such that f_k has trivial character we obtain the interpolation formula

$$\begin{aligned} L_p(f_\infty/K', \chi', k) &= \lambda_B^\epsilon(k)^2 \left(1 - \frac{w_p p^{(k-2)/2}}{a_p(k)} \right)^k \prod_{\ell|D} \frac{\ell-1}{\ell+1} \cdot \frac{(2i)^k \sqrt{d_K}^{k-1}}{4\epsilon} \\ &\quad \times \frac{\|\varphi_{f_k}^{\text{JL}, \#}\|^2}{\|\varphi_{f_k^\#}\|^2} \cdot L_K(1/2, \pi_{f_k^\#} \times \pi_\chi). \end{aligned}$$

Therefore we get the factorization

$$L_p(f_\infty/K', \chi', k) = \eta(k) L_p(f_\infty, \chi'_1, k, k/2) L_p(f_\infty, \chi'_2, k, k/2)$$

where η is the function appearing in (47).

With these choices in force, $L_p(f_\infty/K', \chi', k)$, $L_p(f_\infty, \chi'_1, k, k/2)$ and $L_p(f_\infty, \chi'_2, k, k/2)$ do not vanish at $k = 2$, so there exists a neighbourhood \mathbb{U} of 2 such that they do not vanish at any point of \mathbb{U} . Recall that $\lambda^\epsilon(2)$ is chosen to be equal to 1. Thus the quotient

$$\eta(k, \chi'_1, \chi'_2) := \frac{L_p(f_\infty/K', \chi', k)}{L_p(f_\infty, \chi'_1, k, k/2) L_p(f_\infty, \chi'_2, k, k/2)}$$

is an analytic function on \mathbb{U} which does not vanish at any point of \mathbb{U} , and we define $\eta(k) := \eta(k, \chi'_1, \chi'_2)$. (Observe that $\eta(k, \chi'_1, \chi'_2)$ does not depend on the choice of χ'_1 and χ'_2 made above: any other choice defines a function which coincides with $\eta(k, \chi'_1, \chi'_2)$ on the dense subset of points $k \in \mathbb{U}$ with trivial character because, for these points, $\eta(k)$ does not depend on χ'_1, χ'_2 .)

To conclude the proof, we need only show the statement about $\eta(2)$. For this, we use the following argument. Fix a prime $d|D$ and consider the set of Dirichlet characters χ'_3 and χ'_4 of conductor prime to MDp satisfying the following conditions:

- $\chi'_1(-1) = \chi'_2(-1) = -\chi'_3(-1) = \chi'_4(-1)$;
- $\chi'_1(\ell) = \chi'_2(\ell) = \chi'_3(\ell) = \chi'_4(\ell)$ for all primes $\ell|pM$;
- $\chi'_1(d) = -\chi'_2(d) = -\chi'_3(d) = -\chi'_4(d)$;
- $\chi'_1(\ell) = -\chi'_2(\ell) = \chi'_3(\ell) = -\chi'_4(\ell)$ for all primes $\ell|(D/d)$.

To simplify notations, set $\Lambda_j := L_p(f_\infty, \chi'_j, 2, 1)$ for $j = 1, \dots, 4$. Note that $\chi'_3(-MD) = \chi'_4(-MD) = -w_{MD}$ and $\chi'_3(p) = \chi'_4(p) = w_p$. Thus, as above, the order of vanishing of $L(f, \chi'_3, s)$ and $L(f, \chi'_4, s)$ at $s = 1$ is even and $L_p(f, \chi'_3, s)$ and $L_p(f, \chi'_4, s)$ do not have an exceptional zero at $s = 1$. Using [22] again, we choose χ'_3 and χ'_4 so that $\Lambda_3 \neq 0$ and $\Lambda_4 \neq 0$.

For the pairs of indices (1, 3), (3, 4) and (2, 4) we consider quadratic fields $K'_{i,j}$ and genus characters $\chi'_{i,j}$ of $K'_{i,j}$ such that the associated Dirichlet characters are (χ'_i, χ'_j) ; in particular, if d'_j is the conductor of χ'_j then the discriminant of $K'_{i,j}$ is $d'_i d'_j$. Note that

- $K'_{1,3}$ and $K'_{3,4}$ are imaginary, while $K'_{2,4}$ is real;
- the primes dividing Mp are split in $K'_{1,3}$ and $K'_{3,4}$ and the number of primes dividing D which are inert in these fields is odd;
- the primes dividing MDp are split in $K'_{2,4}$.

Thanks to [3, Proposition 5.1], we have

$$L_p(f_\infty/K'_{i,j}, \chi'_{i,j}, 2) = \langle \phi_2, \phi_2 \rangle \Lambda_i \Lambda_j \quad \text{for } (i, j) = (1, 3) \text{ and } (i, j) = (3, 4),$$

where we use the notation in [3]. In particular, recall that the p -adic L -function in the equation above is defined in [3, §3.2] and that $\langle \phi_2, \phi_2 \rangle \in \mathbb{Q}^\times$ is defined in [3, p. 395]. Since, by construction, $L_p(f_\infty/K'_{i,j}, \chi'_{i,j}, 2)$ is a square in \mathbb{Q}^\times , it follows that $\Lambda_i \Lambda_j \in \langle \phi_2, \phi_2 \rangle \cdot (\mathbb{Q}^\times)^2$ for $(i, j) = (1, 3)$ and $(i, j) = (3, 4)$.

By [4, Theorem 3.6], there is also an equality

$$L_p(f_\infty/K'_{2,4}, \chi'_{2,4}, 2) = \Lambda_2 \Lambda_4$$

where, this time, $L_p(f_\infty/K'_{2,4}, \chi'_{2,4}, 2)$ is defined in [4, Definition 3.4]. Again, by construction, $L_p(f_\infty/K'_{2,4}, \chi'_{2,4}, 2)$ is a square in \mathbb{Q}^\times , so the same is true of the product $\Lambda_2 \Lambda_4$. Now the factorization

$$\prod_{i=1}^4 \Lambda_i = (\Lambda_1 \Lambda_2) \cdot (\Lambda_3 \Lambda_4) = (\Lambda_1 \Lambda_3) \cdot (\Lambda_2 \Lambda_4) \in (\mathbb{Q}^\times)^2$$

shows that $\Lambda_1 \Lambda_2$ is a square in \mathbb{Q}^\times . Thanks to the interpolation formula

$$L_p(f_\infty/K', \chi', 2) = \eta(2) \Lambda_1 \Lambda_2$$

proved before and the fact that, by construction, $L_p(f_\infty/K', \chi', 2)$ is a square in \mathbb{Q}^\times (keep in mind the normalization chosen in part (1) of Proposition 3.2), it follows that the same is true of $\eta(2)$, as was to be shown. \square

Remark 4.34. The function $L_p(f_\infty/K', \chi', k)$ appearing in (48) can, of course, be interpreted as a p -adic L -function attached to the Hida family f_∞ , the real quadratic field K' and the genus character χ' . In particular, observe the following facts.

1) Since p is split in K' , the set $\mathbb{Z}_p^\times \times \mathbb{Z}_p^\times$ is stable under the action of the elements γ_{ψ_σ} , hence, thanks to Lemma 4.18 and the density argument in the proof of Proposition 4.21, the function $L_p(f_\infty/K', \chi', k)$ is well defined independently of the choices of the representative $\tilde{\mu}_f^\epsilon$ and of the set $\{\psi_\sigma\}_{\sigma \in G_{K'}^+}$.

2) As in Remark 4.20, one can show that for all $k \in \mathbb{U}_f$ there is an equality

$$\int_{\mathbb{Z}_p^\times \times \mathbb{Z}_p^\times} Q_\psi^{(k)}(x, y) d\tilde{\mu}_{f, \gamma_\psi}^\pm = \int_{\mathbb{Z}_p^\times \times \mathbb{Z}_p^\times} \langle Q_\psi(x, y) \rangle^{\frac{k-2}{2}} d\tilde{\mu}_{f, \gamma_\psi}^\pm.$$

To show this, one proceeds as in the proof of [3, Lemma 3.7] upon noticing that ψ , $\mathbb{Z}_p \times \mathbb{Z}_p$ and $\mathbb{Z}_p^\times \times \mathbb{Z}_p^\times$ correspond to Ψ , L_Ψ and L'_Ψ in the notations of [3]. Since we need the function in (48) only as an auxiliary object, we will not develop the theory of p -adic L -functions in this context any further.

5. RATIONALITY RESULTS FOR DARMON POINTS ON ELLIPTIC CURVES

Set $N := MDp$, let E/\mathbb{Q} be an elliptic curve of conductor N and let K be a real quadratic field as in §2.1. Recall that $D > 1$ is a square-free product of an *even* number of primes.

Theorem 5.1. *Let χ be a genus character of K corresponding to a pair (χ_1, χ_2) such that $\chi_i(-MD) = -w_{MD}$ for $i = 1, 2$.*

(1) *There exists a point $\mathbf{P}_\chi \in E(H_\chi)^\times$ and a rational number $c \in \mathbb{Q}^\times$ such that*

$$J_\chi = c \log_E(\mathbf{P}_\chi).$$

(2) *The point \mathbf{P}_χ is of infinite order if and only if $L'(E/K, \chi, 1) \neq 0$.*

(3) *A suitable integral multiple of P_χ belongs to the natural image of $E(H_\chi)^\times$ in $E(K_p)$. In particular, P_χ coincides with the image of a global point in $E(K_p) \otimes_{\mathbb{Z}} \mathbb{Q}$.*

Proof. As in the proof of [4, Theorem 4.3], order χ_1, χ_2 in such a way that $\text{sign}(E, \chi_1) = -1$. Then $\chi_1(p) = -w_p = a_p$ and, thanks to [3, Theorem 5.4], $L_p(f_\infty, \chi_1, k, k/2)$ vanishes of order at least 2 at $k = 2$. Furthermore, there are a global point $\mathbf{P}_{\chi_1} \in E(\mathbb{Q}(\sqrt{d_1}))^{\chi_1}$, which is of infinite order if and only if $L'(E/\mathbb{Q}, \chi_1, 1) \neq 0$, and an integer s such that

$$(49) \quad \frac{d^2}{dk^2} L_p(f_\infty, \chi_1, k, k/2)_{k=2} = s \log_E(\mathbf{P}_{\chi_1})^2$$

and

$$(50) \quad s \equiv L^*(f, \psi, 1) \pmod{(\mathbb{Q}^\times)^2}$$

for *any* primitive Dirichlet character ψ for which $L(f, \psi, 1) \neq 0$, $\psi(p) = -\chi(p)$ and $\psi(\ell) = \chi(\ell)$ for all primes $\ell \nmid MD$. Here the *algebraic part* $L^*(f, \psi, 1)$ of $L(f, \psi, 1)$ is as in [4, eq. (24)].

From Theorem 4.33 and the fact that $L_p(f_\infty, \chi_1, k, k/2)$ has order of vanishing at $k = 2$ greater than or equal to 2, it follows that

$$(51) \quad \frac{d^2}{dk^2} L_p(f_\infty/K, \chi, k)_{k=2} = \eta(2) \frac{d^2}{dk^2} L_p(f_\infty, \chi_1, k, k/2)_{k=2} L_p(f_\infty, \chi_2, 2, 1).$$

First suppose that $L'(E/K, \chi, 1) \neq 0$. In this case $L_p(f_\infty, \chi_2, 2, 1) = 2L^*(f, \chi_2, 1) \in \mathbb{Q}^\times$. Set $m := L^*(f, \chi_2, 1)$. By (50), there exists $r \in \mathbb{Q}^\times$ such that $r^2 = s/m$. Define $\mathbf{P}_\chi := \mathbf{P}_{\chi_1}$ in this situation. Setting $v := ntrm \in \mathbb{Q}^\times$, a combination of (49), (51) and Corollary 4.32 yields

$$J_\chi^2 = \eta(2)(nt)^2 sm \log_E(\mathbf{P}_{\chi_1})^2 = (v\sqrt{\eta(2)})^2 \log_E(\mathbf{P}_\chi)^2,$$

from which we obtain the desired result for $c := \pm v\sqrt{\eta(2)} \in \mathbb{Q}^\times$ (recall that, by Theorem 4.33, $\eta(2)$ is a square in \mathbb{Q}^\times).

Now suppose that $L'(E/K, \chi, 1) = 0$. In this case $L_p(f_\infty, \chi_2, 2, 1) = 2L^*(f, \chi_2, 1) = 0$, and the result is an immediate consequence of (51) and Corollary 4.32 on setting $\mathbf{P}_\chi := 0$. This completes the proof of parts (1) and (2).

Finally, since $J_\chi = \log_E(P_\chi)$, if $c = a/b$ with $a, b \in \mathbb{Z}$ then part (2) gives the equality

$$\log_E(bP_\chi) = \log_E(a\mathbf{P}_\chi),$$

which implies part (3) since the kernel of \log_E is the torsion subgroup of $E(K_p)$. \square

REFERENCES

- [1] A. Ash, G. Stevens, Cohomology of arithmetic groups and congruences between systems of Hecke eigenvalues, *J. Reine Angew. Math.* **365** (1986), 192–220.
- [2] B. Balamasubramanyam, M. Longo, Λ -adic modular symbols over totally real fields, *Comment. Math. Helv.*, to appear.
- [3] M. Bertolini, H. Darmon, Hida families and rational points on elliptic curves, *Invent. Math.* **168** (2007), no. 2, 371–431.
- [4] M. Bertolini, H. Darmon, The rationality of Stark–Heegner points over genus fields of real quadratic fields, *Ann. of Math. (2)* **170** (2009), no. 1, 343–370.
- [5] J.-F. Boutot, H. Carayol, Uniformisation p -adique des courbes de Shimura : les théorèmes de Čerednik et de Drinfeld, *Astérisque* **196-197** (1991), no. 7, 45–158.
- [6] H. Cohn, *A classical invitation to algebraic numbers and class fields*, Universitext **9**, Springer-Verlag, New York, 1978.
- [7] H. Darmon, Integration on $\mathcal{H}_p \times \mathcal{H}$ and arithmetic applications, *Ann. of Math. (2)* **154** (2001), no. 3, 589–639.
- [8] S. Dasgupta, Stark–Heegner points on modular Jacobians, *Ann. Sci. École Norm. Sup. (4)* **38** (2005), no. 3, 427–469.
- [9] S. Dasgupta, M. Greenberg, \mathcal{L} -invariants and Shimura curves, submitted (2009).
- [10] S. S. Gelbart, *Automorphic forms on adèle groups*, Annals of Mathematics Studies **83**, Princeton University Press, Princeton, NJ, 1975.
- [11] S. S. Gelbart, An elementary introduction to the Langlands program, *Bull. Amer. Math. Soc. (N. S.)* **10** (1984), no. 2, 177–219.
- [12] J. González, V. Rotger, Non-elliptic Shimura curves of genus one, *J. Math. Soc. Japan* **58** (2006), no. 4, 927–948.
- [13] M. Greenberg, Stark–Heegner points and the cohomology of quaternionic Shimura varieties, *Duke Math. J.* **147** (2009), no. 3, 541–575.
- [14] R. Greenberg, G. Stevens, p -adic L -functions and p -adic periods of modular forms, *Invent. Math.* **111** (1993), no. 2, 407–447.
- [15] H. Hida, Galois representations into $\mathrm{GL}_2(\mathbf{Z}_p[[X]])$ attached to ordinary cusp forms, *Invent. Math.* **85** (1986), no. 3, 545–613.
- [16] B. W. Jordan, *On the Diophantine arithmetic of Shimura curves*, Harvard University Ph. D. thesis, 1981.
- [17] M. Longo, S. Vigni, On the vanishing of Selmer groups for elliptic curves over ring class fields, *J. Number Theory* **130** (2010), no. 1, 128–163.
- [18] M. Longo, S. Vigni, Quaternion algebras, Heegner points and the arithmetic of Hida families, *Manuscripta Math.*, to appear.
- [19] M. Longo, V. Rotger, S. Vigni, On rigid analytic uniformizations of Jacobians of Shimura curves, *Amer. J. Math.*, to appear.
- [20] M. Longo, V. Rotger, S. Vigni, Special values of L -functions and the arithmetic of Darmon points, submitted.
- [21] Y. Matsushima, G. Shimura, On the cohomology groups attached to certain vector valued differential forms on the product of the upper half planes, *Ann. of Math. (2)* **78** (1963), no. 3, 417–449.
- [22] M. R. Murty, V. K. Murty, *Non-vanishing of L -functions and applications*, Progress in Mathematics **157**, Birkhäuser, Basel, 1997.
- [23] J. Nekovář, A. Plater, On the parity of ranks of Selmer groups, *Asian J. Math.* **4** (2000), no. 2, 437–498.
- [24] A. Ogg, Real points on Shimura curves, in *Arithmetic and Geometry I*, M. Artin and J. Tate (eds.), Progress in Mathematics **35**, Birkhäuser, Boston, 1983, 277–307.
- [25] A. Popa, Central values of Rankin L -series over real quadratic fields, *Compos. Math.* **142** (2006), no. 4, 811–866.
- [26] J.-P. Serre, *Trees*, corrected 2nd printing, Springer-Verlag, Berlin, 2003.
- [27] G. Shimura, *Introduction to the arithmetic theory of automorphic functions*, Princeton University Press, Princeton, NJ, 1971.

DIPARTIMENTO DI MATEMATICA PURA E APPLICATA, UNIVERSITÀ DI PADOVA, VIA TRIESTE 63, 35121 PADOVA, ITALY

E-mail address: mlongo@math.unipd.it

DEPARTAMENT DE MATEMÀTICA APLICADA II, UNIVERSITAT POLITÈCNICA DE CATALUNYA, C. JORDI GIRONA
1-3, 08034 BARCELONA, SPAIN

E-mail address: `stefano.vigni@upc.edu`